A counterexample to the four state Mpemba triplet conjecture

Research report and reproducible mathematical certificate

11 October 2026

Abstract

We disprove the four-state case of the finite-temperature triplet conjecture of Avitan, Factor and Gelbwaser-Klimovsky (arXiv:2603.04567v1). An explicit reversible generator with strictly positive rates and four distinct energies fails the necessary-condition test of Avitan, Factor and Gelbwaser-Klimovsky on every original-rate triplet, but has an ordinary asymptotic cooling Mpemba effect between two finite positive initial temperatures. The bath inverse temperature is one; the initial inverse temperatures are 1/1000 and 1/100. Rational interval arithmetic certifies the thermal-overlap inequality. A second standalone verification uses exact inertia counts and spectral residual bounds. The example has a simple slow mode and strict margins, hence belongs to an open family of counterexamples. We connect the triplet inequalities to established stochastic monotonicity, obtaining corrected sufficient conditions for absence. The supporting order argument is presented as an application of existing theory, with explicit comparison to Ohga, Hayakawa and Ito. Overlap plots and a certified finite-time distance crossing show the physical effect, whose amplitude advantage is about 0.1102%.

1 Outcome

The proposed triplet test can miss a genuine four-state relaxation reversal. In the example, the hotter preparation begins farther from equilibrium, yet later becomes slightly closer. The reversal is mathematically rigorous and physically small. Every input rate and bath weight is rational. The energies are logarithms of those weights. A cubic with integer coefficients specifies the slow eigenvalue, and cofactor polynomials specify its left eigenvector exactly. Thermal probabilities at the two selected temperatures are algebraic numbers bounded by rational numbers whose correctness is checked by integer powers.

A companion example has an exactly rational slow eigenvalue and left eigenvector. It gives a second, simpler spectral certificate, at the expense of two less compact rates. Both examples are strictly inside the required positive-rate and simple-slow-mode domain. No assertion about a strong Mpemba effect is needed: the two displayed slow-mode overlaps are nonzero.

2 Problem and literature status

The target is the inference that failure of Eq. (3) on every energy-ordered triplet of a four-state reversible thermalizing generator precludes the Mpemba effect. The triplets retain the original off-diagonal rates. This is the conjecture in Avitan, Factor and Gelbwaser-Klimovsky, arXiv:2603.04567v1, submitted 4 March 2026. Its main text and supporting information, particularly S2–S14, were checked on 11 October 2026. The version history listed only v1. Its finite-temperature claim is numerical; the analytic theorem assumes zero bath temperature and stronger triplet inequalities.

The later Gelbwaser-Klimovsky, arXiv:2609.09988v1, submitted 9 September 2026, repeats the inference without providing a general proof. Its examples without an exact three-state mechanism

concern a different, stronger predicate. Searches by title, identifier, conjecture, correction and counterexample located no other primary-source resolution. This establishes the scope of the literature check; it does not exclude unindexed or unpublished work. The source-to-counterexample correspondence is made explicit in Section 2.3. Publication priority or exhaustive novelty is not claimed.

For supporting no-Mpemba results we additionally checked Ohga, Hayakawa and Ito, arXiv:2410.06623v2, version-history date 19 October 2024, including Result 2 and SI B–D, on 11 October 2026. Their theorem is a prior application of monotone coupling to absence of mean-energy crossings. Section 4.4 compares the assumptions and observables precisely. We do not claim a new general monotonicity principle.

For definitions we use Lu and Raz, PNAS 114, 5083–5088 (2017), and distinguish the strong effect of Klich et al., Phys. Rev. X 9, 021060 (2019). The asymptotic conclusion is supplemented below by certified finite-time crossing brackets.

2.1 Model and reductions

With column probabilities, $\dot{p} = Mp$, $M_{ij} = a_{ij}$ for $i \neq j$, and $M_{jj} = -\sum_{i\neq j} a_{ij}$. The rate a_{ij} goes from j to i. Write

$$q_i = e^{-\beta_b E_i}, \qquad Q = \sum_i q_i, \qquad \pi_b = q/Q.$$

Detailed balance is $M_{ij}q_j = M_{ji}q_i$. An energy shift sets $E_1 = 0$; measuring energies in bath units sets $\beta_b = 1$. A common positive multiplier of M changes only the time unit. We use $a_{12} = 1$. We neither assume equally spaced energies nor normalize the total energy span. These would impose additional restrictions.

If $D = \mathrm{diag}(\pi_b)$, detailed balance makes $S = D^{-1/2}MD^{1/2}$ symmetric: $MD = DM^\top$. Hence the spectrum is real and the generator is diagonalizable. Its Dirichlet form is

$$-f^\top DM^\top f = \sum_{i<j} c_{ij}(f_i - f_j)^2, \qquad c_{ij} = M_{ij}\pi_{bj} = c_{ji}.$$

In the positive-rate domain the stationary eigenvalue is simple. The slowest nonzero eigenvalue is additionally assumed simple and is certified below.

2.2 Exact predicate and its negation

For $i < j < k$, put $A = a_{ik}$, $B = a_{jk}$, $C = a_{ij}$, $t = q_j/q_i$, $u = q_k/q_j$. The source's $(-, 0, +)$ correspond to (i, j, k). Its exact test is

$$\kappa = \frac{A+B}{C(1+t)}, \qquad G = At - B,$$
$$F = A(1 + tu/2) - B(1 + u/2) - C(1 - t),$$
$$P_{ijk} = (\kappa \geq 1 \ \wedge \ F > 0) \ \vee \ (\kappa \leq 1 \ \wedge \ G > 0).$$

Both strict comparisons and both inclusive branch comparisons match Eq. (3) and S9. This is a direct transcription of the published test, using destination-first rate indices.

Lemma 1 (The branch logic reduces to a convex cone)**.** *For positive C and $0 < t < 1$, $\neg P_{ijk}$ is equivalent to $F \leq 0$ and $G \leq 0$.*

Proof. Let $H = A + B - C(1 + t)$, $d = (1 - t)/(1 + t) > 0$, and $h = 2/(1 + t) + u/2 > 0$. Direct expansion gives $F = hG + dH$. If $H > 0$, failure is $F \leq 0$, which forces $G < 0$. If $H < 0$, failure is $G \leq 0$, which forces $F < 0$. At $H = 0$ both branches apply, but $F = hG$, so their strict tests agree. These cases prove both directions. □

For fixed energies the eight linear inequalities define one closed polyhedral cone. The admissible rates are its intersection with the strictly positive orthant. This conclusion does not discard the original branch logic; it proves an equivalent formulation including $\kappa = 1$.

The supporting-information fast-vector formulas divide by $-uG$. We do not use that division. When $G = 0$, direct triplet diagonalization gives nonzero eigenvalues $-A(1 + t + tu)$ and $-[C(1 + t) + Atu]$. If also $H = 0$, they coincide. The predicate remains well-defined whenever $C > 0$. Isolated triplet generators require their diagonals to be recomputed, not inherited as principal submatrices.

2.3 Precise published claim and the contradiction

The target is the finite-temperature conjecture in Avitan, Factor and Gelbwaser-Klimovsky, rather than an extrapolation supplied by this report. Their N-level protocol states: "if all triplets violate the conditions, the N-level system will not exhibit the Mpemba effect." The non-zero-temperature subsection explicitly identifies this as a conjecture. The protocol and SI Eqs. (S10)–(S14) specify the original rates.

Let $\mathsf{ME}_{\mathrm{cool}}(M, E, \beta_b)$ mean that there exist finite $0 < \beta_h < \beta_c < \beta_b$ with $|b_2(\beta_h)| < |b_2(\beta_c)|$, where the bath and generator are fixed and the slow nonzero eigenvalue is simple. Section 2.4 shows that this implies an initial-to-eventual KL-distance reversal. A necessary consequence of the published conjecture, restricted to four states and this regular domain, is

$$\boxed{\bigwedge_{i<j<k} \neg P_{ijk} \quad \Longrightarrow \quad \neg\mathsf{ME}_{\mathrm{cool}}(M, E, \beta_b).} \tag{$\mathcal{C}_4$}$$

This is a universally quantified implication over distinct ordered energies, positive reversible rates and $0 < \beta_b < \infty$. Requiring positive rates and a simple slow mode restricts the domain; it does not enlarge the authors' claim. Restricting the conclusion to cooling makes it weaker. A counterexample to this restricted consequence therefore suffices to refute the unrestricted finite-temperature conjecture, independently of any heating or spectral-boundary convention.

Requirement in ($\mathcal{C}_4$)	Witness in Theorem 1
Four ordered distinct energies	$E = (0, \log(200/199), \log(5/2), \log 400)$.
Finite positive bath	$\beta_b = 1$.
Conservative reversible thermal dynamics	All twelve off-diagonal rates are positive; $\mathbf{1}^\top M = 0$ and $M\,\mathrm{diag}(q)$ is symmetric.
Simple slow nonzero eigenvalue	One root in each of $(-20, -19)$, $(-60, -59)$ and $(-337, -336)$.
Original-rate triplet failures	Exact evaluations below give $F < 0$ and $G < 0$ on all four triplets, with both κ branches represented.
Conclusion contradicted	$\beta_h = 1/1000 < \beta_c = 1/100 < 1$ and $\lvert b_2(\beta_c)\rvert - \lvert b_2(\beta_h)\rvert > 0.0001034076$.

The table records the proof obligations, which the exact construction below discharges. No effective rate, principal-submatrix diagonal, infinite initial temperature or equality case is used. Shifting all energies by the same positive constant also makes $E_1 > 0$, if desired, without changing any rate test or Gibbs distribution.

Three distinct statements. The March paper's zero-temperature theorem assumes $\beta_b \to \infty$, a nondegenerate spectrum and $C > A + B$ for every triplet (SI Eq. S20). Our bath is finite and the latter inequality fails already for 123: $1 < 28 + 1799/8$. Thus the counterexample does not challenge that theorem or the three-state necessity of Eq. (3).

The September paper, Gelbwaser-Klimovsky, Section II, distinguishes Eq. (3) from its exact three-state mechanisms. Write $\mathcal{E}_{ijk}$ for satisfaction of one of those mechanisms. In their nondegenerate three-state setting, necessity gives $\mathcal{E}_{ijk} \Rightarrow P_{ijk}$, but not conversely. Consequently

$$\bigwedge \neg P_{ijk} \implies \bigwedge \neg \mathcal{E}_{ijk}.$$

Necessity supplies no converse to this implication. Its reported four-state effects with no triplet mechanism therefore do not, by that fact alone, establish failure of $(\mathcal{C}_4)$. Section II separately reiterates the original-rate Eq. (3) absence inference. Our witness verifies the stronger premise $\bigwedge \neg P_{ijk}$ explicitly.

General conjecture versus sampled parameter ranges. The March Fig. 2 uses rates from 0.05 to 1 and $\beta_b E_i$ from 0 to 1; the September Fig. 5 uses rates from 0.001 to 1 and $\beta_b E_i$ from 0 to 5. These are sampling ranges, not hypotheses attached to the general inference. Our invariant energy span is $\beta_b(E_4 - E_1) = \log 400 > 5$, outside both boxes. Even the downward-rate ratio $a_{12}/a_{23} = 8/1799 < 0.05$ prevents a common time rescaling from placing all downward rates in the March box. We therefore refute the unrestricted conjecture; we neither reproduce nor contradict the finite sample counts, and make no claim of a counterexample within either sampled box.

2.4 What the cooling witness proves

Choose $Mv_2 = \lambda_2 v_2$, $w_2^\top M = \lambda_2 w_2^\top$ and $w_2^\top v_2 = 1$. Since $\lambda_2 \neq 0$, $w_2^\top \pi_b = 0$. Thus

$$b_2(\beta) = w_2^\top [\pi(\beta) - \pi_b] = w_2^\top \pi(\beta).$$

All initial temperatures here are finite and positive. Cooling has $0 < \beta_h < \beta_c < \beta_b$, with anomalous ordering $|b_2(\beta_h)| < |b_2(\beta_c)|$. Heating instead has $\beta_b < \beta_h < \beta_c$ and requires $|b_2(\beta_c)| < |b_2(\beta_h)|$.

With a simple slow mode,

$$p(t;\beta) - \pi_b = e^{\lambda_2 t} b_2(\beta) v_2 + O(e^{\lambda_3 t}).$$

Consequently a strict overlap comparison gives the same eventual comparison in every fixed norm. For relative entropy,

$$D_{\mathrm{KL}}(p(t;\beta)\|\pi_b) = \tfrac{1}{2} e^{2\lambda_2 t} b_2(\beta)^2 \sum_i v_{2i}^2/\pi_{bi} + o(e^{2\lambda_2 t}).$$

The hotter initial Gibbs state is initially farther in this distance because

$$\frac{d}{d\beta} D_{\mathrm{KL}}(\pi(\beta)\|\pi_b) = (\beta - \beta_b)\,\mathrm{Var}_{\pi(\beta)}(E).$$

Thus the example gives a genuine initial-to-eventual distance reversal. It does not claim reversal at every time or for arbitrary distance functionals without appropriate hypotheses. A strong effect would require a nonbath preparation with zero slow overlap; that is not the witness used here.

Holding the bath, generator and eigenvectors fixed gives

$$b_2'(\beta) = -\,\mathrm{Cov}_{\pi(\beta)}(w_2, E) = -\sum_{i<j} \pi_i(\beta)\pi_j(\beta)(w_{2j} - w_{2i})(E_j - E_i).$$

A stationary point alone does not prove a maximum. We instead certify two strict overlap values on the same bath side. Since $b_2(1) = 0$, their increasing absolute ordering also forces an interior maximum somewhere between β_h and the bath.

3 Main result and exact counterexample

Theorem 1 (Counterexample to the published finite-temperature inference)**.** *There exist four distinct ordered energies, a finite positive bath temperature and a strictly positive reversible generator with a simple slowest nonzero eigenvalue such that all four original-rate predicates P_{ijk} fail, yet two finite cooling preparations obey $|b_2(\beta_h)| < |b_2(\beta_c)|$. Thus the antecedent of $(\mathcal{C}_4)$ holds and its conclusion is false.*

Proof. Set

$$\beta_b = 1, \quad q = (1, 199/200, 2/5, 1/400), \quad E = (0, \log(200/199), \log(5/2), \log 400).$$

These energies are distinct and strictly ordered. The downward rates in the order $(12, 13, 14, 23, 24, 34)$ are

$$(1, 28, 249/10, 1799/8, 4983/200, 251/25).$$

Set upward rates by $a_{ji} = a_{ij}q_j/q_i$ for $i < j$. The full generator is

$$M = \begin{pmatrix} -49029/4000 & 1 & 28 & 249/10 \\ 199/200 & -7280583/79600 & 1799/8 & 4983/200 \\ 56/5 & 17990/199 & -1011751/4000 & 251/25 \\ 249/4000 & 4983/79600 & 251/4000 & -11971/200 \end{pmatrix}.$$

Every off-diagonal entry is positive. Its columns sum to zero and $M\,\mathrm{diag}(q)$ is symmetric. Since $Q = 959/400$, the equilibrium is $\pi_b = (400, 398, 160, 1)/959$.

Exact original-rate triplet evaluations are:

Triplet	κ	F	G
123	7225/57	−1176493/4975	−39403/200
124	3321/133	−32119/1592000	−279/2000
134	1747/1960	−7761/4000	−2/25
234	1391209/12548025	−190400119/1592000	−119/4975

All four predicates fail strictly, with two upper branches and two lower branches. No boundary convention is needed.

The characteristic polynomial is $xC(x)/1273600000000$, where

$$C(x) = 1273600000000x^3 + 530473008000000x^2 + 35732953114072400x + 510539126345027563.$$

Exact opposite endpoint signs show one root in each of $(-337, -336)$, $(-60, -59)$ and $(-20, -19)$. These three disjoint sign-changing intervals exhaust a cubic's roots; all are simple. The largest, denoted r, is λ_2. Approximately,

$$(\lambda_2, \lambda_3, \lambda_4) = (-19.86836952145, -59.91747997894, -336.72876105237).$$

In particular the slow-mode spectral gap exceeds 39 in these time units.

Let $P_i(x)$ be the explicit cofactor polynomials listed in Appendix A. Their common factor corresponds to the first column of $\mathrm{adj}(M^\top - xI)$. Put $w_i = P_i(r)/P_1(r)$. The nonvanishing denominator is certified by the rational isolating interval below. The adjugate identity proves $M^\top w = rw$. It also follows that $q^\top w = 0$. To impose the required biorthogonal normalization set

$$N = \sum_i q_i w_i^2 > 0, \qquad v_i = q_i w_i / N.$$

Detailed balance gives $Mv = rv$ and $w^\top v = 1$. Thus $w = w_2$, $v = v_2$ is a valid consistently normalized pair. Approximately,

$$w = (1, -0.75893729814493, -0.61211930644096, -0.0038663077658171).$$

The relevant overlap is exactly

$$b_2(\beta) = \frac{\sum_i w_i q_i^\beta}{\sum_i q_i^\beta}.$$

For $\beta_h = 1/1000$ and $\beta_c = 1/100$, rational interval certification gives the following deliberately widened enclosures:

$$-0.0937454280 < b_2(\beta_h) < -0.0937454279,$$
$$-0.0938488357 < b_2(\beta_c) < -0.0938488356.$$

Every displayed terminating decimal denotes an exact rational number. In particular

$$|b_2(\beta_c)| - |b_2(\beta_h)| > 0.0001034076 > 0.$$

Both preparations have finite temperatures, $T_h = 1000$ and $T_c = 100$ in units with $k_B = 1$, versus $T_b = 1$. The strict overlap comparison and spectral expansion establish the ordinary cooling Mpemba effect. The verification construction below proves the enclosures without relying on floating-point acceptance tests. This completes the counterexample. □

4 Mathematical explanation and corrected theorem

4.1 Why the triplet inference fails

The highest-energy state's slow-mode value is almost zero, while the intermediate states have strongly negative values. At very high initial temperature, occupation of state 4 therefore dilutes the negative slow overlap. The slightly cooler preparation depopulates state 4 enough to increase the magnitude of that overlap. At still lower temperatures, transfer toward the positive ground-state component returns the overlap to zero at the bath. This explanation concerns projection onto a relaxation mode, not merely mean energy.

An isolated triplet has a different eigenvector and a different equilibrium normalization from the full four-state system. Pairwise detailed balance and the inequalities $F, G < 0$ do not control the orientation of the full slow eigenvector relative to the entire Gibbs curve. In particular, nonmonotone slow eigenvectors are permitted. This explains the missing implication in a triplet-only argument.

All rates, F and G are strict; the slow eigenvalue is separated; the overlap gap is strict. Continuous dependence of a simple eigenprojection, Gibbs probabilities and the predicates on rates and energies therefore yields an open neighborhood of counterexamples, including varying energies. This is not an equality-surface artifact. No numerical radius for that neighborhood is claimed.

There is also an explicit one-parameter family. Let $R_0 = \pi_b \mathbf{1}^\top - I$ and $M_\rho = M + \rho R_0$ for $\rho \geq 0$. Its off-diagonal reset rates are $\rho\pi_{bi}$, for which $F = G = 0$ on every triplet. Thus F, G remain unchanged. On the zero-sum subspace $R_0 = -I$, so every nonzero eigenvalue shifts by $-\rho$ and every mode and overlap is preserved. Positivity and the counterexample persist for all $\rho \geq 0$. A subsequent common positive time rescaling also preserves the result. More explicitly, for every $\alpha > 24$, the same q and the downward rates

$$(\alpha - 24, \alpha + 3, \alpha - 1/10, 199\alpha/200 + 200, 199\alpha/200 + 1/25, 2\alpha/5 + 1/25)$$

form a counterexample. Their generator equals $M + (\alpha - 25)(q\mathbf{1}^\top - QI)$. All nonzero eigenvalues shift by $-(\alpha - 25)Q$, while their ordering, modes, overlaps and all F, G remain unchanged. Positivity follows directly from $\alpha > 24$.

4.2 Visible reversal and its size

Figures 1 and 2 separate two facts: the hotter initial state has a smaller slow-mode amplitude, and the full time-dependent distance actually changes ordering. These use the primary generator in Theorem 1, with the bath and generator fixed throughout. The high-temperature amplitude reaches its numerical maximum near $\beta = 0.0237422$.




Figure 1: Thermal overlap of the exact primary generator, with $w_1 = 1$ and $T_b = 1$. The broad curve conceals a small high-temperature maximum; the enlargement displays it. The blue preparation has a larger slow amplitude than the hotter red preparation. The enlarged ordinate uses the hot amplitude as its reference; the quoted 0.1102% reduction uses the cold amplitude. Numerical curves use 80-digit arithmetic; the two selected overlaps are independently certified exactly.

Initially $D_h(0) \simeq 1.210137886$ and $D_c(0) \simeq 1.155489557$, so the hotter preparation is farther away in relative entropy. The two distances cross numerically near $t = 0.14967233$. Exact rational bounds prove a crossing in $(0.14, 0.16)$: at the endpoints,

$$D_h(0.14) - D_c(0.14) \in (5.9361291891, 5.9361291893)\, 10^{-8},$$
$$D_h(0.16) - D_c(0.16) \in (-1.3100324358, -1.3100324356)\, 10^{-8}.$$

Total variation also reverses, with a certified crossing in $(0.10, 0.11)$ and numerical location 0.10436410. The difference between these times is expected: finite-time crossings depend on the distance. Continuity plus the certified signs proves existence; no uniqueness or earliest-crossing theorem is asserted.

The relevant fractional amplitude reduction is

$$\epsilon_b = 1 - \frac{|b_h|}{|b_c|} \simeq 0.0011018531 = 0.11018531\%.$$

For relative entropy the leading term is quadratic, so

$$1 - \lim_{t\to\infty} \frac{D_h(t)}{D_c(t)} = 1 - \frac{b_h^2}{b_c^2} \simeq 0.22024921\%.$$

Both preparations have the same decay exponent λ_2. This is an ordinary effect, not the exponentially faster strong effect. For an increasingly small common distance threshold, the asymptotic time advance is

$$\Delta t = \frac{\log(|b_c|/|b_h|)}{|\lambda_2|} \simeq 5.5488\, 10^{-5}\, a_{12}^{-1}.$$




Figure 2: Actual evolution $p(t) = e^{tM}\pi(\beta)$ for the same two preparations. Relative entropy decreases for each, and the hotter state's distance becomes smaller. The ratio panel resolves the crossing that is difficult to see on the logarithmic plot. The horizontal dotted line is the asymptotic ratio difference, -0.2202492%. Rational uniformization and logarithm bounds certify opposite KL orderings at $t = 0.14$ and 0.16; the displayed root location is numerical, with no claim of uniqueness.

The slow relaxation time is $1/|\lambda_2| \simeq 0.0503313\, a_{12}^{-1}$, so this advance is only about 0.11025% of one slow relaxation time. It is not a percentage reduction in the full cooling time, which depends on the chosen threshold. The time unit has no assigned experimental value, and $T_h/T_b = 1000$, $T_c/T_b = 100$. These data establish a strict mathematical failure of the triplet inference; they do not establish a practically useful cooling protocol.

4.3 A corrected sufficient condition

Let $L = M^\top$ act on functions, $(Lf)_i = \sum_{j\neq i} a_{ji}(f_j - f_i)$. Define adjacent differences $y_i = f_{i+1} - f_i$, the difference map $Bf = y$, and $(H_0 y)_j = \sum_{\ell<j} y_\ell$. Then $BL = KB$ with $K = BLH_0$, since $L\mathbf{1} = 0$. In the constant-plus-differences basis, L is block triangular with blocks 0 and K, so K has precisely the nonzero generator eigenvalues. For its off-diagonal entries,

$$K_{i\ell} = \begin{cases} \sum_{j\le\ell}(a_{j,i} - a_{j,i+1}), & \ell < i, \\ \sum_{j>\ell}(a_{j,i+1} - a_{j,i}), & \ell > i. \end{cases}$$

Theorem 2 (Order-preserving sufficient condition)**.** *For any finite number of states, strictly ordered energies, a fixed $0 < \beta_b < \infty$, positive reversible rates and a simple slowest nonzero eigenvalue, if every off-diagonal entry of K is nonnegative, there is no asymptotic cooling or heating Mpemba effect at positive initial temperatures.*

Proof. Such a matrix K is called Metzler. After a sufficiently large scalar diagonal shift it is nonnegative. Perron theory gives a nonzero nonnegative eigenvector y at its spectral abscissa, which is λ_2. This does not require irreducibility of K. Choose w with $Bw = y$ and $\pi_b^\top w = 0$. Then $B(Lw - \lambda_2 w) = 0$, so $Lw - \lambda_2 w$ is constant. Its bath expectation is zero, hence it vanishes. Thus w is the slow left eigenfunction, nondecreasing and nonconstant in energy.

The covariance identity now gives $b_2'(\beta) < 0$ for every finite positive β: all its pairwise summands are nonnegative and at least one is strictly positive. Because $b_2(\beta_b) = 0$, its absolute value decreases toward the bath on the cooling side and increases away from the bath on the heating side. These are exactly the normal asymptotic orderings. □

In four states, the six off-diagonal entries are

$$K_{12} = a_{32} + a_{42} - a_{31} - a_{41}, \qquad K_{13} = a_{42} - a_{41},$$
$$K_{21} = a_{12} - a_{13}, \qquad K_{23} = a_{43} - a_{42},$$
$$K_{31} = a_{13} - a_{14}, \qquad K_{32} = a_{13} + a_{23} - a_{14} - a_{24}.$$

The original failures already ensure the three upper entries are nonnegative, since $G \leq 0$ orders upward rates. A corrected, directly checkable sufficient condition is therefore the four original failures together with

$$a_{12} \geq a_{13}, \qquad a_{13} \geq a_{14}, \qquad a_{13} + a_{23} \geq a_{14} + a_{24}.$$

This is a proved sufficient condition, not a proposed necessary characterization.

Corollary 1. *If every triplet has $\kappa \leq 1$ and fails the original predicate, then the absence conclusion is true, for any state count under the theorem's assumptions.*

Proof. $H \leq 0$ and $G \leq 0$ give $A + B \leq C(1 + t)$ and $B \geq At$, so $A \leq C$. These comparisons make each downward summand in K nonnegative; $G \leq 0$ and detailed balance do the same for each upward summand. Apply the theorem. □

A second subfamily consists of downward gap laws $a_{ij} = f(E_j - E_i)$, $i < j$, with positive nonincreasing f. Both $f(x)$ and $e^{-\beta_b x} f(x)$ are nonincreasing, so every required summand is nonnegative. This establishes the same absence conclusion directly, without the false general triplet conjecture. The following comparison locates this argument within established order theory; no general monotonicity novelty is claimed.

4.4 Relation to earlier no-Mpemba theorems

Ohga, Hayakawa and Ito (OHI), Result 2 and SI D already use monotone coupling to exclude Mpemba behavior. Their SI D2 explicitly treats a total order. Our matrix condition is precisely a finite-chain expression of stochastic monotonicity, not a separate physical principle. The relevant distinctions are:

	OHI Result 2	The preceding spectral criterion
State order	Finite partial order; energy respects it	Total order by distinct energies
Dynamics	Irreducible and Gibbs-stationary; reversibility unnecessary	Positive reversible rates
Order condition	Comparable jumps and a monotone coupling	$K = BLH_0$ is Metzler
Bath range	Hypotheses throughout a temperature interval	One fixed finite positive bath
Spectrum	No simplicity requirement	Simple slowest nonzero eigenvalue
Conclusion	No cooling mean-energy crossing at any time	No asymptotic cooling/heating reversal in fixed norms or KL

OHI SI B extends their Result 1, the microscopic characterization, to KL divergence; it does not state Result 2 as an all-time KL theorem. Thus the different observable and time scopes must not be conflated.

Indeed $BL = KB$ gives $Be^{tL} = e^{tK}B$. The Metzler condition is equivalent to e^{tL} preserving increasing functions: sufficiency follows from $e^{tK} \geq 0$, and necessity by differentiation at zero on coordinate vectors of adjacent differences. This also gives a monotone coupling. Choose r large enough that $I + L/r$ is stochastic and $I + K/r$ is entrywise nonnegative. Its transition rows are stochastically ordered. A common Poisson clock of rate r and common-uniform inverse-CDF updates

preserve the order of two copies. On a total order, energy compatibility and comparability of jumps are automatic.

For fixed bath, $e^{tL}E$ is then increasing, so

$$\partial_\beta\{\pi(\beta)^\top e^{tL}E\} = -\operatorname{Cov}_{\pi(\beta)}(e^{tL}E, E) \le 0.$$

This gives all-time energy ordering without reversibility or spectral simplicity. It explains the direct relation to OHI while avoiding an inference from a single observable to all possible distances.

The likelihood-ratio and stochastic-order methods below also have precedent in Fill and Kahn, Ann. Appl. Probab. 23 (2013), 1778–1816, particularly their comparison framework and Corollary 3.3. Their setup compares different kernels under suitable initial laws; the thermal preparations here differ instead. We provide the following consequence for clarity, without a priority claim.

Corollary 2 (All-time divergence ordering under reversible monotonicity)**.** *Let a finite irreducible generator be reversible with positive Gibbs equilibrium, on a total energy order, and let K be Metzler. For any convex Φ finite on positive arguments, with $\Phi(1) = 0$, define*

$$D_\Phi(p\|\pi_b) = \sum_i \pi_{bi}\Phi(p_i/\pi_{bi}).$$

For same-side Gibbs preparations, the farther initial temperature has at least as large a divergence at every $t \ge 0$. This includes KL and total variation, and requires no simple eigenvalue. For KL the inequality is strict at every finite time for distinct initial temperatures.

Proof. Write $p \succeq_{\mathrm{st}} q$ when $\sum_i p_i f_i \ge \sum_i q_i f_i$ for every increasing f. Gibbs laws are stochastically ordered by temperature, by the pairwise covariance identity. A monotone semigroup preserves this order. Thus for cooling $p_h(t) \succeq_{\mathrm{st}} p_c(t) \succeq_{\mathrm{st}} \pi_b$.

Reversibility implies $p_c(t)/\pi_b = e^{tL}[\pi(\beta_c)/\pi_b]$ componentwise. The initial ratio is proportional to $e^{(\beta_b-\beta_c)E_i}$ and is increasing; it stays increasing. Choose a nondecreasing subgradient g of Φ. Convexity and stochastic order give

$$D_\Phi(p_h\|\pi_b) - D_\Phi(p_c\|\pi_b) \ge \sum_i (p_{hi} - p_{ci})g(p_{ci}/\pi_{bi}) \ge 0.$$

For heating, $p_c \preceq_{\mathrm{st}} p_h \preceq_{\mathrm{st}} \pi_b$ and p_h/π_b is decreasing. Apply the same supporting-line inequality at p_h/π_b to obtain $D_\Phi(p_c\|\pi_b) \ge D_\Phi(p_h\|\pi_b)$. KL is strictly convex, and invertibility of e^{tM} keeps two distinct initial distributions distinct at finite t, proving strictness. □

This does not assert all-time ordering for every arbitrary norm. The original spectral proof supplies the every-fixed-norm conclusion asymptotically with a simple slow mode. Our counterexample violates monotonicity explicitly: $K_{21} = a_{12} - a_{13} = -27$. It therefore challenges neither OHI's theorem nor this corollary.

The defensible contribution claim is the exact counterexample to the unrestricted triplet inference, its robust family and mechanism, and the explicit triplet-rate connection to an established sufficient condition. We do not present the general order-preservation mechanism or the elementary divergence consequence as newly discovered principles. The all-$\kappa \le 1$ statement is an application; its broader publication priority remains unestablished.

5 Approaches explored

The initial route comparison, including explicit continue/change criteria, is recorded in `research/approaches.md`. The decisive work covered both theorem construction and adversarial falsification.

1. **Reversible spectral geometry.** Symmetrization and the Dirichlet form gave reliable eigenvalue ordering and an equilibrium-orthogonal search space. The obstacle was that rate inequalities do not make the slow-mode orientation convex. Spectral certificates remain essential; this route did not yield the full absence theorem.

2. **Order and covariance.** The first calculation was $K = BLH_0$. Its Metzler subfamily yielded the theorem above. An exact auxiliary example in `research/theorem_notes.md` disproves the intermediate claim that all triplet failures force a monotone slow eigenvector. This identified the missing implication rather than assuming it.

3. **Exponential-polynomial zero counting.** Multiplying by $Z(\beta)^2$ gives

$$Z^2 b_2' = -\sum_{i<j} (w_j - w_i)(E_j - E_i) e^{-\beta(E_i+E_j)}.$$

   For four states there are at most six distinct exponential terms. Chebyshev/variation arguments could bound turning points, but cannot exclude the certified turning behavior. No new general zero-count theorem is claimed. This route became explanatory, not decisive.

4. **Feasible-rate geometry.** The identity $F = hG + dH$ converted apparent branch enumeration into eight linear inequalities. A search enumerated extremal rays after removing the reset direction, then sampled positive mixtures. It found the first root-search candidate after 67,000 systems. Extremal rays were used for discovery only: we did not assume eigenvector properties are inherited by mixtures.

5. **Backward eigenvector design.** Starting from the discovered thermal mode, we prescribed rational q and rational w with $q^\top w = 0$, then solved the linear eigenvector equations for rates. This produced the rational-mode companion and an exact eigenvalue. The remaining spectrum was checked independently; realization of an arbitrary faster eigenmode would not suffice.

6. **Lumping and perturbation.** The near-neutral highest state explains the observed mode. No adiabatically eliminated rate was substituted into the original predicate. Strict-margin continuity supplies a rigorous open family; a separate singular-perturbation theorem was unnecessary once a finite positive example was certified.

7. **Difficult regimes and targeted sampling.** Broad positive-rate sampling missed the effect. Signed-cone directions plus reset restoration found an independent witness. This concentrated the search on unequal energy gaps and narrow high-temperature windows, where a coarse temperature grid is weak.

8. **Algebraic proof.** Rational bath weights made the selected Gibbs probabilities algebraic even though the energy gaps are not commensurate. Exact root brackets and rational interval arithmetic proved the two-temperature inequality. Commensurate energies were therefore not needed.

9. **Additional idea: reset lineality.** The reset generator lies in the lineality space of all the triplet inequalities and acts as a scalar on nonstationary modes. This allowed signed directions to be made physical without changing their mode geometry, and gives the explicit counterexample family above. The September paper also studies the fully degenerate reset point; general novelty of that point is not claimed.

10. **Additional idea: thermal dual cones.** The cone spanned by Gibbs tangent vectors would have a dual consisting of modes with controlled derivative sign. The first calculation was the covariance expansion. The counterexample shows that the original rate cone's slow modes cross

the required thermal dual. A complete dual-cone characterization remains an extension, not a premise used in the proof.

6 Verification and reproducibility

6.1 Exact certificate construction

Run from the workspace root:

```
python3 research/adversarial_certificate.py
python3 research/adversarial_rational_mode_certificate.py
```

The primary script uses SymPy rational/integer arithmetic to check positivity, column sums, detailed balance and the original Boolean predicates. It derives the characteristic polynomial and cofactor vector and verifies its eigenvector residual and bath expectation modulo $C(x)$. The actual slow root is enclosed between

$$-\frac{42820190741029}{2155193997917} \quad \text{and} \quad -\frac{135549509489200}{6822377112669}.$$

Opposite exact polynomial signs certify this interval. Two disjoint negative intervals similarly contain the other roots; degree three proves completeness. The interval evaluation of P_1 excludes zero, so division by it is justified.

For each $q_i^{1/n}$, $n = 1000, 100$, the script proposes adjacent grid points L_i, U_i of mesh 10^{-24}, then checks exactly

$$0 \le L_i, \qquad L_i^n \le q_i \le U_i^n.$$

Strict monotonicity of the positive nth-power map proves the enclosure. Approximate arithmetic only proposes the endpoints; it never accepts an inequality. Polynomial Horner evaluation, products and division use exact interval endpoints. All denominators are verified to exclude zero. The final interval comparison is also exact. Full rational endpoints, rather than merely rounded decimals, are saved in the JSON certificates.

The tighter certified margin in the primary normalization is greater than

$$0.00010340762800011847331527.$$

The deliberately widened report bounds are conservative. Scaling w consistently changes the displayed numbers but never the conclusion; search logs may use a unit weighted norm, whereas the final report fixes $w_1 = 1$.

6.2 A separate proof using inertia and residual bounds

The primary example also admits a verification that does not use its cubic coefficients, cofactor polynomials, previously computed root brackets or certificate JSON files. The standalone script `research/standalone_primary_check.py` starts only from the six printed downward rates and four bath weights. It uses Python's standard-library integers and fractions, with explicit rejection checks that remain active under optimization. It rebuilds the generator and evaluates the original two-branch predicates directly, without invoking Lemma 1. The following argument explains its complete spectral and thermal certificate.

Exact identification of the slow eigenspace. Put $L = M^\top$, $D_q = \mathrm{diag}(q)$ and $H = D_q L$. Detailed balance makes H symmetric. For rational ξ, exact symmetric elimination of $H - \xi D_q$ into a unit triangular factor, a diagonal factor and the transpose triangular factor gives the following pivot signs, eliminating coordinates in the order $1, 2, 3, 4$:

ξ	Pivot signs	Number of eigenvalues of L above ξ
-337	$+,+,+,+$	4
-336	$+,+,-,+$	3
-60	$+,-,+,+$	3
-59	$+,-,+,-$	2
-20	$+,-,+,-$	2
-19	$+,-,-,-$	1

All pivots are nonzero; the script records their exact rational values. Each elimination step replaces the trailing block by $A_{ij} - A_{ik}A_{kj}/A_{kk}$ and is a congruence. Congruence preserves the counts of positive and negative eigenvalues. Also $H - \xi D_q$ is congruent to $D_q^{1/2} L D_q^{-1/2} - \xi I$. Thus the counts drop by one across each of $(-337, -336)$, $(-60, -59)$ and $(-20, -19)$. These account for three simple negative eigenvalues; $L\mathbf{1} = 0$ accounts for the fourth. In particular the slow nonzero eigenspace is one-dimensional and lies in $(-20, -19)$. No approximate eigensolver is used in this identification.

Residual control replaces an exact eigenvector formula. Define the following exact rational trial values (each decimal terminates):

$$\widehat{r} = -19.86836952145135, \qquad \widehat{w}_1 = 1, \quad \widehat{w}_2 = -0.75893729814493, \quad \widehat{w}_3 = -0.61211930644096,$$

$$\widehat{w}_4 = -\frac{q_1\widehat{w}_1 + q_2\widehat{w}_2 + q_3\widehat{w}_3}{q_4} = -\frac{193315388213}{50000000000000}.$$

Use the weighted norm $\|f\|_q^2 = \sum_i q_i f_i^2$. Direct rational multiplication verifies

$$q^\top \widehat{w} = 0, \qquad \|L\widehat{w} - \widehat{r}\widehat{w}\|_q^2 < 10^{-16}.$$

These trial values are not accepted as eigenvalues or eigenvectors: only their explicitly checked residual bound is used.

Let z be the orthogonal projection of $\widehat{w}$ onto the true slow eigenspace in this weighted inner product. The operator L is self-adjoint there. Its stationary component in $\widehat{w}$ vanishes because $q^\top \widehat{w} = 0$. The two faster eigenvalues are below -59, whereas $\widehat{r} > -20$, so their distances from $\widehat{r}$ exceed 39. Expanding in a weighted orthonormal eigenbasis gives

$$\|L\widehat{w} - \widehat{r}\widehat{w}\|_q^2 \geq 39^2 \|\widehat{w} - z\|_q^2, \qquad \|\widehat{w} - z\|_q < \frac{10^{-8}}{39}.$$

Since $\|\widehat{w}\|_q \geq 1$, the projection is nonzero. The minimum bath weight is $1/400$, hence each coordinate error is less than $20\,10^{-8}/39$. For every probability vector p this implies

$$|z^\top p - \widehat{w}^\top p| < \delta, \qquad \delta = \frac{1}{195000000}. \tag{1}$$

Independent thermal enclosures. For $n = 1000$ and 100, integer binary search finds m_i satisfying

$$\left(\frac{m_i}{10^{12}}\right)^n \le q_i < \left(\frac{m_i+1}{10^{12}}\right)^n.$$

These comparisons use integer powers and positive denominators. Signed sums of the endpoint bounds enclose $\sum_i \widehat{w}_i q_i^{1/n}$; positive sums enclose its partition function. Taking all numerator/denominator endpoint quotients and adding the error in Eq. (1) proves the deliberately widened rational intervals

$$z^\top \pi(1/1000) \in (-0.093745434, -0.093745422),$$
$$z^\top \pi(1/100) \in (-0.093848841, -0.093848830).$$

Both overlaps are negative, and their absolute-value difference exceeds $0.000103396 > 10^{-4}$. This proves the cooling reversal for the actual simple slow mode, rather than for the trial vector. The paired right mode is $D_q z/(z^\top D_q z)$, giving unit left-right overlap. Here the left mode is z, whose first coordinate is in $(1-\delta, 1+\delta)$, rather than exactly one. Dividing by this positive coordinate recovers the main report's normalization; the checker verifies that the normalized advantage still exceeds 10^{-4}.

Independence and its limits. The original certificate uses symbolic characteristic polynomials and cofactor intervals; this checker uses rational congruence and a spectral residual bound. Its standalone implementation requires no research data files or third-party packages. Ten adversarial tests include an isolated execution with only the checker present, identical results under normal and optimized Python, rejection of a transposed generator, a false triplet premise, a wrong mode, a poor eigenvector and reversed temperatures, plus analytically known arithmetic examples. These tests exercise failure paths; they do not themselves prove the checker correct.

This is a separate implementation and a different proof within the same model-assisted project, not independent human authorship, formal proof-assistant verification or external peer review. It still trusts Python's integer/fraction arithmetic and the mathematical argument above. It independently verifies the primary asymptotic counterexample; the finite-time crossing brackets remain supported by their separate uniformization certificate. All methods still require review of the correspondence to the published claim in Section 2.3. A reviewer can run the standalone checker in an empty directory and audit the argument above without trusting the discovery search or any stored numerical output.

6.3 Figure and finite-time verification

Run `research/physics_plots.py` for the figures and data, and `research/finite_time_certificate.py` for exact finite-time bounds. Plot values use 80-digit mpmath arithmetic. Direct matrix exponentiation and reversible symmetric spectral propagation agree within $6\,10^{-81}$ at ten checks spanning $t = 0$ to 0.4. The plot CSV files and numerical metrics are saved under `output/figures/`. Those numerical comparisons are distinct from the exact certificates.

The finite-time certificate uses uniformization with $\nu = 253$ and the exactly stochastic matrix $P = I + M/\nu$:

$$e^{tM}p(0) = e^{-\nu t}\sum_{k=0}^{\infty}\frac{(\nu t)^k}{k!}P^k p(0).$$

It retains terms through $k = 250$. Each omitted probability component is at most one; the scalar positive-series tail is bounded by its first omitted term divided by $1 - \nu t/252$. The same positive Taylor bounds give a rational enclosure of $e^{-\nu t}$. Initial Gibbs probabilities use the already-certified algebraic brackets. For logarithms it uses

$$\log r = 2\sum_{k=0}^{59}\frac{z^{2k+1}}{2k+1} + R, \quad z = \frac{r-1}{r+1}, \quad |R| \le \frac{2|z|^{121}}{121(1-z^2)}.$$

All interval operations round outward onto a rational grid of mesh 10^{-50}. Exact rational endpoint signs prove the displayed KL and TV crossing brackets. The script uses floating point only to print readable values. The root estimates shown on the figures are numerical; their enclosing crossing intervals are exact.

6.4 Executed discovery calculations

The discovery environment used Python with NumPy 1.26.4, SciPy 1.13.1 and SymPy 1.14.0. Seeds and actual parameters are retained in the scripts and output files. Floating-point discovery outputs are not mathematical certificates.

`research/cone_search.py` used seed 10411, energy gaps $\exp U[-4, 2]$, 1,000 ray mixtures per energy set, and 150 cooling inverse temperatures from 0.001 to 0.999. It stopped after 67 energy sets, 67,000 systems, on a strict candidate. Coefficients of the ray mixtures were $\exp U[-9, 3]$; a reset direction restored positivity. SVD acceptance tolerances for rays were 10^{-8} and 10^{-9}, and deduplication tolerance 10^{-6}. These are discovery tolerances only.

The independent positive-rate search used seed 934256 and 20,000,000 draws. Gaps were $10^{U[-3,1]}$, rates $10^{U[-5,5]}$ rescaled by their maximum; 2,020,118 draws passed the predicates. It found no accepted witness on its 18-point cooling/heating grid, under the margins and spectral filters in `research/adversarial_search.py`. The signed search used seed 742908, gaps $10^{U[-3,0.7]}$, and normal signed variables scaled by $10^{U[-3,3]}$, alternating an additional destination-weight factor. It stopped at 9,950,000 draws (573,427 feasible) with a witness. Full grids, filters and stopping records are retained in the scripts and `research/adversarial_notes.md`. Neither a failed search nor its sample count was used as proof. A separate model-assisted verification checked the primary candidate at 70 digits, then reviewed the exact certificate's matrix orientation, branch logic, root exhaustion and interval operations. The rational-mode companion was checked separately by symbolic equations and root counts.

No experiments, empirical measurements, messages to authors, or external publication were performed. All generator tests use original rates. All promised executed certificate checks passed.

6.5 Boundary audit

The disproof does not require zero rates, $\beta_b = 0$, a zero-temperature bath, equality of energies or spectral degeneracy. The source numerically discusses equal and zero rates but does not supply a general finite-temperature boundary proof. If zero rates are allowed, irreducibility must still be checked; otherwise bath equilibrium need not be unique. The printed κ quotient is undefined if $C = 0$. A closure defined through F, G requires a stated extension, not silent division by zero.

For a degenerate slow eigenspace the scalar b_2 must be replaced by the full slow spectral projection. Its norm or entropy contribution can depend on several coefficients. Reversibility prevents Jordan blocks but not this ambiguity. Continuity of a selected scalar mode cannot be assumed across an eigenvalue crossing. Our strict spectral separation avoids these issues. The spectral theorem assumes a simple slowest nonzero eigenvalue; no unstated limiting argument is used. The all-time divergence corollary separately covers degenerate spectra under its reversible monotonicity assumptions.

7 Remaining gap

The explicit witness contradicts the published unrestricted finite-temperature implication ($\mathcal{C}_4$) under its stated original-rate interpretation. This conclusion concerns neither the separate zero-temperature theorem nor the simulation boxes discussed in Section 2.3. What remains open in this work is a sharp replacement characterization, a complete phase diagram of admissible slow-mode orientations, and a

minimal structural description of all counterexamples. We have not proved that a particular number of upper-κ triplets is necessary, nor that all effects under the failed premise share the near-neutral-state mechanism.

The corrected order-preserving conditions are sufficient, not necessary, and are explicitly attributed to established monotonicity methods. The report proves an asymptotic effect and certifies finite crossing brackets for KL and total variation. It does not prove uniqueness of those crossings or characterize every transient distance crossing. The literature search is bounded and does not establish priority over unpublished work.

8 Best next step

A concrete next calculation is to fix the rational bath weights above, impose $w_1 = 1$ and $q^\top w = 0$, and solve the linear rate feasibility problem as (w_2, w_4) vary. Combine each feasible solution with an exact slow-mode spectral check and the sign of $b_2(\beta)b_2'(\beta)$ on $0 < \beta < 1$. This would map the boundary between the corrected order-preserving region and the open counterexample family, and test whether one upper-κ triplet can suffice. The existing rational-mode companion provides a fully specified starting point.

For external review, the immediate useful package is this report, the standalone inertia/residual checker and its adversarial tests, the original spectral/overlap certificates, the finite-time certificate, figures, and the hypothesis-by-hypothesis literature comparison. Section 6.2 gives the alternative mathematical argument and distinguishes algorithmic independence from external peer review. Any eventual publication should separately check priority, polish the structural theorem and have a researcher independently run the certificates. The mathematical result here does not depend on that future step.

Appendix A: Explicit slow-mode cofactor polynomials

The primary counterexample's exact left eigenvector is $w_i = P_i(r)/P_1(r)$, where r is the simple slow root specified in Theorem 1 and

$$\begin{aligned}
P_1(x) &= 3184000000x^3 + 1287155436000x^2 \\
&\quad + 74562002576870x + 532366138003157, \\
P_2(x) &= 3184000000x^2 + 9060415184000x + 532366138003157, \\
P_3(x) &= 89152000000x^2 + 14211422840400x + 532366138003157, \\
P_4(x) &= 79281600000x^2 + 28279185628400x + 532366138003157.
\end{aligned}$$

Appendix B: A rational slow-mode companion

Keep the same q, a_{12}, a_{13}, a_{14} and a_{24}, but set

$$a_{23} = \frac{330154675693}{1470375000}, \qquad a_{34} = \frac{1527327066}{151990625}.$$

Use detailed balance and conservative diagonals as before. Then

$$w = \left(1, -\frac{759}{1000}, -\frac{48957}{80000}, -\frac{1}{250}\right), \qquad \lambda_2 = -\frac{4966671}{250000}$$

obey $M^\top w = \lambda_2 w$ and $q^\top w = 0$ by rational arithmetic. Here $N = 27567998009/16000000000$ and $v = \mathrm{diag}(q)w/N$.

The characteristic polynomial factors as

$$\frac{x(250000x + 4966671)R(x)}{632507162050000000000000},$$

where

$$\begin{aligned}
R(x) &= 2530028648200000000x^2 \\
&\quad + 1002356769715761942000x + 50981800562345565497767.
\end{aligned}$$

One root of R lies in $(-337, -336)$ and the other in $(-60, -59)$, as exact endpoint signs show. The rational eigenvalue is therefore the simple slow mode. The companion certificate checks all original predicates strictly and proves

$$\begin{aligned}
b_2(1/1000) &\in (-0.093755177882, -0.093755177881), \\
b_2(1/100) &\in (-0.093857287853, -0.093857287852).
\end{aligned}$$

This is an independent exact realization of the same thermal mechanism, constructed backward from a prescribed mode.