

MATH 4150: Introduction to Number Theory

Frank Qiang
Instructor: Joshua Stucky

Georgia Institute of Technology
Fall 2025

Contents

1	Aug. 18 — Divisibility	3
1.1	Basic Properties of Divisibility	3
1.2	The Division Algorithm	3
2	Aug. 20 — Prime Numbers	5
2.1	Prime Numbers	5
2.2	Sieve of Eratosthenes	5
2.3	Gaps in Primes	6
2.4	Other Open Problems	6
3	Aug. 25 — Greatest Common Divisors	8
3.1	Greatest Common Divisors	8
3.2	The Euclidean Algorithm	9
4	Aug. 27 — Fundamental Theorem of Arithmetic	11
4.1	The Fundamental Theorem of Arithmetic	11
4.2	Least Common Multiples	12
5	Sept. 3 — Congruences	14
5.1	Dirichlet's Theorem	14
5.2	Congruences	14
6	Sept. 8 — Congruences, Part 2	17
6.1	More on Congruences	17
6.2	Linear Congruences in One Variable	18
7	Sept. 10 — Chinese Remainder Theorem	20
7.1	More on Linear Congruences	20
7.2	The Chinese Remainder Theorem	20
7.3	Wilson's Theorem	21
8	Sept. 15 — Fermat's Little Theorem	23
8.1	Fermat's Little Theorem	23
8.2	Euler's Theorem	24
8.3	Arithmetic Functions and Multiplicativity	24
9	Sept. 17 — Arithmetic Functions	26
9.1	Properties of Multiplicative Functions	26
9.2	Properties of the Euler Phi Function	27

10 Sept. 22 — Exam 1 Review	29
10.1 Practice Problems	29
11 Sept. 29 — Arithmetic Functions, Part 2	31
11.1 The Divisor Function	31
11.2 The Sum of Divisors Function	31
11.3 Perfect Numbers	32
11.4 The Möbius Function	33
12 Oct. 1 — Quadratic Residues	35
12.1 Möbius Inversion	35
12.2 Quadratic Residues	36
12.3 The Legendre Symbol	37
13 Oct. 8 — The Legendre Symbol	38
13.1 More on the Legendre Symbol	38
13.2 Particular Cases of the Legendre Symbol	39
14 Oct. 13 — Quadratic Reciprocity	41
14.1 Applications of Gauss's Lemma	41
14.2 Quadratic Reciprocity	42
15 Oct. 15 — Quadratic Reciprocity, Part 2	45
15.1 Proof of Quadratic Reciprocity	45
15.2 Applications of Quadratic Reciprocity	45
16 Oct. 20 — Primitive Roots	47
16.1 Orders	47
16.2 Primitive Roots	48

Lecture 1

Aug. 18 — Divisibility

Something something pair a' docks. (I forgot to write it down oops.)

1.1 Basic Properties of Divisibility

Definition 1.1. Let $a, b \in \mathbb{Z}$. We say that a *divides* b , and we write $a \mid b$, if there exists $c \in \mathbb{Z}$ such that $b = ac$. We also say that a is a *divisor* (or *factor*) of b . We write $a \nmid b$ if a does not divide b .

Example 1.1.1. We have the following:

1. We have $3 \mid 6$ since $6 = 3 \cdot 2$, and $3 \mid -6$ since $-6 = 3 \cdot (-2)$.
2. For any $a \in \mathbb{Z}$, we have $a \mid 0$ since $0 = a \cdot 0$.
3. Technically, we have $0 \mid 0$, but do not confuse this with the indeterminate form $0/0$.

Proposition 1.1. Let $a, b, c \in \mathbb{Z}$. If $a \mid b$ and $b \mid c$, then $a \mid c$. In particular, divisibility is transitive.

Proof. Since $a \mid b$ and $b \mid c$, there exist integers e, f such that $b = ae$ and $c = bf$. We can write

$$c = bf = (ae)f = a(ef),$$

so that a divides c by definition. □

Proposition 1.2. Let $a, b, c, m, n \in \mathbb{Z}$. If $c \mid a$ and $c \mid b$, then $c \mid (am + bn)$. In other words, c divides any integral linear combination of a and b .

Proof. Since $c \mid a$ and $c \mid b$, we have $a = ce$ and $b = cf$ for some $e, f \in \mathbb{Z}$. Then

$$am + bn = (ce)m + (cf)n = c(em + fn),$$

so that c divides $am + bn$ by definition. □

1.2 The Division Algorithm

Definition 1.2. Let $x \in \mathbb{R}$. The *greatest integer function* (or *floor function*) of x , denoted $[x]$ (or $\lfloor x \rfloor$), is the greatest integer less than or equal to x .

Example 1.2.1. We have the following:

1. If $a \in \mathbb{Z}$, then $[a] = a$. The converse is also true: If $[a] = a$ for $a \in \mathbb{R}$, then $a \in \mathbb{Z}$.
2. We have $[\pi] = 3$, $[e] = 2$, $[-1.5] = -2$, and $[-\pi] = -4$.

Lemma 1.1. *Let $x \in \mathbb{R}$. Then $x - 1 < [x] \leq x$.*

Proof. The upper bound is obvious. To show the lower bound, suppose to the contrary that $[x] \leq x - 1$. Then $[x] < [x] + 1 \leq x$, which contradicts the maximality of $[x]$ as $[x] + 1$ is an integer. \square

Example 1.2.2. We can write $5 = 3 \cdot 1 + 2$ and $26 = 6 \cdot 4 + 2$; this is the *division algorithm*.

Theorem 1.1 (Division algorithm). *Let $a, b \in \mathbb{Z}$ with $b > 0$. Then there exist unique $q, r \in \mathbb{Z}$ such that*

$$a = bq + r, \quad 0 \leq r < b.$$

Call q the quotient and r the remainder of the division.

Proof. First we show existence. Let $q = [a/b]$ and $r = a - b[a/b]$. By construction, $a = bq + r$. To check that $0 \leq r < b$, note that by Lemma 1.1, we have $a/b - 1 < [a/b] \leq a/b$. Multiplying by $-b$ gives

$$-a \leq -b[a/b] < b - a,$$

and adding a gives the desired inequality $0 \leq a - b[a/b] = r < b$.

Now we prove uniqueness. Assume there are $q_1, q_2, r_1, r_2 \in \mathbb{Z}$ such that

$$a = bq_1 + r_1 = bq_2 + r_2, \quad 0 \leq r_1, r_2 < b.$$

Then $0 = (bq_1 + r_1) - (bq_2 + r_2) = b(q_1 - q_2) + (r_1 - r_2)$, so we find that

$$r_2 - r_1 = b(q_1 - q_2).$$

So $b \mid r_2 - r_1$. But $0 \leq r_1, r_2 < b$ implies $-b < r_2 - r_1 < b$, so we must have $r_2 - r_1 = 0$, i.e. $r_1 = r_2$. This then implies $0 = b(q_1 - q_2)$, which gives $q_1 - q_2 = 0$ since $b > 0$, so $q_1 = q_2$ as well. \square

Remark. In the division algorithm, we have $r = 0$ if and only if $b \mid a$.

Example 1.2.3. Suppose $a = -5$, $b = 3$. Then $q = [a/b] = -2$ and $r = a - b[a/b] = 1$, i.e.

$$-5 = 3 \cdot (-2) + 1.$$

Note that $-5 = 3 \cdot (-1) + (-2)$ also, but this does not contradict uniqueness since $-2 \notin [0, 3)$.

Definition 1.3. Let $n \in \mathbb{Z}$. Then n is *even* if $2 \mid n$, and *odd* otherwise.

Lecture 2

Aug. 20 — Prime Numbers

Two fish are in a tank. One says to the other, “Ha, how do you drive this thing?”

2.1 Prime Numbers

Definition 2.1. Let $p \in \mathbb{Z}$ with $p > 1$. Then p is *prime* if the only positive divisors of p are 1 and p . If $n \in \mathbb{Z}$, $n > 1$ and n is not prime, then n is *composite*.

Remark. The number 1 is neither prime nor composite.

Example 2.1.1. The following are prime numbers: 2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43, 47, \dots

Lemma 2.1. *Every integer greater than 1 has a prime divisor.*

Proof. Assume to the contrary that there exists $n > 1$ that has no prime divisor. By the well-ordering principle,¹ we may take n to be the smallest such positive integer. Since n has no prime divisors, n cannot be prime. Thus n has a divisor a with $1 < a < n$. Since $1 < a < n$, a must have a prime divisor p by the minimality of n . But then $p \mid a$ and $a \mid n$, so $p \mid n$ by transitivity, a contradiction. \square

Theorem 2.1 (Euclid). *There are infinitely many prime numbers.*

Proof. Assume to the contrary that there are only finitely many primes p_1, p_2, \dots, p_n . Consider

$$N = p_1 p_2 \cdots p_n + 1.$$

By Lemma 2.1, N has a prime divisor $p = p_j$ for some $1 \leq j \leq n$. Since p divides N and p divides $p_1 p_2 \cdots p_n$, p also divides $N - p_1 p_2 \cdots p_n = 1$, which is a contradiction. \square

Exercise 2.1. Modify the proof and construct infinitely many problematic N .

2.2 Sieve of Eratosthenes

Proposition 2.1. *If n is composite, then n has a prime divisor that is less than or equal to \sqrt{n} .*

Proof. Since n is composite, $n = ab$ where $1 < a, b < n$. Without loss of generality, assume $a \leq b$. We claim $a \leq \sqrt{n}$. To see this, suppose to the contrary that $a > \sqrt{n}$. Then $n = ab \geq a^2 > n$, a contradiction. By Lemma 2.1, a has a prime divisor $p \leq a \leq \sqrt{n}$. But then $p \mid a$ and $a \mid n$, so $p \mid n$. \square

¹The *well-ordering principle* says that every nonempty subset of the positive integers contains a least element.

Remark. The proposition implies that if all the prime divisors of an integer n are greater than \sqrt{n} , then n is prime. So to check the primality of n , it suffices to check divisibility by primes $\leq \sqrt{n}$.

Example 2.1.2. The *sieve of Eratosthenes* proceeds as follows. To find primes ≤ 50 , we can delete multiples of primes $\leq \sqrt{50} \approx 7.07$. To start, we know that 2 is prime. Then cross out all multiples of 2. The smallest number remaining is 3, which we now know must be prime. Then cross out all multiples of 3. Continue this process until we cross out all multiples of 7, and then all remaining numbers are prime.

2.3 Gaps in Primes

Proposition 2.2. *For any positive integer n , there are at least n consecutive composite positive integers.*

Proof. Consider the following list of n consecutive numbers:

$$(n+1)! + 2, \quad (n+1)! + 3, \quad (n+1)! + 4, \quad \dots, \quad (n+1)! + (n+1).$$

Note that for any $2 \leq m \leq n+1$, we have $m \mid m$ and $m \mid (n+1)!$, so m divides $(n+1)! + m$. Thus each number in the above list is composite, so we have at least n consecutive composite integers. \square

Remark. With some modifications to this proof (namely a more “efficient” construction), one can find asymptotic lower bounds for the length of long prime gaps.

Conjecture 2.1.1. *There are infinitely many pairs of primes that differ by exactly 2.*

Remark. Zhang (2013) was able to show that there are infinitely many pairs of primes whose difference is $\leq 70,000,000$. This has been lowered to 246 by the Polymath project, which included Tao and Maynard. Assuming other strong conjectures (Elliot-Halberstam), we can get down to 6.

Remark. In addition to long and short prime gaps, we can also consider the average length of prime gaps. Gauss conjectured that as $x \rightarrow \infty$, the number of primes $\leq x$, denoted $\pi(x)$, satisfies

$$\pi(x) \sim \frac{x}{\log x},$$

i.e. $\pi(x)$ is asymptotic to $x/\log x$. Said differently, this says that the “probability” that an integer $\leq x$ is prime is $\pi(x)/x \sim 1/\log x$. This conjecture was proved independently in 1896 by de la Vallée-Poussin and Hadamard, and is now known as the *prime number theorem*.

Definition 2.2. Let $x \in \mathbb{R}$. Define $\pi(x) = |\{p : p \text{ prime}, p \leq x\}|$.

Theorem 2.2 (Prime number theorem). *As $x \rightarrow \infty$, $\pi(x)$ is asymptotic to $x/\log x$, i.e.*

$$\lim_{x \rightarrow \infty} \frac{\pi(x)}{x/\log x} = 1.$$

2.4 Other Open Problems

Conjecture 2.2.1 (Goldbach). *Every even integer ≥ 4 is a sum of two primes.*

Theorem 2.3 (Ternary Goldbach). *Every odd integer ≥ 7 is a sum of three primes.*

Remark. Goldbach’s conjecture implies ternary Goldbach (subtract 3), but not vice versa.

Definition 2.3. Primes of the form $p = 2^n - 1$ are called *Mersenne primes*, and primes of the form $p = 2^{2^n} + 1$ are called *Fermat primes*.

Conjecture 2.3.1. *There are infinitely many Mersenne primes but only finitely many Fermat primes.*

Lecture 3

Aug. 25 — Greatest Common Divisors

*What do you call a root vegetable, fresh off the oven, and a pig that you throw off the balcony?
One is a heated yam, and the other is a yeeted ham.*

3.1 Greatest Common Divisors

Remark. Given $a, b \in \mathbb{Z}$, not both zero, we can consider the set

$$S = \{c \in \mathbb{Z} : c \mid a \text{ and } c \mid b\},$$

of common divisors of both a and b . Note that $\pm 1 \in S$, so S is nonempty, and S is also finite as at least one of a, b is nonzero. Thus S has a maximal element.

Definition 3.1. Let $a, b \in \mathbb{Z}$, not both zero. Then the *greatest common divisor* of a and b , denoted (a, b) , is the largest integer d such that $d \mid a$ and $d \mid b$. If $(a, b) = 1$, then we say that a, b are *relatively prime* (or *coprime*).

Remark. Note that $(0, 0)$ is not defined. Also note that if $(a, b) = d$, then

$$(a, b) = (-a, b) = (a, -b) = (-a, -b) = d.$$

Example 3.1.1. We will compute $(24, 60)$. The list of positive divisors of 24 and 60 are

$$24 : 1, 2, 3, 4, 6, 8, 12, 24;$$

$$60 : 1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60.$$

We can then see that $(24, 60) = 12$.

Remark. In general, we have $(a, 0) = |a|$.

Proposition 3.1. Let $(a, b) = d$. Then $(a/d, b/d) = 1$.

Proof. Let $d' = (a/d, b/d) > 0$. Then $d' \mid (a/d)$ and $d' \mid (b/d)$, so there exist e, f such that $a/d = ed'$ and $b/d = fd'$. We can write this as $a = ed'd$ and $b = fd'd$. Thus $d'd$ is a common divisor of a and b , so we must have $d' = 1$ by the maximality of d . \square

Proposition 3.2. Let $a, b \in \mathbb{Z}$, not both zero, and let

$$T = \{ma + nb : m, n \in \mathbb{Z}, ma + nb > 0\}.$$

Then $\min T$ exists and is equal to (a, b) .

Proof. Without loss of generality, we can assume $a \neq 0$. Note that $|a| \in T$, so T is nonempty. Thus by the well-ordering principle, T has a minimal element d . Then $d = m'a + n'b$ for some $m', n' \in \mathbb{Z}$. We will show that $d \mid a$, a similar argument shows that $d \mid b$. By the division algorithm, we may write

$$a = dq + r, \quad 0 \leq r < d.$$

It suffices to show that $r = 0$. We can rewrite the above as

$$r = a - dq = a - (m'a + n'b)q = a(1 - m'q) - b(n'q).$$

So r is an integral linear combination of a, b . Since d is the smallest positive integral linear combination of a, b and $0 \leq r < d$, we must have $r = 0$. So d is a common divisor of a, b .

Now suppose $c \mid a$ and $c \mid b$, then $c \mid (ma + nb)$, so c divides $d = m'a + n'b$. Thus $c \leq d$, so $d = (a, b)$. \square

Remark. If $(a, b) = d$, then $d = ma + nb$ for some $m, n \in \mathbb{Z}$. If $d = 1$, then the converse also holds: If

$$1 = ma + nb,$$

and d' is a common divisor of a, b , then $d' \mid 1$, so $d' = 1$.

Remark. Along the way, we showed that any common divisor of a, b divides (a, b) .

Definition 3.2. Let $a_1, \dots, a_n \in \mathbb{Z}$, with at least one nonzero. Then the *greatest common divisor* of a_1, \dots, a_n , denoted (a_1, \dots, a_n) , is the largest integer d such that $d \mid a_i$ for $1 \leq i \leq n$. If $(a_1, \dots, a_n) = 1$, then we say that a_1, \dots, a_n are *relatively prime*, and if $(a_i, a_j) = 1$ for all $1 \leq i \neq j \leq n$, then we say that a_1, \dots, a_n are *pairwise relatively prime*.

Remark. Pairwise relatively prime implies relatively prime, but the converse is not true (e.g. $\{2, 4, 3\}$).

3.2 The Euclidean Algorithm

Lemma 3.1. If $a, b \in \mathbb{Z}$ with $0 < b \leq a$ and $a = bq + r$ with $q, r \in \mathbb{Z}$, then $(a, b) = (r, b)$.

Proof. It suffices to show that the two sets of common divisors (of a, b and of r, b) are the same. Denote by S_1 and S_2 these two sets, respectively. First let $c \in S_1$, so $c \mid a$ and $c \mid b$. We can write

$$r = a - bq,$$

so we have $c \mid r$. Thus $c \in S_2$, so $S_1 \subseteq S_2$. Now let $c \in S_2$, so $c \mid r$ and $c \mid b$. We have

$$a = bq + r$$

by hypothesis, so $c \mid a$, i.e. $c \in S_1$. Thus $S_1 = S_2$, so $(a, b) = \max S_1 = \max S_2 = (r, b)$. \square

Example 3.2.1. The above lemma allows us to compute greatest common divisors more efficiently. We will compute $(803, 154)$. We can write $803 = 5 \cdot 154 + 33$, so $(803, 154) = (154, 33)$. Continuing, we get

$$(803, 154) = (154, 33) = (33, 22) = (22, 11) = (11, 0) = 11.$$

Theorem 3.1 (Euclidean algorithm). Let $a, b \in \mathbb{Z}$ with $0 < b \leq a$. Set $r_{-1} = a$, $r_0 = b$, and inductively write $r_{i-1} = q_i r_i + r_{i+1}$ by the division algorithm for $n \geq 1$. Then $r_n = 0$ for some $n \geq 1$ and $(a, b) = r_{n-1}$.

Proof. Note that $r_1 > r_2 > r_3 > \cdots$. If $r_n \neq 0$ for all $n \geq 1$, then this is a strictly decreasing infinite sequence of positive integers, which is not possible. So $r_n = 0$ for some $n \geq 1$. The conclusion $(a, b) = r_{n-1}$ follows by repeatedly applying the lemma since $(a, b) = (r_i, r_{i+1}) = (r_{n-1}, 0) = r_{n-1}$. \square

Example 3.2.2. By reversing this process, we can write (a, b) explicitly as an integer linear combination of a, b . Using the previous example of computing $(803, 154)$, we can see that

$$\begin{aligned} (803, 154) &= 11 = 33 - 1 \cdot 22 \\ &= 33 - 1 \cdot (154 - 4 \cdot 33) = 5 \cdot 33 - 1 \cdot 154 \\ &= 5 \cdot (803 - 5 \cdot 154) - 1 \cdot 154 = 5 \cdot 803 - 26 \cdot 154. \end{aligned}$$

Thus we have found that $(803, 154) = 5 \cdot 803 - 26 \cdot 154$. Note that this representation is not unique, e.g. we can also write $11 = 19 \cdot 803 - 99 \cdot 154$. In fact, there are infinitely many such representations.

Lecture 4

Aug. 27 — Fundamental Theorem of Arithmetic

What's the difference between a mediocre clown and a rabbit in the gym? One's a bit funny, the other's a fit bunny.

4.1 The Fundamental Theorem of Arithmetic

Lemma 4.1 (Euclid). *Let $a, b \in \mathbb{Z}$ and let p be a prime. If $p \mid ab$, then $p \mid a$ or $p \mid b$.*

Proof. If $p \mid a$, then we are done, so assume $p \nmid a$. Then $(p, a) = 1$. Thus we can write $1 = ma + np$ for some $m, n \in \mathbb{Z}$. Since $p \mid ab$, we can write $ab = pc$ for some $c \in \mathbb{Z}$. Multiplying by b , we have

$$b = bma + bnp = m(cp) + nbp = p(mc + nb).$$

Thus we see that $p \mid b$, as desired. □

Remark. This fails if p is composite: Take $p = 6$, $a = 2$, and $b = 3$.

Exercise 4.1. Determine where the proof fails if p is composite.

Corollary 4.0.1. *Let $a_1, \dots, a_n \in \mathbb{Z}$ and p a prime. If $p \mid a_1 \cdots a_n$, then $p \mid a_i$ for some $1 \leq i \leq n$.*

Proof. Induct on n . The base case $n = 1$ is trivial. If $n = 2$, then this is just Lemma 4.1. Now suppose $n \geq 2$, and we show the result for $n + 1$. Specifically, assume that if $p \mid a_1 \cdots a_n$, then $p \mid a_i$ for some $1 \leq i \leq n$. Suppose $p \mid a_1 \cdots a_n a_{n+1}$. Then $p \mid (a_1 \cdots a_n) a_{n+1}$. So by Lemma 4.1, we have $p \mid a_1 \cdots a_n$ or $p \mid a_{n+1}$. If $p \mid a_{n+1}$, then we are done. Otherwise, $p \mid a_1 \cdots a_n$, so $p \mid a_i$ for some $1 \leq i \leq n$ by the induction hypothesis. In particular, $p \mid a_i$ for some $1 \leq i \leq n + 1$, as desired. □

Theorem 4.1 (Fundamental theorem of arithmetic). *Every integer $m > 1$ may be expressed in the form $m = p_1^{a_1} \cdots p_n^{a_n}$ where p_1, \dots, p_n are distinct primes and a_1, \dots, a_n are positive integers. This form is called the prime factorization of the integer m . Moreover, this factorization is essentially unique, i.e. unique up to permutations of the factors $p_i^{a_i}$.*

Proof. We first prove existence. Assume to the contrary that there exists $m > 1$ that does not have a prime factorization. Without loss of generality, we can assume m is the smallest such integer by the well-ordering principle. In particular, m cannot be prime. So $m = ab$ for some $1 < a, b < m$. Then a, b have prime factorizations. Thus so too does m , a contradiction.

Now we prove uniqueness. Assume that $m = p_1^{a_1} \cdots p_n^{a_n} = q_1^{b_1} \cdots q_r^{b_r}$. Without loss of generality, we can assume $p_1 < p_2 < \cdots < p_n$ and $q_1 < q_2 < \cdots < q_r$. We need to show that $n = r$, $p_i = q_i$ for each i , and $a_i = b_i$ for each i . Let $p_i \mid m$. Then $p_i \mid q_1^{b_1} \cdots q_r^{b_r}$, so $p_i \mid q_j$ for some $1 \leq j \leq r$. Thus $p_i = q_j$ since both are prime. Similarly, given q_i , we have $q_i = p_j$ for some j . Thus the primes in the two factorizations (as sets) are the same. Thus $n = r$, and by the ordering assumption, we have $p_i = q_i$ for each $1 \leq i \leq n$. So

$$m = p_1^{a_1} \cdots p_n^{a_n} = p_1^{b_1} \cdots p_n^{b_n}.$$

Suppose to the contrary that $a_i \neq b_i$ for some i . Without loss of generality, assume $a_i < b_i$. We have $p_i^{b_i} \mid m$, so $p_i^{b_i} \mid p_1^{a_1} \cdots p_{i-1}^{a_{i-1}} p_i^{a_i} p_{i+1}^{a_{i+1}} \cdots p_n^{a_n}$. Thus $p_i^{b_i - a_i} \mid p_1^{a_1} \cdots p_{i-1}^{a_{i-1}} p_{i+1}^{a_{i+1}} \cdots p_n^{a_n}$. Since $a_i < b_i$, we have $b_i - a_i > 0$, so $p_i \mid p_1^{a_1} \cdots p_{i-1}^{a_{i-1}} p_{i+1}^{a_{i+1}} \cdots p_n^{a_n}$ by the transitivity of divisibility. Then $p_i \mid p_j$ for some $j \neq i$, so $p_i = p_j$, which is a contradiction since the p_i are all distinct primes. This proves uniqueness. \square

Remark. This is one reason why we do not consider 1 to be a prime, as we would lose uniqueness.

Example 4.0.1. We can write $60 = 2^2 \cdot 3 \cdot 5$ and $756 = 2^2 \cdot 3^3 \cdot 7$.

4.2 Least Common Multiples

Definition 4.1. Let $a, b \in \mathbb{Z}$ with $a, b > 0$. The *least common multiple* of a and b , denoted $[a, b]$, is the least positive integer m such that $a \mid m$ and $b \mid m$.

Remark. Since ab is a common multiple of a and b , $[a, b]$ always exists by the well-ordering principle.

Example 4.1.1. We will compute $[6, 7]$. The multiples of 6 and 7 include:

$$\begin{aligned} 6 : 6, 12, 18, 24, 30, 36, 42, 48, \dots; \\ 7 : 7, 14, 21, 28, 35, 42, 49, \dots \end{aligned}$$

So we can see that $[6, 7] = 42 = 6 \cdot 7$. On the other hand, $[6, 8] = 24 \neq 6 \cdot 8$.

Remark. The fundamental theorem of arithmetic can be used to calculate both GCDs and LCMs.

Proposition 4.1. Let $a, b \in \mathbb{Z}$ with $a, b > 1$. Write $a = p_1^{a_1} \cdots p_n^{a_n}$ and $b = p_1^{b_1} \cdots p_n^{b_n}$, where the p_i are distinct primes, and $a_i, b_i \geq 0$. Then we have

$$(a, b) = p_1^{\min\{a_1, b_1\}} \cdots p_n^{\min\{a_n, b_n\}} \quad \text{and} \quad [a, b] = p_1^{\max\{a_1, b_1\}} \cdots p_n^{\max\{a_n, b_n\}}.$$

Proof. Left as an exercise. \square

Example 4.1.2. Calculate $(756, 2205)$ and $[756, 2205]$. We can write

$$756 = 2^2 \cdot 3^3 \cdot 5^0 \cdot 7^1 \quad \text{and} \quad 2205 = 2^0 \cdot 3^2 \cdot 5^1 \cdot 7^2.$$

So we have $(756, 2205) = 2^0 \cdot 3^2 \cdot 5^0 \cdot 7^1 = 63$ and $[756, 2205] = 2^2 \cdot 3^3 \cdot 5^1 \cdot 7^2 = 26460$.

Lemma 4.2. Given $x, y \in \mathbb{R}$, we have $\min\{x, y\} + \max\{x, y\} = x + y$.

Proof. The result is obvious if $x = y$. Otherwise, one is the minimum and the other is the maximum. \square

Theorem 4.2. Let $a, b \in \mathbb{Z}$ with $a, b > 1$. Then $(a, b)[a, b] = ab$.

Proof. Write $a = p_1^{a_1} \cdots p_n^{a_n}$ and $b = p_1^{b_1} \cdots p_n^{b_n}$ with $a_i, b_i \geq 0$ and p_i distinct. By Proposition 4.1,

$$\begin{aligned} (a, b)[a, b] &= p_1^{\min\{a_1, b_1\}} \cdots p_n^{\min\{a_n, b_n\}} p_1^{\max\{a_1, b_1\}} \cdots p_n^{\max\{a_n, b_n\}} \\ &= p_1^{\min\{a_1, b_1\} + \max\{a_1, b_1\}} \cdots p_n^{\min\{a_n, b_n\} + \max\{a_n, b_n\}} = p_1^{a_1 + b_1} \cdots p_n^{a_n + b_n} = ab, \end{aligned}$$

where the third equality follows from Lemma 4.2. □

Lecture 5

Sept. 3 — Congruences

No, Tony's the guy with no shins.

5.1 Dirichlet's Theorem

Theorem 5.1 (Dirichlet's theorem on primes in arithmetic progressions). *Let $a, b \in \mathbb{Z}$ with $a, b > 0$ and $(a, b) = 1$. Then the arithmetic progression $a, a + b, a + 2b, a + 3b, \dots$ contains infinitely many primes.*

Remark. Setting $a = b = 1$ recovers the fact that there are infinitely many primes.

Remark. The general case of Dirichlet's theorem is difficult, but we can use the fundamental theorem of arithmetic to prove some special cases, e.g. when $a = 3$ and $b = 4$.

Lemma 5.1. *Let $a, b \in \mathbb{Z}$. If a and b are expressible as $4n + 1$, then so is their product ab .*

Proof. Let $a = 4m + 1$ and $b = 4n + 1$. Then

$$ab = (4m + 1)(4n + 1) = 16mn + 4m + 4n + 1 = 4(4mn + m + n) + 1,$$

which proves the desired result. □

Proposition 5.1. *There are infinitely many primes of the form $4n + 3$ with $n \geq 0$.*

Proof. Assume to the contrary that there are finitely many primes of the form $4n + 3$, say $3, p_1, \dots, p_r$. Then consider the integer $N = 4p_1 \cdots p_r + 3$. The prime factorization of N must contain a prime of the form $4n + 3$, since otherwise N would be a product of primes of the form $4n + 1$, which must again be of the form $4n + 1$. Thus we have $3 \mid N$ or $p_i \mid N$ for some $1 \leq i \leq r$.

If $3 \mid N$, then $3 \mid N - 3 = 4p_1 \cdots p_r$, which is a contradiction. Otherwise, $p_i \mid N$ for some $1 \leq i \leq r$, and we have $p_i \mid N - 4p_1 \cdots p_r = 3$, which is a contradiction as well. □

Remark. The same proof does not work for primes of the form $4n + 1$, since a product of numbers of the form $4n + 3$ is not necessarily again of the form $4n + 3$.

5.2 Congruences

Definition 5.1. Let $a, b, m \in \mathbb{Z}$ with $m > 0$. Then we say that a is *congruent to b modulo m* , and we write $a \equiv b \pmod{m}$, if $m \mid (a - b)$. The integer m is called the *modulus* of the congruence. We write $a \not\equiv b \pmod{m}$ if a is not congruent to b modulo m .

Example 5.1.1. We have $25 \equiv 1 \pmod{4}$ and $25 \equiv 4 \pmod{7}$.

Proposition 5.2. *Congruence modulo m is an equivalence relation on \mathbb{Z} .*

Proof. Reflexivity is clear since $m \mid 0 = (a - a)$ any $a \in \mathbb{Z}$, so $a \equiv a \pmod{m}$. For symmetry, suppose that $a \equiv b \pmod{m}$. Then $m \mid a - b$. But then $m \mid (-1)(a - b) = b - a$, so $b \equiv a \pmod{m}$ as well.

Finally, for transitivity, suppose that $a \equiv b \pmod{m}$ and $b \equiv c \pmod{m}$. Then $m \mid a - b$ and $m \mid b - c$, so m also divides their sum $m \mid (a - b) + (b - c) = a - c$, i.e. $a \equiv c \pmod{m}$. \square

Remark. A consequence of Proposition 5.2 is that \mathbb{Z} is partitioned into its equivalence classes under congruence modulo m . For $a \in \mathbb{Z}$, we write $[a]$ to denote the equivalence class of a modulo m (not to be confused with the floor function).

Example 5.1.2. The equivalence classes of \mathbb{Z} under congruence modulo 4 are

$$\begin{aligned} [0] &= \{\dots, -8, -4, 0, 4, 8, \dots\}, \\ [1] &= \{\dots, -7, -3, 1, 5, 9, \dots\}, \\ [2] &= \{\dots, -6, -2, 2, 6, 10, \dots\}, \\ [3] &= \{\dots, -5, -1, 3, 7, 11, \dots\}. \end{aligned}$$

Definition 5.2. A set of m integers such that every integer is congruent modulo m to exactly one integer of the set is called a *complete residue system* modulo m .

Example 5.2.1. $\{0, 1, 2, 3\}$ is a complete residue system modulo 4. So is $\{4, 5, -6, -1\}$.

Proposition 5.3. *The set $\{0, 1, \dots, m-1\}$ is a complete residue system modulo m .*

Proof. First we prove that every integer is congruent to one of $0, 1, \dots, m-1$ modulo m . By the division algorithm, for any $a \in \mathbb{Z}$, there exist $q, r \in \mathbb{Z}$ with $0 \leq r \leq m-1$ such that $a = qm + r$. Thus we have $a - r = qm$, so $m \mid a - r$, i.e. $a \equiv r \pmod{m}$. This proves existence since $r \in \{0, 1, \dots, m-1\}$.

Now we show uniqueness. Suppose $a \equiv r_1 \pmod{m}$ and $a \equiv r_2 \pmod{m}$ where $r_1, r_2 \in \{0, 1, \dots, m-1\}$. By transitivity, we have $r_1 \equiv r_2 \pmod{m}$, so $m \mid r_1 - r_2$. But $0 \leq r_1, r_2 \leq m-1$, so

$$-(m-1) \leq r_1 - r_2 \leq m-1,$$

so we must have $r_1 - r_2 = 0$, i.e. $r_1 = r_2$. This proves uniqueness. \square

Definition 5.3. The set $\{0, 1, \dots, m-1\}$ is called the set of *least nonnegative residues* modulo m .

Proposition 5.4. *Let $a, b, c, d, m \in \mathbb{Z}$, $m > 0$ such that $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$. Then*

1. $a + c \equiv b + d \pmod{m}$;
2. $ac \equiv bd \pmod{m}$.

Proof. Since $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$, we have $m \mid b - a$ and $m \mid d - c$. Then m divides

$$(b - a) + (d - c) = (b + d) - (a + c),$$

so we have $a + c \equiv b + d \pmod{m}$. This proves (1).

To prove (2), note that since $m \mid a - b$, we also have $m \mid c(a - b)$. Likewise, $m \mid d - c$ implies $m \mid b(d - c)$. Then m divides the difference

$$c(a - b) - b(d - c) = ac - bd,$$

which shows that $ac \equiv bd \pmod{m}$. This shows (2). \square

Remark. This shows that the congruence classes of \mathbb{Z} modulo m form a *ring*.

Example 5.3.1. Consider the complete residue system $\{0, 1, 2, 3\}$ modulo 4. Their squares mod 4 are

$$\{0^2, 1^2, 2^2, 3^2\} \equiv \{0, 1, 0, 1\} \equiv \{0, 1\} \pmod{4}.$$

Lecture 6

Sept. 8 — Congruences, Part 2

6.1 More on Congruences

Example 6.0.1. Compute a complete residue system modulo 5 using

- only even numbers: $\{0, 2, 4, 6, 8\}$,
- only prime numbers: $\{2, 3, 5, 11, 19\}$.

Example 6.0.2. Compute a complete residue system modulo 5 using only numbers $\equiv 1 \pmod{4}$.

Remark. Recall that the set of equivalence classes of \mathbb{Z} modulo m form a ring. In particular, we can construct addition and multiplication tables. For $m = 4$, this looks like:

+	0	1	2	3	×	0	1	2	3
0	0	1	2	3	0	0	0	0	0
1	1	2	3	0	1	0	1	2	3
2	2	3	0	1	2	0	2	0	2
3	3	0	1	2	3	0	3	2	1

Addition modulo 5 is similar, but the multiplication table for $m = 5$ is:

×	0	1	2	3	4
0	0	0	0	0	0
1	0	1	2	3	4
2	0	2	4	1	3
3	0	3	1	4	2
4	0	4	3	2	1

Recall that a ring with no zero divisors (nonzero elements a, b such that $ab = 0$) is an *integral domain*, in particular we see from the multiplication table that $\mathbb{Z}/5\mathbb{Z}$ is an integral domain. Since a finite integral domain is automatically a *field*, we see that $\mathbb{Z}/5\mathbb{Z}$ is a field.

Proposition 6.1. Let $a, b, c, m, \in \mathbb{Z}$ with $m > 0$. Then

$$ca \equiv cb \pmod{m} \quad \text{if and only if} \quad a \equiv b \pmod{m/(m, c)}.$$

In particular, if m is prime, then $ca \equiv cb \pmod{m}$ if and only if $a \equiv b \pmod{m}$ for $c \not\equiv 0 \pmod{m}$.

Proof. (\Rightarrow) We have $ca \equiv cb \pmod{m}$ if and only if $m \mid ca - cb = c(a - b)$. Let $d = (m, c)$. By the transitivity of divisibility, we have $(m/d) \mid (c/d)(a - b)$. But $(m/d, c/d) = 1$, so $(m/d) \mid a - b$. Then we have $a \equiv b \pmod{m/d}$ by the definition of congruence.

(\Leftarrow) Again let $d = (m, c)$. Then $a \equiv b \pmod{m/d}$, so $(m/d) \mid a - b$. Then $m \mid d(a - b)$, and so

$$m \mid d(a - b)(c/d) = c(a - b) = ca - cb,$$

which means $ca \equiv cb \pmod{m}$ by the definition of congruence. \square

Remark. This shows that the congruence classes modulo m form a field if and only if m is prime.

6.2 Linear Congruences in One Variable

Definition 6.1. Let $a, b \in \mathbb{Z}$. A congruence of the form

$$ax \equiv b \pmod{m}$$

is called a *linear congruence* in the variable x .

Example 6.1.1. Consider the following linear congruences:

- $2x \equiv 3 \pmod{4}$ has no solutions;
- $2x \equiv 4 \pmod{6}$ has $x = 2, 5$ as solutions;
- $3x \equiv 9 \pmod{6}$ has $x = 1, 3, 5$ as solutions.

Theorem 6.1. Let $ax \equiv b \pmod{m}$, and let $d = (a, m)$. If $d \nmid b$, then there are no solutions for x in \mathbb{Z} . If $d \mid b$, then the congruence has exactly d incongruent solutions modulo m in \mathbb{Z} .

Proof. Note that $ax \equiv b \pmod{m}$ if and only if $m \mid ax - b$, if and only if $ax - b = my$ for some integer y . This is equivalent to $ax - my = b$. Thus $ax \equiv b \pmod{m}$ is solvable in x if and only if the equation $ax - my = b$ is solvable in x, y .

Let x, y be a solution of $ax - my = b$. Since $d \mid a$ and $d \mid m$, we must have $d \mid b$. Taking contrapositives, this proves the first part of the theorem.

Assume now that $d \mid b$. We prove the second part in 4 steps:

1. We will show that $ax \equiv b \pmod{m}$ has a solution x_0 .
2. We will show that there are infinitely many solutions of a particular form involving x_0 .
3. We will show that any solution has a particular form involving x_0 . (Note that this combines with (2) to give all possible solutions.)
4. We will show that there are exactly d equivalence classes of solutions.

(1) Since $d = (a, m)$, there exist $r, s \in \mathbb{Z}$ such that $d = ra + sm$. Since $d \mid b$, we can write

$$b = \frac{b}{d} \cdot d = \frac{b}{d}(ra + sm) = \frac{br}{d} \cdot a + \frac{bs}{d} \cdot m.$$

Thus $b - a(br/d) = (bs/d)m$, so $m \mid b - a(br/d)$, so $a(br/d) \equiv b \pmod{m}$. Thus $x_0 = br/d$ is a solution.

(2) Let x_0 be any solution of $ax \equiv b \pmod{m}$. Consider $x_0 + (m/d)n$ for $n \in \mathbb{Z}$. Then

$$a(x_0 + (m/d)n) \equiv ax_0 + a(m/d)n \equiv b + (a/d)mn \equiv b \pmod{m},$$

so $x_0 + (m/d)n$ is also solution for any $n \in \mathbb{Z}$.

(3) Let x_0 be a solution of $ax \equiv b \pmod{m}$. Recall from the beginning of the proof that this is equivalent to there being $y_0 \in \mathbb{Z}$ such that $ax_0 - my_0 = b$. Let x be any other solution. Then $ax - my = b$ for some $y \in \mathbb{Z}$, so

$$0 = b - b = (ax_0 - my_0) - (ax - my) = a(x_0 - x) - m(y_0 - y),$$

which gives $a(x_0 - x) = m(y_0 - y)$. This is equivalent to $(a/d)(x_0 - x) = (m/d)(y_0 - y)$. Note that if $y_0 - y = 0$, then $x_0 - x = 0$ as well since $a/d \neq 0$. So we may assume $y_0 - y \neq 0$. Then

$$(m/d) \mid (a/d)(x_0 - x),$$

and since $(a/d, m/d) = 1$, we have $(m/d) \mid (x_0 - x)$. Thus $x \equiv x_0 \pmod{m/d}$. In particular, all solutions to $ax \equiv b \pmod{m}$ are given by $x = x_0 + (m/d)n$ for $n \in \mathbb{Z}$ and any particular solution x_0 .

(4) Let $x_0 + (m/d)n_1$ and $x_0 + (m/d)n_2$ be solutions. Then we have

$$x_0 + (m/d)n_1 \equiv x_0 + (m/d)n_2 \pmod{m}$$

if and only if $(m/d)n_1 \equiv (m/d)n_2 \pmod{m}$. This happens if and only if $m \mid (m/d)(n_1 - n_2)$, if and only if $(m/d)(n_1 - n_2) = km$ for some $K \in \mathbb{Z}$, if and only if $n_1 - n_2 = kd$. In particular, this is equivalent to $n_1 \equiv n_2 \pmod{d}$. Since there are exactly d congruence classes for n , there are exactly d congruence classes of solutions as well, which completes the proof. \square

Lecture 7

Sept. 10 — Chinese Remainder Theorem

7.1 More on Linear Congruences

Corollary 7.0.1. Consider the linear congruence $ax \equiv b \pmod{m}$ and let $d = (a, m)$. If $d \mid b$, then there are exactly d incongruent solutions modulo m , given by

$$x = x_0 + \frac{m}{d} \cdot n, \quad n = 0, 1, \dots, d - 1$$

where x_0 is any particular solution.

Example 7.0.1. We solve $16x \equiv 8 \pmod{28}$. We compute $d = (16, 28)$ by the Euclidean algorithm:

$$28 = 1 \cdot 16 + 12$$

$$16 = 1 \cdot 12 + 4$$

$$12 = 3 \cdot 4 + 0.$$

So $d = 4$. Since $4 \mid 8$, the congruence has 4 incongruent solutions. Working backwards, we have

$$4 = 2 \cdot 16 + (-1) \cdot 28.$$

Multiplying by 2, we get that $8 = 4 \cdot 16 + (-2) \cdot 28$. Taking this equation modulo 28, we get

$$16 \cdot 4 \equiv 8 \pmod{28},$$

so $x_0 = 4$ is a particular solution. Thus all the incongruent solutions are given by $x = 4 + (28/4)n$ for $n = 0, 1, 2, 3$, that is $x = 4, 11, 18, 25$.

Definition 7.1. Any solution of $ax \equiv 1 \pmod{m}$ is called the *multiplicative inverse* of a modulo m . The multiplicative inverse of a is often denoted \bar{a} .

Corollary 7.0.2. The congruence $ax \equiv 1 \pmod{m}$ has a solution if and only if $(a, m) = 1$. In this case, the congruence has a unique solution. In particular, the multiplicative inverse, if it exists, is unique.

7.2 The Chinese Remainder Theorem

Example 7.1.1. Consider the following problem: Find a positive integer having remainder 2 when divided by 3, remainder 1 when divided by 4, and remainder 3 when divided by 5. The problem can be rephrased as asking for a solution to the system of congruences:

$$\begin{cases} x \equiv 2 \pmod{3} \\ x \equiv 1 \pmod{4} \\ x \equiv 3 \pmod{5}. \end{cases}$$

Theorem 7.1 (Chinese remainder theorem). *Let m_1, \dots, m_n be pairwise relatively prime positive integers, and let $b_1, \dots, b_n \in \mathbb{Z}$. Then the system of congruences*

$$\begin{cases} x \equiv b_1 \pmod{m_1} \\ x \equiv b_2 \pmod{m_2} \\ \vdots \\ x \equiv b_n \pmod{m_n} \end{cases}$$

has a unique solution modulo $M = m_1 \cdots m_n$.

Proof. Let $M = m_1 \cdots m_n$ and $M_i = M/m_i$. Then $(M_i, m_i) = 1$, so there are solutions to each system $M_i x_i \equiv 1 \pmod{m_i}$ given by $x_i = \overline{M}_i$. Consider

$$x = b_1 M_1 \overline{M}_1 + b_2 M_2 \overline{M}_2 + \cdots + b_n M_n \overline{M}_n.$$

Note that $m_i \mid M_j$ for $i \neq j$, so $x \equiv b_i M_i \overline{M}_i \equiv b_i \pmod{m_i}$, so x is a solution to the system.

For uniqueness modulo M , let x' be another solution. Then $x' \equiv b_i \pmod{m_i}$ for each $1 \leq i \leq n$. Then

$$x \equiv x' \pmod{m_i}, \quad 1 \leq i \leq n.$$

Thus $m_i \mid x - x'$, so $M \mid x - x'$ since the m_i are pairwise relatively prime, so $x \equiv x' \pmod{M}$. \square

Example 7.1.2. We now solve Example 7.1.1. Using the notation in the proof, we have

$$(m_1, m_2, m_3) = (3, 4, 5), \quad (b_1, b_2, b_3) = (2, 1, 3), \quad M = 60, \quad (M_1, M_2, M_3) = (20, 15, 12).$$

We still need to compute \overline{M}_i . In general, this can be done via the Euclidean algorithm. In this case,

$$(\overline{M}_1, \overline{M}_2, \overline{M}_3) = (2, 3, 3).$$

Now we can calculate the solution using

$$x = b_1 M_1 \overline{M}_1 + b_2 M_2 \overline{M}_2 + b_3 M_3 \overline{M}_3 = (2 \cdot 20 \cdot 2) + (1 \cdot 15 \cdot 3) + (3 \cdot 12 \cdot 3) = 233.$$

Reducing modulo 60, we get that the unique solution is given by $x \equiv 53 \pmod{60}$.

7.3 Wilson's Theorem

Lemma 7.1. *Let p be a prime and let $a \in \mathbb{Z}$. Then a is its own inverse modulo p (i.e., $a \equiv \overline{a} \pmod{p}$) if and only if $a \equiv \pm 1 \pmod{p}$.*

Proof. (\Rightarrow) Suppose $a \equiv \overline{a} \pmod{p}$. Then $a^2 \equiv a\overline{a} \equiv 1 \pmod{p}$, so $p \mid a^2 - 1 = (a - 1)(a + 1)$. Since p is prime, we have $p \mid a - 1$ or $p \mid a + 1$, so $a \equiv \pm 1 \pmod{p}$.

(\Leftarrow) This is obvious since $(\pm 1)^2 = 1$ in \mathbb{Z} , so they are also equal after reducing modulo p . \square

Theorem 7.2 (Wilson's theorem). *Let p be a prime. Then $(p - 1)! \equiv -1 \pmod{p}$.*

Example 7.1.3. The idea behind the proof is the following: Concretely, if $p = 11$, we have

$$(11 - 1)! = 10 \cdot 9 \cdot 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 \pmod{11}$$

By Lemma 7.1, 10 and 1 are their own inverses modulo 11. For each other integer $2 \leq n \leq 9$, we can pair them with their inverses: (2, 6), (3, 4), (5, 9), (7, 8). Then we can write

$$(11 - 1)! \equiv (9 \cdot 5) \cdot (8 \cdot 7) \cdot (6 \cdot 2) \cdot (4 \cdot 3) \cdot 10 \cdot 1 \equiv 10 \cdot 1 \equiv -1 \pmod{11}.$$

Proof of Theorem 7.2. We can easily check the theorem for $p = 2, 3$, so suppose $p > 3$ is a prime. Then each a with $1 \leq a \leq p - 1$ has a unique inverse modulo p , and this inverse is distinct from a if $2 \leq a \leq p - 2$. Pair each such integer with its inverse modulo p , say a and a' . The product of all of these pairs is $(p - 2)!$, so $(p - 2)! \equiv 1 \pmod{p}$. Thus $(p - 1)! \equiv p - 1 \equiv -1 \pmod{p}$. \square

Proposition 7.1 (Converse of Wilson's theorem). *Let $n \in \mathbb{Z}$ with $n > 1$. If $(n - 1)! \equiv -1 \pmod{n}$, then n is prime.*

Proof. Suppose $n = ab$ with $1 \leq a < n$. It suffices to show that $a = 1$. Since $a < n$, we have $a \mid (n - 1)!$. Also, $n \mid (n - 1)! + 1$ by assumption, so $a \mid (n - 1)! + 1$ also since $a \mid n$. Thus

$$a \mid ((n - 1)! + 1) - (n - 1)! = 1,$$

so we must have $a = 1$. \square

Definition 7.2. A prime p is a *Wilson prime* if $(p - 1)! \equiv -1 \pmod{p^2}$.

Example 7.2.1. The first few Wilson primes are 5, 13, 563. In fact, these are the only known ones.

Lecture 8

Sept. 15 — Fermat's Little Theorem

What do you call it when you have your grandmother on speed dial? It's an insta gram.

8.1 Fermat's Little Theorem

Theorem 8.1 (Fermat's little theorem). *Let p be a prime and $a \in \mathbb{Z}$ such that $p \nmid a$. Then*

$$a^{p-1} \equiv 1 \pmod{p}.$$

Proof. Consider the $p-1$ integers $a, 2a, 3a, \dots, (p-1)a$. Note that $p \nmid a_i$ for any $1 \leq i \leq p-1$. Note also that no two of these integers are congruent modulo p : If $ai \equiv aj \pmod{p}$ for some $i \neq j$, then we can multiply by the inverse \bar{a} of a (which exists since $p \nmid a$) to get $i \equiv j \pmod{p}$, which is impossible. Thus $\{a, 2a, \dots, (p-1)a\}$ is a complete nonzero residue system, so

$$a(2a)(3a) \cdots ((p-1)a) \equiv 1 \cdot 2 \cdot 3 \cdots (p-1) \pmod{p}.$$

Then $a^{p-1}(p-1)! \equiv (p-1)! \pmod{p}$, so $a^{p-1} \equiv 1 \pmod{p}$ since $p \nmid (p-1)!$. □

Corollary 8.1.1. *Let p be prime and $a \in \mathbb{Z}$ with $p \nmid a$. Then a^{p-2} is the inverse of a modulo p .*

Proof. By Fermat's little theorem, $a \cdot a^{p-2} = a^{p-1} \equiv 1 \pmod{p}$. □

Corollary 8.1.2. *Let p be prime and $a \in \mathbb{Z}$. Then $a^p \equiv a \pmod{p}$.*

Proof. If $p \mid a$, then both sides are congruent to 0 modulo p . Otherwise, if $p \nmid a$, then we can write $a^p = a \cdot a^{p-1} \equiv a \cdot 1 = a \pmod{p}$ by Fermat's little theorem. □

Corollary 8.1.3. *Let p be prime. Then $2^p \equiv 2 \pmod{p}$.*

Definition 8.1. If $n \in \mathbb{Z}$ is composite and $2^n \equiv 2 \pmod{n}$, then n is called a *pseudoprime*.

Remark. It is known that there are infinitely many (even and odd) pseudoprimes.

Example 8.1.1. Consider $n = 341 = 11 \cdot 31$. To prove that $2^{341} \equiv 2 \pmod{341}$, it suffices to show that $2^{341} \equiv 2 \pmod{11}$ and $2^{341} \equiv 2 \pmod{31}$ by the Chinese remainder theorem. Note that

$$\begin{aligned} 2^{341} &= (2^{10})^{34} \cdot 2 \equiv 1^{34} \cdot 2 = 2 \pmod{11} \\ 2^{341} &= (2^{30})^{11} \cdot 2^{11} \equiv 1^{11} \cdot (2^5)^2 \cdot 2 \equiv 1^2 \cdot 2 = 2 \pmod{31} \end{aligned}$$

by Fermat's little theorem, so 341 is a pseudoprime.

8.2 Euler's Theorem

Definition 8.2. Let $n \in \mathbb{Z}$, $n > 0$. Euler's phi function, denoted $\varphi(n)$, is the number of positive integers $\leq n$ that are relatively prime to n . In other words,

$$\varphi(n) = \#\{m \in \mathbb{Z} : 1 \leq m \leq n, (m, n) = 1\}.$$

Example 8.2.1. We have $\varphi(4) = 2$, $\varphi(14) = 6$, and $\varphi(p) = p - 1$ for any prime p .

Theorem 8.2 (Euler's theorem). Let $a, m \in \mathbb{Z}$ with $m > 0$. If $(a, m) = 1$, then

$$a^{\varphi(m)} \equiv 1 \pmod{m}.$$

Proof. Let $r_1, r_2, \dots, r_{\varphi(m)}$ be the distinct positive integers not exceeding m such that $(r_i, m) = 1$. Then consider the integers $ar_1, ar_2, \dots, ar_{\varphi(m)}$. Note first that $(ar_i, m) = 1$ since $(r_i, m) = 1$ and $(a, m) = 1$ by assumption. Note also that $ar_i \not\equiv ar_j \pmod{m}$ for $i \neq j$ since \bar{a} exists (since $(a, m) = 1$), and multiplying by \bar{a} implies $r_i \equiv r_j \pmod{m}$, which is impossible. Thus the least nonnegative residues of $\{ar_1, \dots, ar_{\varphi(m)}\}$ coincide with $\{r_1, \dots, r_{\varphi(m)}\}$, so we have

$$(ar_1)(ar_2) \cdots (ar_{\varphi(m)}) \equiv r_1 r_2 \cdots r_{\varphi(m)} \pmod{m},$$

thus $a^{\varphi(m)}(r_1 \cdots r_{\varphi(m)}) \equiv (r_1 \cdots r_{\varphi(m)}) \pmod{m}$. Since $(r_1 \cdots r_{\varphi(m)}, m) = 1$, the inverse of $r_1 \cdots r_{\varphi(m)}$ modulo m exists, and multiplying by the inverse gives $a^{\varphi(m)} \equiv 1 \pmod{m}$. \square

Remark. Taking $m = p$ recovers Fermat's little theorem since $\varphi(p) = p - 1$ for prime p .

Definition 8.3. Let m be a positive integer. A set of $\varphi(m)$ integers such that each integer is relatively prime to m and no two are congruent modulo m is called a *reduced residue system* modulo m .

Example 8.3.1. $\{1, 5, 7, 11\}$ is a reduced residue system modulo 12. So is

$$\{5(1), 5(5), 5(7), 5(11)\} = \{5, 25, 35, 55\}.$$

For a prime p , the set $\{1, 2, \dots, p - 1\}$ is always a reduced residue system modulo p .

Corollary 8.2.1. Let $a, m \in \mathbb{Z}$ with $m > 0$ and $(a, m) = 1$. Then $\bar{a} \equiv a^{\varphi(m)-1} \pmod{m}$.

8.3 Arithmetic Functions and Multiplicativity

Definition 8.4. An *arithmetic function* is a function whose domain is the set of positive integers.

Example 8.4.1. The following are examples of arithmetic functions:

1. Euler's φ function;
2. $\tau(n)$, the number of positive divisors of n ;
3. $\sigma(n)$, the sum of the positive divisors of n ;
4. $\omega(n)$, the number of distinct prime factors of n ;
5. $p(n)$, the number of integer partitions of n ;

6. $\Omega(n)$, the number of total prime factors (counted with multiplicity) of n .

Definition 8.5. An arithmetic function f is *multiplicative* if $f(mn) = f(m)f(n)$ whenever $(m, n) = 1$. We say that f is *completely multiplicative* if $f(mn) = f(m)f(n)$ for all m, n .

Remark. Note that if $n > 1$, then we can write $n = p_1^{a_1} \cdots p_r^{a_r}$. If f is multiplicative, then

$$f(n) = f(p_1^{a_1} \cdots p_r^{a_r}) = f(p_1^{a_1}) \cdots f(p_r^{a_r}).$$

So multiplicative functions are determined by their values at prime powers. If f is completely multiplicative, then $f(n) = f(p_1)^{a_1} \cdots f(p_r)^{a_r}$ and f is determined by its values at primes.

Example 8.5.1. The functions φ, v, σ from Example 8.4.1 are multiplicative, while ω, p, Ω are not.

Example 8.5.2. The functions $f(n) = 1$ and $f(n) = 0$ are completely multiplicative. The function f defined by $f(1) = 1$ and $f(n) > 0$ if $n > 1$ is also completely multiplicative.

Remark. If f is multiplicative and not identically zero, then $f(1) = 1$. To see this, take n such that $f(n) \neq 0$ (since f is not identically zero). Then $f(n) = f(n \cdot 1) = f(n)f(1)$, so $f(1) = 1$ since $f(n) \neq 0$.

Lecture 9

Sept. 17 — Arithmetic Functions

9.1 Properties of Multiplicative Functions

Remark. We write $\sum_{d|n} f(d)$ to denote a sum over the positive divisors of n . For instance,

$$\sum_{d|12} f(d) = f(1) + f(2) + f(3) + f(4) + f(6) + f(12).$$

Theorem 9.1. *Let f be an arithmetic function, and for $n \in \mathbb{Z}$, $n > 0$, define*

$$F(n) = \sum_{d|n} f(d).$$

If f is multiplicative, then so is F .

Proof. Let m, n be relative prime. We need to show that $F(mn) = F(m)F(n)$. We have

$$F(mn) = \sum_{d|mn} f(d).$$

We claim that every divisor d of mn can be written uniquely as $d = d_1 d_2$, where $d_1 | m$ and $d_2 | n$. Moreover, any such product $d_1 d_2$ is a divisor of mn . To see this, write $m = p_1^{a_1} \dots p_r^{a_r}$ and $n = q_1^{b_1} \dots q_s^{b_s}$, where all the $p_1, \dots, p_r, q_1, \dots, q_s$ are distinct. Then if $d | mn$, then

$$d = p_1^{e_1} \dots p_r^{e_r} q_1^{f_1} \dots q_s^{f_s}, \quad 0 \leq e_i \leq a_i, 0 \leq f_j \leq b_j.$$

Then we must choose $d_1 = p_1^{e_1} \dots p_r^{e_r}$ and $d_2 = q_1^{f_1} \dots q_s^{f_s}$, which proves the claim.

Using the claim, we can split the sum into

$$F(mn) = \sum_{d|mn} f(d) = \sum_{d_1|m} \sum_{d_2|n} f(d_1 d_2) = \sum_{d_1|m} \sum_{d_2|n} f(d_1) f(d_2) = F(m)F(n),$$

where we note that $(d_1, d_2) = 1$ since $(m, n) = 1$. □

Example 9.0.1. Let $m = 4$, $n = 3$. Then we can write

$$\begin{aligned} F(3 \cdot 4) &= \sum_{d|12} f(d) = f(1) + f(2) + f(3) + f(4) + f(6) + f(12) \\ &= f(1 \cdot 1) + f(1 \cdot 2) + f(3 \cdot 1) + f(1 \cdot 4) + f(3 \cdot 2) + f(3 \cdot 4) \\ &= f(1)f(1) + f(1)f(2) + f(3)f(1) + f(1)f(4) + f(3)f(2) + f(3)f(4) \\ &= (f(1) + f(3))(f(1) + f(2) + f(4)) = F(3)F(4). \end{aligned}$$

9.2 Properties of the Euler Phi Function

Theorem 9.2. *The Euler φ function is multiplicative.*

Proof. Let $m, n \in \mathbb{Z}$, $m, n > 0$ with $(m, n) = 1$. We need to show that $\varphi(mn) = \varphi(m)\varphi(n)$. Consider the array of positive integers $\leq mn$ organized as follows:

$$\begin{array}{cccccc} 1 & m+1 & 2m+1 & \cdots & (n-1)m+1 \\ 2 & m+2 & 2m+2 & \cdots & (n-1)m+2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ i & m+i & 2m+i & \cdots & (n-1)m+i \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ m & 2m & 3m & \cdots & nm \end{array}$$

Consider the i th row. If $(i, m) > 1$, then no element on the i th row is relatively prime to m (and hence cannot be relatively prime to mn). Thus we may restrict our attention to those i that satisfy $(i, m) = 1$. There are, by definition, $\varphi(m)$ such values of i . The entries in the i th row are

$$i, \quad m+i, \quad 2m+i, \quad \dots, \quad (n-1)m+i.$$

We claim that this is a complete residue system modulo n . To see this, suppose that

$$km+i \equiv jm+i \pmod{n}, \quad 0 \leq k, j \leq n-1.$$

Then $km \equiv jm \pmod{n}$. Since $(m, n) = 1$, this implies $k \equiv j \pmod{n}$. Since $0 \leq k, j \leq n-1$, we must have $k = j$. The claim follows since we have n non-congruent elements (modulo n) in the list. Thus, there are $\varphi(n)$ elements in the i th row that are relatively prime to n . Also, $(km+i, m) = (i, m) = 1$ by the Euclidean algorithm, so they are relatively prime to m as well. Thus $\varphi(mn) = \varphi(m)\varphi(n)$. \square

Theorem 9.3. *Let p be prime, $a \in \mathbb{Z}$, $a > 0$. Then $\varphi(p^a) = p^a - p^{a-1}$.*

Proof. The total number of integers not exceeding p^a is p^a . The only integers not relatively prime to p^a are the multiples of p : $p, 2p, 3p, \dots, (p^{a-1})p$. There are p^{a-1} such integers, so $\varphi(p^a) = p^a - p^{a-1}$. \square

Theorem 9.4. *Let $n \in \mathbb{Z}$, $n > 0$. Then*

$$\varphi(n) = n \prod_{p|n} \left(1 - \frac{1}{p}\right).$$

Proof. Write $n = p_1^{a_1} \cdots p_r^{a_r}$. Then

$$\begin{aligned} \varphi(n) &= \varphi(p_1^{a_1} \cdots p_r^{a_r}) = \varphi(p_1^{a_1}) \cdots \varphi(p_r^{a_r}) = (p_1^{a_1} - p_1^{a_1-1}) \cdots (p_r^{a_r} - p_r^{a_r-1}) \\ &= p_1^{a_1} \cdots p_r^{a_r} \left(1 - \frac{1}{p_1}\right) \cdots \left(1 - \frac{1}{p_r}\right) = n \prod_{p|n} \left(1 - \frac{1}{p}\right). \end{aligned}$$

This proves the desired formula. \square

Remark. One can interpret Theorem 9.4 probabilistically: It says that $\varphi(n)$ is n times the “probability” that an integer is not divisible by any of the primes dividing n .

Example 9.0.2. Consider $n = 504 = 2^3 \cdot 3^2 \cdot 7$. Then $\varphi(n)$ is given by

$$\varphi(504) = 504 \left(1 - \frac{1}{2}\right) \left(1 - \frac{1}{3}\right) \left(1 - \frac{1}{7}\right) = 504 \cdot \frac{1}{2} \cdot \frac{2}{3} \cdot \frac{6}{7} = 144.$$

Theorem 9.5 (Gauss). *Let $n \in \mathbb{Z}$, $n > 0$. Then*

$$\sum_{d|n} \varphi(d) = n.$$

Proof. Let d be a divisor of n . Define the set

$$S_d = \{1 \leq m \leq n : (m, n) = d\}.$$

Note that $(m, n) = d$ if and only if $(m/d, n/d) = 1$. Thus $|S_d| = \varphi(n/d)$. Note also that every integer less than or equal to n belongs to exactly one of the S_d , so

$$n = \sum_{d|n} |S_d| = \sum_{d|n} \varphi(n/d) = \sum_{d|n} \varphi(d),$$

which the last equality follows since $\{d : d \mid n\} = \{n/d : d \mid n\}$. □

Example 9.0.3. Let $n = 12$. We verify that $12 = \sum_{d|12} \varphi(d)$. Write the table

d	S_d
1	$\{1, 5, 7, 11\}$
2	$\{2, 10\}$
3	$\{3, 9\}$
4	$\{4, 8\}$
6	$\{6\}$
12	$\{12\}$

Summing the $|S_d| = \varphi(12/d)$, we indeed get $12 = 4 + 2 + 2 + 2 + 1 + 1$.

Lecture 10

Sept. 22 — Exam 1 Review

Why are Saturday and Sunday the strongest days? The other are week days.

10.1 Practice Problems

Exercise 10.1. Show that φ is multiplicative but not completely multiplicative.

Proof. The idea for the first part is to draw an $m \times n$ table of the first mn integers, see the proof of Theorem 9.2 for the details. For the second part, note that $\varphi(2) = 1$, $\varphi(2)^2 = 1$, but $\varphi(4) = 2$. \square

Alternative proof. Let R_k denote the set of residue classes modulo k that are coprime to k . Note that $|R_k| = \varphi(k)$, so it suffices to show there is a bijection $\psi : R_{mn} \rightarrow R_m \times R_n$ for $(m, n) = 1$. Define

$$\psi(a) = (a \bmod m, a \bmod n).$$

To see that ψ is surjective, let $(b, c) \in R_m \times R_n$. Since $(m, n) = 1$, by the Chinese remainder theorem there exists $a \in \mathbb{Z}$, defined modulo mn , such that $a \equiv b \pmod{m}$ and $a \equiv c \pmod{n}$. So $\psi(a) = (b, c)$. Note that $(a, mn) = 1$ since $(a, m) = (b, m) = 1$ and $(a, n) = (c, n) = 1$, so $a \in R_{mn}$. Injectivity follows since the choice of a is unique modulo mn by the Chinese remainder theorem. \square

Exercise 10.2. Compute $(163, 67)$ by the Euclidean algorithm.

Proof. We compute that

$$\begin{aligned}(163, 67) &= (163 - 2 \cdot 67, 67) = (29, 67) \\ &= (29, 67 - 2 \cdot 29) = (29, 9) \\ &= (29 - 3 \cdot 9, 9) = (2, 9) \\ &= (2, 9 - 4 \cdot 2) = (2, 1),\end{aligned}$$

so we have $(163, 67) = 1$. \square

Exercise 10.3. State and prove Wilson's theorem.

Proof. Wilson's theorem states that $(p-1)! \equiv -1 \pmod{p}$ for prime p (the converse also holds). The idea behind the proof is to note that each residue modulo p other than ± 1 can be paired with its (distinct) additive inverse modulo p . For the details, see the proof of Theorem 7.2. \square

Exercise 10.4. Find the least positive solution x to the congruence $x \equiv 20^{110} \pmod{17}$.

Proof. Use Fermat's little theorem: The division algorithm gives $110 = 6 \cdot 16 + 14$, so

$$\begin{aligned} x &\equiv 20^{6 \cdot 16 + 14} \equiv (20^{16})^6 \cdot 20^{14} \pmod{17} \\ &= 1^6 \cdot 20^{14} \equiv 20^{14} = 3^{14} \pmod{17}. \end{aligned}$$

Multiplying both sides by 3^2 gives $9x \equiv 3^2 x \equiv 3^{16} \equiv 1 \pmod{17}$, so it suffices to find the inverse of 9 modulo 17. Using the Euclidean algorithm, we have

$$(17, 9) = (17 - 9, 9) = (8, 9) = (8, 9 - 8) = (8, 1) = 1,$$

so $1 = 9 - 8 = 9 - (17 - 9) = 2 \cdot 9 - 17$. Thus $\bar{9} \equiv 2 \pmod{17}$, so we can take $x = 2$. \square

Exercise 10.5. Find the least positive solution x to the congruence $x \equiv 38^{110} \pmod{21}$.

Proof. First we compute that $\varphi(21) = \varphi(7)\varphi(3) = 6 \cdot 2 = 12$. By Euler's theorem,

$$x \equiv 38^{110} \equiv 17^{110} \equiv 179 \cdot 12 + 2 \equiv (17^{12})^9 \cdot 17^2 \equiv 17^2 \pmod{21}.$$

Now we notice that $17^2 \equiv (-4)^2 \equiv 16 \pmod{21}$, so we can take $x = 16$. \square

Exercise 10.6. Let $a, m \in \mathbb{Z}$ and $m > 1$. If $(a, m) = 1$, show that $a^{\varphi(m)-1}$ is the multiplicative inverse of a modulo m .

Proof. By Euler's theorem, $a \cdot a^{\varphi(m)-1} = a^{\varphi(m)} \equiv 1 \pmod{m}$, so $\bar{a} \equiv a^{\varphi(m)-1} \pmod{m}$. \square

Exercise 10.7. Prove that for odd primes p , we have $2(p-3)! \equiv -1 \pmod{p}$.

Proof. By Wilson's theorem, $(p-1)! \equiv -1 \pmod{p}$. Then we have

$$(p-3)!(p-2)(p-1) \equiv -1 \pmod{p},$$

so $2(p-3)! \equiv -1 \pmod{p}$ since $(p-2)(p-1) \equiv 2 \pmod{p}$. \square

Exercise 10.8. Find integers a, b such that $(a, b) = 3$ and $a + b = 66$.

Proof. It suffices to write $a = 3a_1$, $b = 3b_1$, where $(a_1, b_1) = 1$. One way to do this is $(a_1, b_1) = (1, 21)$:

$$3(a_1 + b_1) = 3(1 + 21) = 3 \cdot 22 = 66.$$

Thus we may take $a = 3$, $b = 63$. \square

Remark. Recall that a *reduced* residue system modulo m is a set $\{r_1, \dots, r_{\varphi(m)}\}$ of integers coprime to m and pairwise incongruent modulo m . Note that the r_i themselves need not be coprime, in fact they may share arbitrarily large common factors: Take any $r_\ell \neq 1$ and consider

$$\{r_\ell r_1, \dots, r_\ell r_{\varphi(m)}\}.$$

By repeating this, we can get arbitrarily large powers of r_ℓ as a common factor.

Lecture 11

Sept. 29 — Arithmetic Functions, Part 2

11.1 The Divisor Function

Definition 11.1. Let $n \in \mathbb{Z}$. The *number of positive divisors* of n , denoted $\tau(n)$, is defined by

$$\tau(n) = \#\{d \in \mathbb{Z} : d > 0, d \mid n\}.$$

Theorem 11.1. $\tau(n)$ is multiplicative.

Proof. Observe that $\tau(n) = \sum_{d \mid n} 1$ and 1 is multiplicative, so the result follows from Theorem 9.1. \square

Remark. Since $\tau(n)$ is multiplicative, it is determined by its behavior on prime powers.

Theorem 11.2. Let p be prime and let $a \in \mathbb{Z}$, $a > 0$. Then $\tau(p^a) = a + 1$.

Proof. The divisors of p^a are exactly the integers $1, p, p^2, \dots, p^a$. There are $a + 1$ of these. \square

Theorem 11.3. Let $n = p_1^{a_1} \cdots p_r^{a_r}$ with p_1, \dots, p_r distinct primes and a_1, \dots, a_r positive integers. Then

$$\tau(n) = \prod_{i=1}^r (a_i + 1).$$

Proof. This follows from τ being multiplicative and Theorem 11.2. \square

Remark. For some interesting reading about $\tau(n)$, see *Dirichlet's divisor problem*.

Example 11.1.1. Consider $504 = 2^3 \cdot 3^2 \cdot 7$. Then $\tau(504) = (3 + 1)(2 + 1)(1 + 1) = 24$.

11.2 The Sum of Divisors Function

Definition 11.2. Let $n \in \mathbb{Z}$, $n > 0$. The *sum of divisors function*, denoted $\sigma(n)$, is defined by

$$\sigma(n) = \sum_{d \mid n} d.$$

Theorem 11.4. $\sigma(n)$ is multiplicative.

Proof. This follows from $f(d) = d$ being multiplicative and Theorem 9.1. \square

Theorem 11.5. Let p be a prime and $a \in \mathbb{Z}$, $a > 0$. Then $\sigma(p^a) = (p^{a+1} - 1)/(p - 1)$.

Proof. The divisors of p^a are $1, p, p^2, \dots, p^a$, so

$$\sigma(p^a) = 1 + p + p^2 + \dots + p^a = \frac{p^{a+1} - 1}{p - 1}$$

by the formula for a (finite) geometric series. \square

Theorem 11.6. Let $n = p_1^{a_1} \cdots p_r^{a_r}$ with p_1, \dots, p_r distinct primes and a_1, \dots, a_r positive integers. Then

$$\sigma(n) = \prod_{i=1}^r \frac{p_i^{a_i+1} - 1}{p_i - 1}.$$

Example 11.2.1. Consider $504 = 2^3 \cdot 3^2 \cdot 7$. Then

$$\sigma(504) = \frac{2^{3+1} - 1}{2 - 1} \cdot \frac{3^{2+1} - 1}{3 - 1} \cdot \frac{7^{1+1} - 1}{7 - 1} = 15 \cdot 13 \cdot 8 = 1560.$$

11.3 Perfect Numbers

Definition 11.3. Let $n \in \mathbb{Z}$, $n > 0$. Then n is a *perfect number* if $\sigma(n) = 2n$, or $\sigma(n) - n = n$.

Remark. Note that $\sigma(n) - n$ is the sum of *proper* divisors of n , so perfect numbers are those that equal the sum of their proper divisors.

Example 11.3.1. 6 and 28 are perfect numbers.

Conjecture 11.6.1. There are infinitely many perfect numbers.

Conjecture 11.6.2. All perfect numbers are even.

Theorem 11.7. Let $n \in \mathbb{Z}$, $n > 0$. Then n is an even perfect number if and only if

$$n = 2^{p-1}(2^p - 1)$$

for some prime p , and $2^p - 1$ is prime (i.e. $2^p - 1$ is a Mersenne prime).

Proof. (\Rightarrow Euler) Assume n is an even perfect number. Then we can write $n = 2^a b$ with $a, b \in \mathbb{Z}$, $a \geq 1$, and b odd. Then we have

$$\sigma(2^a b) = \sigma(2^a) \sigma(b) = (2^{a+1} - 1) \sigma(b).$$

Also, since n is perfect, we can also write $\sigma(2^a b) = 2 \cdot 2^a b = 2^{a+1} b$, thus

$$(2^{a+1} - 1) \sigma(b) = 2^{a+1} b. \tag{1}$$

This implies that $2^{a+1} \mid (2^{a+1} - 1) \sigma(b)$, so $2^{a+1} \mid \sigma(b)$ since $(2^{a+1}, 2^{a+1} - 1) = 1$. Then $\sigma(b) = 2^{a+1} c$ (2) for some integer $c \geq 1$. Substituting this into (1), we get

$$(2^{a+1} - 1) 2^{a+1} c = 2^{a+1} b,$$

so we have $(2^{a+1} - 1)c = b$ (3). We now show that $c = 1$. Suppose to the contrary that $c > 1$. Then (3) implies that b has at least 3 distinct divisors, namely 1, b , c . Then

$$\sigma(b) \geq 1 + b + c.$$

But (2) implies $\sigma(b) = 2^{a+1}c = (2^{a+1} - 1 + 1)c = (2^{a+1} - 1)c + c = b + c$ by (3), a contradiction. So $c = 1$, and by (3), $b = 2^{a+1} - 1$. Also, (2) implies $\sigma(b) = b + 1$, so b must be prime. One can show that $2^{a+1} - 1$ being prime implies $a + 1$ is prime, so $n = 2^a(2^{a+1} - 1)$ with $2^{a+1} - 1$ and $a + 1$ prime.

(\Leftarrow Euclid) Assume that $n = 2^{p-1}(2^p - 1)$ with p and $2^p - 1$ both prime. Then

$$\sigma(2^{p-1}(2^p - 1)) = \sigma(2^{p-1})\sigma(2^p - 1) = (2^p - 1)(2^p - 1 + 1) = (2^p - 1)2^p = 2 \cdot 2^{p-1}(2^p - 1),$$

which shows that $\sigma(n) = 2n$. Thus n is perfect. \square

Remark. Theorem 11.7 gives a characterization of even perfect numbers and a bijection between even perfect numbers and Mersenne primes.

Example 11.3.2. The first 5 perfect numbers correspond to $p = 2, 3, 5, 7, 13$.

11.4 The Möbius Function

Definition 11.4. Let $n \in \mathbb{Z}$, $n > 0$. The *Möbius function*, denoted $\mu(n)$, is defined by

$$\mu(n) = \begin{cases} 1 & \text{if } n = 1, \\ 0 & \text{if } p^2 \mid n \text{ for some } p, \\ (-1)^r & \text{if } n = p_1 \cdots p_r \text{ with } p_i \text{ distinct primes.} \end{cases}$$

Example 11.4.1. Since $504 = 2^3 \cdot 3^2 \cdot 7$, we have $\mu(504) = 0$. On the other hand,

$$\mu(6) = (-1)^2 = 1 \quad \text{and} \quad \mu(30) = (-1)^3 = -1$$

Theorem 11.8. $\mu(n)$ is multiplicative.

Proof. Let m, n be relatively prime positive integers. We need to show that $\mu(mn) = \mu(m)\mu(n)$. This is clear if $m = 1$ or $n = 1$, so we may assume $m, n > 1$. Note that m or n is divisible by a square if and only if mn is divisible by a square, since $(m, n) = 1$. In this case, both $\mu(m)\mu(n)$ and $\mu(mn)$ are 0.

Now suppose m, n are products of distinct primes, say $m = p_1 \cdots p_r$ and $n = q_1 \cdots q_s$. Since $(m, n) = 1$, we have $p_i \neq q_j$ for any $1 \leq i \leq r$ and $1 \leq j \leq s$. Thus

$$\mu(mn) = \mu(p_1 \cdots p_r q_1 \cdots q_s) = (-1)^{r+s} = (-1)^r (-1)^s = \mu(p_1 \cdots p_r) \mu(q_1 \cdots q_s) = \mu(m) \mu(n),$$

as desired. So μ is multiplicative. \square

Proposition 11.1. Let $n \in \mathbb{Z}$, $n > 0$. Then

$$\sum_{d \mid n} \mu(d) = \begin{cases} 1 & \text{if } n = 1, \\ 0 & \text{if } n > 1. \end{cases}$$

Proof. Since μ is multiplicative, so is $F(n) = \sum_{d|n} \mu(d)$ by Theorem 9.1. Thus it suffices to show that $F(p^a) = 0$ for prime powers p^a . We have

$$F(p^a) = \sum_{d|p^a} \mu(d) = \mu(1) + \mu(p) + \mu(p^2) + \cdots + \mu(p^a).$$

Note that $p^2 \mid p^j$ for $j \geq 2$, so $\mu(p^j) = 0$ for $j \geq 2$. Thus

$$F(p^a) = \mu(1) + \mu(p) = 1 - 1 = 0.$$

It is clear that $F(1) = \sum_{d|1} \mu(d) = \mu(1) = 1$, so the result follows. \square

Example 11.4.2. Let $n = 12$. Then we have

$$\begin{aligned} \sum_{d|12} \mu(d) &= \mu(1) + \mu(2) + \mu(3) + \mu(4) + \mu(6) + \mu(12) \\ &= 1 - 1 - 1 + 0 + 1 + 0 = 0. \end{aligned}$$

Lecture 12

Oct. 1 — Quadratic Residues

What do you call a bear with no ear? A bee.

What do you call a magician who loses his magic? Ian.

12.1 Möbius Inversion

Theorem 12.1 (Möbius inversion). *Let f, g be arithmetic functions. Then*

$$f(n) = \sum_{d|n} g(d) \quad \text{if and only if} \quad g(n) = \sum_{d|n} \mu(d) f(n/d) = \sum_{d|n} \mu(n/d) f(d).$$

Proof. (\Rightarrow) Assume $f(n) = \sum_{d|n} g(d)$. Then we can write

$$\sum_{d|n} \mu(d) f(n/d) = \sum_{d|n} \mu(d) \sum_{a|(n/d)} g(a).$$

Note that $a | (n/d)$ if and only if $d | (n/a)$, so we can switch the order of summation to get

$$\sum_{d|n} \mu(d) f(n/d) = \sum_{a|n} g(a) \sum_{d|(n/a)} \mu(d) = \sum_{a|n} g(a) \begin{cases} 1 & \text{if } n = a \\ 0 & \text{otherwise} \end{cases} = g(n),$$

where the second equality is by Proposition 11.1. This proves the forward direction.

(\Leftarrow) Assume that $g(n) = \sum_{d|n} \mu(n/d) f(d)$. Then we have

$$\sum_{d|n} g(d) = \sum_{d|n} \sum_{a|d} \mu(d/a) f(a) = \sum_{a|n} f(a) \sum_{\substack{d|n \\ a|d}} \mu(d/a) = \sum_{a|n} f(a) \sum_{b|(n/a)} \mu(b)$$

where we let $d = ab$ and noted that $ab | n$ if and only if $b | (n/a)$. Then

$$\sum_{d|n} g(d) = \sum_{a|n} f(a) \begin{cases} 1 & \text{if } n = a \\ 0 & \text{otherwise} \end{cases} = f(n)$$

by Proposition 11.1, which proves the reverse direction. □

Example 12.0.1. Recall that $\sum_{d|n} \varphi(d) = n$ by Theorem 9.5. By Möbius inversion,

$$\varphi(n) = \sum_{d|n} \mu(d) \frac{n}{d} = n \sum_{d|n} \frac{\mu(d)}{d} = n \prod_{p^a|n} \sum_{d|p^a} \frac{\mu(d)}{d} = n \prod_{p|n} \left(1 - \frac{1}{p}\right),$$

where $\mu(d)/d$ is multiplicative since $\mu(d)$ and $1/d$ both are. This recovers the product formula for φ .

Example 12.0.2. We have $\tau(n) = \sum_{d|n} 1$. So by Möbius inversion,

$$1 = \sum_{d|n} \tau(n/d) \mu(d).$$

Example 12.0.3. We have $\sigma(n) = \sum_{d|n} d$. So by Möbius inversion,

$$n = \sum_{d|n} \mu(d) \sigma(n/d).$$

12.2 Quadratic Residues

Remark. So far, we have only studied linear congruences, which take the form $ax \equiv b \pmod{m}$. Now we will be interested in *quadratic* congruences, i.e. congruences of the form $ax^2 + bx \equiv c \pmod{m}$. We will primarily restrict to the case $x^2 \equiv a \pmod{p}$ for p an odd prime (the question is easy for $p = 2$).

Definition 12.1. Let $a, m \in \mathbb{Z}$, $m > 0$, and $(a, m) = 1$. Then a is a *quadratic residue modulo m* if the congruence $x^2 \equiv a \pmod{m}$ has a solution. Otherwise, a is a *quadratic non-residue modulo m* .

Example 12.1.1. The quadratic residues modulo 11 are

$$\{1^2, 2^2, 3^2, 4^2, 5^2, 6^2, 7^2, 8^2, 9^2, 10^2\} \equiv \{1, 4, 9, 5, 3, 3, 5, 9, 4, 1\} \equiv \{1, 3, 4, 5, 9\} \pmod{11}.$$

The quadratic non-residues are $\{2, 6, 7, 8, 10\}$. Note that the sizes of these sets are the same.

Proposition 12.1. Let p be an odd prime and $a \in \mathbb{Z}$, $p \nmid a$. Then $x^2 \equiv a \pmod{p}$ has either 0 or 2 incongruent solutions modulo p .

Proof. Assume $x^2 \equiv a \pmod{p}$ has a solution x_0 . Then $-x_0$ is also a solution. It is also incongruent to p , since if $x_0 \equiv -x_0 \pmod{p}$, then $2x_0 \equiv 0 \pmod{p}$, which implies $p \mid 2x_0$. Since p is odd, we must have $p \mid x_0$, so $x_0 \equiv 0 \pmod{p}$. But then $a \equiv x_0^2 \equiv 0 \pmod{p}$, a contradiction. Thus $x^2 \equiv a \pmod{p}$ has at least two incongruent solutions modulo p if it has a solution at all.

We now show $x^2 \equiv a \pmod{p}$ has at most 2 incongruent solutions. Suppose x_0, x_1 are solutions. Then

$$x_0^2 \equiv x_1^2 \equiv a \pmod{p}.$$

Then $x_0^2 - x_1^2 \equiv 0 \pmod{p}$, so $p \mid x_0^2 - x_1^2 = (x_0 - x_1)(x_0 + x_1)$. Thus $p \mid x_0 - x_1$ or $p \mid x_0 + x_1$. In the first case, $x_0 \equiv x_1 \pmod{p}$, and in the second case, $x_0 \equiv -x_1 \pmod{p}$. So any solution is congruent to either x_0 or $-x_0$, which means that $x^2 \equiv a \pmod{p}$ has at most 2 incongruent solutions. \square

Corollary 12.1.1. Let p be an odd prime and $a \in \mathbb{Z}$, $p \nmid a$. If $x^2 \equiv a \pmod{p}$ is solvable with $x = x_0$, then the two solutions are given by x_0 and $p - x_0$.

Proposition 12.2. *Let p be an odd prime. Then there are exactly $(p-1)/2$ quadratic residues and $(p-1)/2$ quadratic non-residues modulo p .*

Proof. For each $1 \leq x \leq p-1$, if $x^2 \equiv a \pmod{p}$, then $(p-x)^2 \equiv a \pmod{p}$ as well, and these are the only two such residues which square to a . That is, for each pair

$$(1, p-1), \quad (2, p-2), \quad \dots, \quad (i, p-i)$$

for $1 \leq i \leq (p-1)/2$, we get a unique quadratic residue, namely i^2 . Since there are $(p-1)/2$ pairs of residues modulo p formed in this way, there are exactly $(p-1)/2$ quadratic residues modulo p . These are given by $1^2, 2^2, \dots, ((p-1)/2)^2$. The remaining $(p-1)/2$ elements are quadratic non-residues. \square

12.3 The Legendre Symbol

Definition 12.2. Let p be an odd prime and $a \in \mathbb{Z}$, $p \nmid a$. The *Legendre symbol* is

$$\left(\frac{a}{p}\right) = \begin{cases} 1 & \text{if } a \text{ is a quadratic residue modulo } p, \\ -1 & \text{if } a \text{ is a quadratic non-residue modulo } p. \end{cases}$$

Example 12.2.1. Recall that 1, 3, 4, 5, 9 are quadratic residues modulo 11, so

$$\left(\frac{1}{11}\right) = \left(\frac{3}{11}\right) = \left(\frac{4}{11}\right) = \left(\frac{5}{11}\right) = \left(\frac{9}{11}\right) = 1.$$

The quadratic non-residues modulo 11 were 2, 6, 7, 8, 10, so

$$\left(\frac{2}{11}\right) = \left(\frac{6}{11}\right) = \left(\frac{7}{11}\right) = \left(\frac{8}{11}\right) = \left(\frac{10}{11}\right) = -1.$$

Example 12.2.2. Evaluate $\left(\frac{3}{7}\right)$. This asks whether 3 is a quadratic residue modulo 7. That is, whether there is a solution to the quadratic congruence $x^2 \equiv 3 \pmod{7}$. One can check that

$$\{1^2, 2^2, 3^2\} \equiv \{1, 4, 2\} \pmod{7},$$

and these are all of the quadratic residues by Proposition 12.2. Thus $\left(\frac{3}{7}\right) = -1$.

Lecture 13

Oct. 8 — The Legendre Symbol

13.1 More on the Legendre Symbol

Exercise 13.1. Find all the quadratic residues modulo 23.

We know that the quadratic residues modulo 23 are the squares of $1, \dots, 11$, so

$$\begin{aligned}\{1^2, 2^2, 3^2, 4^2, 5^2, 6^2, 7^2, 8^2, 9^2, 10^2, 11^2\} &\equiv \{1, 4, 9, 16, 2, 13, 3, 18, 12, 8, 6\} \pmod{23} \\ &= \{1, 2, 3, 4, 6, 8, 9, 12, 13, 16, 18\} \pmod{23}\end{aligned}$$

is the set of quadratic residues modulo 23.

Theorem 13.1 (Euler's criterion). *Let p be an odd prime with $a \in \mathbb{Z}$ and $p \nmid a$. Then*

$$\left(\frac{a}{p}\right) \equiv a^{(p-1)/2} \pmod{p}.$$

Proof. Suppose first that $\left(\frac{a}{p}\right) = 1$. Then $x^2 \equiv a \pmod{p}$ has a solution, say $x = x_0$. Then

$$a^{(p-1)/2} \equiv (x_0^2)^{(p-1)/2} \equiv x_0^{p-1} \equiv 1 \equiv \left(\frac{a}{p}\right) \pmod{p}$$

by Fermat's little theorem. Now suppose that $\left(\frac{a}{p}\right) = -1$. Since $p \nmid a$, for each $1 \leq i \leq p-1$, the linear congruence $ij \equiv a \pmod{p}$ has a unique solution j with $1 \leq j \leq p-1$. Note that $i \neq j$ since a is not a quadratic residue modulo p . Thus we can pair the residues $1, 2, \dots, p-1$ into $(p-1)/2$ pairs (i, j) such that $ij \equiv a \pmod{p}$. Multiplying these pairs together, we get

$$(p-1)! \equiv 1 \cdot 2 \cdot 3 \cdots (p-1) \equiv a^{(p-1)/2} \pmod{p}.$$

The left-hand side is congruent to $-1 = \left(\frac{a}{p}\right)$ by Wilson's theorem, which completes the proof. \square

Example 13.0.1. We compute $\left(\frac{3}{7}\right)$. Using Euler's criterion,

$$\left(\frac{3}{7}\right) \equiv 3^{(7-1)/2} \equiv 3^3 \equiv 27 \equiv 6 \equiv -1 \pmod{7},$$

so we get that $\left(\frac{3}{7}\right) = -1$.

Proposition 13.1. *Let p be an odd prime and $a, b \in \mathbb{Z}$ with $p \nmid a$ and $p \nmid b$. Then*

1. $\left(\frac{a^2}{p}\right) = 1$;
2. if $b \equiv a \pmod{p}$, then $\left(\frac{b}{p}\right) = \left(\frac{a}{p}\right)$;
3. $\left(\frac{ab}{p}\right) = \left(\frac{a}{p}\right)\left(\frac{b}{p}\right)$.

Proof. (1) The congruence $x^2 \equiv a^2 \pmod{p}$ has a solution $x = a$.

(2) The congruence $x^2 \equiv a \pmod{p}$ is equivalent to the congruence $x^2 \equiv b \equiv a \pmod{p}$.

(3) By Euler's criterion, $\left(\frac{ab}{p}\right) \equiv (ab)^{(p-1)/2} \equiv a^{(p-1)/2}b^{(p-1)/2} \equiv \left(\frac{a}{p}\right)\left(\frac{b}{p}\right) \pmod{p}$. Since $\left(\frac{ab}{p}\right)$, $\left(\frac{a}{p}\right)$, and $\left(\frac{b}{p}\right)$ are each ± 1 , congruence modulo p is equivalent to equality (since $1 \neq -1$ for $p \geq 3$). \square

Example 13.0.2. Calculate $\left(\frac{-11}{7}\right)$. Using the above properties and Euler's criterion, we have

$$\left(\frac{-11}{7}\right) = \left(\frac{-1}{7}\right)\left(\frac{11}{7}\right) = \left(\frac{-1}{7}\right)\left(\frac{4}{7}\right) = \left(\frac{-1}{7}\right) \equiv (-1)^3 \equiv -1 \pmod{7}$$

since 4 is a quadratic residue modulo 7. So $\left(\frac{-11}{7}\right) = -1$.

13.2 Particular Cases of the Legendre Symbol

Remark. If $a = \pm 2^{a_0} p_1^{a_1} \cdots p_r^{a_r}$, then we have

$$\left(\frac{a}{p}\right) = \left(\frac{\pm 1}{p}\right) \left(\frac{2}{p}\right)^{a_1} \left(\frac{p_1}{p}\right)^{a_1} \cdots \left(\frac{p_r}{p}\right)^{a_r}.$$

Thus to evaluate $\left(\frac{a}{p}\right)$, it suffices to understand $\left(\frac{-1}{p}\right)$, $\left(\frac{2}{p}\right)$, and $\left(\frac{q}{p}\right)$ for odd primes q .

Theorem 13.2. Let p be an odd prime. Then

$$\left(\frac{-1}{p}\right) = (-1)^{(p-1)/2} = \begin{cases} 1 & \text{if } p \equiv 1 \pmod{4}, \\ -1 & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

Proof. The first equality follows from Euler's criterion. The second is a direct computation: Note that p can only be congruent to 1 or 3 modulo 4. If $p \equiv 1 \pmod{4}$, then $p = 1 + 4k$ for some $k \in \mathbb{Z}$. Then

$$(-1)^{(p-1)/2} = (-1)^{(1+4k-1)/2} = (-1)^{2k} = 1.$$

Similarly, if $p \equiv 3 \pmod{4}$, then $p = 3 + 4k$ for some $k \in \mathbb{Z}$, and

$$(-1)^{(p-1)/2} = (-1)^{(3+4k-1)/2} = (-1)^{1+2k} = -1.$$

This proves the second equality. \square

Lemma 13.1 (Gauss's lemma). Let p be an odd prime and let $a \in \mathbb{Z}$ with $p \nmid a$. Let n be the number of least positive residues of the integers

$$a, \quad 2a, \quad 3a, \quad \dots, \quad \left(\frac{p-1}{2}\right)a$$

that are greater than $p/2$. Then $\left(\frac{a}{p}\right) = (-1)^n$.

Proof. Let r_1, \dots, r_n be the least positive residues among $a, 2a, \dots, ((p-1)/2)a$ that are greater than $p/2$, and let s_1, \dots, s_m be the residues which are less than $p/2$. Note that none of the r_i, s_j are congruent to 0 modulo p since $p \nmid a$. Now consider the $(p-1)/2$ integers given by

$$p - r_1, \quad p - r_2, \quad \dots, \quad p - r_n, \quad s_1, \quad s_2, \quad \dots, \quad s_m.$$

We claim that this is the set of residues $1, 2, \dots, (p-1)/2$ in some order. All elements are ≥ 1 , and are $\leq (p-1)/2$ since they are $< p/2$ and are integers. So it suffices to show that there are no duplicates.

If $p - r_i \equiv p - r_j \pmod{p}$, then $r_i \equiv r_j \pmod{p}$, so $k_i a \equiv k_j a \pmod{p}$ for some $k_i \neq k_j$. Since $(a, p) = 1$, we can multiply by its inverse \bar{a} to get $k_i \equiv k_j \pmod{p}$, which is a contradiction. By a similar argument, the s_j are all distinct. It only remains to consider $p - r_i \equiv s_j \pmod{p}$. Then

$$-k_i a \equiv k_j a \pmod{p}$$

for some $1 \leq k_i, k_j \leq (p-1)/2$. The congruence then implies $-k_i \equiv k_j \pmod{p}$. But $p - k_i > p/2 \geq (p-1) \geq k_j$, so this congruence is impossible. This proves the claim.

Thus, multiplying all the numbers together gives

$$\begin{aligned} \left(\frac{p-1}{2}\right)! &\equiv (p - r_1) \cdots (p - r_n) s_1 \cdots s_m \equiv (-1)^n a \cdot 2a \cdots \left(\frac{p-1}{2}\right) a \\ &\equiv (-1)^n a^{(p-1)/2} \left(\frac{p-1}{2}\right)! \pmod{p}. \end{aligned}$$

Since $p \nmid ((p-1)/2)!$, we can multiply by its inverse to get $a^{(p-1)/2} \equiv (-1)^n \pmod{p}$. The result then follows since the left-hand side is congruent to $\left(\frac{a}{p}\right)$ by Euler's criterion and $\left(\frac{a}{p}\right), (-1)^n$ are ± 1 . \square

Example 13.0.3. We use Gauss's lemma to calculate $\left(\frac{6}{11}\right)$. We have $\left(\frac{6}{11}\right) = (-1)^n$, where n is the number of least positive residues among

$$6, \quad 2 \cdot 6, \quad 3 \cdot 6, \quad 4 \cdot 6, \quad 5 \cdot 6$$

that are larger than $11/2 = 5.5$. Reducing the above gives $\{6, 1, 7, 2, 8\}$, so $n = 3$. Thus $\left(\frac{6}{11}\right) = -1$.

Lecture 14

Oct. 13 — Quadratic Reciprocity

Why did the man bring his watch to the bank? He wanted to save time.

14.1 Applications of Gauss's Lemma

Exercise 14.1. Calculate the following:

$$\left(\frac{-1}{13}\right), \quad \left(\frac{2}{17}\right), \quad \left(\frac{-14}{11}\right), \quad \left(\frac{18}{23}\right).$$

For the first, since $13 \equiv 1 \pmod{4}$, we have $\left(\frac{-1}{13}\right) = 1$. For the second, we compute

$$\{2, 2(2), 3(2), 4(2), 5(2), 6(2), 7(2), 8(2)\} \equiv \{2, 4, 6, 8, 10, 12, 14, 16\} \pmod{17}.$$

Four of these residues are greater than $17/2 = 8.5$, so $\left(\frac{2}{17}\right) = (-1)^4 = 1$. For the third, write

$$\left(\frac{-14}{11}\right) = \left(\frac{-1}{11}\right) \left(\frac{14}{11}\right) = \left(\frac{-1}{11}\right) \left(\frac{3}{11}\right) = -\left(\frac{3}{11}\right)$$

since $11 \equiv 3 \pmod{4}$. To compute $\left(\frac{3}{11}\right)$, we list

$$\{3, 2(3), 3(3), 4(3), 5(3)\} \equiv \{3, 6, 9, 1, 4\} \pmod{11}.$$

Two of the above are greater than $11/2 = 5.5$, so $\left(\frac{3}{11}\right) = (-1)^2 = 1$. Thus $\left(\frac{-14}{11}\right) = -1$. For the last,

$$\left(\frac{18}{23}\right) = \left(\frac{2}{23}\right) \left(\frac{9}{23}\right) = \left(\frac{2}{23}\right),$$

which we can compute by enumerating

$$\begin{aligned} &\{2, 2(2), 3(2), 4(2), 5(2), 6(2), 7(2), 8(2), 9(2), 10(2), 11(2)\} \\ &\equiv \{2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22\} \pmod{23}. \end{aligned}$$

Six of the above are greater than $23/2 = 11.5$, so $\left(\frac{18}{23}\right) = \left(\frac{2}{23}\right) = (-1)^6 = 1$.

Theorem 14.1. *Let p be an odd prime. Then*

$$\left(\frac{2}{p}\right) = (-1)^{(p^2-1)/8} = \begin{cases} 1 & \text{if } p \equiv 1, 7 \pmod{8}, \\ -1 & \text{if } p \equiv 3, 5 \pmod{8}. \end{cases}$$

Proof. By Gauss's lemma, we have $\left(\frac{2}{p}\right) = (-1)^n$, where n is the number of least positive residues of

$$2, \quad 2(2), \quad 3(2), \quad 4(2), \quad \dots, \quad \left(\frac{p-1}{2}\right) 2.$$

Let $k \in \mathbb{Z}$ with $1 \leq k \leq (p-1)/2$. Note that $2k < p/2$ if and only if $k < p/4$ (we always have $2k < p$), so there are $\lfloor p/4 \rfloor$ values of k for which $2k < p/2$. Thus, there are $(p-1)/2 - \lfloor p/4 \rfloor$ values of k for which $2k > p/2$ (recall that p is odd), so $n = (p-1)/2 - \lfloor p/4 \rfloor$. To show that $(p^2-1)/8$ and $(p-1)/2 - \lfloor p/4 \rfloor$ always have the same parity, we can just check the four cases:

- $p \equiv 1 \pmod{8}$. Then $p = 8m + 1$ for some $m \in \mathbb{Z}$. Then

$$n = \frac{p-1}{2} - \left\lfloor \frac{p}{4} \right\rfloor = \frac{8m+1-1}{2} - \left\lfloor \frac{8m+1}{4} \right\rfloor = 4m - 2m = 2m.$$

Note that this is even. On the other hand, we can check that

$$\frac{p^2-1}{8} = \frac{(8m+1)^2-1}{8} = 8m^2 + 2m.$$

This is also even, so the parity matches in this case.

- Check the cases $p \equiv 3, 5, 7 \pmod{8}$ similarly as an exercise.

Since $(p^2-1)/8$ and n agree modulo 2, we have $(-1)^{(p^2-1)/8} = (-1)^n = \left(\frac{2}{p}\right)$. □

Example 14.0.1. We have $\left(\frac{2}{23}\right) = 1$ since $23 \equiv 7 \pmod{8}$.

14.2 Quadratic Reciprocity

Remark. We will now try to understand $\left(\frac{q}{p}\right)$ for distinct odd primes p, q .

Theorem 14.2 (Law of quadratic reciprocity). *Let p, q be distinct odd primes. Then*

$$\left(\frac{p}{q}\right) \left(\frac{q}{p}\right) = (-1)^{(p-1)(q-1)/4} = \begin{cases} 1 & \text{if } p \equiv 1 \text{ or } q \equiv 1 \pmod{4}, \\ -1 & \text{if } p \equiv q \equiv 3 \pmod{4}. \end{cases}$$

Remark. Quadratic reciprocity allows us to simplify the calculation for $\left(\frac{p}{q}\right)$. For example, consider

Which primes are quadratic residues modulo 17, i.e. evaluate $\left(\frac{p}{17}\right)$?

This is a finite problem: We may just compute all squares modulo 17. Now consider

For which primes p is 17 a quadratic residue, i.e. evaluate $\left(\frac{17}{p}\right)$?

This is a priori an infinite problem, but we can convert it to the previous one by quadratic reciprocity.

Example 14.0.2. Compute $\left(\frac{7}{53}\right)$. We use quadratic reciprocity: $7 \equiv 3 \pmod{4}$ and $53 \equiv 1 \pmod{4}$, so

$$\left(\frac{7}{53}\right) = \left(\frac{53}{7}\right) = \left(\frac{4}{7}\right) = 1$$

since 4 is always a square modulo any prime.

Example 14.0.3. Calculate $\left(\frac{-158}{101}\right)$. We can first write

$$\left(\frac{-158}{101}\right) = \left(\frac{-1}{101}\right) \left(\frac{158}{101}\right) = \left(\frac{158}{101}\right) = \left(\frac{57}{101}\right) = \left(\frac{3}{101}\right) \left(\frac{19}{101}\right)$$

since $101 \equiv 1 \pmod{4}$, $158 \equiv 57 \pmod{101}$, and $57 = 3 \cdot 19$. We can now apply quadratic reciprocity (note that we could not have done this earlier, since 158, 57 are not prime):

$$\left(\frac{-158}{101}\right) = \left(\frac{101}{3}\right) \left(\frac{101}{19}\right) = \left(\frac{2}{3}\right) \left(\frac{6}{19}\right) = \left(\frac{2}{3}\right) \left(\frac{25}{19}\right) = -1 \cdot 1 = 1.$$

Lemma 14.1. Let p be an odd prime number and let $a \in \mathbb{Z}$, $p \nmid a$, a odd. Let

$$N = \sum_{j=1}^{(p-1)/2} \left\lfloor \frac{ja}{p} \right\rfloor.$$

Then $\left(\frac{a}{p}\right) = (-1)^N$.

Proof. Let r_1, r_2, \dots, r_n be the least non-negative residues among $a, 2a, \dots, ((p-1)/2)a$ that are $> p/2$. Likewise, let s_1, \dots, s_m be the remaining residues that are $< p/2$. Note that

$$r_1, \dots, r_n, s_1, \dots, s_m$$

are all distinct modulo p (they come from $a, 2a, 3a, \dots, ((p-1)/2)a$, which are distinct since $p \nmid a$). This means that the fractions r_i/p , s_j/p are also all distinct. Then

$$ja = p \cdot \frac{ja}{p} = p \left(\left\lfloor \frac{ja}{p} \right\rfloor + \frac{\text{remainder}}{p} \right) = p \left\lfloor \frac{ja}{p} \right\rfloor + \text{remainder depending on } j,$$

where the remainders are exactly the numbers $r_1, \dots, r_n, s_1, \dots, s_m$. Then

$$\sum_{j=1}^{(p-1)/2} ja = \sum_{j=1}^{(p-1)/2} p \left\lfloor \frac{ja}{p} \right\rfloor + \sum_{i=1}^n r_i + \sum_{j=1}^m s_j. \quad (1)$$

Note also that

$$\sum_{j=1}^{(p-1)/2} j = \sum_{i=1}^n (p - r_i) + \sum_{j=1}^m s_j = pn - \sum_{i=1}^n r_i + \sum_{j=1}^m s_j. \quad (2)$$

Subtracting (2) from (1) gives the equation

$$\sum_{j=1}^{(p-1)/2} j(a-1) = \sum_{j=1}^{(p-1)/2} p \left\lfloor \frac{ja}{p} \right\rfloor - pn + 2 \sum_{i=1}^n r_i.$$

Taking the above equation modulo 2, since a is odd, we get

$$\sum_{j=1}^{(p-1)/2} p \left\lfloor \frac{ja}{p} \right\rfloor - pn \equiv 0 \pmod{2},$$

so $pN \equiv pn \pmod{2}$, so $N \equiv n \pmod{2}$ since $2 \nmid p$. So $(-1)^N = (-1)^n = \left(\frac{a}{p}\right)$ by Gauss's lemma. \square

Example 14.0.4. We compute $\left(\frac{7}{11}\right)$ using Lemma 14.1. We calculate

$$N = \sum_{j=1}^5 \left\lfloor \frac{7j}{11} \right\rfloor = \left\lfloor \frac{7}{11} \right\rfloor + \left\lfloor \frac{14}{11} \right\rfloor + \left\lfloor \frac{21}{11} \right\rfloor + \left\lfloor \frac{28}{11} \right\rfloor + \left\lfloor \frac{35}{11} \right\rfloor = 0 + 1 + 1 + 2 + 3 = 7,$$

so $\left(\frac{7}{11}\right) = (-1)^7 = -1$ by Lemma 14.1.

Lecture 15

Oct. 15 — Quadratic Reciprocity, Part 2

What do you call a place in America that receives shipments of pollinators? A U.S. bee port.

15.1 Proof of Quadratic Reciprocity

Proof of Theorem 14.2. Without loss of generality, assume $p > q$. Consider a $q \times p$ grid on \mathbb{R}^2 . Let L be the line from $(0, 0)$ to $N = (q, p)$. Let $A = ((p-1)/2, 0)$, $B = (0, (q-1)/2)$. Let M be the intersection of L with the line $x = (p-1)/2$ and D be the intersection of L with the line $y = (q-1)/2$. Also let $C = ((p-1)/2, (q-1)/2)$. We count the number of lattice points in the rectangle $OABC$, not including the axes. This number is clearly $(p-1)(q-1)/4$. Now observe that:

1. The line ON has slope q/p . In particular, ON contains no lattice points.
2. The y -coordinate of M is $((p-1)/2)(q/p) = q/2 - q/2p$. This lies between the consecutive integers $(q-1)/2$ and $(q+1)/2$:

$$\frac{q-1}{2} = \frac{q}{2} - \frac{1}{2} < \frac{q}{2} - \frac{q}{2p} < \frac{q}{2} < \frac{q+1}{2}.$$

So the number of lattice points in $OABC$ excluding the axes, and below the line ON is

$$N_1 = \sum_{j=1}^{(p-1)/2} \left\lfloor \frac{jq}{p} \right\rfloor.$$

Likewise, the number of lattice points above the line ON is

$$N_2 = \sum_{j=1}^{(q-1)/2} \left\lfloor \frac{jp}{q} \right\rfloor.$$

Thus the total number of lattice points in question is $N_1 + N_2 = (p-1)(q-1)/4$. Then

$$\left(\frac{p}{q}\right) \left(\frac{q}{p}\right) = (-1)^{N_2} (-1)^{N_1} = (-1)^{N_1+N_2} = (-1)^{(p-1)(q-1)/4}$$

by Lemma 14.1, which proves the claim. □

15.2 Applications of Quadratic Reciprocity

Remark. Note that we have characterized the primes for which -1 and 2 are quadratic residues.

Example 15.0.1. For what primes p is 3 a quadratic residue? It suffices to compute when $\left(\frac{3}{p}\right) = 1$. By quadratic reciprocity, we have

$$\left(\frac{3}{p}\right) = \begin{cases} \left(\frac{p}{3}\right) & p \equiv 1 \pmod{4}, \\ -\left(\frac{p}{3}\right) & p \equiv 3 \pmod{4}. \end{cases}$$

Note that the only (non-zero) quadratic residue modulo 3 is 1. If $p \equiv 1 \pmod{4}$, then $\left(\frac{p}{3}\right) = 1$ if and only if $p \equiv 1 \pmod{3}$. If $p \equiv 3 \pmod{4}$, then $\left(\frac{p}{3}\right) = -1$ if and only if $p \equiv 2 \pmod{3}$. By the Chinese remainder theorem, we can rewrite the first condition as $p \equiv 1 \pmod{12}$ and the second condition as $p \equiv -1 \pmod{12}$. Thus we see that $\left(\frac{3}{p}\right) = 1$ if and only if $p \equiv \pm 1 \pmod{12}$.

Example 15.0.2. Characterize the primes p for which both 2 and 3 are quadratic residues modulo p . We want p such that $\left(\frac{2}{p}\right) = \left(\frac{3}{p}\right) = 1$. We already know that $\left(\frac{2}{p}\right) = 1$ if and only if $p \equiv \pm 1 \pmod{8}$ and $\left(\frac{3}{p}\right) = 1$ if and only if $p \equiv \pm 1 \pmod{12}$. So by the Chinese remainder theorem, we have $\left(\frac{2}{p}\right) = \left(\frac{3}{p}\right) = 1$ if and only if $p \equiv \pm 1 \pmod{24}$.

Example 15.0.3. Characterize the primes p for which 13 is a quadratic residue modulo p . By quadratic reciprocity, we have $\left(\frac{13}{p}\right) = \left(\frac{p}{13}\right)$. The non-zero quadratic residues modulo 13 are

$$\{1, 3, 4, 9, 10, 12\},$$

So $\left(\frac{13}{p}\right) = \left(\frac{p}{13}\right) = 1$ if and only if $p \equiv 1, 3, 4, 9, 10, 12 \equiv \pm 1, \pm 3, \pm 4 \pmod{13}$.

Example 15.0.4. Characterize the primes p for which 11 is a quadratic residue modulo p . By quadratic reciprocity, we have that

$$\left(\frac{11}{p}\right) = \begin{cases} \left(\frac{p}{11}\right) & p \equiv 1 \pmod{4}, \\ -\left(\frac{p}{11}\right) & p \equiv 3 \pmod{4}. \end{cases}$$

The quadratic residues modulo 11 are 1, 3, 4, 5, 9, and the quadratic non-residues are 2, 6, 7, 8, 10. If $p \equiv 1 \pmod{4}$, then $\left(\frac{p}{11}\right) = 1$ if and only if $p \equiv 1, 3, 4, 5, 9 \pmod{11}$, and if $p \equiv 3 \pmod{4}$, then $\left(\frac{p}{11}\right) = -1$ if and only if $p \equiv 2, 6, 7, 8, 10 \pmod{11}$. One can check each of these cases by the Chinese remainder theorem, and one gets $\left(\frac{11}{p}\right) = 1$ if and only if $p \equiv 1, 5, 7, 9, 19, 25, 35, 37, 39, 43 \pmod{44}$, or

$$p \equiv \pm 1, \pm 5, \pm 7, \pm 9, \pm 19 \pmod{44}.$$

Example 15.0.5. Characterize the primes p for which -1 and 2 are both quadratic residues modulo p . Recall that $\left(\frac{-1}{p}\right) = 1$ if and only if $p \equiv 1 \pmod{4}$ and $\left(\frac{2}{p}\right) = 1$ if and only if $p \equiv \pm 1 \pmod{8}$, so we see that $\left(\frac{-1}{p}\right) = \left(\frac{2}{p}\right) = 1$ if and only if $p \equiv 1 \pmod{8}$.

Example 15.0.6. Characterize the primes p for which both -1 and 3 are quadratic residues modulo p . Recall that $\left(\frac{-1}{p}\right) = 1$ if and only if $p \equiv 1 \pmod{4}$ and $\left(\frac{3}{p}\right) = 1$ if and only if $p \equiv \pm 1 \pmod{12}$. So we have $\left(\frac{-1}{p}\right) = \left(\frac{3}{p}\right) = 1$ if and only if $p \equiv 1 \pmod{12}$.

Exercise 15.1. Characterize the primes p for which 3 and 5 are quadratic residues modulo p .

Lecture 16

Oct. 20 — Primitive Roots

What happens when you step on a grape? Nothing, it lets out a little whine.

16.1 Orders

Remark. Let m be a positive integer and $(a, m) = 1$. By Euler's theorem, we know that

$$a^{\varphi(m)} \equiv 1 \pmod{m}.$$

However, it may happen that $a^g \equiv 1 \pmod{m}$ for some smaller g .

Definition 16.1. Let $a, m \in \mathbb{Z}$ with $m > 0$, $(a, m) = 1$. Then the *order of a modulo m* , denoted $\text{ord}_m a$, is the least positive integer n such that $a^n \equiv 1 \pmod{m}$.

Example 16.1.1. We compute $\text{ord}_7 2$. We can compute that

$$\begin{aligned} 2^1 &\equiv 2 \pmod{7}, \\ 2^2 &\equiv 4 \pmod{7}, \\ 2^3 &\equiv 1 \pmod{7}, \end{aligned}$$

so we see that $\text{ord}_7 2 = 3$. Note that Euler's theorem only guarantees $\text{ord}_7 2 \leq \phi(7) = 6$.

Example 16.1.2. We compute $\text{ord}_7 3$. We can compute that

$$\begin{aligned} 3^1 &\equiv 3 \pmod{7}, \\ 3^2 &\equiv 2 \pmod{7}, \\ 3^3 &\equiv 6 \pmod{7}, \\ 3^4 &\equiv 4 \pmod{7}, \\ 3^5 &\equiv 5 \pmod{7}, \\ 3^6 &\equiv 1 \pmod{7}, \end{aligned}$$

so we see that $\text{ord}_7 3 = 6$.

Proposition 16.1. Let $a, m \in \mathbb{Z}$ with $m > 0$ and $(a, m) = 1$. Then $a^n \equiv 1 \pmod{m}$ for some positive integer n if and only if $\text{ord}_m a \mid n$. In particular, $\text{ord}_m a \mid \varphi(m)$.

Proof. (\Rightarrow) Suppose that $a^n \equiv 1 \pmod{m}$. By the division algorithm, there exists $q, r \in \mathbb{Z}$ such that

$$n = q(\text{ord}_m a) + r, \quad 0 \leq r < \text{ord}_m a.$$

Then $1 = a^n \equiv a^{q(\text{ord}_m a)+r} \equiv (a^{\text{ord}_m a})^q a^r \equiv a^r \pmod{m}$, which can only happen if $r = 0$ by the definition of $\text{ord}_m a$ and $0 \leq r < \text{ord}_m a$. Therefore, $\text{ord}_m a \mid n$.

(\Leftarrow) Suppose that $\text{ord}_m a \mid n$. Then $n = q(\text{ord}_m a)$, so $a^n = a^{q(\text{ord}_m a)} \equiv (a^{\text{ord}_m a})^q \equiv 1 \pmod{m}$. \square

Example 16.1.3. We compute $\text{ord}_{13} 2$. By Proposition 16.1, it suffices to check divisors of $\varphi(13) = 12$:

$$\begin{aligned} 2^1 &\equiv 2 \pmod{13}, \\ 2^2 &\equiv 4 \pmod{13}, \\ 2^3 &\equiv 8 \pmod{13}, \\ 2^4 &\equiv 3 \pmod{13}, \\ 2^6 &\equiv 12 \pmod{13}, \\ 2^{12} &\equiv 1 \pmod{13}, \end{aligned}$$

thus $\text{ord}_{13} 2 = 12$. Note that we did not need to compute $2^7, 2^8$, etc. to verify this.

Proposition 16.2. Let $a, m \in \mathbb{Z}$ with $m > 0$ and $(a, m) = 1$. If i, j are non-negative integers, then $a^i \equiv a^j \pmod{m}$ if and only if $i \equiv j \pmod{\text{ord}_m a}$.

Proof. Without loss of generality, suppose $i \geq j$.

(\Rightarrow) Assume $a^i \equiv a^j \pmod{m}$. Then we can write

$$a^j \equiv a^i \equiv a^j a^{i-j} \pmod{m}.$$

Since $(a, m) = 1$, we can cancel a^j to get $1 \equiv a^{i-j} \pmod{m}$. Then by Proposition 16.1, $\text{ord}_m a \mid i - j$.

(\Leftarrow) Assume $i \equiv j \pmod{\text{ord}_m a}$. Then $\text{ord}_m a \mid i - j$, so there exists $n \in \mathbb{Z}$ such that $i - j = n(\text{ord}_m a)$. Thus $i = j + n(\text{ord}_m a)$, and we have $a^i \equiv a^{j+n(\text{ord}_m a)} \equiv a^j (a^{\text{ord}_m a})^n \equiv a^j \pmod{m}$. \square

Example 16.1.4. We have seen previously that $\text{ord}_7 2 = 3$. So if i, j are non-negative integers, then $2^i \equiv 2^j \pmod{7}$ if and only if $i \equiv j \pmod{3}$ by Proposition 16.2. Note that

$$2000 \equiv 2 \pmod{3},$$

so we can calculate $2^{2000} \equiv 2^2 \equiv 4 \pmod{7}$.

16.2 Primitive Roots

Definition 16.2. Let $r, m \in \mathbb{Z}$ with $m > 0$ and $(r, m) = 1$. Then r is called a *primitive root modulo m* if $\text{ord}_m r = \varphi(m)$.

Remark. See *primitive root diffusers* for an interesting application (also *quadratic residue diffusers*).

Example 16.2.1. We have seen that 3 is a primitive root modulo 7 and 2 is a primitive root modulo 13. On the other hand, 2 is not a primitive root modulo 7.

Example 16.2.2. We prove that there are no primitive roots modulo 8. The reduced residues modulo 8 are 1, 3, 5, 7, and $\varphi(8) = 4$. But $1^2 \equiv 3^2 \equiv 5^2 \equiv 7^2 \equiv 1$, so none of these are primitive roots modulo 8.

In particular, not all integers m possess a primitive root. The *primitive root theorem* (later) tells us that m has a primitive root if and only if $m = 1, 2, 4, p^k, 2p^k$, where p is an odd prime.

Proposition 16.3. *Let r be a primitive root modulo m . Then $\{r, r^2, r^3, \dots, r^{\varphi(m)}\}$ is a complete set of reduced residues modulo m .*

Proof. Since r is a primitive root modulo m , we have $(r, m) = 1$, and so $(r^n, m) = 1$ for any $n \geq 1$. Also, there are $\varphi(m)$ elements in the list, so it remains to show that they are distinct modulo m .

To do this, suppose that $r^i \equiv r^j \pmod{m}$ for some $1 \leq i, j \leq \varphi(m)$. Then Proposition 16.2 implies that $i \equiv j \pmod{\varphi(m)}$, so $i = j$. Thus the r^i are distinct modulo m . \square

Remark. Proposition 16.3 says that a primitive root (when it exists) generates the reduced residues modulo m .

Example 16.2.3. Recall that 3 is a primitive root modulo 7. We saw in Example 16.1.2 that

$$\{3^1, 3^2, 3^3, 3^4, 3^5, 3^6\} \equiv \{3, 2, 6, 4, 5, 1\} \pmod{7},$$

in particular this is a complete set of reduced residues modulo 7.

Example 16.2.4. Recall that 2 is a primitive root modulo 13. We can compute

$$\{2^1, 2^2, 2^3, 2^4, 2^5, 2^6, 2^7, 2^8, 2^9, 2^{10}, 2^{11}, 2^{12}\} \equiv \{2, 4, 8, 3, 6, 12, 11, 9, 5, 10, 7, 1\} \pmod{13},$$

which is a complete set of reduced residues modulo 13.

Remark. If a primitive root exists, it is in general not unique. We will determine how many there are next lecture (we will see that there are $\varphi(\varphi(m))$ of them).

Exercise 16.1. Show there are no primitive roots modulo 12.

To do this, write $\mathbb{Z}/12\mathbb{Z} \cong \mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$, then we have

$$(\mathbb{Z}/12\mathbb{Z})^\times \cong (\mathbb{Z}/3\mathbb{Z})^\times \times (\mathbb{Z}/4\mathbb{Z})^\times = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$$

which is not cyclic. Alternatively, one can just compute directly for $(\mathbb{Z}/12\mathbb{Z})^\times = \{1, 5, 7, 11\}$ that

$$1^2 \equiv 5^2 \equiv 7^2 \equiv 11^2 \equiv 1 \pmod{12},$$

so none of these can be primitive roots modulo 12.