

# MATH 4318 Final Formula Sheet

## Differentiation

**Theorem 1** (Quotient rule). If  $f, g : U \rightarrow \mathbb{R}$  are differentiable and  $g(x_0) \neq 0$ , then

$$(f/g)'(x_0) = \frac{f'(x_0)g(x_0) - f(x_0)g'(x_0)}{g(x_0)^2}.$$

**Theorem 2** (Cauchy's mean value theorem). Let  $f, g \in C([a, b])$  be differentiable in  $(a, b)$ . If  $g'(x) \neq 0$  for any  $x \in (a, b)$ , then there exists  $x_0 \in (a, b)$  such that

$$\frac{f'(x_0)}{g'(x_0)} = \frac{f(b) - f(a)}{g(b) - g(a)}.$$

**Theorem 3.** Let  $R \in (0, \infty)$  and  $f \in C^\infty(x_0 - R, x_0 + R)$ . If there exists  $M > 0$  such that for all  $x \in (x_0 - R, x_0 + R)$ ,  $|f^{(n)}(x)| \leq M$  for all  $n \in \mathbb{N}$ , then

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$$

for all  $x \in (x_0 - R, x_0 + R)$ .

**Theorem 4** (Lagrange remainder). Let  $f \in C^n([a, b])$  and assume that  $f$  is  $(n+1)$ -times differentiable in  $(a, b)$ . Then

$$f(x) - \sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (x-a)^k = \frac{f^{(n+1)}(\xi)}{(n+1)!} (x-a)^{n+1}$$

for some  $\xi \in [a, x]$ .

## Integration

**Definition 1** (Riemann integrability). A function  $f : [a, b] \rightarrow \mathbb{R}$  is Riemann integrable on  $[a, b]$  if there exists  $A \in \mathbb{R}$  such that for all  $\epsilon > 0$ , there exists  $\delta > 0$  such that  $|S - A| < \epsilon$  whenever  $S$  is any Riemann sum on a partition of width  $< \delta$ . We call  $A$  the Riemann integral of  $f$  on  $[a, b]$ .

**Theorem 5** (Cauchy criterion for integrability). A function  $f : [a, b] \rightarrow \mathbb{R}$  is integrable if and only if for any  $\epsilon > 0$ , there exists  $\delta > 0$  such that  $|S_1 - S_2| < \epsilon$  whenever  $S_1$  and  $S_2$  are Riemann sums for partitions of width  $< \delta$ .

**Definition 2** (Step function). We say  $f : [a, b] \rightarrow \mathbb{R}$  is a step function if there exists a partition  $a = x_0 < x_1 < \dots < x_n = b$  such that  $f$  is constant on each interval  $(x_{i-1}, x_i)$ .

**Theorem 6.** A function  $f : [a, b] \rightarrow \mathbb{R}$  is integrable if and only if for any  $\epsilon > 0$ , there exist step functions  $f_1, f_2$  such that  $f_1(x) \leq f(x) \leq f_2(x)$  for all  $x \in [a, b]$  and

$$\int_a^b (f_2 - f_1) dx < \epsilon.$$

**Definition 3** (Oscillation amplitude). Let  $f : [a, b] \rightarrow \mathbb{R}$  be bounded and  $a = x_0 < x_1 < \dots < x_n = b$  be a partition. Then

$$\omega_i(f) = \sup\{|f(x) - f(y)| : x, y \in [x_{i-1}, x_i]\}$$

is the oscillation amplitude of  $f$  on  $[x_{i-1}, x_i]$ .

**Theorem 7.** A function  $f : [a, b] \rightarrow \mathbb{R}$  is integrable if and only if for any  $\epsilon > 0$ , there exists a partition such that

$$\sum_{n=1}^n \omega_i(f)(x_i - x_{i-1}) < \epsilon.$$

**Theorem 8** (Du Bois-Reymond). Let  $f$  be bounded on  $[a, b]$ . Then  $f \in \mathcal{R}([a, b])$  if and only if for any  $\epsilon, a > 0$ , there exists a partition such that the total length of subintervals with  $\omega_i(f) \geq \epsilon$  is  $< a$ .

## Exchange of Limit Operations

**Theorem 9.** Let  $\{f_n\}$  be a sequence of functions on an open interval  $U \subseteq \mathbb{R}$  such that each  $f_n$  has a continuous derivative. Suppose  $\{f'_n\}$  converges uniformly on  $U$  and for some  $a \in U$ ,  $\{f'_n(a)\}$  converges. Then  $\lim_{n \rightarrow \infty} f_n = f$  exists and  $f$  is differentiable. Furthermore, we have  $f' = \lim_{n \rightarrow \infty} f'_n$ .

**Theorem 10.** Let  $f_n : [a, b] \rightarrow \mathbb{R}$  be continuously differentiable. Suppose that  $\sum_{n=1}^{\infty} f_n(x)$  converges pointwise on  $[a, b]$  and  $\sum_{n=1}^{\infty} f'_n(x)$  converges uniformly on  $[a, b]$ . Then  $\sum_{n=1}^{\infty} f_n(x)$  converges uniformly on  $[a, b]$  and

$$\frac{d}{dx} \sum_{n=1}^{\infty} f_n(x) = \sum_{n=1}^{\infty} f'_n(x).$$

## Infinite Series

**Theorem 11.** If  $a_n \geq 0$ , then  $\sum_{n=1}^{\infty} a_n$  either converges or diverges to  $\infty$ .

**Theorem 12** (Alternating series test). Let  $\{a_n\}$  be a decreasing sequence with  $a_n \rightarrow 0$  as  $n \rightarrow \infty$ . Then the series  $\sum_{n=1}^{\infty} (-1)^{n+1} a_n$  converges (to  $S$ , say), and the partial sums  $S_n$  have error  $|S_n - S| \leq a_{n+1}$ .

**Theorem 13.** Let  $f_n \in C([a, b])$ . If  $\sum_{n=1}^{\infty} f_n(x)$  converges uniformly on  $(a, b)$ , then it converges uniformly on  $[a, b]$ .

## Power Series

**Theorem 14.** We have the following:

1. If  $\sum_{n=0}^{\infty} a_n x^n$  converges at  $x = x_1 \neq 0$ , then it converges absolutely for all  $x$  with  $|x| < |x_1|$ .
2. If  $\sum_{n=0}^{\infty} a_n x^n$  diverges at  $x = x_2 \neq 0$ , then it diverges for all  $x$  with  $|x| > |x_2|$ .

**Theorem 15** (Hadamard's formula). For a power series  $\sum_{n=0}^{\infty} a_n x^n$ , let  $L = \limsup_{n \rightarrow \infty} |a_n|^{1/n}$ . Then its radius of convergence is  $R = 1/L$ .

**Theorem 16.** For a series  $\sum_{n=0}^{\infty} a_n x^n$  with  $a_n \neq 0$ , if  $\lim_{n \rightarrow \infty} |a_{n+1}/a_n| = L$ , then its radius of convergence is  $R = 1/L$ .

**Theorem 17.** If  $\sum_{n=0}^{\infty} a_n x^n$  has radius of convergence  $R > 0$ , then for any  $0 < r < R$ , the power series  $\sum_{n=0}^{\infty} a_n r^n$  converges uniformly on  $[-r, r]$ . Moreover, if  $\sum_{n=0}^{\infty} a_n x^n$  converges at  $x = R < \infty$  (or  $x = -R$ ), then  $\sum_{n=0}^{\infty} a_n x^n$  converges uniformly on  $[0, R]$  (or  $[-R, 0]$ ).

**Theorem 18.** If  $\sum_{n=0}^{\infty} a_n x^n$  has radius of convergence  $R > 0$ , then  $f(x) = \sum_{n=0}^{\infty} a_n x^n \in C^\infty(-R, R)$ .

**Theorem 19.** Suppose  $f(x) = \sum_{n=0}^{\infty} a_n x^n$  has radius of convergence  $R > 0$ . Then for any  $x \in (-R, R)$ ,  $f \in \mathcal{R}([0, x])$  and

$$\int_0^x f(t) dt = \sum_{n=0}^{\infty} \frac{a_n}{n+1} x^{n+1}.$$

## Differentiation in $\mathbb{R}^n$

**Theorem 20.** Let  $f : U \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$  where  $U$  is open. Then  $f$  is differentiable at  $x = a$  if and only if there exist functions  $A_1, \dots, A_n$  on  $U$ , continuous at  $x = a$ , such that

$$f(x) - f(a) = A_1(x)(x_1 - a_1) + \dots + A_n(x)(x_n - a_n)$$

for all  $x \in U$ . In this case,  $\partial f / \partial x_i(a) = A_i(a)$ .

**Theorem 21.** Let  $U$  be an open set in  $\mathbb{R}^n$  and suppose that  $f : U \rightarrow \mathbb{R}$  has partial derivatives  $f'_1, \dots, f'_n$  on  $U$  which are continuous on  $x = a$ . Then  $f$  is differentiable at  $x = a$ .

**Theorem 22** (Implicit function theorem). Let  $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$  and  $U \times V \subseteq \mathbb{R}^n \times \mathbb{R}^m$  be a neighborhood of  $(x_0, y_0)$ . Suppose  $f$  and  $\partial f / \partial y$  are continuous on  $U \times V$ , and  $f(x_0, y_0) = 0$  and

$$\det \left( \frac{\partial f}{\partial y}(x_0, y_0) \right) \neq 0.$$

Then there exists a neighborhood  $U_0 \times V_0 \subseteq U \times V$  of  $(x_0, y_0)$  and a unique continuous function  $\varphi : U_0 \rightarrow V_0$  satisfying

$$\begin{cases} f(x, \varphi(x)) = 0, \\ \varphi(x_0) = y_0. \end{cases}$$

**Theorem 23.** Let  $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$  be continuous and suppose that  $\partial f / \partial y$  exists and is continuous on  $[a, b] \times [c, d]$ . Then

$$\frac{d}{dy} \int_a^b f(x, y) dx = \int_a^b \frac{\partial f}{\partial y}(x, y) dx.$$

## Integration in $\mathbb{R}^n$

**Theorem 24** (Lebesgue's criterion for Riemann integrability). Let  $A \subseteq \mathbb{R}^n$  be a set with volume and let  $f : A \rightarrow \mathbb{R}$  be a bounded function that is continuous except on a subset of  $A$  with zero volume. Then  $f$  is integrable on  $A$ .

**Theorem 25.** Suppose  $f(x, y)$  is integrable on  $D = [a, b] \times [c, d]$  and for each  $x \in [a, b]$ ,  $f(x, y)$  is integrable on  $[c, d]$ . Then

$$\int_a^b dx \left[ \int_c^d f(x, y) dy \right] = \iint_D f(x, y) dx dy$$

## Reverse Triangle Inequality

**Proposition 1.** For all  $x, y \in \mathbb{R}$ , we have  $||x| - |y|| \leq |x - y|$ .