

# MATH 6441: Algebraic Topology I

Frank Qiang  
Instructor: John Etnyre

Georgia Institute of Technology  
Spring 2025

# Contents

<b>1</b>	<b>Jan. 6 — CW-Complexes</b>	<b>3</b>
1.1	Introduction and Motivation . . . . .	3
1.2	CW-Complexes . . . . .	4
1.3	Homotopy . . . . .	5
<b>2</b>	<b>Jan. 8 — Homotopy</b>	<b>6</b>
2.1	More on Homotopy . . . . .	6
2.2	Homotopy Groups . . . . .	6
2.3	Homotopy Equivalence . . . . .	7
<b>3</b>	<b>Jan. 13 — Homotopy, Part 2</b>	<b>9</b>
3.1	More on Homotopy Equivalence . . . . .	9
3.2	Homotopy Extension Property . . . . .	10
<b>4</b>	<b>Jan. 15 — Fundamental Group</b>	<b>13</b>
4.1	Fundamental Group . . . . .	13
4.2	Induced Homomorphisms . . . . .	15
<b>5</b>	<b>Jan. 22 — Simple Computations</b>	<b>16</b>
5.1	Fundamental Groups and Homotopy Equivalence . . . . .	16
5.2	Simple Computations of Fundamental Groups . . . . .	17
5.3	Fundamental Group of the Circle . . . . .	18
<b>6</b>	<b>Jan. 27 — Fundamental Group of the Circle</b>	<b>20</b>
6.1	Proof of Path Lifting . . . . .	20
6.2	Applications of the Fundamental Group of $S^1$ . . . . .	20
6.3	Free Products of Groups . . . . .	21
<b>7</b>	<b>Jan. 29 — Some Group Theory</b>	<b>23</b>
7.1	Group Presentations . . . . .	23
7.2	Seifert-van Kampen Theorem . . . . .	25
<b>8</b>	<b>Feb. 3 — Seifert-van Kampen Theorem</b>	<b>26</b>
8.1	Applications of the Seifert-van Kampen Theorem . . . . .	26
8.2	Proof of the Seifert-van Kampen Theorem . . . . .	27
8.3	Covering Spaces . . . . .	28
<b>9</b>	<b>Feb. 5 — Covering Spaces</b>	<b>29</b>
9.1	Examples of Covering Spaces . . . . .	29
9.2	Covering Spaces and Lifting . . . . .	29
9.3	Connections to the Fundamental Group . . . . .	30

<b>10 Feb. 10 — Covering Spaces, Part 2</b>	<b>32</b>
10.1 More Connections to the Fundamental Group . . . . .	32
10.2 Covering Space Isomorphisms . . . . .	34
<b>11 Feb. 12 — Covering Spaces, Part 3</b>	<b>36</b>
11.1 Galois Correspondence . . . . .	36
<b>12 Feb. 17 — Covering Spaces, Part 4</b>	<b>40</b>
12.1 Deck Transformations . . . . .	40
<b>13 Feb. 19 — Singular Homology</b>	<b>42</b>
13.1 Covering Spaces and Group Actions . . . . .	42
13.2 Introduction to Homology . . . . .	42
13.3 Singular Homology . . . . .	43

# Lecture 1

## Jan. 6 — CW-Complexes

### 1.1 Introduction and Motivation

Algebraic topology builds “functions” (actually *functors*)

$$\{\text{topological spaces, continuous maps}\} \longrightarrow \{\text{algebraic things, algebraic maps}\},$$

where “algebraic things” can be groups, vector spaces, etc. The main objective of algebraic topology is to *distinguish topological spaces*, e.g. showing that  $\mathbb{R}^n \not\cong \mathbb{R}^m$  for  $n \neq m$ . More applications are:

1. Studying maps between spaces.<sup>1</sup>

- Does a given space  $M$  embed in  $N$ ? For instance, for what  $m$  does  $\mathbb{R}P^n$  embed in  $\mathbb{R}^m$ ? (This is still not known in general.) Here  $\mathbb{R}P^n$  is the real projective space.
- Lifting maps, i.e. given  $f : A \rightarrow B$  and  $g : E \rightarrow B$ , does there exist a map  $\tilde{f} : A \rightarrow E$  such that  $g \circ \tilde{f} = f$ ? In other words, is there a map  $\tilde{f}$  such that the following diagram commutes:

$$\begin{array}{ccc} & & E \\ & \nearrow \tilde{f} & \downarrow g \\ A & \xrightarrow{f} & B \end{array}$$

- Fixed point problems: Given  $f : X \rightarrow X$ , does  $f$  have a fixed point, i.e.  $x \in X$  such that  $f(x) = x$ ? Such theorems are used to prove the existence of solutions to ordinary differential equations, for instance.

2. Group actions, e.g. which finite groups act freely on  $S^n$ ?
3. Group theory, e.g. showing that every subgroup of a free group is free. Another example is that if  $F_n$  is the free group on  $n$  generators, then its *commutator*  $[F_n, F_n]$  is not finitely generated.
4. Algebra, e.g. proving the fundamental theorem of algebra.

This course will cover the following topics:

1. The *fundamental group*  $\pi_1(X, x_0)$  of a space  $X$  for  $x_0 \in X$ , and *covering spaces*.
2. The *homology groups*  $H_k(X)$  for  $k = 0, 1, 2, \dots$ . These groups are abelian.
3. The *cohomology ring*  $H^*(X) = \bigoplus_{k=0}^{\infty} H^k(X)$ .

But before getting to this, we need to develop some important ideas.

---

<sup>1</sup>All maps and functions in this class are continuous unless otherwise specified.

## 1.2 CW-Complexes

**Definition 1.1.** Let  $D^n \subseteq \mathbb{R}^n$  be the unit disk and  $S^{n-1} = \partial D^n$ . Given a topological space  $Y$  and a continuous map  $a : S^{n-1} \rightarrow Y$ , the space obtained from  $Y$  by *attaching* an  $n$ -cell (via  $a$ ) is

$$Y \cup_a D^n = (Y \sqcup D^n) / \sim,$$

where the equivalence relation  $\sim$  is given by  $x \sim a(x)$  for  $x \in \partial D^n$ . Here  $\sqcup$  denotes disjoint union.

**Definition 1.2.** An  $n$ -complex or an  $n$ -dimensional CW-complex is defined inductively by:

- A  $(-1)$ -complex is the empty set  $\emptyset$ .
- An  $n$ -complex  $X^n$  is a space obtained from an  $(n-1)$ -complex by attaching  $n$ -cells.

An  $n$ -complex is *finite* if it involves only a finite number of cells. The  $k$ -skeleton of  $X$  is the union of all  $n$ -cells in  $X$  with  $n \leq k$ .

**Remark.** Any CW-complex is Hausdorff. See Hatcher for a proof.

**Example 1.2.1.** Here are some examples of CW-complexes:

- A 0-complex is a union of points. This is because  $D^0 = \{\text{pt}\}$  and  $\partial D^0 = \emptyset$ .
- A 1-complex is a graph (points and lines connecting them).
- The torus  $T$  (a square with opposite sides identified) is a 2-complex. Here the 0-skeleton  $T^{(0)}$  is the common corner on the square and the 1-skeleton  $T^{(1)}$  is two sides of the square after taking the quotient. The 2-skeleton  $T = T^{(2)}$  is the entire torus.
- Another example of a 2-complex is the two-holed torus, which is obtained by identifying the edges of an octagon (pairs of every other edge identified with opposite orientation).<sup>2</sup>
- A third example of a 2-complex is  $X^{(1)} \cup_a D^2$  given an attaching map  $D^2 \rightarrow X^{(1)}$ .
- Consider the unit sphere  $S^n \subseteq \mathbb{R}^{n+1}$ . One way to give  $S^2$  a CW-complex structure is to see the sphere as two disks  $D^2$  glued together, resulting in one 0-cell, one 1-cell, and two 2-cells. Another way is to start with two points, attach two 1-cells to get a circle, and then attaching two disks to get  $S^2$ . This results in two 0-cells, two 1-cells, and two 2-cells.

The second idea generalizes to  $S^n$ . We can write

$$S^n = S^{n-1} \cup_{a_1} D^n \cup_{a_2} D^n,$$

where  $S^{n-1}$  inductively has a CW-complex structure. This yields two  $k$ -cells for each  $k \leq n$ .

Another way to put a CW-complex structure on  $S^n$  is to attach  $D^n$  to a point with  $\partial D^n \rightarrow \{\text{pt}\}$ . In particular, notice that a space can in general have several different CW-complex structures.

- Consider the  $n$ -dimensional real projective space

$$\mathbb{R}P^n = \{\text{lines through the origin in } \mathbb{R}^{n+1}\}.$$

Since each line through the origin passes through  $S^n$  twice, we can equivalently think of  $\mathbb{R}P^n$  as the unit sphere  $S^n$  with antipodal points identified.

---

<sup>2</sup>This CW-decomposition of the two-holed torus results in one 0-cell, four 1-cells, and one 2-cell.

We can also think of this as  $D^n$  with antipodal points on  $\partial D^n$  identified. Since  $\partial D^n = S^{n-1}$ , this is simply  $\mathbb{R}P^{n-1} \cup_a D^n$ , where  $a : \partial D^n \rightarrow \mathbb{R}P^{n-1}$  is the quotient map. This gives  $\mathbb{R}P^n$  a CW-complex structure with one  $k$ -cell for each  $k \leq n$ .

- The complex projective space  $\mathbb{C}P^n$  has a similar CW-complex structure with one  $k$ -cell for each even  $k \leq 2n$ . One can verify this as an exercise.
- Any smooth manifold has a CW-complex structure. See Hatcher.

**Exercise 1.1.** Show the product of CW-complexes is a CW-complex.

**Definition 1.3.** A *subcomplex* of a CW-complex  $X$  is a closed subset  $A \subseteq X$  that is a union of cells in  $X$ . In particular,  $A$  is also a CW-complex and  $(X, A)$  is called a *CW-pair*.

## 1.3 Homotopy

**Definition 1.4.** Let  $X$  and  $Y$  be topological spaces. Two maps  $f, g : X \rightarrow Y$  are *homotopic*, denoted  $f \sim g$ , if there exists a continuous map  $\Phi : X \times [0, 1] \rightarrow Y$  such that

$$\Phi(x, 0) = f(x) \quad \text{and} \quad \Phi(x, 1) = g(x)$$

for all  $x \in X$ . In this case,  $\Phi$  is called a *homotopy* between  $f$  and  $g$ .

**Remark.** We note the following:

- A homotopy  $\Phi$  gives a family of maps  $\phi_{t_0} : X \rightarrow Y$  given by  $x \mapsto \phi(x, t_0)$  which is continuous in  $t_0$ . So maps are homotopic if there is a continuous family of maps between them.
- If  $A \subseteq X$  then we say that  $f$  is *homotopic to  $g$  rel  $A$* , denoted  $f \sim_A g$ , if there exists  $\Phi$  as above with the additional property that  $\Phi(x, t) = f(x)$  for all  $x \in A$ , i.e. points in  $A$  are fixed.
- If  $A \subseteq X$  and  $B \subseteq Y$ , then the notation  $f : (X, A) \rightarrow (Y, B)$  means that  $f : X \rightarrow Y$  and  $f(a) \in B$  for each  $a \in A$ . We say that  $f$  is a *map of pairs*. If  $f, g : (X, A) \rightarrow (Y, B)$ , then  $f, g$  are *homotopic as maps of pairs* if each  $\phi_t : (X, A) \rightarrow (Y, B)$  is a map of pairs.

# Lecture 2

## Jan. 8 — Homotopy

### 2.1 More on Homotopy

**Example 2.0.1.** For any space  $X$ , any map  $f : X \rightarrow [0, 1]$  is homotopic to the map  $g : X \rightarrow [0, 1]$  given by  $x \mapsto 0$ . To see this, we have the homotopy  $\Phi : X \times [0, 1] \times [0, 1]$  defined by

$$(x, t) \mapsto (1 - t)f(x).$$

We can see that  $\Phi(x, 0) = f(x)$  and  $\Phi(x, 1) = 0 = g(x)$ .

**Exercise 2.1.** Show that homotopy is an equivalence relation on maps  $X \rightarrow Y$ .

**Definition 2.1.** Let  $C(X, Y) = \{\text{continuous maps from } X \text{ to } Y\}$ . Let  $[X, Y] = C(X, Y)/\sim$ , i.e. homotopic maps are identified with each other.

**Example 2.1.1.** We have the following:

1.  $[X, [0, 1]] = \{g\}$  for any space  $X$ , where  $g$  is the map  $x \mapsto 0$  as above.
2.  $[\{*\}, X] = \{\text{path components of } X\}$ .

### 2.2 Homotopy Groups

**Definition 2.2.** We call a space  $X$  *pointed* if there is a designated “base point”  $x_0 \in X$ . Given two pointed spaces  $(X, x_0)$  and  $(Y, y_0)$ , we define

$$[X, Y]_0 = \{\text{homotopy classes of maps of pairs } (X, \{x_0\}) \rightarrow (Y, \{y_0\})\}.$$

**Definition 2.3.** Let  $y_0$  be the north pole in  $S^n$ , i.e.  $S^n \subseteq \mathbb{R}^{n+1}$  is the unit sphere and  $y_0 = (0, \dots, 0, 1)$ . The  $n$ th homotopy group of a pointed space  $(X, x_0)$  is  $\pi_n(X, x_0) = [S^n, X]_0$ .

**Remark.** The homotopy group  $\pi_n(X, x_0)$  is in fact a group. We will study  $\pi_1(X, x_0)$  next and it is called the *fundamental group* of  $(X, x_0)$ .

**Remark.** For which  $(Y, y_0)$  is  $[Y, X]_0$  “naturally” a group for all  $(X, x_0)$ ? Similarly, for which  $(Y, y_0)$  is  $[X, Y]_0$  a group for all  $(X, x_0)$ ? Here, given a map  $f : (X_1, x_1) \rightarrow (X_2, x_2)$ , there is an obvious *induced map*  $f_* : [Y, X_1]_0 \rightarrow [Y, X_2]_0$  given by  $[g] \mapsto [f \circ g]$ . Similarly, there is a map  $f^* : [X_2, Y]_0 \rightarrow [X_1, Y]_0$  given by  $[g] \mapsto [g \circ f]$ . In the questions above, “naturally” means that  $f_*$  and  $f^*$  are homomorphisms for any  $(X_1, x_1)$  and  $(X_2, x_2)$ . The (perhaps unsatisfying) answer is that a space satisfying the first condition is called an *H-space*, and a space satisfying the second is called an *H'-space*.

## 2.3 Homotopy Equivalence

**Definition 2.4.** We say that  $f : X \rightarrow Y$  is the *homotopy inverse* to a function  $g : Y \rightarrow X$  if  $f \circ g \sim \text{id}_Y$  and  $g \circ f \sim \text{id}_X$ , where  $\text{id}_X$  and  $\text{id}_Y$  are the identity maps on  $X$  and  $Y$ . If  $g$  has a homotopy inverse, then we call  $g$  a *homotopy equivalence* from  $Y$  to  $X$  and we call  $X, Y$  *homotopy equivalent*.<sup>1</sup>

**Exercise 2.2.** Show that homotopy equivalence is an equivalence relation.

**Lemma 2.1.** *The following are equivalent:*

1.  $X$  and  $Y$  are homotopy equivalent.
2. For any space  $Z$ , there is a one-to-one correspondence  $\phi_Z : [X, Z] \rightarrow [Y, Z]$  such that for all continuous maps  $h : Z \rightarrow Z'$ , the following diagram commutes:

$$\begin{array}{ccc} [X, Z] & \xrightarrow{\phi_Z} & [Y, Z] \\ \downarrow h_* & & \downarrow h_* \\ [X, Z'] & \xrightarrow{\phi_{Z'}} & [Y, Z'] \end{array}$$

3. For any space  $Z$ , there is a one-to-one correspondence  $\phi^Z : [Z, X] \rightarrow [Z, Y]$  such for that all continuous maps  $h : Z \rightarrow Z'$ , the following diagram commutes:

$$\begin{array}{ccc} [Z', X] & \xrightarrow{\phi^{Z'}} & [Z', Y] \\ \downarrow h^* & & \downarrow h^* \\ [Z, X] & \xrightarrow{\phi^Z} & [Z, Y] \end{array}$$

*Proof.* This is left as an exercise. □

**Remark.** Based on the previous lemma, two spaces are homotopy equivalent if and only if homotopy classes of maps to and from the space are “naturally equivalent.”

**Example 2.4.1.** We have the following:

- Homeomorphic spaces are homotopy equivalent.
- Let  $X = S^1$  and  $Y = S^1 \times [0, 1]$ . We claim that  $X$  is homotopy equivalent to  $Y$ .

Define the maps  $f : S^1 \rightarrow S^1 \times [0, 1]$  by  $x \mapsto (x, 0)$  and  $g : S^1 \times [0, 1] \rightarrow S^1$  by  $(x, t) \mapsto x$ . Then we can see that  $g \circ f : S^1 \rightarrow S^1$  maps  $x \mapsto x$ , so  $g \circ f = \text{id}_{S^1}$ . On the other hand, the composition  $f \circ g : S^1 \times [0, 1] \rightarrow S^1 \times [0, 1]$  maps  $(x, t) \mapsto (x, 0)$ . Now  $f \circ g \sim \text{id}_{S^1 \times [0, 1]}$  by homotopy. For instance, define  $\Phi : (S^1 \times [0, 1]) \times [0, 1] \rightarrow (S^1 \times [0, 1])$  by  $((x, t), s) \mapsto (x, st)$ , so

$$\Phi((x, t), 1) = (x, t) = \text{id}_{S^1 \times [0, 1]}(x, t) \quad \text{and} \quad \Phi((x, t), 0) = (x, 0) = f \circ g.$$

Thus  $f$  is a homotopy equivalence from  $S^1$  to  $S^1 \times [0, 1]$ . Note that  $S^1 \times [0, 1]$  is the annulus.

**Definition 2.5.** A space is called *contractible* if it is homotopy equivalent to a point.

**Example 2.5.1.** The spaces  $[0, 1]$  and  $\mathbb{R}^n$  are contractible (exercise).

---

<sup>1</sup>We will denote homotopy equivalence by  $X \simeq Y$  or simply  $X \sim Y$ .



**Definition 2.6.** If  $A \subseteq X$ , then a *retraction of  $X$  to  $A$*  is a map  $r : X \rightarrow A$  such that  $r(a) = a$  for every  $a \in A$ . A *deformation retraction* of  $X$  to  $A$  is a retraction  $r : X \rightarrow A$  that is homotopic rel  $A$  to the identity map  $\text{id}_X$ , i.e. we can find  $\phi_t : X \rightarrow X$  for  $t \in [0, 1]$  such that  $\phi_0(x) = x$  and  $\phi_1(X) \subseteq A$  and  $\phi_t(x) = x$  for all  $x \in A$  and  $t \in [0, 1]$ .

**Remark.** If  $X$  deformation retracts to  $A$ , then  $X$  is homotopy equivalent to  $A$ . To see this, suppose we have a homotopy  $\phi_t : X \rightarrow X$  as above, and let  $i : A \rightarrow X$  be the inclusion map. Then  $\phi_1 \circ i = \text{id}_A$  and  $i \circ \phi_1 = \phi_1 \sim \phi_0 = \text{id}_X$ , so  $\phi_1$  is a homotopy equivalence from  $X$  to  $A$ .

**Definition 2.7.** Given two spaces  $X, Y$  and a map  $f : X \rightarrow Y$ , the *mapping cylinder* of  $f$  is the space

$$M_f = ((X \times [0, 1]) \cup Y) / \sim,$$

where the equivalence relation  $\sim$  is given by  $(x, 1) \sim f(x)$  for  $x \in X$ .

**Remark.** The mapping cylinder  $M_f$  deformation retracts to  $Y$ . To see this, consider the map  $\tilde{\phi}_t$  given by  $(x, s) \mapsto (x, (1-t)s + t)$  on  $X \times [0, 1]$  and  $y \mapsto y$  on  $Y$ . Since  $\tilde{\phi}_t$  respects the equivalence relation, it descends to a map  $\phi_t : M_f \rightarrow M_f$  on the quotient space. Note that  $\phi_0 = \text{id}_{M_f}$  and  $\phi_1(M_f) = Y \subseteq M_f$ , and  $\phi_t = \text{id}_Y$  for all  $t$ . Thus  $\phi_1$  is a deformation retraction. In particular, this means that  $M_f \simeq Y$ .

**Remark.** There are obvious inclusions  $i : X \rightarrow M_f$  given by  $x \mapsto (x, 0)$  and  $j : Y \rightarrow M_f$  given by  $y \mapsto y$ . Note that  $\phi_1$  defined above is the homotopy inverse to  $j$ . Now we have the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow i & \downarrow j \\ & & M_f \end{array}$$

where  $j$  is a homotopy equivalence and  $j \circ f \sim i$  (exercise).

**Remark.** The above remark shows the following “slogan” of algebraic topology:

Any map is an inclusion up to homotopy.

**Example 2.7.1.** Let  $X$  be three circles with two enclosed in a third bigger one, and let  $Y$  be two circles enclosing the inner two circles of  $X$  connected by a line segment. Let  $Z$  be the region inside by the outer circle of  $X$  and outside the inner two circles of  $X$ .

Define  $f : X \rightarrow Y$  to be the map which sends  $x \in X$  to the point in  $Y$  at the other end of an interval (points on the inner circles of  $X$  are mapped by radial lines to the circles in  $Y$ , and points on the outer circle of  $X$  are mapped radially to either the circles or the line segment in  $Y$ ). One can write an explicit formula for  $f$  as an exercise.

Then  $Z$  is homeomorphic to  $M_f$ , and in particular  $Z \simeq Y$ . Similarly,  $M_f$  is homotopy equivalent to two circles joined at a point, or a circle with a diameter. Thus by transitivity, these two spaces and  $Z$  are all homotopy equivalent to each other.

# Lecture 3

## Jan. 13 — Homotopy, Part 2

### 3.1 More on Homotopy Equivalence

**Lemma 3.1.** *If  $(X, A)$  is a CW pair and  $A$  is contractible, then  $X/A \simeq X$ .<sup>1</sup>*

**Exercise 3.1.** The following are some applications of this lemma:

1. Let  $X$  be a connected graph (i.e. a 1-complex), and let  $A$  be an edge in  $X$  connecting distinct vertices. Then  $A$  is contractible, so  $X/A \simeq X$ . Continuing this process, let  $A$  be a maximal tree in  $X$ , which will also be contractible. Then  $X \simeq X/A$ , so any connected graph is homotopy equivalent to a *wedge of circles* (with number of circles equal to the number of self-loops in the graph).<sup>2</sup>
2. Consider the space  $X$  obtained by attaching a 1-cell  $A$  connecting the north and south poles on a sphere. Let  $B$  be half of a great circle connecting the endpoints of  $A$ . Clearly  $A$  and  $B$  are both contractible. After collapsing  $A$ , we see that  $X \simeq X/A$ , which is  $S^2$  with the north and south poles identified. On the other hand, by contracting  $B$  instead we see that  $X \simeq X/B$ , which is  $S^2 \vee S^1$ .

Note that after contracting  $A$ , the subset  $B$  is actually no longer contractible.

3. Let  $X$  be a torus with attached disks  $A_1, A_2, A_3$  in the tube of the torus. Then

$$X \simeq ((X/A_1)/A_2)/A_3,$$

which is three spheres lying in a circle, each attached to the next one at a single point.

We can also obtain this space by considering the space  $Y$  of three spheres attached in a line with an extra 1-cell  $B$  attached at the ends of the chain of circles. Also let  $A$  be the union of halves of great circles going through the chain of circles, with the same endpoints as  $B$ . Then we can see that this creates the same space as before, so that  $X \simeq Y/B \simeq Y$ . On the other hand, by contracting  $A$ , we see that  $Y \simeq Y/A = S^2 \vee S^2 \vee S^2 \vee S^1$ .

Of course, all of these spaces are then homotopy equivalent to each other by transitivity.

**Lemma 3.2.** *Let  $(X, A)$  be a CW pair and  $f, g : A \rightarrow Y$  be homotopic maps. Then  $X \cup_f Y \simeq X \cup_g Y$ .*

**Example 3.0.1.** Let  $Y = S^2$ , and  $X = D^2$ , and  $A = \partial D^2$ . Let  $g : A \rightarrow Y$  map  $A$  to a great circle, and let  $f : A \rightarrow Y$  map  $A$  to the north pole. One can show as an exercise that  $f \sim g$  (e.g. by pulling the equator towards the north pole). So the lemma says that  $X \cup_g Y$ , which is a sphere with a disk glued along its equator, is homotopy equivalent to  $X \cup_f Y = S^2 \vee S^2$ .

<sup>1</sup>Here  $X/A$  denotes the quotient of  $X$  obtained by collapsing all of  $A$  to a single point.

<sup>2</sup>A *wedge of pointed spaces*  $(X, x_0) \vee (Y, y_0)$  is the space obtained from  $X \sqcup Y$  by identifying  $x_0$  and  $y_0$ .

## 3.2 Homotopy Extension Property

**Remark.** To prove both of these lemmas, we need the *homotopy extension property* (HEP).

**Definition 3.1.** A space  $X$  and a subspace  $A \subseteq X$  have the *homotopy extension property* if given  $F_0 : X \rightarrow Y$  (for any  $Y$  and  $F_0$ ) and a homotopy  $f_t : A \rightarrow Y$  such that  $f_0 = F_0|_A$ , then there is a homotopy  $F_t : X \rightarrow Y$  such that  $F_t|_A = f_t$  for every  $t$ .

**Lemma 3.3.** A pair  $(X, A)$  has the homotopy extension property if and only if

$$(X \times \{0\}) \cup (A \times [0, 1])$$

is a retract of  $X \times [0, 1]$ .

*Proof.* ( $\Leftarrow$ ) We will assume that  $A$  is closed (not necessarily but makes the proof easier, and almost all examples satisfy this). By assumption, we have a retraction

$$r : (X \times [0, 1]) \rightarrow (X \times \{0\}) \cup (A \times [0, 1]).$$

Given  $F_0 : X \rightarrow Y$  and  $f_t : A \rightarrow Y$  such that  $f_0 = F_0|_A$ , we can define a map

$$\tilde{F} : (X \times \{0\}) \cup (A \times [0, 1]) \rightarrow Y$$

by  $x \mapsto F_0(x)$  on  $X \times \{0\}$  and  $(a, t) \mapsto f_t(a)$  on  $A \times [0, 1]$ . This map  $\tilde{F}$  is continuous since the definitions of  $\tilde{F}$  agree on the intersection and the intersection  $A \times \{0\}$  is closed. Now define

$$F : X \times [0, 1] \rightarrow Y$$

by  $F = \tilde{F} \circ r$ , which is a homotopy of  $F_0$  that extends  $f_t$ .

( $\Rightarrow$ ) Let  $Y = (X \times [0, 1]) \cup (A \times [0, 1])$ . Let

$$F_0 : X \rightarrow (X \times \{0\}) \cup (A \times [0, 1])$$

be given by  $x \mapsto (x, 0)$ , and

$$f_t : A \mapsto (X \times \{0\}) \cup (A \times [0, 1])$$

be given by  $a \mapsto (a, t)$ . Then the homotopy extension property yields an extension

$$F : X \times [0, 1] \rightarrow (X \times \{0\}) \cup (A \times [0, 1]),$$

which is a retraction, as desired. □

**Lemma 3.4.** If  $(X, A)$  is a CW pair, then  $(X \times \{0\}) \cup (A \times [0, 1])$  is a (deformation) retract of  $X \times [0, 1]$ . In particular,  $(X, A)$  satisfies the homotopy extension property.

*Proof.* The main idea is that for any disk  $D^n$ , the space  $(D^n \times \{0\}) \cup (\partial D^n \times [0, 1])$  is a deformation retract of  $D^n \times [0, 1]$ . To see this, let  $D^n \subseteq \mathbb{R}^n$  be the unit disk and  $D^n \times [0, 1] \subseteq \mathbb{R}^{n+1}$ . Let

$$p = (0, \dots, 0, 2).$$

For any  $x \in D^n \times [0, 1]$ , let  $\ell_x$  be the line through  $p$  and  $x$ . Note that

$$\ell_x \cap ((D^n \times [0, 1]) \cup (\partial D^n \times [0, 1]))$$

is a unique point. Define  $\tilde{r}(x)$  to be this point, which yields a map

$$\tilde{r} : D^n \times [0, 1] \rightarrow (D^n \times \{0\}) \cup (\partial D^n \times [0, 1])$$

Note that for  $x \in (D^n \times \{0\}) \cup (\partial D^n \times [0, 1])$ , then  $\tilde{r}(x) = x$  since the point of intersection is unique and  $x$  is already in the intersection. Show as an exercise that  $\tilde{r}$  is continuous. Then setting

$$\tilde{r}_t = t\tilde{r} + (1 - t)\text{id}_{D^n \times [0, 1]}$$

gives a deformation retraction of  $D^n \times [0, 1]$  onto  $(D^n \times \{0\}) \cup (\partial D^n \times [0, 1])$ .

To show the general case that  $X \times [0, 1]$  retracts to  $(X \times \{0\}) \cup (A \times [0, 1])$ , we induct on the dimension of cells. Define  $r$  on  $X^{(0)} \times [0, 1] \rightarrow (X \times \{0\}) \cup (A \times [0, 1])$  by the following:

- if a 0-cell  $D^0 \subseteq A$ , then let  $r$  be the identity on  $D^0 \times [0, 1]$ ;
- if a 0-cell  $D^0$  is not in  $A$ , then send  $D^0 \times [0, 1]$  to  $D^0 \subseteq X \times \{0\}$ .

Now inductively assume  $r$  is defined on  $X^{(k-1)} \times [0, 1] \rightarrow (X \times \{0\}) \cup (A \times [0, 1])$ . For each  $k$ -cell  $D^k$ ,

- if  $D^k \subseteq A$ , then let  $r$  be the identity on  $D^k \times [0, 1]$ ;
- if  $D^k$  is not in  $A$ , then note that  $\partial D^k \times [0, 1] \rightarrow X^{(k-1)} \times [0, 1]$  is defined by induction, and we have an “inclusion” (here  $a : \partial D^k \rightarrow X^{(k-1)}$  is the attaching map for  $D^k$ )

$$\begin{array}{ccc} D^k & \xrightarrow{i} & X^{(k-1)} \cup D^k \xrightarrow{q} (X^{(k-1)} \cup D^k) / \{(x \in \partial D^k) \sim (a(x) \in X^{(k-1)})\} \\ & \searrow j & \nearrow \\ & & \end{array}$$

So we let  $D^k \times \{0\} \rightarrow (X \times \{0\}) \cup (A \times [0, 1])$  be the map  $j$  into  $X \times \{0\}$ . This defines  $r$  on

$$(D^k \times [0, 1]) \cup (\partial D^k \times [0, 1]),$$

which extends to  $D^k \times [0, 1]$  by composing with the map  $\tilde{r}$  from above.

This inductively defines the retraction  $r$  on all of  $X \times [0, 1]$ . The last claim follows by Lemma 3.3.  $\square$

**Remark.** Now we can prove the first two lemmas from the beginning of the day.

*Proof of Lemma 3.1.* We prove that the result holds for any  $(X, A)$  which satisfies the homotopy extension property. We show that the quotient map  $q : X \rightarrow X/A$  has a homotopy inverse. Since  $A$  is contractible, we know there exists a homotopy  $f_t : A \rightarrow A \subseteq X$ , such that  $f_0 = \text{id}_A$  and  $f_1$  is constant. Let  $F_0 : X \rightarrow X$  be the identity (note that  $F_0|_A = \text{id}_A$ ), so that the homotopy extension property gives a homotopy  $F_t : X \rightarrow X$  extending  $f_t$ . Since  $F_t(A) \subseteq A$  for every  $t$ , we get maps

$$\tilde{F}_t : X/A \rightarrow X/A,$$

which are well defined since points in  $A$  are mapped to points in  $A$ . Furthermore,  $F_1(A) = \{\text{pt}\}$ , so  $F_1$  factors through  $X/A$  to give a map  $h : X/A \rightarrow X$  satisfying  $F_1 = h \circ q$ . This gives the diagram:

$$\begin{array}{ccc} X & \xrightarrow{F_1} & X \\ q \downarrow & \nearrow h & \downarrow q \\ X/A & \xrightarrow{\tilde{F}_1} & X/A \end{array}$$

It is easy to check that  $h \circ q = F_1$  and  $q \circ h = \overline{F}_1$ , so that the diagram commutes. But then

$$h \circ q = F_1 \sim F_0 = \text{id}_X \quad \text{and} \quad q \circ h = \overline{F}_1 \sim \overline{F}_0 = \text{id}_{X/A},$$

so  $q$  is a homotopy equivalence. □

*Proof of Lemma 3.2.* Let  $F : A \times [0, 1] \rightarrow Y$  be a homotopy, which extends to  $F : X \times [0, 1] \rightarrow Y$  by the homotopy extension property. Consider the mapping cylinder

$$M_F = (X \times [0, 1]) \cup_F Y,$$

and one can show that  $M_F \simeq X \cup_g Y \simeq X \cup_f Y$ . □

# Lecture 4

## Jan. 15 — Fundamental Group

### 4.1 Fundamental Group

**Remark.** The basic idea of the *fundamental group* is to study the topology of a space with loops mapped into the space. For instance, intuitively, any loop in  $S^2$  can be “pulled back” to (i.e. is homotopic to) a constant loop. On the other hand, a loop wrapping around the hole in  $T^2$  gets stuck and cannot be “pulled back” to a constant loop. The same issue happens for a loop in  $T^2$  around the cylindrical part. The fundamental group is a way to make this intuition precise and measure the “holes” in a space.

**Definition 4.1.** The *fundamental group* of a based space  $(X, x_0)$  is

$$\pi_1(X, x_0) = [(S^1, n), (X, x_0)]_0,$$

i.e. the homotopy classes of loops in  $X$  based at  $x_0$ . Here  $n = (0, 1)$  is the north pole of  $S^1$ .

**Exercise 4.1.** Let  $S^1 \subseteq \mathbb{R}^2$  be the unit circle and let  $p : [0, 1] \rightarrow S^1$  be given by

$$t \mapsto (\cos 2\pi t, \sin 2\pi t).$$

Show that  $p$  is a quotient map, so we can think of  $S^1$  as  $[0, 1]$  with  $0, 1$  identified. Moreover, show that there is a one-to-one correspondence between maps of the form<sup>1</sup>

$$\gamma : ([0, 1], \{0, 1\}) \rightarrow (X, x_0) \quad \text{and} \quad \tilde{\gamma} : (S^1, \{(1, 0)\}) \rightarrow (X, x_0)$$

given by  $\tilde{\gamma} \mapsto \tilde{\gamma} \circ p = \gamma$ , and that homotopies of  $\tilde{\gamma}$  rel  $\{(1, 0)\}$  correspond to homotopies of  $S^1$  rel  $\{(1, 0)\}$ .

**Remark.** Using the above exercise, we can think of  $\pi_1(X, x_0) = [S^1, X]_0$  instead as

$$\pi_1(X, x_0) = [[0, 1], \{0, 1\}], (X, x_0)]_0.$$

Given a based loop  $\gamma : [0, 1] \rightarrow X$ , we denote its equivalence class in  $\pi_1(X, x_0)$  by  $[\gamma]$ .

**Definition 4.2.** If  $\gamma_1, \gamma_2$  are loops in  $X$  based at  $x_0 \in X$ , their *concatenation*  $\gamma_1 * \gamma_2 : [0, 1] \rightarrow X$  is

$$t \mapsto \begin{cases} \gamma_1(2t) & \text{if } 0 \leq t \leq 1/2, \\ \gamma_2(2t - 1) & \text{if } 1/2 \leq t \leq 1. \end{cases}$$

**Remark.** Concatenation of loops indeed yields another loop since  $\gamma_1 * \gamma_2(0) = \gamma_1 * \gamma_2(1) = x_0$  and  $\gamma_1 * \gamma_2$  is continuous since the definitions agree on the closed set  $\{1/2\}$ .

---

<sup>1</sup>Such a loop  $\gamma$  is called a *based loop*.

**Remark.** We can clearly see that  $\gamma_1 * \gamma_2$  is well-defined given  $\gamma_1$  and  $\gamma_2$ , but can we define  $[\gamma_1] * [\gamma_2]$  for homotopy classes of loops in a well-defined manner? We need to check that if  $\gamma_1 \sim \gamma_2$  and  $\delta_1 \sim \delta_2$  (i.e.  $\gamma_1, \gamma_2 \in [\gamma_1]$  and  $\delta_1, \delta_2 \in [\delta_1]$ ), then we also have  $\gamma_1 * \delta_1 \sim \gamma_2 * \delta_2$ .

To do this, let  $H : [0, 1] \times [0, 1] \rightarrow X$  be the homotopy from  $\gamma_1$  to  $\gamma_2$  and  $G : [0, 1] \times [0, 1] \rightarrow X$  be the homotopy from  $\delta_1$  to  $\delta_2$ . We need to construct a homotopy  $\tilde{H} : [0, 1] \times [0, 1] \rightarrow X$  from  $\gamma_1 * \delta_1$  to  $\gamma_2 * \delta_2$ .

Note that if we think of  $[0, 1] \times [0, 1]$  as the unit square, then  $\tilde{H}$  is already defined on the boundary: the left and right sides are constantly  $x_0$ , the top side is  $\gamma_2 * \delta_2$ , and the bottom side is  $\gamma_1 * \delta_1$ . So we only need to define it on the interior. For this, note that the vertical line in the middle of the square is also constantly  $x_0$  by construction: This creates two rectangles on each half, which we can fill with  $H$  and  $G$ .

More formally, we can construct the homotopy  $\tilde{H} : [0, 1] \times [0, 1] \rightarrow X$  explicitly via

$$(t, s) \mapsto \begin{cases} H(2t, s) & \text{if } 0 \leq t \leq 1/2, \\ G(2t - 1, s) & \text{if } 1/2 \leq t \leq 1. \end{cases}$$

This is continuous since the definitions agree on the closed set  $\{t = 1/2\}$ . Thus, setting

$$[\gamma_1] * [\delta_1] = [\gamma_1 * \delta_1]$$

gives a well-defined binary operation by the above discussion.

**Lemma 4.1.** *The pair  $(\pi_1(X, x_0), *)$  is a group.*

*Proof.* For the identity, let  $e : [0, 1] \rightarrow X$  be the constant loop  $t \mapsto x_0$ . We will show that

$$[e] * [\gamma] = [\gamma] = [\gamma] * [e].$$

The picture is that  $[0, 1] \times [0, 1]$  has  $\gamma$  on the top side and  $\gamma * e$  on the bottom. By drawing a line from the midpoint of the bottom side and the top-right corner, we see that we can fill the right portion with just  $x_0$  and the left portion with  $\gamma$ . The equation of this line is  $s = 2t - 1$ , so  $t = (s + 1)/2$ . Thus from the picture, we can write the explicit homotopy  $H : [0, 1] \times [0, 1] \rightarrow X$  via

$$H(t, s) = \begin{cases} \gamma(2/(s + 1), t) & \text{if } 0 \leq t \leq (s + 1)/2, \\ x_0 & \text{if } (s + 1)/2 \leq t \leq 1. \end{cases}$$

One can use a similar construction to show that  $[e] * [\gamma] = [\gamma]$ , so that  $[e]$  is an identity element.

Now we show the existence of inverses. Given a loop  $\gamma$ , define  $\bar{\gamma}$  via  $\bar{\gamma}(t) = \gamma(1 - t)$ , i.e.  $\gamma$  backwards. Set  $\gamma_s(t) = \gamma(st)$ . Note that as  $t$  goes from 0 to 1,  $\gamma_s$  goes from  $\gamma(0)$  to  $\gamma(s)$ , and also that  $\bar{\gamma}_s(t) = \gamma(s - st)$ . So we can write the homotopy  $H : [0, 1] \times [0, 1] \rightarrow X$  between  $\gamma * \bar{\gamma}$  and  $e$  by

$$H(t, s) = \begin{cases} \gamma_s(2t) & \text{if } 0 \leq t \leq 1/2, \\ \bar{\gamma}_s(2t - 1) & \text{if } 1/2 \leq t \leq 1 \end{cases} = \begin{cases} \gamma(2st) & \text{if } 0 \leq t \leq 1/2, \\ \gamma(s - s(2t - 1)) & \text{if } 1/2 \leq t \leq 1. \end{cases}$$

Thus setting  $[\gamma]^{-1} = [\bar{\gamma}]$  gives us inverses.

Finally, for associativity, we need to see that  $(\gamma_1 * \gamma_2) * \gamma_3 \sim \gamma_1 * (\gamma_2 * \gamma_3)$ . Again by drawing a picture, we see that we can draw two diagonal lines connecting the starting points of  $\gamma_2$  on top and bottom and the ending points of  $\gamma_1$  on top and bottom. Write an explicit formula for the homotopy as an exercise.  $\square$

## 4.2 Induced Homomorphisms

**Remark.** If  $f : X \rightarrow Y$  and  $x_0 \in X$ , let  $y_0 = f(x_0)$ . Then given a based loop  $\gamma : [0, 1] \rightarrow X$ , note that the composition  $f \circ \gamma : [0, 1] \rightarrow Y$  is a based loop in  $Y$ . Also, if  $\gamma \sim \delta$ , then  $f \circ \gamma \sim f \circ \delta$  (if  $H$  is a homotopy from  $\gamma \sim \delta$ , then  $f \circ H$  is a homotopy from  $f \circ \gamma$  to  $f \circ \delta$ ). In particular,  $f$  induces a map

$$f_* : \pi_1(X, x_0) \rightarrow \pi_1(Y, y_0).$$

**Lemma 4.2.** *The induced map  $f_* : \pi_1(X, x_0) \rightarrow \pi_1(Y, y_0)$  is a homomorphism.*

*Proof.* Note that

$$\gamma_1 * \gamma_2(t) = \begin{cases} \gamma_1(2t) & \text{if } 0 \leq t \leq 1/2, \\ \gamma_2(2t - 1) & \text{if } 1/2 \leq t \leq 1 \end{cases}$$

and that

$$(f \circ \gamma_1) * (f \circ \gamma_2)(t) = \begin{cases} f(\gamma_1(2t)) & \text{if } 0 \leq t \leq 1/2, \\ f(\gamma_2(2t - 1)) & \text{if } 1/2 \leq t \leq 1, \end{cases}$$

so  $f \circ (\gamma_1 * \gamma_2) = (f \circ \gamma_1) * (f \circ \gamma_2)$ , which implies  $f_*([\gamma_1 * \gamma_2]) = f_*([\gamma_1]) * f_*([\gamma_2])$ .  $\square$

**Exercise 4.2.** Check the following as an exercise:

1.  $(f \circ g)_* = f_* \circ g_*$ ;
2. if  $f : X \rightarrow Y$  is homotopic rel base point to  $g : X \rightarrow Y$ , then  $f_* = g_* : \pi_1(X, x_0) \rightarrow \pi_1(Y, y_0)$ .

**Remark.** How does  $\pi_1$  depend on the base point? Let  $x_0, x_1 \in X$ , and suppose that there exists a path  $h : [0, 1] \rightarrow X$  with  $h(0) = x_0$  and  $h(1) = x_1$ . Then if  $\gamma$  is a loop based at  $x_1$ , we can get a loop based at  $x_0$  by going from  $x_0$  to  $x_1$  along  $h$ , taking  $\gamma$ , and then going back to  $x_0$ . More explicitly, this is

$$h * \gamma * \bar{h}(t) = \begin{cases} h(3t) & \text{if } 0 \leq t \leq 1/3, \\ \gamma(3t - 1) & \text{if } 1/3 \leq t \leq 2/3, \\ \bar{h}(3t - 2) & \text{if } 2/3 \leq t \leq 1. \end{cases}$$

**Lemma 4.3.** *The path  $h$  induces an isomorphism  $\phi_h : \pi_1(X, x_1) \rightarrow \pi_1(X, x_0)$  by  $[\gamma] \mapsto [h * \gamma * \bar{h}]$ .*

*Proof.* Check as an exercise that  $\phi_h$  is a well-defined homomorphism. To show that  $\phi_h$  is an isomorphism, we claim that  $\phi_{\bar{h}}$  is an inverse of  $\phi_h$ . To see this, let  $[\gamma] \in \pi_1(X, x_0)$ . Then

$$\phi_h \circ \phi_{\bar{h}}([\gamma]) = [h * \bar{h} * \gamma * h * \bar{h}] = [h * \bar{h}] * [\gamma] * [h * \bar{h}] = [e] * [\gamma] * [e] = [\gamma],$$

where the second equality follows by the same proof for associativity of  $*$ . This proves the result.  $\square$

**Remark.** Note the following based on the above lemma:

1. The isomorphism class of  $\pi_1(X, x_0)$  only depends on the path component of  $X$  containing  $x_0$ .
2. The isomorphism *depends on*  $h$ . One needs to be careful about using the correct identification.



# Lecture 5

## Jan. 22 — Simple Computations

### 5.1 Fundamental Groups and Homotopy Equivalence

**Lemma 5.1.** *Suppose  $f_0, f_1 : X \rightarrow Y$  are homotopic by the homotopy  $H : X \times [0, 1] \rightarrow Y$ . Let  $x_0 \in X$  be a basepoint and define  $h : [0, 1] \rightarrow Y$  by  $t \mapsto H(x_0, t)$ . Then the following diagram commutes:*

$$\begin{array}{ccc} \pi_1(X, x_0) & \xrightarrow{(f_0)_*} & \pi_1(Y, f_0(x_0)) \\ & \searrow (f_1)_* & \uparrow \phi_h \\ & & \pi_1(Y, f_1(x_0)) \end{array}$$

*Proof.* Fix an arbitrary  $[\gamma] \in \pi_1(X, x_0)$ , and we construct a homotopy from  $h * (f_1 \circ \gamma) * \bar{h}$  to  $f_0 \circ \gamma$ . The picture is the following: we have  $h * (f_1 \circ \gamma) * \bar{h}$  at the top and  $f_0 \circ \gamma$  at the bottom, where  $t$  parametrizes the horizontal direction and  $s$  parametrizes the vertical direction. Draw a trapezoidal shape by connecting the middle two points on the top edge to the two bottom corners.

Define the homotopy as follows: For a fixed  $s$ , define  $H'(t, s)$  first by  $h_s(t)$  in the first third,  $H(\gamma(t), s)$  in the second third, and then  $\bar{h}_s(t)$  in the last third. Construct the explicit homotopy as an exercise.  $\square$

**Theorem 5.1.** *If  $f : X \rightarrow Y$  is a homotopy equivalence, then the induced map*

$$f_* : \pi_1(X, x_0) \rightarrow \pi_1(Y, f(x_0))$$

*on fundamental groups is an isomorphism.*

*Proof.* Let  $g$  be a homotopy inverse to  $f$ , so we have the composition:

$$\pi_1(X, x_0) \xrightarrow{f_*} \pi_1(Y, f(x_0)) \xrightarrow{g_*} \pi_1(X, g(f(x_0)))$$

But we know  $g \circ f = \text{id}_X$ , so by the lemma there exists a path  $h : X \rightarrow X$  from  $x_0$  to  $g(f(x_0))$  such that  $g_* \circ f_* = \phi_h$  is an isomorphism. So  $f_*$  is injective. Similarly,  $f_* \circ g_*$  is an isomorphism, and therefore  $f_*$  is surjective. Since we already know  $f_*$  is a homomorphism, this shows that  $f_*$  is an isomorphism.  $\square$

**Remark.** Recall that we have defined a “functor”

$$\{\text{pointed topological spaces, pointed maps}\} \rightarrow \{\text{groups, homomorphisms}\},$$

where homotopy equivalent spaces are mapped to isomorphic groups and homotopic maps give rise to the “same” homomorphism. We will finally make some computations of fundamental groups next.

## 5.2 Simple Computations of Fundamental Groups

**Lemma 5.2.** *If  $X$  is contractible, then  $\pi_1(X, x_0) = \{1\}$  for all  $x_0 \in X$ , where  $\{1\}$  is the trivial group.*

*Proof.* If  $Y = \{y_0\}$  is a one-point space, then there exists a unique loop  $\gamma : [0, 1] \rightarrow X$  given by  $t \mapsto y_0$ . So  $\pi_1(Y, y_0) = \{1\}$ . Since  $X$  is contractible, it is homotopy equivalent to  $Y$  and so  $\pi_1(X, x_0) = \{1\}$ .  $\square$

**Definition 5.1.** We say that a space  $X$  is *simply connected* if

1.  $X$  is path-connected, and
2.  $\pi_1(X, x_0) = \{1\}$  for some  $x_0 \in X$ .

**Remark.** Simply connected means that “points in  $X$  are connected in a very simple way.”

**Lemma 5.3.** *A space  $X$  is simply connected if and only if every two points in  $X$  are connected by a unique homotopy class of paths in  $X$ .*

*Proof.* ( $\Leftarrow$ ) Clearly  $X$  is path-connected. Furthermore, any loop based at  $x_0$  is a path from  $x_0$  to itself, and the constant is as well. Thus any loop is homotopic to the constant loop, i.e.  $\pi_1(X, x_0) = \{1\}$ .

( $\Rightarrow$ ) For any  $a, b \in X$ , there exists a path from  $a$  to  $b$  (since  $X$  is path-connected). Now suppose  $\gamma, \delta : [0, 1] \rightarrow X$  are paths from  $a$  to  $b$ . By  $\pi_1(X, a) = \{1\}$  we know that  $\gamma * \bar{\delta} \sim e_a$ , so

$$\gamma \sim \gamma * (\bar{\delta} * \delta) \sim (\gamma * \bar{\delta}) * \delta \sim e_a * \delta \sim \delta,$$

i.e.  $\gamma$  and  $\delta$  are in the same homotopy class.  $\square$

**Lemma 5.4.** *Let  $X = A \cup B$ , where  $A, B, A \cap B$  are open and path-connected. Let  $x_0 \in A \cap B$ . Then any loop  $\gamma : [0, 1] \rightarrow X$  based at  $x_0$  can be written as*

$$\gamma \sim \gamma_1 * \gamma_2 * \cdots * \gamma_n,$$

where each  $\gamma_i$  is a loop in  $A$  or  $B$  based at  $x_0$ .

*Proof.* We first claim that there exist  $0 = t_0 < t_1 < \cdots < t_n = 1$  such that  $\text{im } \gamma|_{[t_{i-1}, t_i]} \subseteq A$  or  $B$ , and  $\gamma(t_i) \in A \cap B$  for every  $i$ . The proof of this will use the following topology fact:

**Lemma** (Lebesgue number lemma). Let  $X$  be a compact metric space and  $\{U_\alpha\}_{\alpha \in A}$  be an open cover. Then there exists a *Lebesgue number*  $\delta > 0$  such that for all sets  $S$  with  $\text{diam}(S) = \sup_{x, y \in S} d(x, y) < \delta$ , there exists  $\alpha \in A$  such that  $S \subseteq U_\alpha$ .

To prove the claim, let  $U_1 = \gamma^{-1}(A)$  and  $U_2 = \gamma^{-1}(B)$ , which is an open cover of  $[0, 1]$ . So there exists  $\delta > 0$  such that if  $|b - a| < \delta$ , then  $[a, b] \subseteq U_i$  for  $i = 1$  or  $2$ . Thus  $\gamma([a, b]) \subseteq A$  or  $B$ . Now let  $n$  be a positive integer such that  $1/n < \delta$ , so that for each  $i$  we have

$$\text{im } \gamma|_{[i/n, (i+1)/n]} \subseteq A \text{ or } B.$$

So we start with  $t_i = i/n$  for  $i = 0, \dots, n$ . Now if  $\gamma|_{[t_{i-1}, t_i]}$  and  $\gamma|_{[t_i, t_{i+1}]}$  both have image in  $A$  (or both in  $B$ ), then throw out  $t_i$ . Then  $\gamma|_{[t_{i-1}, t_i]}$  will have image in  $A$  or  $B$  and  $\gamma(t_i) \in A \cap B$ , as desired.

Now given the claim, let  $\delta_i : [0, 1] \rightarrow A \cap B$  connect  $x_0$  to  $\gamma(t_i)$ , and set  $\gamma_i = \gamma|_{[t_{i-1}, t_i]}$ . Then

$$\gamma \sim \gamma_1 * \gamma_2 * \cdots * \gamma_n \sim (\gamma_1 * \bar{\delta}_1) * (\delta_1 * \gamma_2 * \bar{\delta}_2) * \cdots * (\delta_{n-1} * \gamma_n),$$

where each of the above loops is either in  $A$  or  $B$ .  $\square$

**Theorem 5.2.** *We have  $\pi_1(S^n, x_0) = \{1\}$  for all  $n \geq 2$ .*

*Proof.* We have  $\pi_1(S^n, x_0) = \{1\}$  for all  $n \geq 2$ . Let  $A = S^n \setminus \{(0, \dots, 0, 1)\}$  and  $B = S^n \setminus \{(0, \dots, 0, -1)\}$ . Note that  $A \cong B \cong \mathbb{R}^n$ , so they are path-connected. Furthermore,  $A \cap B = S^{n-1} \times \mathbb{R}$ , which is also path-connected if  $n \geq 2$ .<sup>1</sup> Now take any  $x_0 \in A \cap B$ . Then any  $[\gamma] \in \pi_1(S^n, x_0)$  can be written as

$$[\gamma] = [\gamma_1] * [\gamma_2] * \cdots * [\gamma_n],$$

where  $[\gamma_i] \in \pi_1(A, x_0)$  or  $\pi_1(B, x_0)$  by the lemma. But we have

$$\pi_1(A, x_0) = \pi_1(B, x_0) = \{1\},$$

so  $[\gamma] = [e_{x_0}]$  and hence  $\pi_1(S^n, x_0) = \{1\}$ . □

**Theorem 5.3.** *Given two spaces  $X$  and  $Y$ , we have  $\pi_1(X \times Y, (x_0, y_0)) \cong \pi_1(X, x_0) \times \pi_1(Y, y_0)$ .*

*Proof.* The map  $\Phi : \pi_1(X, x_0) \times \pi_1(Y, y_0) \rightarrow \pi_1(X \times Y, (x_0, y_0))$  given by

$$([\gamma], [\delta]) \mapsto [\gamma \times \delta],$$

where  $(\gamma \times \delta)(t) = (\gamma(t), \delta(t))$ , is an isomorphism. Check as an exercise that  $\Phi$  is well-defined and a bijection (for the second part, consider the projections  $p_X : X \times Y \rightarrow X$  and  $p_Y : X \times Y \rightarrow Y$ ). □

## 5.3 Fundamental Group of the Circle

Our next objective is the following computation:

**Theorem 5.4.** *We have  $\pi_1(S^1, (1, 0)) \cong \mathbb{Z}$ . In particular, the map sending  $n \in \mathbb{Z}$  to*

$$\gamma_n : [0, 1] \rightarrow S^1 : t \mapsto (\cos 2\pi nt, \sin 2\pi nt)$$

*is an isomorphism  $\mathbb{Z} \rightarrow \pi_1(S^1, (1, 0))$ .*

**Remark.** The proof is an example of a very important technique that we will see again soon. The proof involves studying the map

$$p : \mathbb{R} \rightarrow S^1 : t \mapsto (\cos 2\pi t, \sin 2\pi t).$$

Note that  $p^{-1}((1, 0)) = \mathbb{Z}$ . This is a particular example of a *covering map*, which we will study later.

**Definition 5.2.** If  $\gamma : [0, 1] \rightarrow S^1$  is a path based at the point  $(1, 0)$ , then a *lift of  $\gamma$  based at  $n \in \mathbb{Z}$*  is a map  $\tilde{\gamma}_n : [0, 1] \rightarrow \mathbb{R}$  such that  $\tilde{\gamma}_n(0) = n$  and  $p \circ \tilde{\gamma}_n = \gamma$ .

**Lemma 5.5.** *We have the following:*

- (a) *For each  $n \in \mathbb{Z}$ , each loop  $\gamma : [0, 1] \rightarrow S^1$  based at  $(1, 0)$  lifts to a unique path  $\tilde{\gamma}_n$  based at  $n$ .*
- (b) *If  $\gamma \sim \gamma'$  are loops in  $S^1$  based at  $(1, 0)$  and  $\tilde{\gamma}_n, \tilde{\gamma}'_n$  are their lifts based at  $n$ , then  $\tilde{\gamma}_n \sim \tilde{\gamma}'_n \text{ rel } \{0, 1\}$ .*

*The above properties are called path lifting and homotopy lifting.*

*Proof.* We will prove this next class. □

---

<sup>1</sup>Note that this does not work for  $n = 1$ : the intersection  $S^0 \times \mathbb{R} = \{\pm 1\} \times \mathbb{R}$  is not path-connected.

*Proof of Theorem 5.4.* Given  $[\gamma] \in \pi_1(S^1, (1, 0))$ , part (a) of Lemma 5.5 says that there is a unique lift  $\tilde{\gamma}_0 : [0, 1] \rightarrow \mathbb{R}$ . Since  $\tilde{\gamma}_0(1) \in p^{-1}((1, 0)) = \mathbb{Z}$ , we can define  $\Phi : \pi_1(S^1, (1, 0)) \rightarrow \mathbb{Z}$  by  $[\gamma] \mapsto \tilde{\gamma}_0(1)$ . Part (b) of Lemma 5.5 says that  $\Phi$  is well-defined. We need to show the following:

1.  $\Phi$  is surjective:

Let  $\tilde{\delta}^n(t) = nt$  for  $t \in [0, 1]$  and  $\delta^n = p \circ \tilde{\delta}^n$ . Then  $\tilde{\delta}^n$  is the lift of  $\delta^n$  based at 0, and  $\Phi([\delta^n]) = n$ .

2.  $\Phi$  is injective:

Suppose  $\gamma, \gamma'$  are loops in  $S^1$  such that

$$\Phi([\gamma]) = \tilde{\gamma}_0(1) = \tilde{\gamma}'_0(1) = \Phi([\gamma']).$$

Set  $\tilde{H}(s, t) = (1 - t)\tilde{\gamma}_0(s) + t\tilde{\gamma}'_0(s)$  and  $H = p \circ \tilde{H}$ . Then  $H$  is a homotopy from  $\gamma$  to  $\gamma'$  (exercise).

3.  $\Phi$  is a homomorphism:

Let  $[\gamma], [\gamma'] \in \pi_1(S^1, (1, 0))$  and let  $\tilde{\gamma}_0, \tilde{\gamma}'_0$  be their lifts based at 0. Then

$$\Phi([\gamma]) = \tilde{\gamma}_0(1) = n \in \mathbb{Z} \quad \text{and} \quad \Phi([\gamma']) = \tilde{\gamma}'_0(1) = m \in \mathbb{Z}.$$

Then note the following:

(a)  $t \mapsto n + \tilde{\gamma}'_0(t)$  is a lift of  $\gamma'$  and starts at  $n$ , so by uniqueness it is  $\tilde{\gamma}'_n$ , and

(b)  $\tilde{\gamma}_0 * \tilde{\gamma}'_n$  is a lift of  $\gamma * \gamma'$ .

So  $\Phi([\gamma] * [\gamma']) = \tilde{\gamma} * \tilde{\gamma}'_0(1) = n + m = \Phi([\gamma]) + \Phi([\gamma'])$ , i.e.  $\Phi$  is a homomorphism.

Thus  $\Phi$  is an isomorphism, and we have  $\pi_1(S^1, (1, 0)) \cong \mathbb{Z}$ . □

# Lecture 6

## Jan. 27 — Fundamental Group of the Circle

### 6.1 Proof of Path Lifting

*Proof of Lemma 5.5.* (a) Let  $A = S^1 \setminus \{(1, 0)\}$ , and note that

$$p^{-1}(A) = \bigcup_{i \in \mathbb{Z}} (i, i+1) = \bigcup_{i \in \mathbb{Z}} A_i.$$

Notice that each restriction  $p|_{A_i} : A_i \rightarrow A$  is a homeomorphism. Now let  $B = S^1 \setminus \{(-1, 0)\}$ , so

$$p^{-1}(B) = \bigcup_{i \in \mathbb{Z}} \left(i - \frac{1}{2}, i + \frac{1}{2}\right) = \bigcup_{i \in \mathbb{Z}} B_i.$$

Similarly, each  $p|_{B_i} : B_i \rightarrow B$  is a homeomorphism. Now if  $\gamma : [0, 1] \rightarrow S^1$  is contained in  $A$  (or  $B$ ), we can choose any  $i \in \mathbb{Z}$  and let  $\tilde{\gamma} = (p|_{A_i})^{-1} \circ \gamma$ , giving a lift of  $\gamma$ . Then for a general  $\gamma : [0, 1] \rightarrow S^1$  with  $\gamma(0) = (1, 0)$ , the set  $\{\gamma^{-1}(A), \gamma^{-1}(B)\}$  is an open cover of the compact metric space  $[0, 1]$ , so there exists a Lebesgue number  $\delta > 0$  such that any interval  $[a, b]$  with  $b - a < \delta$  lies in either  $\gamma^{-1}(A)$  or  $\gamma^{-1}(B)$ . Choose  $n$  such that  $1/n < \delta$ . If  $t_n = i/n$  for  $i = 0, \dots, n$ , then

$$\gamma([t_i, t_{i+1}]) \subseteq A \text{ or } B$$

for every  $i$ . Again for convenience, if  $[t_{i-1}, t_i]$  and  $[t_i, t_{i+1}]$  are both in  $\gamma^{-1}(A)$  or  $\gamma^{-1}(B)$ , then discard  $t_i$ . So we have a partition  $0 = t_0 < t_1 < \dots < t_k = 1$  such that (note that  $\gamma$  starts at  $(1, 0) \notin A$ )

$$\gamma([t_i, t_{i+1}]) \subseteq \begin{cases} A & \text{if } i \text{ is odd,} \\ B & \text{if } i \text{ is even.} \end{cases}$$

Then we want to build  $\tilde{\gamma}_n$ . Define  $\tilde{\gamma}_n$  on  $[t_0, t_1]$  to be  $(p|_{B_n})^{-1} \circ \gamma|_{[t_0, t_1]}$ . Now  $\tilde{\gamma}_n(t_1) \in A_i$  for a unique  $i$ , so define  $\tilde{\gamma}_n$  on  $[t_1, t_2]$  by  $(p|_{A_i})^{-1} \circ \gamma|_{[t_1, t_2]}$ . Note that  $\tilde{\gamma}_n$  is continuous on  $[t_0, t_2]$  since the two definitions agree at  $t = t_1$ . Inductively continue to define the lift  $\tilde{\gamma}_n$  on all of  $[0, 1]$ .

(b) The proof is very similar to path lifting. Given a homotopy  $H : [0, 1] \times [0, 1] \rightarrow S^1$ , we can find a Lebesgue number  $\delta > 0$  for  $\{H^{-1}(A), H^{-1}(B)\}$ . Pick  $n$  such that  $\sqrt{2}/n < \delta$  and break  $[0, 1] \times [0, 1]$  into  $n^2$  squares of side length  $1/n$ . The diameter of each square is at most  $\sqrt{2}/n$ , so each square can be lifted as above. Finish the construction as an exercise to lift  $H$  on all of  $[0, 1] \times [0, 1]$ .  $\square$

### 6.2 Applications of the Fundamental Group of $S^1$

**Corollary 6.0.1.** *There is no retraction  $D^2 \rightarrow \partial D^2$ .*

*Proof.* Suppose there was a retraction  $r : D^2 \rightarrow \partial D^2$ , and let  $i : S^1 \rightarrow D^2$  be the inclusion of  $S^1$  as the boundary of  $D^2$ . Then we have the composition:

$$S^1 \xrightarrow{i} D^2 \xrightarrow{r} S^1$$

Noting that  $r \circ i = S^1 \rightarrow S^1$  is the identity, so  $(r \circ i)_* : \pi_1(S^1, (1, 0)) \rightarrow \pi_1(S^1, (1, 0))$  is the identity map. In particular,  $r_* \circ i_* = (r \circ i)_*$  is the identity map, hence  $i_*$  must be injective. But

$$i_* : \pi_1(S^1, (1, 0)) \rightarrow \pi_1(D^2, (1, 0))$$

where  $\pi_1(S^1, (1, 0)) \cong \mathbb{Z}$  and  $\pi_1(D^2, (1, 0)) = \{1\}$ , so  $i_*$  cannot be injective. Contradiction.  $\square$

**Corollary 6.0.2.** *Any map  $f : D^2 \rightarrow D^2$  has a fixed point, i.e.  $x \in D^2$  such that  $f(x) = x$ .*

*Proof.* Suppose otherwise that  $f : D^2 \rightarrow D^2$  has no fixed points. Then for each  $x \in D^2$ , there is a unique ray  $R_x$  starting at  $f(x)$  and going through  $x$ . Note that  $R_x \cap \partial D^2$  in a unique point (on the interior of  $R_x$ ). Define  $r : D^2 \rightarrow S^1$  by  $x \mapsto R_x \cap \partial D^2$ . Show that  $r$  is continuous as an exercise (e.g. parametrize the line). But then  $r$  is a retraction  $D^2 \rightarrow \partial D^2$ , a contradiction.  $\square$

**Remark.** There are more applications such as the fundamental theorem of algebra, the ham sandwich theorem, and the Borsuk-Ulam theorem. See Hatcher for more details.

## 6.3 Free Products of Groups

**Definition 6.1.** Let  $G_1$  and  $G_2$  be groups. A *word* in  $G_1 \sqcup G_2$  is a finite sequence

$$x = (x_1, x_2, \dots, x_n)$$

for some  $n$ , where each  $x_i$  is in  $G_1$  or  $G_2$ . Define an equivalence relation on words in  $G_1 \sqcup G_2$  which is generated by (show as an exercise that this is in fact an equivalence relation):

1. replace  $a, b$  in a sequence by  $ab$  if  $a, b$  are in the same group (or the reverse of this), and
2. if  $e_i$  (the identity in  $G_i$ ) is in a sequence, then remove it (or add it in any place in a sequence).

Denote the equivalence class of a word  $x$  by  $[x]$ . Call a word  $x = (x_1, \dots, x_n)$  *reduced* if

1.  $x_j \neq e_i$  for any  $j$  or  $i$ , and
2.  $x_i$  and  $x_{i+1}$  are from different groups.

Show that each  $[x]$  contains a unique reduced word (note: uniqueness is hard). The *free product* of  $G_1$  and  $G_2$  is the group  $G_1 * G_2$  of all equivalence classes of words in  $G_1 \sqcup G_2$ , with multiplication

$$[x_1, \dots, x_n] \cdot [y_1, \dots, y_m] = [x_1, \dots, x_n, y_1, \dots, y_m].$$

**Remark.** Note that in  $G_1 * G_2$ , the identity  $e$  is the empty word and the inverse is given by

$$[x_1, \dots, x_n]^{-1} = [x_n^{-1}, \dots, x_1^{-1}].$$

Check as an exercise that  $G_1 * G_2$  is in fact a group (really only need to check associativity).

**Proposition 6.1.** *Let  $j_i : G_i \rightarrow G_1 * G_2$  be the inclusion of  $G_i$  into  $G_1 * G_2$ . Given any homomorphisms  $\phi_i : G_i \rightarrow H$  where  $H$  is any group, there exists a unique homomorphism  $\phi : G_1 * G_2 \rightarrow H$  such that  $\phi \circ j_i = \phi_i$ , i.e. the following diagram commutes:*

$$\begin{array}{ccccc}
 & & G_1 & & \\
 & & \downarrow j_1 & \searrow \phi_1 & \\
 G_2 & \xrightarrow{j_2} & G_1 * G_2 & \xrightarrow{\phi} & H \\
 & \searrow \phi_2 & & & 
 \end{array}$$

*Proof.* If the  $x_i$  are reduced and  $x_1 \in G_1$ , then define

$$\phi_1(x_1, x_2, \dots, x_n) = \phi_1(x_1) \cdot \phi_2(x_2) \cdot \phi_1(x_3) \cdot \dots$$

Then check the following as an exercise:

1. Show such that  $\phi$  exists and is unique.
2. Show this property *defines* the free product, i.e. if  $D$  is another group satisfying the property in the proposition, then  $D \cong G_1 * G_2$ .

The second part above says that this is the *universal property* of the free product. □

**Example 6.1.1.** Represent  $\mathbb{Z}$  in product notation via  $\{x^n\}$ , where  $x^n x^m = x^{n+m}$ . Then

$$\mathbb{Z} * \mathbb{Z} = \{x^n\} * \{y^m\} = \{e, x^{n_1} y^{m_1} \dots x^{n_k}, x^{n_1} y^{m_1} \dots y^{m_k}, y^{m_1} x^{n_1} \dots y^{m_k}, y^{m_1} x^{n_1} \dots x^{n_k}\}.$$

This group is called the *free group on two generators*, and  $\mathbb{Z}$  is the *free group on one generator*.

# Lecture 7

## Jan. 29 — Some Group Theory

### 7.1 Group Presentations

**Definition 7.1.** The *free group* on  $n$  generators, denoted  $F_n$ , is defined inductively via

$$F_n = F_{n-1} * \mathbb{Z},$$

where  $F_1 = \mathbb{Z}$ . (One can also consider  $F_\infty$ .)

**Remark.** Note that any homomorphism  $\phi : \mathbb{Z} \rightarrow G$  (for any group  $G$ ) is determined by  $\phi(1)$ . Moreover, given any  $g \in G$ , there is a unique homomorphism which maps  $1 \mapsto g$ . Thus by the universal property of the free product, a homomorphism  $F_n \rightarrow G$  is determined uniquely by a choice of  $g_1, \dots, g_n \in G$ .

**Definition 7.2.** A *group presentation* is a group  $\langle X|R \rangle$  defined as follows:

- $X$  is some set (of generators);
- $R$  is a set of words (relations) in  $X \cup X^{-1}$  (formally denote  $x \in X$  as  $x^{-1} \in X^{-1}$ );
- let  $n = |X|$  and  $F_n$  be the free group on  $n$  generators, so that we can think of  $R \subseteq F_n$ ;
- let  $\langle R \rangle$  be the smallest normal subgroup of  $F_n$  containing  $R$ ;
- define the group  $\langle X|R \rangle = F_n / \langle R \rangle$ .

We say that  $\langle X|R \rangle$  is a *presentation* of a group  $G$  if  $G \cong \langle X|R \rangle$ .

**Example 7.2.1.** The group  $\langle g|g^n \rangle$  is all the words in  $g, g^{-1}$ :

$$\dots, g^{-2}, g^{-1}, e, g, g^2, g^3, \dots,$$

but  $g^n = e$ , so  $g^{n+1} = g^n g = eg = g$  and thus we have  $g^{-1} = eg^{-1} = g^n g^{-1} = g^{n-1} g g^{-1} = g^{n-1}$ . So there is a one-to-one correspondence between elements of  $\langle g|g^n \rangle$  and  $g^k$  for  $k = 0, \dots, n-1$ .

**Exercise 7.1.** Show that  $\langle g|g^n \rangle \cong \mathbb{Z}/n$ , so that  $\langle g|g^n \rangle$  is a presentation of  $\mathbb{Z}/n$ .

**Lemma 7.1.** *Every group has a presentation.*

*Proof.* Let  $G$  be a group. Let  $X \subseteq G$  be a collection of elements of  $G$  that generate  $G$  (e.g. take  $X = G$  itself). Let  $n = |X|$ , so there exists a unique  $\phi : F_n \rightarrow G$  sending the generators of  $F_n$  to the  $g_i \in X$ . Let  $N = \ker \phi$ , so the first isomorphism theorem says that  $G \cong F_n / N$  (note that  $\phi$  is clearly surjective). Let  $R$  be a subset of  $N$  that generates  $N$  (e.g. take  $R = N$ ). Then  $G \cong \langle X|R \rangle$ .  $\square$

**Remark.** Using fewer generators to write  $G$  or  $N$  may give more less complicated presentations of  $G$ .



**Definition 7.3.** We say that  $G$  is *finitely generated* if  $G \cong \langle X | R \rangle$  such that  $|X| < \infty$ . We say that  $G$  is *finitely presented* if both  $|X|, |R| < \infty$ .

**Exercise 7.2.** Show the following:

1. If  $G = \langle g_1, \dots, g_n | r_1, \dots, r_m \rangle$ , then for any group  $H$  and any map  $h : \{g_1, \dots, g_n\} \rightarrow H$  satisfying  $h(r_i) = e_H$  (the notation  $h(r_i)$  means to replace any letters  $g_j$  in  $r_i$  by  $h(g_j)$ ), there exists a unique homomorphism  $\phi_h : G \rightarrow H$  such that  $\phi_h(g_i) = h(g_i)$ .
2. If  $G_1 = \langle g_1, \dots, g_n | r_1, \dots, r_m \rangle$  and  $G_2 = \langle h_1, \dots, h_k | s_1, \dots, s_\ell \rangle$ , then

$$G_1 * G_2 = \langle g_1, \dots, g_n, h_1, \dots, h_k | r_1, \dots, r_m, s_1, \dots, s_\ell \rangle.$$

**Definition 7.4.** Given groups  $G_1, G_2$  and  $K$ , and homomorphisms  $\psi_i : K \rightarrow G_i$ , the *free product with amalgamation* is

$$G_1 *_K G_2 = \frac{G_1 * G_2}{\langle \{\psi_1(k)\psi_2(k)^{-1}\}_{k \in K} \rangle},$$

where  $\langle \{\psi_1(k)\psi_2(k)^{-1}\}_{k \in K} \rangle$  is the smallest normal subgroup of  $G_1 * G_2$  containing  $\{\psi_1(k)\psi_2(k)^{-1}\}_{k \in K}$ .

**Remark.** The idea is that  $G_1 *_K G_2$  is the set of all words in  $G_1 \cup G_2$  but if we see  $\psi_1(k)$  in a word, we can replace it by  $\psi_2(k)$  and vice versa. In terms of group presentations, if

$$\begin{aligned} G_1 &= \langle g_1, \dots, g_n | r_1, \dots, r_m \rangle, \\ G_2 &= \langle g'_1, \dots, g'_{n'} | r'_1, \dots, r'_{m'} \rangle, \\ K &= \langle h_1, \dots, h_k | s_1, \dots, s_\ell \rangle, \end{aligned}$$

then we can write

$$G_1 *_K G_2 = \langle g_1, \dots, g_n, g'_1, \dots, g'_{n'} | r_1, \dots, r_m, r'_1, \dots, r'_{m'}, \psi_1(h_1)(\psi_2(h_1))^{-1}, \dots, \psi_1(h_k)(\psi_2(h_k))^{-1} \rangle.$$


Note that the  $s_i$  show up in the above since  $\psi_1, \psi_2$  are homomorphisms.

**Exercise 7.3.** Show the following:

1. Check that the above presentation for  $G_1 *_K G_2$  is correct.
2. Let  $\iota_i : G_i \rightarrow G_1 * G_2$  be the inclusions and  $j_i : G_i \rightarrow G_1 *_K G_2$  be the induced maps. Then given any homomorphisms  $\phi_i : G_i \rightarrow H$  (where  $H$  is any group) such that

$$\phi_1 \circ \psi_1(k) = \phi_2 \circ \psi_2(k) \quad \text{for all } k \in K,$$

then there exists a unique homomorphism  $\Phi : G_1 *_K G_2 \rightarrow H$  such that  $\Phi \circ j_i = \phi_i$ , i.e.



Show that this is the *universal property* for the free product with amalgamation.

## 7.2 Seifert-van Kampen Theorem

**Theorem 7.1** (Seifert-van Kampen). *Let  $X$  be a topological space with base point  $x_0$ . Let  $A, B \subseteq X$  be open sets with  $X = A \cup B$  such that  $A, B, A \cap B$  are path-connected and  $x_0 \in A \cap B$ . Let*

$$\psi_A : \pi_1(A, x_0) \rightarrow \pi_1(X, x_0) \quad \text{and} \quad \psi_B : \pi_1(B, x_0) \rightarrow \pi_1(X, x_0)$$

*be the homomorphisms induced from the inclusions  $A \cap B \rightarrow A$  and  $A \cap B \rightarrow B$ . Then*

$$\pi_1(X, x_0) \cong \pi_1(A, x_0) *_{\pi_1(A \cap B, x_0)} \pi_1(B, x_0).$$

**Remark.** There is a more general version where  $X = \bigcup_{\alpha \in A} U_\alpha$ . See Hatcher for more details.

**Example 7.4.1.** Consider  $W_2 = S^1 \vee S^1$ , a wedge of two circles. Let  $x_0$  be the point of intersection of the circles. Let  $A$  be an open neighborhood of the left circle and  $B$  be an open neighborhood of the right circle. Note that  $A \simeq B \simeq S^1$  and  $A \cap B \simeq \{\text{pt}\}$ . So we see that

$$\pi_1(A, x_0) \cong \mathbb{Z} \cong \langle g_1 | \rangle, \quad \pi_1(B, x_0) \cong \mathbb{Z} \cong \langle g_2 | \rangle, \quad \pi_1(A \cap B, x_0) = \{e\},$$

and so  $\psi_A : \pi_1(A \cap B, x_0) \rightarrow \pi_1(A, x_0)$  and  $\psi_B : \pi_1(A \cap B, x_0) \rightarrow \pi_1(B, x_0)$  must both map  $e \mapsto e$ . Thus

$$\pi_1(W_2, x_0) \cong \pi_1(A, x_0) *_{\pi_1(A \cap B, x_0)} \pi_1(B, x_0) \cong \langle g_1, g_2 | \psi_A(e)(\psi_B(e))^{-1} \rangle \cong \langle g_1, g_2 | \rangle \cong F_2,$$

by the Seifert-van Kampen theorem.

**Exercise 7.4.** Show the following:

1. If  $W_n$  is a wedge of  $n$  circles, then  $\pi_1(W_n, x_0) \cong F_n$ .
2. We have  $\pi_1(\text{any connected graph}) = F_n$  for some  $n$ .

**Example 7.4.2.** Consider the torus  $T^2 = S^1 \times S^1$ . Recall that for a product, we know

$$\pi_1(T^2) \cong \pi_1(S^1) \times \pi_1(S^1) \cong \mathbb{Z} \times \mathbb{Z}.$$

We can also use van Kampen's theorem to see this. Think of  $T^2$  as a square with opposite sides identified. Let  $A$  be the square with a circle missing in the circle, which deformation retracts to the boundary of the square. Let  $B$  be a big disk in the middle, so that  $A \cap B$  deformation retracts to a circle. So

$$A \simeq W_2, \quad B \simeq \{\text{pt}\}, \quad A \cap B \simeq S^1.$$

Thus by our previous computations, we have

$$\pi_1(A, x_0) \cong \langle g_1, g_2 | \rangle, \quad \pi_1(B, x_0) \cong \{e\}, \quad \pi_1(A \cap B, x_0) \cong \langle h | \rangle.$$

The inclusion  $\psi_B : \pi_1(A \cap B, x_0) \rightarrow \pi_1(B, x_0)$  sends  $h \rightarrow e$ , and  $\psi_A : \pi_1(A \cap B, x_0) \rightarrow \pi_1(A, x_0)$  sends  $h \mapsto g_1 g_2 g_1^{-1} g_2^{-1}$ . To see the last claim, push a loop in  $A \cap B$  to the boundary of the square (note that under the homotopy equivalence of  $A$  and  $W_2$ , we can assume  $x_0$  maps to a corner point), where it follows the four edges of the square. These are the loops  $g_1, g_2, g_1^{-1}, g_2^{-1}$ , where the last two loops are oriented in the opposite direction of the first two. Thus by van Kampen's theorem,

$$\pi_1(T^2, x_0) \cong \pi_1(A, x_0) *_{\pi_1(A \cap B, x_0)} \pi_1(B, x_0) \cong \langle g_1, g_2 | \psi_A(h)(\psi_B(h))^{-1} \rangle = \langle g_1, g_2 | g_1 g_2 g_1^{-1} g_2^{-1} \rangle.$$

**Exercise 7.5.** Check the following:

1. Show that  $\mathbb{Z} \times \mathbb{Z} \cong \langle g_1, g_2 | g_1 g_2 g_1^{-1} g_2^{-1} \rangle$ , so that  $\pi_1(T^2) \cong \mathbb{Z} \times \mathbb{Z}$  again.
2. Compute  $\pi_1(\Sigma_g, x_0)$ , where  $\Sigma_g$  is the surface of genus  $g$ . Show that  $\pi_1(\Sigma_g)$  is not abelian if  $g > 1$ .

# Lecture 8

## Feb. 3 — Seifert-van Kampen Theorem

### 8.1 Applications of the Seifert-van Kampen Theorem

**Theorem 8.1.** *Let  $X$  be a path-connected space  $f : \partial D^n \rightarrow X$  be continuous with  $x_0 \in \partial D^n$ . Set*

$$Y = X \cup_f D^n = X \sqcup D^n / \{(x \in D^n) \sim (f(x) \in X)\}.$$

*Then (for  $n = 1$ , we need  $X$  to have a base point  $f(x_0)$  with an open neighborhood  $U \simeq \{f(x_0)\}$ )*

$$\pi_1(Y, y_0) = \begin{cases} \pi_1(X, f(x_0)) * \mathbb{Z} & \text{if } n = 1, \\ \pi_1(X, f(x_0)) / \langle r \rangle & \text{if } n = 2, \\ \pi_1(X, f(x_0)) & \text{if } n \geq 3, \end{cases}$$

where  $r = f_*(g)$  where  $g$  generates  $\pi_1(\partial D^2, x_0) \cong \mathbb{Z}$ .

*Proof.* We proof this in the case  $n = 2$ . Let

$$A = X \cup_f (D^2 \setminus \{0\}) = X \cup_f (S^1 \times (0, 1]) \simeq X$$

and  $B$  be the interior of  $D^2$  (so  $B \simeq \{\text{pt}\}$ ). Then we can see that

$$A \cap B = (\text{int } D^2) \setminus \{0\} = S^1 \times (0, 1) \simeq S^1.$$

Note that we can choose  $y_0$  to be  $f(x_0) \in X$  because the that is where it is sent under the deformation retraction from  $A$  to  $X$ . Now  $\psi_A : \pi_1(A \cap B, y_0) \rightarrow \pi_1(A, y_0)$

$$\pi_1(A \cap B, y_0) \cong \langle g \rangle \quad \text{and} \quad \pi_1(A, y_0) \cong \pi_1(X, f(x_0))$$

is given by  $g \mapsto f_*(g)$ , and  $\psi_B(g) = e$ . Thus the Seifert-van Kampen theorem implies

$$\pi_1(Y, y_0) \cong \pi_1(A, y_0) *_{\pi_1(A \cap B, y_0)} \pi_1(B, y_0) \cong \frac{\pi_1(A, y_0) * \{e\}}{\langle \psi_A(g)(\psi_B(g))^{-1} \rangle} \cong \frac{\pi_1(A, y_0)}{\langle r \rangle} \cong \frac{\pi_1(X, f(x_0))}{\langle r \rangle},$$

which is the desired result for  $n = 2$ . The  $n \geq 3$  case is similar (except the intersection is now contractible in this case). Check the  $n = 1$  case as an exercise.  $\square$

**Remark.** This allows us to compute the fundamental group of any CW complex (hence any manifold).

**Corollary 8.1.1.** *Let  $G$  be a finitely presented group. Then there exists a topological space  $X$  (in fact, a compact CW complex) such that  $\pi_1(X, x_0) \cong G$ .*

*Proof.* Let  $G = \langle g_1, \dots, g_n | r_1, \dots, r_m \rangle$  and  $W_n$  be the wedge of  $n$  circles, so that

$$\pi_1(W_n, x_0) \cong F_n \cong \langle g_1, \dots, g_n \rangle.$$

Now for each  $r_i$  let  $f_i : S^1 \rightarrow W_n$  be a map such that  $(f_i)_*(g) = r_i$  (show as an exercise that such  $f_i$  exists; essentially for each word, take the loops corresponding to each letter in order). Let

$$X = W_n \cup_{f_i} \left( \bigsqcup_{i=1}^m D^2 \right),$$

and the previous theorem tells us that  $\pi_1(X, x_0) \cong \langle g_1, \dots, g_n | r_1, \dots, r_m \rangle \cong G$ .  $\square$

**Remark.** The topological space realizing  $G$  as its fundamental group is not unique. For instance, we can take the above construction and add a 5-cell, which does not change the fundamental group.

## 8.2 Proof of the Seifert-van Kampen Theorem

*Proof of Theorem 7.1.* We have the inclusions  $A \subseteq X$  and  $B \subseteq X$ , which induce maps

$$\phi_A : \pi_1(A, x_0) \rightarrow \pi_1(X, x_0) \quad \text{and} \quad \phi_B : \pi_1(B, x_0) \rightarrow \pi_1(X, x_0).$$

By the universal property of free products, we get a map  $\Phi : \pi_1(A, x_0) * \pi_1(B, x_0) \rightarrow \pi_1(X, x_0)$  by

$$([\gamma_1], [\delta_1], [\gamma_2], \dots) \mapsto \phi_A([\gamma_1])\phi_B([\delta_1])\phi_A([\gamma_2]) \dots$$

Note that if  $[\gamma] \in \pi_1(A \cap B, x_0)$ , then  $\psi_A([\gamma]) = [\gamma] = \psi_B([\gamma])$ , so

$$\phi_A \circ \psi_A([\gamma]) = [\gamma] = \phi_B \circ \psi_B([\gamma]).$$

This tells us that  $\Phi(\psi_A([\gamma])(\psi_B([\gamma]))^{-1}) = e$ , so we see that

$$K = \langle \psi_A([\gamma])(\psi_B([\gamma]))^{-1} \rangle_{[\gamma] \in \pi_1(A \cap B, x_0)}$$

lies in the kernel of  $\Phi$ . This gives us an induced map (still called it  $\Phi$ )

$$\Phi : \pi_1(A, x_0) *_{\pi_1(A \cap B, x_0)} \pi_1(B, x_0) \rightarrow \pi_1(X, x_0).$$

Lemma 5.4 says that  $\Phi$  is surjective, so it suffices to check injectivity. To do this, let  $[\gamma_i] \in \pi_1(A, x_0)$  and  $[\eta_i] \in \pi_1(B, x_0)$  with

$$\Phi([\gamma_1][\eta_1] \dots [\gamma_n][\eta_n]) = [\gamma_1 * \eta_1 * \dots * \gamma_n * \eta_n] = e. \quad (*)$$

We need to see that we can get from the word  $[\gamma_1][\eta_1] \dots [\gamma_n][\eta_n]$  to the empty word by a sequence of:

1. replace  $a, b$  by  $a \cdot b$  if  $a, b$  are in the same group (and the reverse of this);
2. if we see  $\psi_A(k)$  in the word, we can replace it with  $\psi_B(k)$  (and the reverse of this).

We will prove the theorem only for  $n = 2$ . Now  $(*)$  says that there exists a homotopy  $H$  between  $x_0$  and  $\gamma_1 * \eta_1$ . As before, we can use the Lebesgue number lemma to find  $n$  such that squares of side length  $1/n$  are mapped by  $H$  into either  $A$  or  $B$  (we can assume that the number of  $\gamma_i, \eta_i$  divides  $n$ ). Check as an exercise that we can assume  $H(i/n, j/n) = x_0$ , i.e. that we can change  $H$  and  $\gamma_i, \eta_i$  by a homotopy

such that the homotopy of  $\gamma_i$  is in  $A$  and the homotopy of  $\eta_i$  is in  $B$  (hint: consider radial lines around  $(i/n, j/n)$ ). So we have an  $n \times n$  grid where each corner point is  $x_0$ , and the bottom edges are

$$\gamma_1 \sim \gamma'_1 * \gamma''_1 \quad \text{and} \quad \eta_1 \sim \eta'_1 * \eta''_1,$$

where these homotopies take place in  $A$  or  $B$ , respectively. Let  $a_1, a_2, a_3, a_4$  be the four edges lying above  $\gamma'_1, \gamma''_1, \eta'_1, \eta''_1$  on the grid (recall that  $n = 2$  in this case) We will show that we can go from  $[\gamma_1][\eta_1]$  to  $[a_1][a_2][a_3][a_4]$  using (1) and (2). Then we can inductively push this to the top, which is the empty word.

Let  $\delta_1, \delta_2, \delta_3$  be the three interior edges which connect the bottom row and the second-to-last row. Since each square maps to  $A$  or  $B$ , the first three squares lie in  $A$  and the last one lies in  $B$ . Let

$$G = \pi_1(A, x_0) \quad \text{and} \quad H = \pi_1(B, x_0).$$

Note that we have

$$\delta_1 * \eta'_1 \sim \delta_1 * \eta' * \delta_3 * \bar{\delta}_3 \text{ in } G, \quad \delta_1 * \eta'_1 * \delta_3 \sim a_1 * a_2 * a_3 \text{ in } G, \quad \delta_3 * \eta''_1 \sim a_4 \text{ in } H,$$

and thus we can write

$$\begin{aligned} [\gamma_1]^G [\eta_1]^H &= [\gamma_1]^G ([\eta'_1][\eta''_1])^H \stackrel{(1)}{=} [\gamma_1]^G [\eta'_1]^H [\eta''_1]^H \\ &\stackrel{(2)}{=} [\gamma_1]^G [\eta'_1]^G [\eta''_1]^H \\ &\stackrel{(1)}{=} ([\gamma_1][\eta'_1])^G [\eta''_1]^H = ([\gamma_1][\eta'_1][\delta_3][\bar{\delta}_3])^G [\eta''_1]^H \\ &\stackrel{(2)}{=} ([\gamma_1][\eta'_1][\delta_3])^G [\bar{\delta}_3]^H [\eta''_1]^H \\ &\stackrel{(1)}{=} ([a_1][a_2][a_3])^G ([\bar{\delta}_3][\eta''_1])^H = ([a_1][a_2][a_3])^G [a_4]^H \\ &\stackrel{(1)}{=} [a_1]^G [a_2]^G [a_3]^G [a_4]^H, \end{aligned}$$

which proves the statement for  $n = 2$ . See Hatcher for the general case.  $\square$

### 8.3 Covering Spaces

**Definition 8.1.** A *covering space* of a space  $X$  is a pair  $(\tilde{X}, p)$  where  $\tilde{X}$  is a space and  $p : \tilde{X} \rightarrow X$  such that every point  $x \in X$  has an *evenly covered neighborhood*. An open set  $U$  is *evenly covered* if

$$p^{-1}(U) = \text{disjoint union of open sets } \{U_\alpha\} \text{ in } \tilde{X}$$

such that  $p|_{U_\alpha} : U_\alpha \rightarrow U$  is a homeomorphism for every  $\alpha$ .

**Example 8.1.1.** The following are examples of covering maps:

1. If  $p : \tilde{X} \rightarrow X$  is a homeomorphism, then it is a covering map.
2. We saw that  $p : \mathbb{R} \rightarrow S^1$  given by  $t \mapsto (\cos 2\pi t, \sin 2\pi t)$  is a covering map.

**Exercise 8.1.** If  $(\tilde{X}, p)$  is a covering space of  $X$  and  $(\tilde{Y}, p')$  is a covering space of  $Y$ , then show that

$$p \times p' : \tilde{X} \times \tilde{Y} \rightarrow X \times Y, \quad (x, y) \mapsto (p(x), p'(y))$$

is a covering map. This gives a covering map  $\mathbb{R}^2 \rightarrow T^2$  by  $(t, s) \mapsto (\cos 2\pi t, \sin 2\pi t, \cos 2\pi s, \sin 2\pi s)$ .

# Lecture 9

## Feb. 5 — Covering Spaces

### 9.1 Examples of Covering Spaces

**Example 9.0.1.** The following are more examples of covering maps:

3. Define  $p_n : S^1 \rightarrow S^1$  by  $\theta \mapsto n\theta$ . These are covering maps for each  $n \in \mathbb{Z}$ .
4. Let  $X$  be a wedge of two circles (corresponding to  $a, b$ ), and let  $\tilde{X}$  be a circle with three outer circles attached, evenly spaced (the inner circle corresponding to  $a_1, a_2, a_3$  and the three outer circles corresponding to  $b_1, b_2, b_3$ ). Let  $p$  map  $a_i$  to  $a$  and  $b_i$  to  $b$ . This is a covering map.

Write out a formula for  $p$  and really check that  $p$  is a covering map as an exercise.

5. Again let  $X$  be a wedge of two circles, labeled  $a, b$ . Let  $\tilde{X}$  be a wedge of two circles, with an extra circle attached on either side. Let  $a_3, b_3$  be the extra circles on the left and right, let  $b_1, b_2$  be the top and bottom halves of the circle on the left, and let  $a_1, a_2$  be the top and bottom halves of the circles on the right. Let  $p$  map  $a_i$  to  $a$  and  $b_i$  to  $b$ . Then  $p$  is a covering map.
6. Consider the quotient map  $p : S^2 \rightarrow \mathbb{R}P^2$ . This map is a covering map. Note that each point in  $\mathbb{R}P^2$  has a neighborhood whose preimage is two open sets on opposite sides of  $S^2$ .

### 9.2 Covering Spaces and Lifting

**Lemma 9.1.** *Let  $(\tilde{X}, p)$  be a covering space of a connected space  $X$ . Then the cardinality  $|p^{-1}(x)|$  is independent of  $x \in X$ .*

*Proof.* Fix  $x_0 \in X$  and let  $n = |p^{-1}(x_0)|$ . Let  $A = \{x \in X : |p^{-1}(x)| = n\}$ , and note that  $A \neq \emptyset$  since  $x_0 \in A$ . We will show that  $A$  is both open and closed, which implies that  $A = X$  by connectedness.

To see that  $A$  is open, let  $x \in A$ . By the definition of a covering space, there exists an open set  $U$  in  $X$  such that  $x \in U$  and  $p^{-1}(U) = \{U_1, \dots, U_n\}$ . Thus for any  $x' \in U$ , we have

$$p^{-1}(x') \subseteq p^{-1}(U) = \{U_1, \dots, U_n\},$$

and  $p^{-1}(x') \cap U_i = \{\text{pt}\}$  since  $p|_{U_i} : U_i \rightarrow U$  is a homeomorphism. Thus  $|p^{-1}(x')| = n$ , so  $x' \in A$ . This holds for each  $x' \in U$ , so  $U \subseteq A$ , i.e.  $A$  is an open set.

One can make a similar argument to check that if  $x \notin A$ , then there exists a open set  $U$  about  $x$  such that  $U \cap A = \emptyset$ . This shows that  $X \setminus A$  is open, i.e.  $A$  is closed, which completes the proof.  $\square$

**Definition 9.1.** We call  $|p^{-1}(x)|$  the *degree* of the covering space.

**Definition 9.2.** If  $(\tilde{X}, p)$  is a covering space for  $X$  and  $f : Y \rightarrow X$  is a continuous map, then a *lift* of  $f$  to  $\tilde{X}$  is a map  $\tilde{f} : Y \rightarrow \tilde{X}$  such that  $p \circ \tilde{f} = f$ , i.e. the following diagram commutes:

$$\begin{array}{ccc} & & \tilde{X} \\ & \nearrow \tilde{f} & \downarrow p \\ Y & \xrightarrow{f} & X \end{array}$$

If  $f(y_0) = x_0$  and  $\tilde{x}_0 \in \tilde{X}$  with  $p(\tilde{x}_0) = x_0$ , then  $\tilde{f}$  is a *lift of  $f$  based at  $\tilde{x}_0$*  if  $\tilde{f}$  is a lift and  $\tilde{f}(y_0) = \tilde{x}_0$ .

**Lemma 9.2.** Let  $(\tilde{X}, p)$  be a covering space of  $X$ ,  $x_0 \in X$ , and  $\tilde{x}_0 \in p^{-1}(x_0)$ . Then

- (a) each path  $\gamma : [0, 1] \rightarrow X$  based at  $x_0$  has a unique lift  $\tilde{\gamma} : [0, 1] \rightarrow \tilde{X}$  based at  $\tilde{x}_0$ .
- (b) if  $H : Y \times [0, 1] \rightarrow X$  is a homotopy with  $h_0(y) = H(y, 0)$  and  $\tilde{h}_0 : Y \rightarrow \tilde{X}$  a lift of  $h_0$ , then there is a unique lift  $\tilde{H} : Y \times [0, 1] \rightarrow \tilde{X}$  of  $H$  such that  $\tilde{H}(y, 0) = \tilde{h}_0(y)$ .

The above properties are called *path lifting* and *homotopy lifting*.

*Proof.* (a) The proof of this part is exactly the proof of part (a) in Lemma 5.5.

(b) The proof from Lemma 5.5 works if  $Y = [0, 1]$ . For the general case, see proof of Theorem 10.1.  $\square$

## 9.3 Connections to the Fundamental Group

**Lemma 9.3.** If  $(\tilde{X}, p)$  is a path connected covering space of  $X$  and  $x_0 \in X$ ,  $\tilde{x}_0 \in p^{-1}(x_0)$ , then the map  $p_* : \pi_1(\tilde{X}, \tilde{x}_0) \rightarrow \pi_1(X, x_0)$  satisfies the following:

- 1.  $p_*$  is injective;
- 2. the image of  $p_*$  is the set of loops in  $\pi_1(X, x_0)$  that when lifted are loops in  $\tilde{X}$  based at  $\tilde{x}_0$ ;
- 3. the index  $[\pi_1(X, x_0) : p_*(\pi_1(\tilde{X}, \tilde{x}_0))]$  is the degree of  $(\tilde{X}, p)$ .

*Proof.* (1) Suppose that  $p_*([\gamma]) = [e]$ , so there exists a homotopy  $H$  in  $X$  between  $x_0$  and  $p \circ \gamma$ . Note that  $\gamma$  is a lift of  $H(t, 0)$ , so Lemma 9.2 says  $H$  lifts to a homotopy  $\tilde{H}$  starting at  $\gamma$  in  $\tilde{X}$ . Note that  $H|_{\{0\} \times [0, 1]}$  is a constant loop and the loop  $t \mapsto \tilde{x}_0$  is a lift of  $H|_{\{0\} \times [0, 1]}$  based at  $\tilde{x}_0$ , so by uniqueness we see that  $\tilde{H}|_{\{0\} \times [0, 1]} = \tilde{x}_0$ . Similarly, we see that  $\tilde{H}|_{\{1\} \times [0, 1]} = \tilde{x}_0$  and  $\tilde{H}|_{[0, 1] \times \{1\}} = \tilde{x}_0$ . Thus  $\tilde{H}$  is a homotopy of loops based at  $\tilde{x}_0$  from  $\gamma$  to the constant loop, i.e.  $[\gamma] = [e_{x_0}]$ . So  $p_*$  is injective.

(2) Clearly if  $[\gamma] \in \pi_1(X, x_0)$  lifts to a loop  $\tilde{\gamma}$  based at  $\tilde{x}_0$ , then  $[\gamma] = p_*([\tilde{\gamma}])$ , so  $[\gamma]$  is in the image of  $p_*$ . Now if  $[\eta] = p_*([\gamma])$ , then  $\eta \sim p \circ \gamma$  in  $X$ . Let  $\tilde{\eta}$  be the lift of  $\eta$  based at  $\tilde{x}_0$ . By Lemma 9.2, the homotopy  $\eta \sim p \circ \gamma$  lifts to a homotopy  $\tilde{\eta} \sim \gamma$  rel endpoints. But  $\gamma$  is a loop, so  $\tilde{\eta}$  must be a loop.

(3) Let  $H = p_*(\pi_1(\tilde{X}, \tilde{x}_0)) \leq \pi_1(X, x_0)$ . If  $[\gamma] \in \pi_1(X, x_0)$  and  $[\delta] \in H$ , then note that by part (2),  $\delta$  lifts to a loop  $\tilde{\delta}$  based at  $\tilde{x}_0$ . Let  $\tilde{\delta} * \gamma$  be a lift of  $\delta * \gamma$  based at  $\tilde{x}_0$ , and note that  $\tilde{\gamma}(1) = \tilde{\delta} * \gamma(1) = \tilde{\delta} * \tilde{\gamma}(1)$ . This allows us to define

$$\phi : \{\text{right cosets of } H\} \rightarrow p^{-1}(x_0)$$

by  $H[\gamma] \mapsto \tilde{\gamma}(1)$ , which is well-defined by the above arguments.

If  $\tilde{x}_1 \in p^{-1}(x_0)$ , then let  $\tilde{\gamma}$  be a path in  $\tilde{X}$  from  $\tilde{x}_0$  to  $\tilde{x}_1$ . Let  $\gamma = p \circ \tilde{\gamma}$ , which is a loop in  $X$  based at  $x_0$ . Clearly  $\phi(H[\gamma]) = \tilde{\gamma}(1) = \tilde{x}_1$ , so  $\phi$  is onto. Now suppose that  $\phi(H[\gamma]) = \phi(H[\eta])$ . If  $\tilde{\gamma}, \tilde{\eta}$  are lifts of  $\gamma$  based at  $\tilde{x}_0$ , then  $\tilde{\gamma}(1) = \tilde{\eta}(1)$ . Thus  $\tilde{\gamma} * \tilde{\eta}^{-1}$  is a loop in  $\tilde{X}$ , so

$$p_*([\tilde{\gamma} * \tilde{\eta}^{-1}]) = [\gamma] * [\eta]^{-1} = [\gamma] * [\eta]^{-1} \in H.$$

This gives  $H[\gamma] = H[\eta]$ , so  $\phi$  is injective. Thus  $\phi$  is a bijection, so  $[\pi_1(X, x_0) : H] = |p^{-1}(x_0)|$ .  $\square$

**Example 9.2.1.** Recall the following examples from before:

1. For the covering map  $p : \mathbb{R} \rightarrow S^1$ , we have  $p_* : \pi_1(\mathbb{R}, 0) \rightarrow \pi_1(S^1, (1, 0))$  which sends  $e \mapsto 0$ , if we view  $p_*$  as  $p_* : \{e\} \rightarrow \mathbb{Z}$ . One can see all three properties of the above lemma in this example, in particular that  $[\pi_1(S^1) : \pi_1(\mathbb{R})] = \infty = |p^{-1}((1, 0))|$ .
2. Let  $p_n : S^1 \rightarrow S^1$  send  $\theta \mapsto n\theta$ . Then  $(p_n)_* : \pi_1(S^1, (1, 0)) \rightarrow \pi_1(S^1, (1, 0))$ , which we can view as a map  $(p_n)_* : \mathbb{Z} \rightarrow \mathbb{Z}$ , or  $(p_n)_* : \langle g \rangle \rightarrow \langle h \rangle$ . Then  $(p_n)_*$  maps  $g \mapsto h^n$ , so the subgroup of  $\mathbb{Z}$  corresponding to this covering space is  $n\mathbb{Z}$ . Again we can see all three properties of the above lemma, in particular that we have  $[\mathbb{Z} : n\mathbb{Z}] = n = |p_n^{-1}((1, 0))|$ .

**Exercise 9.1.** Check the properties of the above lemma explicitly for Example 9.0.1(4). Note that picking different base points can yield different images of  $p_*$ .



# Lecture 10

## Feb. 10 — Covering Spaces, Part 2

### 10.1 More Connections to the Fundamental Group

**Lemma 10.1.** *If  $(\tilde{X}, p)$  is a path connected covering space of  $X$  and  $x_0 \in X$ , then*

$$\{p_*(\pi_1(\tilde{X}, \tilde{x}))\}_{\tilde{x} \in p^{-1}(x_0)}$$

*is a conjugacy class of subgroups of  $\pi_1(X, x_0)$ .*

*Proof.* Let  $\tilde{x}_0, \tilde{x}_1 \in p^{-1}(x_0)$  and  $H_i = p_*(\pi_1(\tilde{X}, \tilde{x}_i))$ . Since  $\tilde{X}$  is path connected, we can find a path  $h$  in  $\tilde{X}$  from  $\tilde{x}_0$  to  $\tilde{x}_1$ . Clearly  $\gamma = p \circ h$  is a loop in  $X$  based at  $x_0$ . If  $[\eta] \in H_1$ , then by Lemma 9.3  $\eta$  lifts to a loop  $\tilde{\eta}$  based at  $\tilde{x}_1$ . Note that  $h * \tilde{\eta} * \bar{h}$  is a loop based at  $\tilde{x}_0$  in  $\tilde{X}$ . Then

$$[\gamma] \cdot [\eta] \cdot [\gamma]^{-1} = [(p \circ h) * (p \circ \tilde{\eta}) * (p \circ \bar{h})] = p_*([h * \tilde{\eta} * \bar{h}]) \in H_0.$$

This says that  $[\gamma]H_1[\gamma]^{-1} \subseteq H_0$ . The same argument says  $[\gamma]^{-1}H_0[\gamma] \subseteq H_1$ , so

$$H_0 = [\gamma][\gamma]^{-1}H_0[\gamma][\gamma]^{-1} \subseteq [\gamma]H_1[\gamma]^{-1}.$$

This implies that  $H_0 = [\gamma]H_1[\gamma]^{-1}$ , so  $H_0, H_1$  are conjugate.

We still need to show we get the full conjugacy class. For this, suppose  $H \leq \pi_1(X, x_0)$  such that there exists  $[\alpha] \in \pi_1(X, x_0)$  with  $[\alpha]H[\alpha]^{-1} = H_0$ . If  $[\alpha] \in H_0$ , then  $H = H_0$  and we are done. If  $[\alpha] \notin H_0$ , then  $\alpha$  lifts to a path  $\tilde{\alpha}$  (not necessarily a loop) based at  $\tilde{x}_0$ . Let  $\tilde{x}_2 = \tilde{\alpha}(1)$ , and set  $H_2 = p_*(\pi_1(\tilde{X}, \tilde{x}_2))$ . From above, we see that  $H = H_2$ , which completes the proof.  $\square$

**Definition 10.1.** A space  $X$  is *locally path connected* if for every  $y \in Y$  and open set  $U$  containing  $y$ , there exists an open set  $V$  such that  $y \in V \subseteq U$  and  $V$  is path connected.

**Example 10.1.1.** An example of a space which is path connected but not locally path connected is

$$C = (\{1/n\} \times [0, 1])_{n \in \mathbb{N}} \times (\{0\} \times [0, 1]) \cup ([0, 1] \times \{0\}),$$

also known as the *comb space*. We can see that  $C$  is path connected, but it is not locally path connected, e.g. take  $y = (0, 1)$  and  $U$  to be any small enough open neighborhood containing  $y$ .

**Theorem 10.1.** *Let  $(\tilde{X}, p)$  be a covering space of  $X$ . Let  $x_0 \in X$  and  $\tilde{x}_0 \in p^{-1}(x_0)$ . Suppose  $f : Y \rightarrow X$  is any map with  $Y$  path connected and locally path connected, with  $y_0 \in Y$  such that  $f(y_0) = x_0$ . Then there exists a lift  $\tilde{f} : Y \rightarrow \tilde{X}$  of  $f$  such that  $\tilde{f}(y_0) = \tilde{x}_0$  if and only if  $f_*(\pi_1(Y, y_0)) \subseteq p_*(\pi_1(\tilde{X}, \tilde{x}_0))$ .*

*Proof.* ( $\Rightarrow$ ) If  $\tilde{f} : Y \rightarrow \tilde{X}$  exists, then  $p \circ \tilde{f} = f$ , and so

$$f_*(\pi_1(Y, y_0)) = (p \circ \tilde{f})_*(\pi_1(Y, y_0)) = p_*(\tilde{f}_*(\pi_1(Y, y_0))) \subseteq p_*(\pi_1(\tilde{X}, \tilde{x}_0))$$

since  $f_*(\pi_1(Y, y_0)) \subseteq \pi_1(X, x_0)$ . This proves the first direction.

( $\Leftarrow$ ) Suppose that  $f_*(\pi_1(Y, y_0)) \subseteq p_*(\pi_1(\tilde{X}, \tilde{x}_0))$ . Since  $Y$  is path connected, for any  $y \in Y$ , there exists a path  $\gamma : [0, 1] \rightarrow Y$  from  $y_0$  to  $y$ . Then there is a unique lift  $\widetilde{f \circ \gamma} : [0, 1] \rightarrow \tilde{X}$  based at  $\tilde{x}_0$ . Define

$$\tilde{f}(y) = \widetilde{f \circ \gamma}(1).$$

Note that if  $\tilde{f}$  is well-defined, then it is clear that  $p \circ \tilde{f} = f$ .

To see that  $\tilde{f}$  is well-defined, we must show that  $\widetilde{f \circ \gamma_1}(1)$  is independent of the choice of path  $\gamma$ . Let  $\gamma, \eta$  be paths from  $y_0$  to  $y$ . Note that  $\gamma * \bar{\eta}$  is a loop in  $Y$  based at  $y_0$ , so

$$(f \circ \gamma) * (f \circ \bar{\eta}) = f_*(\gamma * \bar{\eta}) \in p_*(\pi_1(\tilde{X}, \tilde{x}_0)).$$

Thus by Lemma 9.3, this loop lifts to a loop in  $\tilde{X}$  based at  $\tilde{x}_0$ . Then

$$(f \circ \gamma) * (f \circ \bar{\eta}) = \widetilde{(f \circ \gamma) * (f \circ \bar{\eta})},$$

where  $\widetilde{f \circ \gamma}$  is a lift based at  $\tilde{x}_0$  and  $\widetilde{f \circ \bar{\eta}}$  is a lift based at  $\widetilde{f \circ \gamma}(1)$ . But  $(f \circ \gamma) * (f \circ \bar{\eta})$  is a loop, so

$$\widetilde{f \circ \bar{\eta}}(1) = \tilde{x}_0.$$

Then  $\widetilde{f \circ \bar{\eta}}$  is the lift of  $\eta$  based at  $\tilde{x}_0$ , so  $\widetilde{f \circ \bar{\eta}}(1) = \widetilde{f \circ \bar{\eta}}(0) = \widetilde{f \circ \gamma}(1)$ . Thus,  $\widetilde{f \circ \eta}(1) = \widetilde{f \circ \gamma}(1)$ .

Now it just remains to show that  $\tilde{f}$  is continuous. To see this, take any open set  $U \subseteq \tilde{X}$ , and we will show that for all  $y \in \tilde{f}^{-1}(U)$ , there exists an open set  $V$  in  $Y$  such that  $y \in V \subseteq \tilde{f}^{-1}(U)$  (this implies that  $\tilde{f}^{-1}(U)$  is open). Let  $W$  be an evenly covered open set containing  $f(y)$ , and  $\tilde{W} \subseteq p^{-1}(W)$  such that  $p|_{\tilde{W}} : \tilde{W} \rightarrow W$  is a homeomorphism (by possibly shrinking  $W$ , e.g. by intersecting  $\tilde{W}$  with  $U$  and taking the image under  $p$ , we can assume that  $\tilde{W} \subseteq U$ ). Since  $Y$  is locally path connected, there exists a path connected open set  $V$  in  $Y$  containing  $y$  and  $V \subseteq f^{-1}(W)$ .

Fix a path  $\gamma$  from  $y_0$  to  $y$ . For any point  $y' \in V$ , there exists a path  $\eta$  from  $y$  to  $y'$  in  $V$ . By definition,

$$\tilde{f}(y') = \widetilde{f \circ (\gamma * \eta)}(1).$$

But if  $\widetilde{f \circ \eta}$  is a lift of  $f \circ \eta$  based at  $\widetilde{f \circ \gamma}(1) = \tilde{f}(y)$ , then

$$f \circ (\gamma * \eta)(1) = \widetilde{f \circ \eta}(1).$$

We know that  $\widetilde{f \circ \eta} = (p|_{\tilde{W}})^{-1} \circ f \circ \eta$ , so we see that

$$\tilde{f}(y') = \widetilde{f \circ \eta}(1) \in \tilde{W} \subseteq U.$$

This shows that  $V \subseteq \tilde{f}^{-1}(U)$ , so  $\tilde{f}$  is continuous, which completes the proof.  $\square$

**Remark.** It is fairly common in algebraic topology for a topological hypothesis to imply some type of algebraic conclusion. The opposite, however, as in the above theorem, is much rarer.

**Lemma 10.2.** *Let  $(\tilde{X}, p)$  be a covering space of  $X$ , and let  $\tilde{f}_1, \tilde{f}_2 : Y \rightarrow \tilde{X}$  be two lifts of  $f : Y \rightarrow X$ . If  $Y$  is connected and  $\tilde{f}_1$  and  $\tilde{f}_2$  agree at one point, then  $\tilde{f}_1 = \tilde{f}_2$ .*

*Proof.* Let  $A = \{y \in Y : \tilde{f}_1(y) = \tilde{f}_2(y)\}$ . Note that  $A \neq \emptyset$  by assumption. If  $y \in A$ , then let  $U$  be an evenly covered neighborhood of  $f(y)$  and  $\tilde{U} \subseteq p^{-1}(U)$  such that  $\tilde{f}_1(y) = \tilde{f}_2(y) \in \tilde{U}$  and  $p|_{\tilde{U}} : \tilde{U} \rightarrow U$  is a homeomorphism. Since  $f$  is continuous, there exists an open neighborhood  $V$  of  $y$  in  $Y$  such that  $f(V) \subseteq U$ . Now we have

$$\tilde{f}_1|_V = (p|_{\tilde{U}})^{-1} \circ f|_V = \tilde{f}_2|_V,$$

so  $V \subseteq A$ , which shows that  $A$  is open. A similar argument shows that  $Y \setminus A$  is open, so  $A$  is closed. Since  $Y$  is connected and  $A$  is open, closed, nonempty, we must have  $A = Y$ , hence  $\tilde{f}_1 = \tilde{f}_2$ .  $\square$

## 10.2 Covering Space Isomorphisms

**Definition 10.2.** We say that two covering spaces  $(\tilde{X}_1, p_1)$  and  $(\tilde{X}_2, p_2)$  of  $X$  are *isomorphic* if there exists a homeomorphism  $h : \tilde{X}_1 \rightarrow \tilde{X}_2$  such that  $p_2 \circ h = p_1$ , i.e. the following diagram commutes:

$$\begin{array}{ccc} \tilde{X}_1 & \xrightarrow{\quad h \quad} & \tilde{X}_2 \\ & \searrow p_1 & \downarrow p_2 \\ & & X \end{array}$$

**Corollary 10.1.1.** *Suppose  $(\tilde{X}_1, p_1)$  and  $(\tilde{X}_2, p_2)$  are path connected, locally path connected covering spaces of  $X$  and  $x_0 \in X$ ,  $\tilde{x}_i \in p_i^{-1}(x_0)$ , then*

- (a) *if  $(p_1)_*(\pi_1(\tilde{X}_1, \tilde{x}_1)) \subseteq (p_2)_*(\pi_1(\tilde{X}_2, \tilde{x}_2))$ , then  $p_1$  lifts to a covering map  $p : \tilde{X}_1 \rightarrow \tilde{X}_2$  which takes the base point  $\tilde{x}_1$  to the base point  $\tilde{x}_2$ ;*
- (b)  *$(\tilde{X}_1, \tilde{x}_1)$  and  $(\tilde{X}_2, \tilde{x}_2)$  are isomorphic covering spaces taking  $\tilde{x}_1$  to  $\tilde{x}_2$  if and only if*

$$(p_1)_*(\pi_1(\tilde{X}_1, \tilde{x}_1)) = (p_2)_*(\pi_1(\tilde{X}_2, \tilde{x}_2));$$

- (c)  *$(\tilde{X}_1, \tilde{x}_1)$  and  $(\tilde{X}_2, \tilde{x}_2)$  are isomorphic covering spaces of  $X$  if and only if  $(p_1)_*(\pi_1(\tilde{X}_1, \tilde{x}_1))$  is conjugate to  $(p_2)_*(\pi_1(\tilde{X}_2, \tilde{x}_2))$ .*

*Proof.* (a) By Theorem 10.1, we get a lift  $p : \tilde{X}_1 \rightarrow \tilde{X}_2$  of  $p_1$  taking  $\tilde{x}_1$  to  $\tilde{x}_2$ . We need to show that  $p$  is a covering map. Let  $x \in \tilde{X}_2$ , and we show that  $x$  has an evenly covered neighborhood. Let  $U$  be a neighborhood of  $p_2(x)$  in  $X$  which is evenly covered by both  $p_1$  and  $p_2$  (e.g. take the intersection of the two neighborhoods evenly covered by  $p_1$  and  $p_2$ ), so there exists a unique  $\tilde{U} \subseteq \tilde{X}_2$  such that  $x \in \tilde{U}$  and  $p_2|_{\tilde{U}} : \tilde{U} \rightarrow U$  is a homeomorphism. Write  $p^{-1}(\tilde{U}) = \bigcup_{\alpha} U_{\alpha}$ . Clearly  $\bigcup_{\alpha} U_{\alpha} \subseteq p^{-1}(U)$ , so  $p_1|_{U_{\alpha}} : U_{\alpha} \rightarrow U$  is a homeomorphism. Thus  $p|_{U_{\alpha}} : p_2^{-1}|_{\tilde{U}} \circ p_1|_{U_{\alpha}}$  is a homeomorphism  $U_{\alpha} \rightarrow \tilde{U}$ , so  $\tilde{U}$  is evenly covered.

(b)  $(\Rightarrow)$  This is clear.  $(\Leftarrow)$  Let  $\tilde{p}_1 : \tilde{X}_1 \rightarrow \tilde{X}_2$  and  $\tilde{p}_2 : \tilde{X}_2 \rightarrow \tilde{X}_1$  be the lifts from part (a). We have:

$$\begin{array}{ccccc} \tilde{X}_2 & \xrightarrow{\tilde{p}_2} & \tilde{X}_1 & \xrightarrow{\tilde{p}_1} & \tilde{X}_2 \\ & \searrow p_2 & \downarrow p_1 & \swarrow p_2 & \\ & & X & & \end{array}$$

Note that  $\tilde{p}_1 \circ \tilde{p}_2 : \tilde{X}_2 \rightarrow \tilde{X}_2$  takes  $\tilde{x}_2$  to  $\tilde{x}_2$  and is a lift of  $p_2$  to  $\tilde{X}_2$ . But  $\text{id}_{\tilde{X}_2}$  is also a lift of  $p_2$  and agrees with  $\tilde{p}_1 \circ \tilde{p}_2$  on  $\tilde{x}_2$ , so by Lemma 10.2,  $\tilde{p}_1 \circ \tilde{p}_2 = \text{id}_{\tilde{X}_2}$ . Similarly, we can show that  $\tilde{p}_2 \circ \tilde{p}_1 = \text{id}_{\tilde{X}_1}$ , so  $\tilde{p}_2$  is a homeomorphism and  $\tilde{X}_1, \tilde{X}_2$  are isomorphic.

(c) This follows from Lemma 10.1 and part (b). □

# Lecture 11

## Feb. 12 — Covering Spaces, Part 3

### 11.1 Galois Correspondence

**Definition 11.1.** A space  $X$  is *semi-locally simply connected* if each point  $x \in X$  has a neighborhood  $U$  such that  $\pi_1(U, x) \rightarrow \pi_1(X, x)$  is trivial.

**Example 11.1.1.** Let  $X = \bigcup_{n=1}^{\infty} S_{1/n}(1/n, 0)$ , where  $S_{1/n}(1/n, 0)$  is a circle of radius  $1/n$  about the point  $(1/n, 0)$ . Note that any neighborhood of  $(0, 0)$  has non-trivial fundamental group that injects into  $\pi_1(X, (0, 0))$ . So  $X$  is not semi-locally simply connected. On the other hand, let  $X'$  be the cone on  $X$ :

$$X' = (X \times [0, 1]) / (X \times \{0\}).$$

Then  $X'$  is semi-locally simply connected but not locally simply connected.

**Remark.** All CW-complexes and manifolds are semi-locally simply connected.

**Theorem 11.1.** Let  $X$  be a path connected, locally path connected, semi-locally simply connected space, and fix  $x_0 \in X$ . Then there is a one-to-one correspondence

$$\left\{ \begin{array}{c} \text{basepoint preserving isomorphism classes} \\ \text{of covering spaces } (\tilde{X}, p, \tilde{x}_0) \end{array} \right\} \longleftrightarrow \{ \text{subgroups of } \pi_1(X, x_0) \}$$

given by  $(\tilde{X}, p, \tilde{x}_0) \mapsto p_*(\pi_1(\tilde{X}, \tilde{x}_0))$  with inverse  $H \mapsto (\tilde{X}_H, p_H, \tilde{x}_0)$ , which satisfies:

1. If  $H \leq K$ , then  $(\tilde{X}_H, p_H, \tilde{x}_H)$  is also a covering space of  $(\tilde{X}_K, p_K, \tilde{x}_K)$ .
2. If  $p_1$  in  $(\tilde{X}_1, p_1, \tilde{x}_1)$  lifts to a cover of  $(\tilde{X}_2, p_2, \tilde{x}_2)$ , then  $p_{1*}(\pi_1(\tilde{X}_1, \tilde{x}_1)) \leq p_{2*}(\pi_1(\tilde{X}_2, \tilde{x}_2))$ .
3.  $[\pi_1(X, x_0) : H] = n$  if and only if  $(\tilde{X}_H, p_H, \tilde{x}_H)$  is degree  $n$ .

In addition, there is a one-to-one correspondence

$$\{ \text{covering spaces } (\tilde{X}, p) \text{ of } X \} \longleftrightarrow \{ \text{conjugacy classes of subgroups of } \pi_1(X, x_0) \}.$$

*Proof.* Note that (1) and (2) follow from Corollary 10.1.1 once we have the one-to-one correspondence, and (3) follows from Lemma 9.3. Also, once we have a well-defined map  $H \mapsto (\tilde{X}_H, p_H, \tilde{x}_H)$  such that  $(p_H)_*(\pi_1(\tilde{X}_H, \tilde{x}_H)) = H$ , then the first one-to-one correspondence follows from Corollary 10.1.1(b). The second one-to-one correspondence follows from Corollary 10.1.1(c).

So we are left to construct  $(\tilde{X}_H, p_H, \tilde{x}_H)$  given  $H \leq \pi_1(X, x_0)$ . Given  $H$ , we define an equivalence relation on paths based at  $x_0$ : Call two based paths  $\gamma, \eta : [0, 1] \rightarrow X$  (i.e.  $\gamma(0) = \eta(0) = x_0$ )  $H$ -equivalent if:

1.  $\gamma(1) = \eta(1)$ , and
2.  $[\gamma * \bar{\eta}] \in H$ .

Denote  $H$ -equivalence by  $\gamma \sim_H \eta$  and its equivalence classes by  $\langle \gamma \rangle$ . Show as an exercise that this is an equivalence relation, and if  $H = \{e\}$ , then  $\gamma \sim_H \eta$  if and only if  $\gamma, \eta$  are homotopic rel endpoints. Let

1.  $\tilde{X}_H = \{\langle \gamma \rangle : \gamma \text{ is a path in } X \text{ based at } x_0\}$ .

Note that this is a set: We will need to put a topology on  $\tilde{X}_H$  later.

2.  $p_H : \tilde{X}_H \rightarrow X : \langle \gamma \rangle \mapsto \gamma(1)$ .
3.  $\tilde{x}_H = \langle e_{x_0} \rangle$ , where  $e_{x_0}$  is the constant path  $x \mapsto x_0$ . Clearly  $p_H(\tilde{x}_H) = x_0$ .

Note that  $p_H$  is onto, since  $X$  is path connected. In order to put a topology on  $\tilde{X}_H$ , we need to study the topology of  $X$  in terms of paths. We claim that

$$\mathcal{U} = \left\{ U \subseteq X : \begin{array}{l} U \text{ is open, path connected, and} \\ \pi_1(U, x) \rightarrow \pi_1(X, x) \text{ is trivial for some } x \in U \end{array} \right\}$$

is a basis for the topology on  $X$ .<sup>1</sup> To see this, note that if  $\pi_1(U, x) \rightarrow \pi_1(X, x)$  is trivial for some  $x \in U$ , then this is true for all  $y \in U$  since  $U$  is path-connected:

$$\begin{array}{ccc} \pi_1(U, x) & \xrightarrow{\text{triv}} & \pi_1(X, x) \\ \downarrow \phi_h & & \uparrow \phi_h \\ \pi_1(U, y) & \xrightarrow{i_*} & \pi_1(X, y) \end{array}$$

Also if  $U \in \mathcal{U}$  and  $V \subseteq U$  is open and path connected, then:

$$\begin{array}{ccccc} \pi_1(V, x) & \longrightarrow & \pi_1(U, x) & \xrightarrow{\text{triv}} & \pi_1(X, x) \\ & \searrow \text{triv} & & & \end{array}$$

so  $V \in \mathcal{U}$  too. Now let  $x \in X$  and  $U$  be any open set in  $X$  containing  $x$ . Since  $X$  is semi-locally simply connected, there exists an open set  $V$  containing  $x$  such that  $\pi_1(V, x) \rightarrow \pi_1(X, x)$  is trivial. Then  $U \cap V$  is an open set containing  $x$  and  $\pi_1(U \cap V, x) \rightarrow \pi_1(X, x)$  is trivial. Now  $X$  being locally path connected implies that there exists an open set  $W$  such that  $x \in W \subseteq U \cap V$  and  $W$  is path connected. Clearly  $\pi_1(W, x) \rightarrow \pi_1(X, x)$  is trivial, so  $W \in \mathcal{U}$ . Thus  $\mathcal{U}$  is a basis for the topology on  $X$ .

Now for each  $U \in \mathcal{U}$  and a path  $\gamma$  from  $x_0$  to a point in  $U$ , set

$$U_\gamma = \{\langle \gamma * \eta \rangle : \eta \text{ a path in } U \text{ such that } \eta(0) = \gamma(1)\}.$$

Note that  $U_\gamma$  is a subset of  $\tilde{X}_H$ . We claim that  $\{U_\gamma\}_{U \in \mathcal{U}, \gamma \text{ a path from } x_0 \text{ to } x \in U}$  forms a basis for a topology on  $\tilde{X}_H$ .<sup>2</sup> To see this, note the following:

<sup>1</sup>Recall that a collection of open sets in  $X$  is a *basis for the topology on  $X$*  if for every  $x \in X$  and open set  $U$  containing  $x$ , there exists a set  $O$  in the collection such that  $x \in O \subseteq U$ .

<sup>2</sup>Recall that a collection of subsets of a set  $X$  form a *basis for a topology on  $X$*  if for each pair of sets  $U, V$  in the collection and  $x \in U \cap V$ , there is another set  $W$  in the collection such that  $x \in W \subseteq U \cap V$ , and  $X$  is the union of the sets in the collection.

1. If  $\langle \gamma \rangle = \langle \delta \rangle$ , then  $U_\gamma = U_\delta$  (so we can write  $U_{\langle \gamma \rangle}$ ).

This is because if  $\gamma \sim_H \delta$ , then  $\gamma * \eta \sim_H \delta * \eta$  for all paths  $\eta$  in  $U$  with  $\eta(0) = \eta(1)$ :

$$\delta * \eta * \overline{(\gamma * \eta)} = \gamma * \eta * \bar{\eta} * \bar{\delta} \sim \gamma * \bar{\delta}.$$

2.  $p_H : U_{\langle \gamma \rangle} \rightarrow U$  is onto since  $U$  is path connected.

3.  $p_H : U_{\langle \gamma \rangle} \rightarrow U$  is injective.

This is because if  $p_H(\langle \gamma * \eta \rangle) = p_H(\langle \gamma * \eta' \rangle)$ , then  $\gamma * \eta(1) = \gamma * \eta'(1)$ , so  $\eta * \bar{\eta}'$  is a loop in  $U$  based at  $x = \gamma(1)$ . So  $\eta * \bar{\eta}'$  is homotopic to a constant loop in  $X$ . Thus  $\gamma * \eta \sim \gamma * \eta'$  rel endpoints, so

$$[(\langle \gamma * \eta \rangle * \overline{(\gamma * \eta')})] = [e_{x_0}] \in H.$$

This gives  $\langle \gamma * \eta \rangle = \langle \gamma * \eta' \rangle$ , i.e.  $p_H$  is injective.

4. If  $\langle \gamma' \rangle \in U_{\langle \gamma \rangle}$ , then  $U_{\langle \gamma' \rangle} = U_{\langle \gamma \rangle}$ .

This is because by hypothesis, there exists a path  $\eta$  in  $U$  such that  $\langle \gamma' \rangle = \langle \gamma * \eta \rangle$ , so we can take  $\gamma * \eta$  to represent  $\gamma'$  by note (1). So if  $\langle \delta \rangle \in U_{\langle \gamma' \rangle}$ , then

$$\delta = (\gamma * \eta) * \eta' \sim \gamma * (\eta * \eta'),$$

so  $\langle \delta \rangle \in U_{\langle \gamma \rangle}$ . This shows that  $U_{\langle \gamma' \rangle} \subseteq U_{\langle \gamma \rangle}$ , and similarly one can prove that  $U_{\langle \gamma \rangle} \subseteq U_{\langle \gamma' \rangle}$ .

Now if  $\langle \delta \rangle \in U_{\langle \gamma \rangle} \cap V_{\langle \gamma' \rangle}$ , then  $U_{\langle \gamma \rangle} = U_{\langle \delta \rangle}$  and  $V_{\langle \gamma' \rangle} = V_{\langle \delta \rangle}$ . by (4). So if  $W$  is any element of  $\mathcal{U}$  such that  $\delta(1) \in W \subseteq U \cap V$ , then

$$\langle \delta \rangle = W_{\langle \delta \rangle} \subseteq U_{\langle \delta \rangle} \cap V_{\langle \delta \rangle} = U_\gamma \cap U_{\gamma'}.$$

Clearly  $\tilde{X}_H$  is the union of the sets in  $\{U_\gamma\}$ , so we have a basis for a basis on  $\tilde{X}_H$ .

We claim that with this topology,  $(\tilde{X}_H, p_H)$  is a covering space of  $(X, x_0)$ . For any  $U \in \mathcal{U}$  and a path  $\gamma$  from  $x_0$  to a point in  $U$ ,  $p|_{U_{\langle \gamma \rangle}} : U_{\langle \gamma \rangle} \rightarrow U$  is a homeomorphism. To see this, first note  $p_H$  is a bijection. Next,  $p_H|_{U_{\langle \gamma \rangle}}$  is continuous since for any basic open set  $V \subseteq U$  and path  $\delta$  from  $x_0$  to a point in  $V$ ,

$$(p_H|_{U_{\langle \gamma \rangle}})^{-1}(V) = V_{\langle \delta \rangle}$$

which is open. This also shows that the image of a basic open set  $V_{\langle \delta \rangle}$  is an open set  $V$ , so  $p_H|_{U_{\langle \gamma \rangle}}$  is a homeomorphism. Note that this implies  $p_H : \tilde{X}_H \rightarrow X$  is continuous. Now if  $x \in X$  is any point, let  $U \in \mathcal{U}$  be an open set containing  $x$ . Then we have

$$p^{-1}(U) = \bigcup_{\substack{\gamma \text{ a path from } x_0 \\ \text{to a point in } U}} U_{\langle \gamma \rangle},$$

and  $p_H|_{U_{\langle \gamma \rangle}} : U_{\langle \gamma \rangle} \rightarrow U$  is a homeomorphism, so  $U$  is evenly covered, i.e.  $(\tilde{X}_H, p_H)$  is a covering space.

It only remains to show  $(p_H)_*(\pi_1(\tilde{X}_H, \tilde{x}_H)) = H$  in  $\pi_1(X, x_0)$ . If  $[\gamma] \in H$ , let  $\gamma_t(s)$  be the path

$$\gamma_t : [0, 1] \rightarrow X : s \mapsto \gamma(ts).$$

Note that  $\tilde{\gamma} : [0, 1] \rightarrow \tilde{X}_H$  given by  $t \mapsto \langle \gamma_t \rangle$  is a loop in  $\tilde{X}_H$  since  $\tilde{\gamma}_0 = \gamma_0$  (the constant path in  $\tilde{X}_H$ ) and  $\tilde{\gamma}(1) = \langle \gamma \rangle = \langle e_{x_0} \rangle$  since  $[\gamma] \in H$ . We also have  $(p_H \circ \tilde{\gamma})(t) = \gamma_t(1) = \gamma(t)$ , so  $p_H \circ \tilde{\gamma} = \gamma$ . So  $\tilde{\gamma}$  is a lift of  $\gamma$  and a loop, therefore  $[\gamma] \in \text{im}((p_H)_*)$  by Lemma 9.3(b). Now if  $\gamma \notin H$ , then the path  $\tilde{\gamma}$  is not a loop since  $\tilde{\gamma}(1) = \langle \gamma \rangle \notin H$ , so  $\tilde{\gamma}(1) \neq \langle e_{x_0} \rangle$ . So  $[\gamma] \notin \text{im}((p_H)_*)$  by Lemma 9.3(b). These two properties imply that  $(p_H)_*(\pi_1(\tilde{X}_H, \tilde{x}_H)) = H$ , which completes the proof of the theorem.  $\square$

**Remark.** There is a “lattice” of subgroups of  $\pi_1(X, x_0)$  and a “lattice” of covering spaces of  $X$ . The above theorem says that these are the same (similar as in Galois theory).

**Remark.** By the above correspondence, there is a unique covering space of  $X$  corresponding to the trivial subgroup  $\{e\} \subseteq \pi_1(X, x_0)$ . This is called the *universal cover* of  $X$ .



# Lecture 12

## Feb. 17 — Covering Spaces, Part 4

### 12.1 Deck Transformations

**Definition 12.1.** Let  $p : \tilde{X} \rightarrow X$  be a covering space. A *deck transformation*, or a *covering transformation*, is a covering space isomorphism, i.e. a homeomorphism  $f : \tilde{X} \rightarrow \tilde{X}$  such that  $p \circ f = p$ . The set  $G(\tilde{X})$  of deck transformations is clearly a group under composition.

**Example 12.1.1.** Consider the covering map  $p_4 : S^1 \rightarrow S^1$  given by  $\theta \mapsto 4\theta$ . Then each  $\phi_k : S^1 \rightarrow S^1$  given by  $\theta \mapsto \theta + (2\pi k)/4$  satisfies  $p_4 \circ \phi_k = p_4$ , so the  $\phi_k$  are deck transformations. Now suppose  $f : S^1 \rightarrow S^1$  is another deck transformation, so  $f(\tilde{x}_1) = \tilde{x}_i$  for some  $i$  (let the  $\tilde{x}_i$  be the four preimages of  $\tilde{x}_1$  under  $p_4$ ). But there exists  $k$  such that  $\phi_k(\tilde{x}_1) = \tilde{x}_i$ , and these are both lifts of  $p$  which agree at  $\tilde{x}_1$ , so  $f = \phi_k$  by Lemma 10.2. Thus  $G(S^1 \xrightarrow{p_4} S^1) = \{\phi_0, \phi_1, \phi_2, \phi_3\} \cong \mathbb{Z}/4\mathbb{Z}$  (a generator is  $\phi_1$ ).

**Example 12.1.2.** Let  $p : \tilde{X} \rightarrow X$  be as in Example 9.0.1(5). Let  $\tilde{x}_1$  be the center intersection point in  $\tilde{X}$  and  $\tilde{x}_2, \tilde{x}_3$  be the two intersection points at the sides. Let  $x_0$  be the intersection point in  $X$ . Note that if we lift  $a$  to  $\tilde{X}$  based at  $\tilde{x}_1$ , then we get a path, but if we lift  $a$  to  $\tilde{X}$  based at  $\tilde{x}_3$ , then we get a loop. So if  $\phi : \tilde{X} \rightarrow \tilde{X}$  is a deck transformation, then it cannot take  $\tilde{x}_1$  to  $\tilde{x}_3$ , since  $\phi$  applied to a lift is a lift. Similarly, there are no deck transformations taking  $\tilde{x}_1$  to  $\tilde{x}_2$ . So any deck transformation takes  $\tilde{x}_1$  to  $\tilde{x}_1$ . Thus Lemma 10.2 says that any deck transformation is the identity, i.e.  $G(\tilde{X}) = \{1\}$ .

**Remark.** In the first example, a covering map of degree 4 gave a group of order 4, whereas in the second example, a covering map of degree 3 gave a group of order 1.

**Definition 12.2.** A covering space  $p : \tilde{X} \rightarrow X$  is called *normal* if for every  $x \in X$  and  $\tilde{x}, \tilde{x}' \in p^{-1}(x)$ , there exists  $\phi \in G(\tilde{X})$  such that  $\phi(\tilde{x}) = \tilde{x}'$ .

**Remark.** The first example is normal, while the second one is not.

**Theorem 12.1.** Let  $p : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$  be a path connected, locally path connected covering space of  $X$ . Let  $H = p_*(\pi_1(\tilde{X}, \tilde{x}_0)) \leq \pi_1(X, x_0)$ . Then

1.  $(\tilde{X}, p)$  is normal if and only if  $H$  is a normal subgroup of  $\pi_1(X, x_0)$ ;
2.  $G(\tilde{X}) \cong N(H)/H$ , where  $N(H)$  is the normalizer of  $H$ , i.e. the largest subgroup of  $\pi_1(X, x_0)$  in which  $H$  is normal.

*Proof.* (1)  $(\Rightarrow)$  Let  $[\gamma] \in \pi_1(X, x_0)$  and let  $\tilde{\gamma}$  be a lift of  $\gamma$  based at  $\tilde{x}_0$ . Set  $\tilde{x}_1 = \tilde{\gamma}(1)$ . From the proof of Lemma 10.1, we have

$$[\gamma]p_*(\pi_1(\tilde{X}, \tilde{x}_0))[\gamma]^{-1} = p_*(\pi_1(\tilde{X}, \tilde{x}_1)).$$

Now  $\tilde{X}$  normal implies that there exists  $\phi \in G(\tilde{X})$  such that  $\phi_1(\tilde{x}_0) = \tilde{x}_1$ , so we get an isomorphism

$$\phi_* : \pi_1(\tilde{X}, \tilde{x}_1) \rightarrow \pi_1(\tilde{X}, \tilde{x}_0).$$

Then since  $p_* \circ \phi_*$  (as  $\phi$  is a deck transformation), we have

$$p_*(\pi_1(\tilde{X}, \tilde{x}_0)) = p_* \circ \phi_*(\pi_1(\tilde{X}, \tilde{x}_1)) = p_*(\pi_1(\tilde{X}, \tilde{x}_1)),$$

so that  $[\gamma]p_*(\pi_1(\tilde{X}, \tilde{x}_0))[\gamma]^{-1} = p_*(\pi_1(\tilde{X}, \tilde{x}_0))$ , i.e.  $p_*(\pi_1(\tilde{X}, \tilde{x}_0))$  is normal.

( $\Leftarrow$ ) Let  $\tilde{x}_0, \tilde{x}_1$  be two points in  $p^{-1}(x_0)$ , and set  $H_1 = p_*(\pi_1(\tilde{X}, \tilde{x}_1))$ . Let  $h$  be a path from  $\tilde{x}_0$  to  $\tilde{x}_1$ , and let  $\gamma = p \circ h$ . By Lemma 10.1, we have  $[\gamma]H_1[\gamma]^{-1} = H$ . Since  $H$  is normal, we have  $H = H_1$ , i.e.

$$p_*(\pi_1(\tilde{X}, \tilde{x}_1)) = p_*(\pi_1(\tilde{X}, \tilde{x}_0)).$$

Now by Theorem 10.1, there are lifts of  $p$  to  $\phi_1, \phi_2$  such that

$$\begin{array}{ccccc} (\tilde{X}, \tilde{x}_1) & \xrightarrow{\phi_1} & (\tilde{X}, \tilde{x}_0) & \xrightarrow{\phi_2} & (\tilde{X}, \tilde{x}_0) \\ & \searrow p & \downarrow p & \swarrow p & \\ & & (X, x_0) & & \end{array}$$

so  $\phi_1, \phi_2$  are deck transformations taking  $\tilde{x}_0$  to  $\tilde{x}_1$  or the reverse.

(2) From above, if  $[\gamma]H[\gamma]^{-1} = H$ , there exists  $\phi \in G(\tilde{X})$  such that  $\phi(\tilde{x}_0) = \tilde{\gamma}(1)$ , where  $\tilde{\gamma}$  is the lift of  $\gamma$  based at  $\tilde{x}_0$ . So we get a map  $\Phi : N(H) \rightarrow G(\tilde{X})$ . First, we claim that  $\Phi$  is a homomorphism. Suppose  $\phi_i = \Phi([\gamma_i])$  for  $i = 1, 2$  and  $[\gamma_i] \in N(H)$ . Then  $\phi_i(\tilde{x}_0) = \tilde{x}_i$ , so  $\tilde{\phi}_i$  is a path from  $\tilde{x}_0$  to  $\tilde{x}_i$ . Note  $\phi_1(\tilde{x}_2)$  is a lift of  $\gamma$  based at  $\tilde{x}_1$ , and  $\tilde{\gamma}_1 * \phi_1(\tilde{\gamma}_2)$  is a path from  $\tilde{x}_0$  to  $\phi_1(\tilde{x}_2)$ . Then

$$[p \circ (\tilde{\gamma}_1 * (\phi_1 \circ \tilde{\gamma}_2))] = [\gamma_1 * \gamma_2] = [\gamma_1] \cdot [\gamma_2],$$

so  $\phi_1 \circ \phi_2 = \Phi([\gamma_1] \cdot [\gamma_2])$ , i.e.  $\Phi$  is a homomorphism. It only remains to show (as an exercise) that  $\Phi$  is surjective and that  $\ker \Phi = H$ , so  $G(\tilde{X}) \cong N(H)/H$  by the first isomorphism theorem.  $\square$

**Remark.** If  $H$  is normal in  $\pi_1(X, x_0)$ , then  $G(\tilde{X}) = \pi_1(X, x_0)/H$ . In particular, if  $\tilde{X}$  is the universal cover (so  $\pi_1(\tilde{X})$  is trivial), then  $G(\tilde{X}) = \pi_1(X, x_0)$ .

# Lecture 13

## Feb. 19 — Singular Homology

### 13.1 Covering Spaces and Group Actions

**Definition 13.1.** A *group action* on a topological space  $X$  is a pair  $(G, \rho)$  where  $G$  is a group and  $\rho : G \rightarrow \text{Homeo}(X)$  is a group homomorphism.<sup>1</sup>

**Remark.** If  $G$  acts on  $X$ , then we can form the quotient space  $X/G$  where two points  $x_1, x_2 \in X$  are identified if there exists  $g \in G$  such that  $\rho(g)(x_1) = x_2$ . Then  $X/G$  is called the *orbit space* of the action.

**Theorem 13.1.** Let  $G$  be a group acting on  $X$  such that  $(*)$  for all  $x \in X$ , there is a neighborhood  $U$  such  $x$  such that  $g_1 U \cap g_2 U \neq \emptyset$  implies  $g_1 = g_2$ . Then:

1.  $\rho : X \rightarrow X/G$  is a normal covering space;
2.  $G \cong G(X \rightarrow X/G)$  if  $X$  is path-connected ( $G(X \rightarrow X/G)$  is the group of deck transformations);
3.  $G \cong \pi_1(X/G)/\rho_*(\pi_1(X))$  if  $X$  is path connected and locally path connected.

*Proof.* This is left as an exercise (or see Hatcher). Note that the hypothesis implies each point in  $X/G$  has an evenly covered neighborhood, so  $(X, \rho)$  is a covering space. Also, any two preimages of a point in  $X/G$  are related by the group action, i.e. is a deck transformation of  $X$ , so  $(X, \rho)$  is normal.  $\square$

**Exercise 13.1.** Show that if  $G$  is a finite group acting freely on a Hausdorff space  $X$  (i.e. no element of  $G$  has a fixed point), then the action satisfies  $(*)$ .

**Example 13.1.1.** Let  $X$  be the four-holed torus, arranged in a triangular shape. Then rotation by  $2\pi/3$  is an action of  $G = \mathbb{Z}/3\mathbb{Z}$  on  $X$  with no fixed points. Then  $X/G$  is a two-holed torus, and by the above theorem  $X$  is a three-fold cover of  $X/G$ .

**Example 13.1.2.** The group  $G = \mathbb{Z}/2\mathbb{Z}$  acts on  $S^n$  by sending  $x \mapsto -x$ , so  $S^n$  is a covering space of  $S^n/G = \mathbb{R}P^n$ . Note that we have already seen this before without appealing to the above theorem.

### 13.2 Introduction to Homology

**Remark.** There are many types of (ordinary) homology, i.e. singular, simplicial, cubical, cellular, etc. All of these are isomorphic on CW-complexes. We will define the most general homology theory, called *singular homology*, and then derive *cellular homology*, which is easier to compute.

---

<sup>1</sup>Here  $\text{Homeo}(X)$  is the group of homeomorphisms of  $X$ .

There are also *generalized homology* theories, such as *bordism theory*, *K-theory*, etc. These are in many ways similar to ordinary homology, but they do not satisfy the dimension axiom, which can give them wildly different behavior. We will briefly mention bordism theory later on in the class.

Underlying these homology theories is *homological algebra*, which is the study of *chain complexes*. Such objects show up in many places and they give rise to more “homology theories,” such as *Floer homology*, *Heegaard-Floer homology*. These are *not* “real homology theories” in the sense of algebraic topology.

## 13.3 Singular Homology

**Definition 13.2.** The *standard  $p$ -simplex* is

$$\Delta^p = \left\{ \sum_{i=0}^p t_i e_i \in \mathbb{R}^{p+1} : \sum_{i=0}^p t_i = 1, t_i \geq 0 \right\},$$

where  $\{e_0, \dots, e_p\}$  are the standard basis vectors for  $\mathbb{R}^{p+1}$ .

**Example 13.2.1.** The following are examples of  $p$ -simplices:

1. For  $p = 0$ , we have  $\Delta^0 = \{t_1 e_1 \in \mathbb{R} : t_1 = 1, t_1 \geq 0\} = e_1$ , which is a point.
2. For  $p = 1$ , we have  $\Delta^1 = \{t_0 e_0 + t_1 e_1 \in \mathbb{R}^2 : t_0 + t_1, t_i \geq 0\}$  is the convex hull of the points  $e_0$  and  $e_1$  in  $\mathbb{R}^2$ , which is a line segment.
3. For  $p = 2$ , the standard 2-simplex  $\Delta^2$  is the convex hull of  $e_0, e_1, e_2 \in \mathbb{R}^3$ , which is a triangle.

**Definition 13.3.** Given any vectors  $v_0, \dots, v_p \in \mathbb{R}^n$ , by  $[v_0, \dots, v_p]$  we denote the map

$$\begin{aligned} \Delta^p &\longrightarrow \mathbb{R}^n \\ (t_0, \dots, t_p) &\longmapsto \sum_{i=0}^p t_i v_i \end{aligned}$$

**Example 13.3.1.** Consider the following examples:

1.  $[e_0, \dots, e_p]$  parametrizes the standard  $p$ -simplex.
2.  $[e_0, \dots, \widehat{e}_i, \dots, e_p]$  (where the hat means to remove  $e_i$ ) is the  $i$ th *face map*, which parametrizes the  $i$ th *face* of  $\Delta^p$ . Denote this map by  $F_i^p = [e_0, \dots, \widehat{e}_i, \dots, e_p]$ .

**Remark.** If  $i > j$ , then

$$F_i^p \circ F_j^{p-1} = [e_0, \dots, \widehat{e}_j, \dots, \widehat{e}_i, \dots, e_p],$$

so  $F_2^2 \circ F_1^1$  identifies  $[e_0]$  with  $[e_0]$  in  $\Delta^2$ . On the other hand, if  $j \geq i$ , then

$$F_i^p \circ F_j^{p-1} = [e_0, \dots, \widehat{e}_i, \dots, \widehat{e}_{j+1}, \dots, e_p].$$

For instance,  $F_0^3 \circ F_1^2$  identifies  $[e_0, e_1]$  with  $[e_1, e_3]$  in  $\Delta^3$ .

**Definition 13.4.** A *singular  $p$ -simplex* in a space  $X$  is a continuous map  $\sigma : \Delta^p \rightarrow X$ . The *singular group of  $p$ -chains* in  $X$ , denoted  $C_p(X)$ , is the free abelian group generated by singular  $p$ -simplices.

**Remark.** An element in  $C_p(X)$  is a finite formal sum  $\sum_{i=1}^k n_i \sigma_i$ , where  $n_i \in \mathbb{Z}$  and  $\sigma_i$  is a singular  $p$ -simplex. There is an obvious way to add elements in  $C(X)$ , and this makes  $C(X)$  into a group

**Definition 13.5.** An element in  $C_p(X)$  is called a *singular  $p$ -chain*. Given a singular  $p$ -simplex  $\sigma$ , the  $i$ th *face* of  $\sigma$  is  $\sigma^{(i)} = \sigma \circ F_i^p$ , and the *boundary* of  $\sigma$  is

$$\partial\sigma = \sum_{i=0}^p (-1)^i \sigma^{(i)}.$$

**Remark.** If  $\sigma$  is a singular  $p$ -simplex, then  $\partial\sigma \in C_{p-1}(X)$ .

**Example 13.5.1.** Let  $\Delta^2 = [e_0, e_1, e_2]$  and  $\sigma : \Delta^2 \rightarrow X$ . Then  $\partial\sigma = \sigma^{(0)} - \sigma^{(1)} + \sigma^{(2)}$  traces the edges  $[\sigma(e_0), \sigma(e_1)]$ ,  $[\sigma(e_1), \sigma(e_2)]$ , and  $[\sigma(e_2), \sigma(e_0)]$ , which forms the boundary of  $\sigma(\Delta^2) \subseteq X$ .

**Definition 13.6.** Define the  $p$ th *boundary map* to be

$$\begin{aligned} \partial_p : C_p(X) &\longrightarrow C_{p-1}(X) \\ \sum_{i=1}^k n_i \sigma_i &\longmapsto \sum_{i=1}^k n_i (\partial\sigma_i). \end{aligned}$$

**Lemma 13.1.** We have  $\partial_{p-1} \circ \partial_p = 0$ , i.e.  $\partial^2 = 0$ .

*Proof.* We can calculate

$$\begin{aligned} \partial_{p-1} \circ \partial_p \sigma &= \partial_{p-1} \left( \sum_{i=0}^p (-1)^i (-1)^i \sigma \circ F_i^p \right) \\ &= \sum_{i=0}^p (-1)^i \left( \sum_{j=0}^{p-1} (-1)^j \sigma \circ F_i^p \circ F_j^{p-1} \right) \\ &= \sum_{0 \leq j < i \leq p} (-1)^{i+j} \sigma \circ F_i^p \circ F_j^{p-1} + \sum_{0 \leq i \leq j \leq p-1} (-1)^{i+j} \sigma \circ F_i^p \circ F_j^{p-1} \\ &= \sum_{0 \leq j < i \leq p} (-1)^{i+j} \sigma \circ [e_0, \dots, \widehat{e}_j, \dots, \widehat{e}_i, \dots, e_p] \\ &\quad + \sum_{0 \leq i \leq j \leq p-1} (-1)^{i+j} \sigma \circ [e_0, \dots, \widehat{e}_i, \dots, \widehat{e}_{j+1}, \dots, e_p]. \end{aligned}$$

Relabel  $k = j + 1$  in the second sum, we find that

$$\begin{aligned} \partial_{p-1} \circ \partial_p \sigma &= \sum_{0 \leq j < i \leq p} (-1)^{i+j} \sigma \circ [e_0, \dots, \widehat{e}_j, \dots, \widehat{e}_i, \dots, e_p] \\ &\quad + \sum_{0 \leq i < k \leq p-1} (-1)^{i+k+1} \sigma \circ [e_0, \dots, \widehat{e}_i, \dots, \widehat{e}_k, \dots, e_p] = 0 \end{aligned}$$

since these are the same but with an extra  $-1$  on the second sum, so they cancel out to 0.  $\square$

**Remark.** The lemma implies that  $\text{im } \partial_{p+1} \subseteq \ker \partial_p$ .

**Definition 13.7.** We define the  $p$ th (singular) *homology group* of  $X$  to be

$$H_p(X) = \frac{\ker \partial_p}{\text{im } \partial_{p+1}}.$$

An element of  $\ker \partial_p$  is called a  $p$ -*cycle*, and an element of  $\text{im } \partial_{p+1}$  is called a  $p$ -*boundary*. We say that two chains  $c_1, c_2 \in C(X)$  are *homologous* if there exists some  $d \in C_{p+1}(X)$  such that  $c_1 - c_2 = \partial d$ .

**Remark.** Writing  $c_1 \sim c_2$  if  $c_1, c_2$  are homologous and letting the equivalence classes be denoted by  $[c]$ , we can also write  $H_p$  as

$$H_p(X) = \{[c] : c \text{ is a } p\text{-cycle}\}.$$

**Remark.** There are no singular  $p$ -simplices for  $p < 0$ , so  $C_p(X) = \{0\}$  for  $p < 0$ .

**Lemma 13.2.** *If  $X$  is a one-point space, then we have*

$$H_p(X) = \begin{cases} \mathbb{Z} & \text{if } p = 0, \\ 0 & \text{otherwise.} \end{cases}$$

**Lemma 13.3.** *We have  $H_0(X) \cong \bigoplus_n \mathbb{Z}$ , where  $X$  has  $n$  path components.*