


=====

ECLIPSING BINARY CLASSIFIER

=====

Target: TIC 52357
Mission: TESS
Period: 0.5480 days
Primary depth: 0.0122
Secondary depth: 0.0022
Depth ratio: 0.180

Classification: EA — Algol-li



Correct predicted


=====

ECLIPSING BINARY CLASSIFIER

=====

Target: KIC 2983113
Mission: KEPLER
Period: 0.5927 days
Primary depth: 0.1339
Secondary depth: 0.1252
Depth ratio: 0.935

Classification: EW — W UMa-li



Correctly predicted


=====

ECLIPSING BINARY CLASSIFIER

=====

Target: KIC 9832227
Mission: KEPLER
Period: 0.4585 days
Primary depth: 0.1401
Secondary depth: 0.0925
Depth ratio: 0.661

Classification: EB – Beta Ly



Correct predicted


=====

ECLIPSING BINARY CLASSIFIER

=====

Target: KIC 2305372
Mission: KEPLER
Period: 1.4013 days
Primary depth: 0.7070
Secondary depth: 0.0951
Depth ratio: 0.135

Classification: EA – Algol-li



Correct predicted


=====

ECLIPSING BINARY CLASSIFIER

=====

Target: TIC 2490641
Mission: TESS
Period: 0.5253 days
Primary depth: 0.7983
Secondary depth: 0.1514
Depth ratio: 0.190

Classification: EA – Algol-li



Incorrect prediction. Stress test.

The classifier employs a simplified rule-based approach mainly relying on eclipse depths and light-curve morphology. It can misclassify systems with extra astrophysical variability, like starspots, indicating that more advanced modelling or machine-learning techniques are needed for reliable classification.


=====

ECLIPSING BINARY CLASSIFIER

=====

Target: KIC 9832227
Mission: KEPLER
Period: 0.4585 days
Primary depth: 0.1401
Secondary depth: 0.0925
Depth ratio: 0.661

Classification: EB — Beta Lyr



Correct predicted