

EULAR recommendations for the management of systemic lupus erythematosus: 2023 update

Antonis Fanouriakis ¹, Myrto Kostopoulou ¹, Jeanette Andersen,² Martin Aringer ³, Laurent Arnaud ⁴, Sang-Cheol Bae ⁵, John Boletis,⁶ Ian N Bruce,⁷ Ricard Cervera,⁸ Andrea Doria ⁹, Thomas Dörner ¹⁰, Richard A Furie ¹¹, Dafna D Gladman ¹², Frederic A Houssiau ¹³, Luís Sousa Inês ¹⁴, David Jayne ¹⁵, Marios Kouloumas,¹⁶ László Kovács,¹⁷ Chi Chiu Mok ¹⁸, Eric F Morand ¹⁹, Gabriella Moroni,²⁰ Marta Mosca,²¹ Johanna Mucke ²², Chetan B Mukhtyar ²³, György Nagy ^{24,25,26}, Sandra Navarra,²⁷ Ioannis Parodis ^{28,29,30}, José M Pego-Reigosa,³¹ Michelle Petri ³², Bernardo A Pons-Estel,³³ Matthias Schneider,²² Josef S Smolen,³⁴ Elisabet Svenungsson ^{28,29}, Yoshiya Tanaka ³⁵, Maria G Tektonidou ³⁶, YK Onno Teng ³⁷, Angela Tincani ³⁸, Edward M Vital ³⁹, Ronald F van Vollenhoven ⁴⁰, Chris Wincup ⁴¹, George Bertsias ⁴², Dimitrios T Boumpas ^{1,43,44}

Handling editor David S Pisetsky

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/ard-2023-224762>).

For numbered affiliations see end of article.

Correspondence to

Dr Dimitrios T Boumpas, Rheumatology and Clinical Immunology Unit, 4th Department of Internal Medicine, "Attikon" University Hospital, Joint Academic Rheumatology Program, National and Kapodistrian University of Athens, Athens 12462, Greece; boumpasd@uoc.gr

Received 22 July 2023

Accepted 18 September 2023

Published Online First

12 October 2023



© Author(s) (or their employer(s)) 2024. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Fanouriakis A, Kostopoulou M, Andersen J, et al. *Ann Rheum Dis* 2024;**83**:15–29.

ABSTRACT

Objectives To update the EULAR recommendations for the management of systemic lupus erythematosus (SLE) based on emerging new evidence.

Methods An international Task Force formed the questions for the systematic literature reviews (January 2018–December 2022), followed by formulation and finalisation of the statements after a series of meetings. A predefined voting process was applied to each overarching principle and recommendation. Levels of evidence and strengths of recommendation were assigned, and participants finally provided their level of agreement with each item.

Results The Task Force agreed on 5 overarching principles and 13 recommendations, concerning the use of hydroxychloroquine (HCQ), glucocorticoids (GC), immunosuppressive drugs (ISDs) (including methotrexate, mycophenolate, azathioprine, cyclophosphamide (CYC)), calcineurin inhibitors (CNIs, cyclosporine, tacrolimus, voclosporin) and biologics (belimumab, anifrolumab, rituximab). Advice is also provided on treatment strategies and targets of therapy, assessment of response, combination and sequential therapies, and tapering of therapy. HCQ is recommended for all patients with lupus at a target dose 5 mg/kg real body weight/day, considering the individual's risk for flares and retinal toxicity. GC are used as 'bridging therapy' during periods of disease activity; for maintenance treatment, they should be minimised to equal or less than 5 mg/day (prednisone equivalent) and, when possible, withdrawn. Prompt initiation of ISDs (methotrexate, azathioprine, mycophenolate) and/or biological agents (anifrolumab, belimumab) should be considered to control the disease and facilitate GC tapering/discontinuation. CYC and rituximab should be considered in organ-threatening and refractory disease, respectively. For active lupus nephritis, GC, mycophenolate or low-dose intravenous CYC are recommended as anchor drugs, and add-on therapy with belimumab or CNIs (voclosporin or tacrolimus) should

be considered. Updated specific recommendations are also provided for cutaneous, neuropsychiatric and haematological disease, SLE-associated antiphospholipid syndrome, kidney protection, as well as preventative measures for infections, osteoporosis, cardiovascular disease.

Conclusion The updated recommendations provide consensus guidance on the management of SLE, combining evidence and expert opinion.

INTRODUCTION

Since 2008, when the first EULAR recommendations for the management of systemic lupus erythematosus (SLE) were published and widely adopted,¹ a series of specific recommendations on disease monitoring, neuropsychiatric SLE and lupus nephritis (LN), pregnancy and women's health issues in SLE have been developed.^{2–5} More recently, updated recommendations on the management of general SLE and LN were published in 2019 and 2020, respectively, the latter jointly with the European Renal Association/European Dialysis and Transplant Association (ERA/EDTA).

Since the 2019 update, the pace of new developments in SLE has accelerated. A second biological drug, anifrolumab, has been approved for the management of the disease, while the field of LN has also witnessed major breakthroughs; the approval of belimumab and voclosporin, a novel calcineurin inhibitor (CNI), for patients with active LN has inspired discussions on a 'paradigm shift' in the treatment of LN, moving from the traditional 'induction-maintenance' regimen to the early use of combination therapies.^{6,7} These advances created the impetus for an update of the recommendations, to provide guidance on an evolving landscape and capitalise on the experience gained thus far. Given

the drug pipeline and lessons learnt regarding trial design from previous trial failures, it is likely that new therapeutic options will continue to emerge, and SLE may finally enter the era of more frequent updates of its management recommendations, similar to other diseases.^{8,9}

METHODS

Per the EULAR standard operating procedures (SOPs)¹⁰ and the AGREE II document,¹¹ the convenor (DTB) submitted a proposal for an update of the recommendations for the management of SLE, which was approved by the EULAR Quality of Care Subcommittee and the EULAR Council. Following approval, a Task Force was assembled to form the research questions for the systematic literature review (SLR), that consisted of 35 rheumatologists, five nephrologists and two patient representatives (JA, MKou), also including two methodologists (GB, CBM) and two fellows responsible for the SLR (AF, MK). Importantly, the Task Force also included non-European experts, four from the Americas (two from the USA (RF, MP), one from Canada (DDG), one from Argentina (BAPE)), four from Asia (SCB, CCM, SVN, YT) and one from Australia (EM). EMerging Eular NETwork was also represented by two members (JM, CW). All members completed a COI form.

Before the first meeting, an outline of the proposed methodology, a set of the main research questions and the respective Population, Intervention, Comparison and Outcomes (PICO) were circulated among the panellists, who were encouraged to comment, edit and propose additional topics for the SLR. Through this process, it was decided that the SLRs would focus on five domains: (1) management of general and organ-specific SLE, (2) targets of treatment, (3) tapering/withdrawal of treatment, (4) management of patients with SLE and antiphospholipid syndrome (APS), and (5) the efficacy and safety of vaccination against herpes zoster and SARS-CoV2 viruses (a general SLR for infection prevention was not decided, as there are dedicated EULAR recommendations on this topic). Separate PICO and search strings were developed for each domain, resulting in five SLRs. The two fellows, supervised by the methodologists, performed the PICO-based SLRs using two different databases (PubMed and Central). Importantly, since the previous recommendations on general SLE had included papers through December 2017, the current SLRs were limited to English language publications published between January 2018 and December 2022. Pertinent articles identified by manual search within the reference list of the originally retrieved publications were also included. A risk of bias assessment was performed using the Cochrane Risk of Bias tool and Newcastle-Ottawa scale for randomised controlled trials (RCT) and observational studies, respectively. The final level of evidence and grading of recommendations, according to the Oxford Evidence-based Medicine grading levels,¹² considered also the body of evidence that had informed the 2019 EULAR recommendations.¹³

The results of the SLRs, focusing on new evidence and data quality, were presented during the first meeting (29 April 2023), held virtually to facilitate the attendance of the international Task Force members. The participants reviewed the evidence, and an initial draft of the statements/recommendations was formulated following open deliberations. Then, suggestions and edits by all members were incorporated by the convenor, methodologists and the fellows responsible for the SLR to a new modified set of recommendations, which was again circulated to the panel members to propose any additional changes. To achieve consensus, a second meeting was held (7 May 2023), in

which panellists discussed in detail each overarching principle and recommendation and came up with the final version which served as basis for the voting. During the third and last meeting (14 May 2023), all Task Force members who were present, were asked to vote per bullet point whether they agreed or not in principle with the respective statement. As per the EULAR SOPs, a recommendation was immediately accepted if more than 75% of those present voted in favour. In cases where consensus was not met, possible causes of disagreement were discussed, amendments to the statements were made and a second round of voting requiring more than 66% agreement from participants took place. Following approval of the recommendation, every Task Force member provided their level of agreement (LoA) for each statement in a scale from 0 (no agreement at all) to 10 (100% agreement).

Of note, the 2019 recommendations counted a total of 33 recommendations belonging to 4 domains (goals of treatment, general principles of treatment, specific manifestations and comorbidities). The current update aligns with the EULAR SOPs to include a maximum 15 bullet point statements. To this end, recommendations from the previous set were either omitted or merged, and new recommendations were formulated. Because of this change from the previous update, all overarching principles and individual recommendations are hereby presented as new, even for individual recommendations where the essence has remained unchanged, because a detailed description of ‘merged’, ‘omitted’ and ‘rephrased’ statements was considered impractical.

RESULTS

Following the meetings mentioned above, the Task Force formulated a final number of 5 overarching principles and 13 recommendations (table 1). The detailed results of the SLR are summarised in the online supplemental appendix; however, parts of the data are also presented herein, to provide an explanation of the results.

Overarching principles

The overarching principles contain general information on the management of SLE, reflect common sense and are not accompanied by a respective level of evidence. They are nevertheless important to set an overall framework for the approach to a patient with SLE and highlight the role of physician–patient interaction. The overarching principles were voted as a group of principles (ie, not individually) and agreement was 100%.

A. SLE requires multidisciplinary, individualised management with patient education and shared decision-making, taking into consideration the costs to patient and society.

While rheumatologists are the specialists who should primarily care for patients with SLE, the multisystem nature of the disease often mandates the involvement of other disciplines (eg, nephrologists, dermatologists etc), and treatment decisions should be individualised considering patient preferences and patient education. Comparative costs of different treatments should be weighed against the cost of illness and the societal impact of SLE, in terms of social and work participation. Mean (SD) LoA was 9.88 (0.40).

B. SLE disease activity should be assessed at each clinic visit (the frequency depending on physician’s discretion), with evaluation of organ damage (at least annually), using validated instruments.

Task Force members agreed on the need to formally assess lupus disease activity at each visit for the need of therapy adaptation. The phrasing ‘frequency depending on physician’s discretion’ was decided following deliberations, in which it became

Table 1 EULAR Recommendations for the management of patients with systemic lupus erythematosus—2023 update

	Level of agreement	
	Mean (SD)	% with score ≥8
Overarching principles		
A. SLE requires multidisciplinary, individualised management with patient education and shared decision-making, taking into consideration the costs to patient and society.	9.88 (0.40)	100
B. SLE disease activity should be assessed at each clinic visit (the frequency depending on physician's discretion), with evaluation of organ damage (at least annually), using validated instruments.	9.74 (0.63)	100
C. Non-pharmacological interventions, including sun protection, smoking cessation, healthy, balanced diet, regular exercise and measures to promote bone health are important to improve long-term outcomes	9.90 (0.37)	100
D. Pharmacological interventions are directed by patient characteristics, type and severity of organ involvement, treatment-related harms, comorbidities, risk for progressive organ damage, and patient preferences.	10 (0)	100
E. Early SLE diagnosis (including serological assessment), regular screening for organ involvement (especially nephritis), prompt initiation of treatment aiming at remission (or low disease activity if remission is not possible) and strict adherence to treatment are essential to prevent flares and organ damage, improve prognosis and enhance quality of life.	9.81 (0.51)	100
Recommendation/statement		
1. Hydroxychloroquine is recommended for all patients (1b/A), unless contraindicated, at a target dose of 5 mg/kg real body weight/day (2b/B) but individualised based on risk for flare (2b/B) and retinal toxicity.	9.21 (1.35)	90.4
2. Glucocorticoids, if needed, are dosed based on the type and severity of organ involvement (2b/C), and should be reduced to maintenance dose of ≤5 mg/day (prednisone equivalent) (2a/B) and, when possible, withdrawn; in patients with moderate-to-severe disease, pulses of intravenous methylprednisolone (125–1000 mg/day, for 1–3 days) (3b/C) can be considered.	9.57 (0.77)	97.6
3. In patients not responding to hydroxychloroquine (alone or in combination with glucocorticoids) or patients unable to reduce glucocorticoids below doses acceptable for chronic use, addition of immunomodulating/immunosuppressive agents (eg, methotrexate (1b/B), azathioprine (2b/C) or mycophenolate (2a/B)) and/or biological agents (eg, belimumab (1a/A) or anifrolumab (1a/A)) should be considered.	9.32 (0.91)	95.2
4. In patients with organ-threatening or life-threatening disease, intravenous cyclophosphamide (2b/C) should be considered; in refractory cases, rituximab (2b/C) may be considered.	9.38 (0.99)	95.2
5. Treatment of active skin disease should include topical agents (glucocorticoids, calcineurin inhibitors) (2b/B), antimalarials (hydroxychloroquine, chloroquine) (1a/A), and/or systemic glucocorticoids (4/C) as needed, with methotrexate (1b/B), mycophenolate (4/C), anifrolumab (1a/A), or belimumab (1a/B) considered as second-line therapy.	9.35 (1.06)	95.2
6. In active neuropsychiatric disease attributed to SLE, glucocorticoids and immunosuppressive agents for inflammatory manifestations (1b/A) and antiplatelet agents/anticoagulants for atherothrombotic/aPL-related manifestations (2b/C) should be considered.	9.68 (0.81)	97.6
7. For acute treatment of severe autoimmune thrombocytopenia, high-dose glucocorticoids (including pulses of intravenous methylprednisolone) (4/C), with or without intravenous immunoglobulin G (4/C), and/or rituximab (2b/B), and/or high-dose intravenous cyclophosphamide (4/C), followed by maintenance therapy with rituximab (2b/B), azathioprine (2b/C), mycophenolate (2b/C), or cyclosporine (4/C) should be considered.	9.48 (0.86)	97.6
8. Patients with active proliferative lupus nephritis should receive low-dose (EuroLupus) intravenous cyclophosphamide (1a/A) or mycophenolate (1a/A) and glucocorticoids (pulses of intravenous methylprednisolone followed by lower oral doses); combination therapy with belimumab (either with cyclophosphamide or mycophenolate (1b/A)) or calcineurin inhibitors (especially voclosporin or tacrolimus, combined with mycophenolate, 1b/A) should be considered.	9.36 (1.06)	92.8
9. Following renal response, treatment of lupus nephritis should continue for at least 3 years (2b/B); patients initially treated with mycophenolate alone or in combination with belimumab or a calcineurin inhibitor should remain on these drugs (1a/A), whereas azathioprine or mycophenolate should replace cyclophosphamide for those initially treated with cyclophosphamide alone (1a/A) or in combination with belimumab (1a/A).	9.56 (0.81)	95.2
10. In patients at high-risk for renal failure (defined as reduced GFR, histological presence of cellular crescents or fibrinoid necrosis, or severe interstitial inflammation), high-dose (NIH regimen) intravenous cyclophosphamide (1a/A) in combination with pulse intravenous methylprednisolone, can be considered.	9.57 (0.86)	95.2
11. In patients with SLE achieving sustained remission, gradual tapering of treatment should be considered, with withdrawal of glucocorticoids first (2a/B).	9.89 (0.38)	100
12. SLE associated with thrombotic antiphospholipid syndrome (APS) should be managed with long-term vitamin K antagonists after the first arterial or unprovoked venous thrombotic event (1b/B); low dose aspirin (75–100 mg/day) should be considered in patients with SLE without APS but with high-risk aPL profile (2a/B).	9.57 (0.83)	97.6
13. Immunisations for the prevention of infections (herpes zoster virus, human papillomavirus, influenza, COVID-19 and pneumococcus), management of bone health, nephroprotection and cardiovascular risk, and screening for malignancies, should be performed (5/D).	9.85 (0.36)	100

Levels of evidence according to the Oxford Evidence-based Medicine grading levels (<https://www.cebm.net/wp-content/uploads/2014/06/CEBM-Levels-of-Evidence-2.1.pdf>). aPL, antiphospholipid antibodies; GFR, glomerular filtration rate; NIH, National Institutes of Health; SLE, systemic lupus erythematosus.

clear that optimal frequency of patient visits may vary, from a few days in a patient with active LN to up to 6 months in patients with long-standing quiescent disease; thus, judgement of the treating physician cannot be substituted by a fixed range of intervals. The most frequently used validated instruments for measuring disease activity are the various versions of the SLE Disease Activity Index (SELENA-SLEDAI or SLEDAI-2K) and the British Isles Lupus Assessment Group (BILAG). Annual assessment of irreversible damage is important because damage accrual has significant prognostic value. Mean (SD) LoA for this principle was 9.74 (0.63).

C. Non-pharmacological interventions, including sun protection, smoking cessation, healthy, balanced diet, regular exercise and measures to promote bone health are important to improve long-term outcomes.

These interventions are not specific to SLE and pertain also to the general population, although patients with lupus should particularly avoid sun exposure due to the characteristic photosensitivity of the disease. The importance of smoking cessation should be emphasised, as smoking may also interfere with the efficacy of antimalarials¹⁴ and biologics (belimumab)¹⁵ among its other detrimental sequelae. Importantly, a dedicated set of EULAR recommendations on non-pharmacological management of SLE and systemic sclerosis were published recently; the

reader is referred there for more relevant information.¹⁶ Mean (SD) LoA was 9.90 (0.37).

D. Pharmacological interventions are directed by patient characteristics, type and severity of organ involvement, treatment-related harms, comorbidities, risk for progressive organ damage and patient preferences.

The phenotypic heterogeneity and variable severity of organ involvement, as well as differential response to drugs based on patient characteristics, mandate an individualised approach. Pharmacological treatment of lupus may range from hydroxychloroquine (HCQ) monotherapy for patients with mild skin and/or joint symptoms, to highly potent immunosuppressive medications like high-dose glucocorticoids (GC) and cyclophosphamide (CYC) in patients with organ-threatening or life-threatening disease. When choosing therapy, immutable characteristics, such as race and ethnicity, as well as socioeconomic determinants and access to different drugs should be taken into account. For example, black patients with LN may be more responsive to mycophenolate than CYC.¹⁷ Patient research partners in the Task Force highlighted the importance of patient preferences in the treatment decisions, which form the basis of shared decision-making. This principle received the highest mean LoA, mean (SD) 10 (0), indicating complete agreement of all individual panellists.

E. Early SLE diagnosis (including serological assessment), regular screening for organ involvement (especially nephritis), prompt initiation of treatment aiming at remission (or low disease activity if this is not possible), and strict adherence to treatment are essential to prevent flares and organ damage, improve prognosis and enhance quality of life.

The final overarching principle highlights four pillars in the management of SLE: (1) need for early diagnosis, because despite increased awareness in combination with the new, more sensitive classification criteria,¹⁸ recent studies support that patients with SLE still face diagnostic delay (median 2 years from onset of symptoms)^{19,20}; (2) vigilant monitoring for new organ involvement, mainly LN, especially during the first years of the disease, but also thereafter. This need for increased awareness for signs of new-onset kidney involvement was emphasised by several Task Force members, because LN represents a major milestone in the natural history of the disease and delaying its diagnosis has profound prognostic repercussions; (3) pursuing a treatment target, which should ideally be remission, as defined by the recent Definition Of Remission In SLE (DORIS) criteria,²¹ or alternatively, a state of low disease activity, such as the Lupus Low Disease Activity state (LLDAS).²² Both remission and LLDAS have been extensively validated and proven to reduce the risk for damage and other adverse outcomes in patients with SLE (a detailed analysis of the favourable outcomes associated with remission and LLDAS is given in the online supplemental appendix); and (4) the importance of patient adherence to treatment. Specific reference to the issue of adherence in the overarching principles was emphasised by several panellists, including the patient research partners, because medication non-adherence, despite reported wide variations, is considered a major cause of treatment failure.²³ A trusting relationship between the physician and patient forms the basis for the minimisation of the risk of non-adherence. Mean (SD) LoA for the final overarching principle was 9.81 (0.51).

Individual recommendations

1. Hydroxychloroquine is recommended for all patients (1b/A), unless contraindicated, at a target dose of 5 mg/kg real body weight/day (2b/B), but individualised based on risk for flare (2b/B) and retinal toxicity.

HCQ is the mainstay of treatment for patients with SLE and the current SLR extended the existing body of evidence regarding the multiple beneficial effects of HCQ in various aspects of the disease. In the 2019 recommendations, emphasis was placed on the specification that HCQ dose 'should not exceed 5 mg/kg real body weight/day', in view of data which suggested a higher than previously thought risk for retinal toxicity by the use of more sensitive screening techniques.²⁴ A recent observational study assessed the risk for flares in relation to HCQ dose during the previous 6-month period and found an almost twofold risk for any flare for doses ≤ 5 mg/kg/day (vs > 5 mg/kg/day), increased to more than sixfold for moderate or severe flares, with a threshold dose calculated near 5 mg/kg/day.²⁵ Until more data become available regarding benefit-risk relationship of different doses, it was decided that the HCQ target dose remains at 5 mg/kg/day; however, this should be individualised based on risk for flare and retinal toxicity, with patients at higher risk for retinal toxicity (kidney disease, preexisting macular or retinal disease, tamoxifen use) being candidates for closer ophthalmologic follow-up. In selected patients and circumstances (eg, moderate or severe disease), initial HCQ dose higher than 5 mg/kg/day (but not exceeding 400 mg/day) may be used, followed by lowering of the

dose to within range once the patient has improved. In addition, Task Force members suggested the use of monitoring HCQ blood levels to guide the optimal dose for each patient and assess for possible non-adherence to therapy, based on studies suggesting that HCQ whole blood levels may reflect patient adherence to treatment.²⁶ Although a universal recommendation for HCQ blood level monitoring would be impractical, it can nevertheless be used to guide dosage adaptations, in settings where it is available. Finally, the use of antimalarials other than HCQ was discussed mainly by non-European panellists, to address the issue of potential limited availability or higher cost of HCQ in some countries. In such settings, chloroquine may be used as an alternative, bearing in mind that it may be more toxic than HCQ (mainly for retinal toxicity).²⁷ Finally, quinacrine can be considered in patients with cutaneous manifestations and HCQ-induced retinopathy. The statement on HCQ was agreed on by 77.8% of participants following one round of amendments (the only statement where this was needed) and mean (SD) LoA was 9.21 (3.35).

2. Glucocorticoids, if needed, are dosed based on the type and severity of organ involvement (2b/C), and should be reduced to maintenance dose of ≤ 5 mg/day (prednisone equivalent) (2a/B) and, when possible, withdrawn; in patients with moderate-to-severe disease, pulses of intravenous methylprednisolone (125–1000 mg per day, for 1–3 days) (3b/C) can be considered.

Minimisation of GC use, in view of their detrimental effects, was a major theme of discussion during the Task Force meetings. Numerous studies in the current SLR confirmed associations of different cut-offs for daily prednisone dose with adverse outcomes, most of which pointed to the threshold of 5 mg/day. Although a controlled trial of different GC tapering regimens or maintenance doses is still lacking in SLE, the Task Force elected to lower the 'acceptable' threshold of daily prednisone dose for maintenance treatment to maximum 5 mg/day prednisone equivalent, as compared with 7.5 mg/day in the 2019 recommendations. Ideally, one could envision the use of GC only as 'bridging therapy' in SLE, similar to rheumatoid arthritis (lowest possible dose for the shortest possible period), and the complete withdrawal of GC is the optimal target.

Intravenous pulses of methylprednisolone (MP) of various doses (depending on disease severity and patient weight) capitalise on the immediate non-genomic effects of GC,²⁸ and may allow for a faster tapering of per os (PO) GC.²⁹ Importantly, pulse IV MP has not been linked to certain established GC-related harms, like avascular necrosis.³⁰ Initial PO dose also depends on disease severity; a retrospective study in 206 patients with LN using propensity score matching found higher rates of 1-year complete response in patients who started with ≥ 40 mg/day compared with those who started with ≤ 30 mg/day, without increased risk for GC-related damage.³¹ A smaller study in non-renal lupus had found similar reduction in disease activity at 1 year and higher risk of damage with starting dose > 30 mg/day versus ≤ 30 mg/day.³² Despite these discrepancies, most panellists agreed that it is the chronic exposure to GC that confers the major risk, and the statement received 96.3% agreement and mean (SD) LoA was 9.57 (0.77).

3. In patients not responding to hydroxychloroquine (alone or in combination with glucocorticoids) or patients unable to reduce glucocorticoids below doses acceptable for chronic use, addition of immunomodulating/immunosuppressive agents (eg, methotrexate (1b/B), azathioprine (2b/C) or mycophenolate (2a/B)) and/or biological agents (eg, belimumab (1a/A) or anifrolumab (1a/A)) should be considered.

This statement emphasises the value of conventional and biological immunomodulatory/immunosuppressive drugs for the control of the disease and facilitation of GC tapering and withdrawal. Since no new, high-quality data emerged in the past 4 years regarding conventional immunosuppressive drugs, deliberations regarding this statement focused on two main issues: (1) inclusion of anifrolumab, following its approval in 2021,^{33 34} as well as belimumab,³⁵ as biological agents with proven efficacy in controlling disease activity, reducing flares, and allowing for GC dose reduction. In the recommendation, there is no hierarchy in the choice between anifrolumab and belimumab, as the two drugs have not been compared in a head-to-head trial and their approval was the result of RCTs in similar extrarenal SLE populations. The panel noted that there are more than 10 years of real-life clinical experience with belimumab, while no real-life data for anifrolumab had been published by the time of the SLR completion. (2) The positioning of biological agents in relation to conventional immunosuppressive drugs for the treatment of SLE. For the latter point, while considerations from specific countries, healthcare settings and biological reimbursement policies have to be taken into account, most panelists agreed that prior use of a conventional immunosuppressive drug (MTX, AZA, mycophenolate mofetil or mycophenolic acid (henceforth combined referred to as 'mycophenolate', see online supplemental table 1 for details), leflunomide³⁶ or others) should not be mandatory for initiating anifrolumab or belimumab. Of note, this is unchanged from the 2019 recommendations. The rationale driving this statement was that, despite their substantially higher cost, approved biological drugs have proven their efficacy in high-quality RCTs, while such data are lacking for conventional immunosuppressive drugs, which continue to be used based on rheumatologists' long-term real-life experience. This recommendation received 84.6% agreement and mean (SD) LoA was 9.32 (0.91).

4. In patients with organ-threatening or life-threatening disease, intravenous cyclophosphamide (2b/C) should be considered; in refractory cases, rituximab (2b/C) may be considered.

Due to its gonadal toxicity, high-dose IV CYC is reserved for severe cases of lupus, in which it may be considered owing to its high efficacy. Rituximab (RTX) is used off-label in SLE and is recommended in circumstances³⁷ where other drugs have failed, with notable exceptions (eg, immune cytopenias, see below) where it can be used earlier. Although a universal definition of refractory disease is lacking in SLE, it is conceived as disease not responding to different classes of immunosuppressive medications. The combination of RTX with CYC had been used in early studies of RTX,³⁸ but additional benefit has not been confirmed³⁹ and this combination comes at the expense of an increased risk for infections. Sequential strategies of RTX followed by belimumab are being explored, but more data are needed.^{40 41}

Patients not responding to any of the aforementioned therapies might be offered other options, such as plasma exchange,⁴² haematopoietic stem cell transplantation⁴³ or experimental therapies. In 2022, the use of CAR T-cells in five patients with severe, refractory SLE was published with encouraging results, yet RCTs and more long-term data are needed.⁴⁴ Agreement for this recommendation was 100%, with mean (SD) LoA 9.38 (0.99).

5. Treatment of active skin disease should include topical agents (glucocorticoids, calcineurin inhibitors) (2b/B), antimalarials (hydroxychloroquine, chloroquine) (1a/A), and/or systemic glucocorticoids (4/C) as needed, with anifrolumab (1a/A), belimumab (1a/B), methotrexate (1b/B), or mycophenolate (4/C), considered as second-line therapy.

For the treatment of active skin disease in SLE, few new data have emerged since the 2019 recommendations, and a significant body of evidence continues to originate from studies in patients with cutaneous lupus erythematosus. Recommended first-line treatment (topical agents, antimalarials and/or systemic GC) has not changed in the statement. HCQ is the antimalarial of choice, but chloroquine may be used in the settings discussed earlier.⁴⁵ Quinacrine (mepacrine) may also be used in cases of inadequate response or toxic retinopathy, as add-on to HCQ or alternative therapy, respectively,⁴⁶ but its use is limited by frequent intolerance and unavailability in many countries.

For the ~40% of patients not responding to first-line therapy,⁴⁷ comparative studies among existing immunosuppressive drugs are lacking. Despite this paucity, recommended second-line drugs have partly changed from 2019, because the Task Force decided to recommend drugs more familiar to rheumatologists (such as MTX or mycophenolate, instead of dapsone or retinoids). A small retrospective study in 73 patients with refractory CLE to first-line therapy found similar response rates (~65%) between MTX and mycophenolate.⁴⁸ Anifrolumab and belimumab have both shown efficacy in mucocutaneous manifestations of SLE,^{49 50} although only anifrolumab has used the Cutaneous Lupus Area and Severity Index in its clinical programme, whereas belimumab has reported responses according to the general instruments SLEDAI and BILAG (hence, the designation B in the Grading of Recommendation, despite positive RCT data). Importantly, the list of recommended drugs is indicative and other treatments may be considered as second-line or third-line options, including dapsone, retinoids, CNI, AZA, CYC and RTX, ideally in collaboration with dermatologists experienced in the treatment of CLE. Finally, thalidomide and, more recently, lenalidomide, are effective in various subtypes of cutaneous lupus,^{51 52} and lenalidomide has a lower risk for polyneuropathy than thalidomide; however, they should both be reserved for patients that have failed multiple previous agents, and with the utmost caution in women of reproductive age. This recommendation was agreed on by 96.3%, mean LoA (SD) was 9.35 (1.06).

6. In active neuropsychiatric disease attributed to SLE, glucocorticoids and immunosuppressive agents for inflammatory manifestations (1b/A) and antiplatelet agents/anticoagulants for atherothrombotic/antiphospholipid antibodies (aPL)-related manifestations (2b/C) should be considered.

This recommendation has remained unchanged, since no new data have emerged during the last 5 years that would deem a modification of the recommendation appropriate.⁵³ Approach to a patient with possible neuropsychiatric SLE should follow the general principles as in the general population and symptomatic treatment as per individual manifestation should be considered. Attribution of a neuropsychiatric manifestation to SLE is challenging and published attribution models may be used in doubtful cases.^{54 55} For severe inflammatory manifestations (eg, myelopathy, acute confusional state), potent immunosuppressive agents, like CYC or RTX,⁵⁶ should be preferred. For the approved biological drugs anifrolumab and belimumab, there is a paucity of evidence regarding their efficacy in neuropsychiatric manifestations, because patients with active, severe forms of such manifestations were excluded from the RCTs of both drugs, and under-represented in real-life use of belimumab. Anticoagulant treatment is mainly indicated in cases of cerebrovascular disease, such as ischaemic stroke associated with aPL, because its value in other manifestations is not clear. Agreement on this recommendation was 96.3% and mean (SD) LoA was 9.68 (0.81).

7. For acute treatment of severe autoimmune thrombocytopenia, high-dose glucocorticoids (including pulses of intravenous

methylprednisolone (4/C), with or without intravenous immunoglobulin G (4/C), and/or rituximab (2b/B), and/or high-dose intravenous cyclophosphamide (4/C), followed by maintenance therapy with rituximab (2b/B), azathioprine (2b/C), mycophenolate (2b/C), or cyclosporine (4/C), should be considered.

Similar to neuropsychiatric SLE, treatment of autoimmune thrombocytopenia in SLE also did not witness major developments since 2019; thus, the current recommendation reflects the principles outlined in the previous update. A platelet number of 20–30 000/mm³ is typically used as the cut-off, below which therapy is indicated. Therapy includes: (1) acute phase treatment with GC (including pulses of intravenous MP, followed by 0.5–0.7 mg/kg/day prednisone equivalent with tapering) with or without IVIG; RTX may also be used early in this setting,⁵⁷ based also on the drug's documented efficacy in idiopathic immune thrombocytopenia, (2) early use of immunosuppressive medications as GC-sparing agents; a small retrospective study showed that patients with SLE with immune cytopenias who relapsed had less often received concomitant immunosuppressive agents following treatment of the initial episode.⁵⁸ More importantly, mycophenolate was shown to reduce relapse when used as first line in a RCT in patients with immune thrombocytopenia.⁵⁹ While a similar RCT has not been performed in SLE, this study provided proof-of-concept for the first-line use of immunosuppressive medications in immune thrombocytopenia, a practice commonly followed by most rheumatologists. Regarding choice of drug for maintenance therapy, there is no hierarchy between the recommended drugs, and this is left to the treating physician's discretion. In cases refractory to these drugs, thrombopoietin receptor (TPO) agonists and splenectomy are options; although the two modalities have not been formally compared in SLE and data mainly come from observational studies,⁶⁰ it seems reasonable to use TPO agonists prior to splenectomy, given the possible complications and long-term sequelae of the latter. However, it should be considered that TPO agonists have been associated with a higher risk of thromboembolic events, therefore their use should be avoided in aPL-positive patients.⁶¹ Fostamatinib, a spleen tyrosine kinase inhibitor, is approved for the treatment of chronic immune thrombocytopenia, but has not been tested in SLE. Similar treatment options (excluding TPO agonists) pertain to SLE autoimmune haemolytic anaemia. This recommendation received 92.6% agreement and mean (SD) LoA was 9.48 (0.86).

8. Patients with active proliferative lupus nephritis should receive low-dose (EuroLupus) intravenous cyclophosphamide (1a/A) or mycophenolate (1a/A) and glucocorticoids (pulses of intravenous methylprednisolone followed by lower oral doses); combination therapy with belimumab (either with cyclophosphamide or mycophenolate (1b/A)) or calcineurin inhibitors (especially voclosporin or tacrolimus, combined with mycophenolate, 1b/A) should be considered.

The recommendation regarding treatment of active LN received the highest attention, in light of the recent approvals of belimumab and voclosporin. Discussions revolved mainly around the position of these drugs in the therapeutic algorithm of LN, that is whether they should be used upfront in an early combination therapy with standard-of-care (SoC, low-dose CYC⁶² or mycophenolate in combination with GC, see online supplemental table 1 for usual drug doses in SLE), or whether they should be reserved for non-responding or relapsing disease. In this regard, deliberations towards the formulation of this recommendation focused on the following facts: (1) LN is by default severe disease, accompanied by increased morbidity and mortality, and leading to gradual nephron loss and chronic

kidney disease (CKD),⁶³ (2) rates of complete response at 1–2 years with SoC therapy (ie, control arms) in recent clinical trials (including the phase 3 trials of belimumab and voclosporin, respectively)^{64 65} are consistently low, in the range of 20%–30%, and (3) both belimumab and voclosporin based on their RCTs have been approved for *all patients with active LN*, meaning that all patients can potentially receive them, including as first line. Of note, in a *post-hoc* analysis of the BLISS-LN, belimumab in combination with SoC was found to reduce the risk for flares by 55% compared with SoC alone, and preserve glomerular filtration rate (GFR) better than SoC.^{66 67} To this end, the possibility for a universal recommendation of early combination therapy aiming to increase renal response rates was intensively discussed among panellists. Counterarguments included the high cost of these therapies and the potential of overtreating some patients who would respond to treatment with mycophenolate or low-dose intravenous CYC alone, as well as respective cost or risk considerations, particularly relevant for long-term use of a CNI. Indeed, some patients with LN present with clinically and histologically milder disease, while others with risk factors for progression to end-stage kidney disease; at present, it is unclear which patients will benefit more from early combination therapies. Of note, recent real-life studies have reported higher response rates with SoC, compared with rates reported in RCTs.^{68 69} Based on all the above, it was proposed that early combination therapy 'should be considered' in all adult patients with active LN, emphasising the fact that treating physicians have the option to decide if and when combination therapy should be used. In the case of CNIs, combination refers to voclosporin as well as tacrolimus (TAC), since the current SLR confirmed the superiority of TAC+mycophenolate over SoC (mainly high-dose CYC), although based on evidence exclusively from Asian populations.⁷⁰

Additional points regarding the recommendation for LN warrant further clarification: first, no distinction between LN histological classes is made in the recommendation. It should be noted that neither belimumab nor voclosporin induced higher renal response rates versus placebo in a post-hoc RCT analysis of the small subgroup of patients with class V LN.^{66 71} Nevertheless, patients with pure class V were underrepresented in these trials (less than 20%), and also there were fewer flares with belimumab in the pure V subgroup. Collectively, the Task Force opined that more data are needed to decide on the optimal treatment of class V LN; to provide a succinct message, it was elected that the statement considers patients with *any* class of LN that needs treatment. Of note, the recommended proteinuria cut-off for immunosuppressive treatment in class V LN remains 1 g/day, as per the 2019 EULAR/ERA-EDTA recommendations.⁷² Second, the secondary analysis of the BLISS-LN trial found that belimumab was more efficacious in patients with baseline proteinuria below 3 g/day.⁶⁶ Third, voclosporin provided a rapid reduction in proteinuria, which may be preferable in patients with a high baseline urine protein in the nephrotic range.⁶⁵ Fourth, in AURORA-1, patients with baseline GFR <45 mL/min were excluded, thus the safety of voclosporin in patients with a low baseline GFR (30–45 mL/min) is as yet unclear. The final decision for the treatment of active LN should depend on the individual patient characteristics as outlined above (histological class, baseline GFR, proteinuria), presence of extrarenal manifestations, comorbidities, risk for toxicity, access to drugs and cost issues, and patient preferences. If a combination therapy is not opted for in patients with treatment-naïve LN, add-on treatment with belimumab or voclosporin should be considered in patients with inadequate response by 3–6 months, or those who

flare. In these cases, physicians should consider to acquire expert consultation.

Regarding dosing of GC, pulses of intravenous MP (eg, 250–1000 mg for 1–3 days) are recommended as part of the initial (or ‘remission induction’) regimen, unless there are major safety concerns (eg, infection). For oral therapy, large controlled trials comparing different GC regimens have not been performed in LN. The 2019 EULAR/ERA-EDTA recommendations endorsed a lower dose GC regimen for initial therapy compared with traditional practices (starting dose 0.3–0.5 mg/kg/day and 20 mg/day for proliferative classes and class V, respectively), though acknowledging that this was not based on high-quality data.⁷³ In the current version of recommendations, the principle for lower cumulative GC exposure was maintained. Importantly, belimumab demonstrated superior GC-sparing potential than SoC in the BLISS-LN study, while the AURORA study of voclosporin used significantly lower GC doses than earlier studies (20–25 mg/day starting oral prednisone rapidly tapered to 5 mg by 12 weeks), lending further support to their use in LN. A recommended GC initial dose and tapering strategy is shown in online supplemental table 1. The overall agreement for this recommendation was 92.8%, with a mean (SD) LoA of 9.36 (1.06).

Thrombotic microangiopathy (TMA) may be evident in up to 20% of LN biopsies, particularly in the presence of aPL, and is associated with an adverse impact on prognosis. Although not mentioned in the statement, treatment options in TMA-LN were discussed, even though high-quality data are lacking. In addition to anticoagulation therapy,⁷⁴ there is emerging evidence for the use of eculizumab, a monoclonal antibody against complement protein C5 which is efficacious in cases of complement-mediated TMA, for patients with LN and histological evidence of TMA.⁷⁵

9. Following renal response, treatment of LN should continue for at least 3 years (2b/B); patients initially treated with mycophenolate alone or in combination with belimumab or a calcineurin inhibitor should remain on these drugs (1a/A), whereas mycophenolate or azathioprine should replace cyclophosphamide for those initially treated with cyclophosphamide alone (1a/A) or in combination with belimumab (1a/A).

Following the choice of initial treatment, renal response should be monitored according to the 2019 EULAR/ERA-EDTA targets (reduction in proteinuria $\geq 25\%$ and 50% at 3 and 6 months, respectively, and below 500–700 mg/day at 12 months, all with GFR within 10% from baseline). These therapeutic goals have now been validated.⁷⁶ Provided that response is achieved, subsequent (or ‘maintenance’) therapy should depend on the initial regimen. If the initial regimen included mycophenolate (either monotherapy or in combination with belimumab or voclosporin), then the same regimen should continue for at least 3 years; on the other hand, if low-dose CYC had been initially used, alone or in combination with belimumab, it should be replaced by mycophenolate or AZA while belimumab should be continued (if used initially).

Duration of immunosuppressive therapy was also intensively discussed. Immunosuppressive treatment in LN, particularly in proliferative classes, should continue for at least 3 years.⁷⁷ Of note, in case of initial therapy with mycophenolate/CNI combination, there is a concern regarding the duration of therapy, as long-term use of ‘legacy’ CNIs (tacrolimus, cyclosporine A) has been associated with nephrotoxicity and GFR decline. Several Asian RCTs have investigated CNI combination therapy as remission induction, but not as a long-term maintenance therapy.^{78 79} In this regard, it is reassuring that the long-term AURORA-2 study extending to 3 years use of voclosporin/mycophenolate

combination recently reported stable levels of GFR throughout the 3-year period.⁸⁰ This recommendation was agreed on by 96.4% of participants and mean (SD) LoA was 9.56 (0.81).

10. In patients at high risk for kidney failure (defined as reduced glomerular filtration rate, histological presence of cellular crescents or fibrinoid necrosis, or severe interstitial inflammation), high-dose (National Institutes of Health regimen) intravenous cyclophosphamide (1a/A) in combination with pulse intravenous methylprednisolone can be considered.

A subset of patients with LN present with baseline clinical and histological characteristics associated with an adverse long-term prognosis. Such patients can still be treated as in recommendation nr. 9, but it should be noted that patients with such characteristics are underrepresented or excluded in all recent trials in LN (eg, the BLISS-LN and AURORA-1 excluded patients with $\text{GFR} < 30$ and $\leq 45 \text{ mL/min/1.73 m}^2$, respectively). Thus, the relative efficacy of these regimens in patients at high risk for kidney failure is currently unclear. A small (32 patients) post hoc analysis of the Aspreva Lupus Management Study found similar response rates (proteinuria and serum creatinine) between mycophenolate and high-dose intravenous CYC.⁸¹ Thus, for patients presenting with impaired kidney function or histological evidence of crescentic glomerulonephritis and/or severe interstitial inflammation, the traditional high-dose intravenous CYC regimen (0.5–0.75 g/m² monthly for 6 months) can also be considered, since it is the most extensively studied therapeutic regimen in severe LN, in the early studies from the NIH. Importantly, this recommendation received 100% agreement among Task Force members, and a mean (SD) LoA of 9.57 (0.86).

Figures 1 and 2 outline the existing treatment options for the management of extrarenal SLE and LN, respectively.

11. In patients with SLE achieving sustained remission, gradual tapering of treatment should be considered, with withdrawal of glucocorticoids first (2a/B).

The possibility of tapering immunosuppressive treatment in patients with SLE with quiescent disease was a specific research question for the SLR, and concerned GC, immunosuppressive drugs (conventional and biological), and finally, HCQ (in this sequence). Regarding GC, a meta-analysis calculated a 24% pooled incidence of SLE flares following GC discontinuation, but relative risk for major flares was not increased compared with patients who continued GC.⁸² In an investigator-initiated study from France, discontinuation of prednisone in patients who had clinically quiescent disease for more than 1 year and received stable 5 mg/day proved inferior to continuation of the same dose, in terms of risk of disease flares.⁸³ Caveats of this study were discussed between Task Force members, mainly the abrupt discontinuation of prednisone and the fact that biological agents like belimumab were not received by any patient. It was decided that these limitations, together with observational studies that have shown no increased risk of flares with gradual tapering of GC to complete withdrawal^{84 85} and the detrimental effects of long-term GC use, allow for a recommendation that gradual GC tapering to discontinuation should be attempted in SLE, in line with current recommendation 2.

Regarding immunosuppressive agents, a second investigator-initiated RCT (Weaning of Immunosuppression in Lupus, WIN-LUPUS) tested whether withdrawal of mycophenolate or AZA after 2–3 years of therapy in LN would be non-inferior to continuation for the occurrence of renal relapses. The study failed to show non-inferiority, as patients in the discontinuation group had more relapses of LN and more extrarenal flares.⁷⁷ On the contrary, similar to GC, uncontrolled observational studies have reported successful withdrawal of immunosuppressive therapy

Treatment of Non-Renal Systemic Lupus Erythematosus




Figure 1 Treatment of non-renal systemic lupus erythematosus. Top-to-bottom sequence does not imply order of preference (eg, MTX, AZA and MMF are equal options for second-line therapy in mild disease or first-line therapy in moderate disease). *Mild disease: constitutional symptoms; mild arthritis; rash ≤ 9% body surface area; platelet count (PLTs) 50–100 × 10⁹/L; SLEDAI ≤ 6; BILAG C or ≤ 1 BILAG B manifestation. *Moderate disease: moderate–severe arthritis ('RA-like'; rash 9%–18% BSA; PLTs 20–50 × 10⁹/L; serositis; SLEDAI 7–12; ≥ 2 BILAG B manifestations). *Severe disease: major organ threatening disease (cerebritis, myelitis, pneumonitis, mesenteric vasculitis); thrombocytopenia with platelets < 20 × 10⁹/L; TTP-like disease or acute haemophagocytic syndrome; rash > 18% BSA SLEDAI > 12; ≥ 1 BILAG A manifestations. [†]Recommendation of belimumab and anifrolumab as first-line therapy in severe disease refers to cases of extrarenal SLE with non-major organ involvement, but extensive disease from skin, joints, and so on. The use of anifrolumab as add-on therapy in severe disease refers mainly to severe skin disease. For patients with severe neuropsychiatric disease, anifrolumab and belimumab are not recommended. ANI, anifrolumab; aPL, antiphospholipid antibodies; APS, antiphospholipid syndrome; AZA, azathioprine; BEL, belimumab; BILAG, British Isles Lupus Assessment Group; CNI, calcineurin inhibitor; CYC, cyclophosphamide; GC, glucocorticoids; HCQ, hydroxychloroquine; IV, intravenous; MMF, mycophenolate mofetil; MTX, methotrexate; PO, per os; RTX, rituximab; SLEDAI, SLE Disease Activity Index; VKA, vitamin K antagonists.

in LN. Of note, the total duration of therapy as well as of remission prior to discontinuation of immunosuppressive drugs are particularly important in the LN setting^{86 87}; patients should have received at least 3–5 years of therapy and be in remission for at least 2 years before withdrawal can be attempted. Prior to withdrawal, tapering should be undertaken very gradually. A repeat kidney biopsy-guided decision for therapy withdrawal in patients in clinical remission, in order to assess for residual histological activity predictive of a subsequent flare, is supported by recent observational studies and could be considered, although this has not been formally tested in a RCT.^{88 89}

Contrary to GC and immunosuppressive drugs, HCQ should not be discontinued in patients with SLE in the absence of unacceptable side-effects; in addition to its multiple benefits including survival,⁹⁰ the SLR concluded that HCQ discontinuation is associated with increased risk for flares (data from observational studies).^{91–93} Additionally, HCQ therapy is a protective factor against disease relapse in patients discontinuing GC or immunosuppressive agents.^{86 94} Although complete discontinuation is discouraged (with the exception of adverse effects), data on tapering/dose reduction are equivocal^{91 92}; thus, a decision for HCQ dose reduction in patients in remission should be taken on an individualised basis.

Importantly, the statement on tapering treatment in quiescent SLE received the highest LoA between members, 100%, and mean (SD) LoA 9.89 (0.38).

12. SLE associated with thrombotic antiphospholipid syndrome (APS) should be managed with long-term vitamin K antagonists after the first arterial or unprovoked venous thrombotic event (1b/B); low-dose aspirin (75–100 mg/day) should be considered in patients with SLE/without APS with high-risk aPL profile (2a/B).

Patients with SLE with concomitant APS represent a challenging endotype of the lupus spectrum. To this end, although EULAR has published specific recommendations for the management of APS in 2019,⁹⁵ it was decided that the significance of SLE-APS merits a specific question for the SLR. Management of definite SLE-aPL/APS should follow the same principles of therapy as primary APS, including the long-term use of vitamin K antagonists in patients with unprovoked venous and those with arterial thrombotic events.⁹⁵ After the first RCT on novel direct oral anticoagulants (DOACs) in APS (Trial of Rivaroxaban in AntiPhospholipid Syndrome trial), based on which the 2019 EULAR recommendations for the management of APS recommended against their use in patients with triple aPL positivity or prior arterial thrombosis, three additional RCTs compared

Treatment of Lupus Nephritis




Figure 2 Treatment of lupus nephritis. Top-to-bottom sequence does not imply order of preference (similar to figure 1). #In addition to general protective measures, as outlined in figure 1. §BEL should always be given in combination with MMF or low-dose CYC as initial therapy, and with MMF or AZA as maintenance therapy. ^CNIs should be given in combination with MMF. *Particularly recommended in the presence of poor prognostic factors: reduced eGFR, histological presence of cellular crescents or fibrinoid necrosis, or severe interstitial inflammation. †Extension of high-dose CYC to subsequent phase refers to severe LN cases, in which bimonthly or quarterly CYC pulses may be given following six monthly pulses. ‡In relapsing/refractory disease, especially after failure to CYC-based regimens. ACEi, angiotensin-converting enzyme inhibitors; APS, antiphospholipid syndrome; ARB, angiotensin receptor blockers; AZA, azathioprine; BEL, belimumab; CNI, calcineurin inhibitor; CYC, cyclophosphamide; eGFR, estimated glomerular filtration rate; GC, glucocorticoids; HCQ, hydroxychloroquine; IV, intravenous; MMF, mycophenolate mofetil; MP, methylprednisolone; PO, per os; RTX, rituximab; SGLT2i, sodium glucose transporter 2 inhibitors; TAC, tacrolimus; Upr, urine protein; VKA, vitamin K antagonists; VOC, voclosporin.

vitamin K antagonists versus DOACs in patients with thrombotic APS.⁹⁵ A recent meta-analysis of these trials showed that DOAC use was associated with increased risk of subsequent arterial thrombotic events (OR: 5.43), especially stroke (OR: 10.74), and the composite of arterial or venous thrombotic events (OR: 4.46).⁹⁶ In patients who have not experienced a thrombotic event, primary prophylaxis with low-dose aspirin should be considered in those with a high risk aPL profile, defined as lupus anticoagulant positivity, or double (any combination of lupus anticoagulant, anticardiolipin antibodies or anti-beta2glycoprotein I antibodies) or triple aPL (all three aPL). Apart from its other beneficial effects, HCQ has also potential antithrombotic effects and may reduce aPL levels,⁹⁷ and is particularly recommended in patients with SLE-aPL or SLE/APS.^{98,99}

Catastrophic APS (CAPS) is a rare complication of APS, with concomitant or successive thrombosis in ≥3 organs. Although a detailed overview of the therapeutic options for CAPS was outside the scope of the SLR, several Task Force members deemed it important to cover this issue in the manuscript text. High-quality studies for the treatment of CAPS in SLE are lacking. Precipitating conditions (eg, infections) should be aggressively sought for and treated, to minimise the risk for the development of CAPS. Triple therapy with full anticoagulation, high-dose GC and plasma exchange and/or IVIG is recommended for

patients with CAPS; more recently, the complement inhibitor eculizumab has shown promise, especially in patients with CAPS with features of complement-mediated TMA (microangiopathic haemolytic anaemia, thrombocytopenia, acute kidney injury).¹⁰⁰ The 2019 EULAR recommendations for the management of APS stated that complement inhibitors or RTX may be considered in refractory CAPS. Agreement for this recommendation was 96.4% and mean (SD) LoA was 9.57 (0.83).

13. Immunisations for the prevention of infections (herpes zoster virus, human papillomavirus, influenza, COVID-19 and pneumococcus), management of bone health, nephroprotection and cardiovascular risk, and screening for malignancies, should be performed (5/D).

The final recommendation is a statement regarding major comorbidities in SLE, reflecting expert opinion based on the evidence for the general benefit of the mentioned measures. As on APS, EULAR has issued specific recommendations regarding vaccinations¹⁰¹ and cardiovascular risk management¹⁰² in patients with autoimmune rheumatic diseases, including SLE, and the reader is referred to the respective manuscripts for further details. In view of the COVID-19 pandemic and the burden of herpes zoster in patients with lupus,¹⁰³ the current SLR included a research question regarding the efficacy and safety of vaccines against SARS-CoV-2 and herpes zoster virus

Box 1 Future research agenda in systemic lupus erythematosus (SLE)

Existing therapies

- ⇒ Utility of measurement of drug blood levels (hydroxychloroquine, mycophenolate, etc)
- ⇒ Randomised trials testing different initial glucocorticoid doses and different tapering regimens
- ⇒ Optimal duration of therapy and timing of immunosuppression discontinuation (both for renal and extrarenal disease)
- ⇒ Value of repeat kidney biopsy prior to immunosuppression discontinuation
- ⇒ Value of per-protocol repeat kidney biopsy after one year of treatment to guide further treatment
- ⇒ Prediction of flare in patients who taper drugs
- ⇒ Prediction of flare in patients who attain the treatment target
- ⇒ Use of statins, low-dose aspirin and other conventional therapies to prevent cardiovascular disease

Pathophysiology and biomarkers

- ⇒ Pre-SLE cohort initiatives to delineate who is at risk to develop SLE and what are the sequential 'hits' that lead to disease development
- ⇒ Involvement of particular organ systems over others, multisystem versus organ-dominant disease
- ⇒ Prediction of response to specific therapies (by clinical, cellular and/or molecular markers)
- ⇒ Biomarkers for response to different biological drugs (pharmacogenetics, transcriptomics, etc)

Clinical trial design and new drug development

- ⇒ Optimisation of clinical trial design and study endpoints to maximise probability of new drug approval
- ⇒ Handling of background medication to avoid polypharmacy and 'dilution' of positive effects of drugs under study
- ⇒ Inclusion of organ-specific endpoints and better and more sensitive measurements of disease activity
- ⇒ Increase in number of adequately trained trial sites (recruitment, infrastructure and training)
- ⇒ Academia versus industry-driven clinical trials
- ⇒ Evaluation and standardisation of patient-reported measures of disease activity/outcomes

(HZV) in lupus. Both the live attenuated and the more efficacious recombinant zoster vaccine have been used in patients with SLE and, although studies are limited, they are considered safe.^{104 105} Similarly, several observational studies have proven the immunogenicity and safety of SARS-CoV-2 vaccines, which are recommended for patients with SLE.¹⁰⁶ Prompt identification and management of infections/sepsis are essential in SLE, and vigilant monitoring for opportunistic infections is warranted in selected patients receiving potent immunosuppressive drugs (eg, high-dose GC, CYC, RTX).¹⁰⁷ In patients with LN, adjunct treatment with nephroprotective agents is of utmost importance to decelerate nephron loss, in combination with immunosuppressive therapy. Renin-angiotensin-aldosterone blockade is required, unless not tolerated, to control hypertension (target level below 130/80 mm Hg).¹⁰⁸ More recently, novel classes of agents, mainly sodium glucose transport 2 (SGLT-2) inhibitors ('flozins'), have gained attention as kidney protective drugs for any case of CKD; SGLT-2 is expressed in kidney biopsies of

patients with LN,¹⁰⁹ and its targeting seems reasonable. A preliminary study of dapagliflozin in a small number of patients with SLE (18 with LN) found no difference in proteinuria following therapy.¹¹⁰ Until more data are available, SGLT-2 inhibitors may be considered in patients with LN with reduced GFR below 60–90 mL/min or proteinuria more than 0.5–1 g/day, on top of ACE/ARB during the maintenance phase. The final recommendation received 92.8% agreement and mean (SD) LoA was 9.85 (0.36).

DISCUSSION

For these recommendations, we assembled a Task Force of world-leading experts in the field of SLE from four continents, to assure the widest possible representation and broad expertise. This is particularly important, because SLE across the world is very different in terms of presentation and severity. To this end, the current set of recommendations sought to ensure a fair balance, and guarantee equal representation of mild, moderate and severe disease, the relative frequency of which may well vary between different countries and continents.

A major modification to the previous sets of EULAR recommendations for SLE was the reduction of the number of individual recommendations. Indeed, to align with the EULAR SOPs but also streamline and simplify the recommendations and facilitate their dissemination, the Task Force managed to condense the recommendations to 13 (with 5 overarching principles). Similar to the 2019 version, the first recommendations (1–4) refer to optimal use of commonly used drugs, recommendations 5–12 deal with specific organ manifestations, and the final recommendation covers the issue of adjunct treatments and comorbidities.

Regarding the use of individual drugs, the emphasis on a more restricted use of GC evolved further from the 2019 recommendations. Thus, GC should be used only if needed, as for mild forms of SLE HCQ alone may suffice. A maximum recommended maintenance dose of 5 mg/day prednisone equivalent is now recommended, stricter than the 7.5 mg/day in the previous recommendations. Of note, this change did not come in view of new high-quality data or RCTs, although we found observational studies linking mean prednisone doses 5 mg/day with adverse sequelae. Nevertheless, there was unanimous agreement among Task Force members that a strong recommendation for a lower dose of GC should be given in view of the detrimental effects of their long-term use and the approval of new agents with GC-sparing effects.

To avoid long-term exposure to GC, early use of immunosuppressive drugs is recommended in SLE. The sequence between conventional and biological drugs was a matter of extensive debate within the Task Force. Nevertheless, in 2019, it was already stated that add-on treatment with belimumab (then the only approved biologic) 'should be considered in patients not responding to combinations of HCQ and GC with or without immunosuppressive agents'. Thus, it was agreed that placing biological drugs (now, belimumab and anifrolumab) after failure of conventional drugs would constitute a step backwards from 2019. To this end, the current recommendations do not require prior failure to one or more conventional drugs before initiating a biological agent, although for the majority cases it may be prudent to try at least one conventional immunosuppressive.

Since 2019, anifrolumab was approved for the treatment of extrarenal SLE in 2021. On the other hand, there is now more than 10 years real-life experience with belimumab, with results confirming good control of disease activity, reduction of flares and halting of damage accrual.^{111 112} The two drugs

have a different mechanism of action and have not been directly compared. Two indirect treatment comparison studies (each supported by each one of the manufacturing companies and using a different patient database) reported conflicting results at low levels of evidence and high risk of bias and, therefore, cannot be interpreted as comparative efficacy studies.^{113 114} Thus, with current evidence, the two drugs are recommended with no hierarchy between them. Of note, both drugs seem to have better efficacy in serologically active patients at baseline, although this should not limit their use to this subset of patients.^{115 116}

Undoubtedly, the most expected outcome of the current update was the verdict regarding the positioning of belimumab and voclosporin in the treatment of LN. The rationale behind the phrasing that early combination therapy with either belimumab or a CNI 'should be considered' reflects the fact that worldwide treatment recommendations for SLE should take into account different patient characteristics, but also variable access to drugs in high-income versus low-income countries. If an early combination therapy is chosen, specific patient characteristics may favour belimumab over a CNI or vice versa, for instance, the presence of extrarenal disease activity for belimumab, or nephrotic-range proteinuria for CNI. Importantly, an update of dedicated EULAR recommendations for the management of LN is currently being scheduled.

Patients with SLE experience a wide variety of symptoms, which extend beyond the classical manifestations that require immunosuppressive therapy. Indeed, symptoms such as fatigue, non-inflammatory pain, mood disturbance and cognitive dysfunction are among the ones most frequently referred and valued by patients. A recently proposed system categorised symptoms of SLE in two types: the typical inflammatory symptoms ('type 1') requiring immunosuppression, and symptoms such as those mentioned above ('type 2'), which do not respond to immunosuppressive therapy, yet often dominate patient-reported outcomes.¹¹⁷ We acknowledge that the current recommendations mainly address classic inflammatory SLE manifestations, in part because the Task Force felt that the data on type 2 symptoms are not so robust to justify specific management recommendations. Nevertheless, a holistic care of patients with SLE should value and address all symptoms mentioned by patients, both those requiring immunosuppressive therapy, as well as those in need of complimentary approaches.¹¹⁸

A crucial point pertaining to any set of management recommendations is their implementation in real-life clinical practice, often not self-explanatory because recommendations by definition cannot capture all aspects of everyday clinical practice. To tackle this need, the EULAR SOPs suggest the definition of quality indicators in tabular form, at least for the most relevant recommendations, to serve as a checklist for treating physicians and facilitate rate of adherence after a reasonable period of time. Importantly, following the issue of the 2019 EULAR recommendations for the management of SLE, such a set of quality indicators was published,¹¹⁹ and subsequently tested independently in relation to quality of life of patients.¹²⁰ A similar initiative for the current recommendations would be valuable for their wider dissemination and implementation.

In conclusion, the 2023 recommendations for the management of SLE provide current state-of-the-art guidance for treating physicians around the world. Further issues for the future research agenda in SLE are shown in [box 1](#). This updated version will inform rheumatologists and nephrologists, health professionals, patients, regulators, payers and other stakeholders on the way modern treatment of SLE is perceived from experts in the field spanning four continents. It is the hope of this

Task Force that the developments in the treatment of SLE will continue at such pace to mandate a new update of the EULAR recommendations within the next 3 years.

Author affiliations

- ¹Rheumatology and Clinical Immunology Unit, "Attikon" University Hospital, National and Kapodistrian University of Athens, Athens, Greece
- ²Lupus Europe, Copenhagen, Denmark
- ³Division of Rheumatology, Department of Medicine III, University Medical Center & Faculty of Medicine Carl Gustav Carus at the TU Dresden, Dresden, Germany
- ⁴Department of Rheumatology, Hôpitaux Universitaires de Strasbourg, INSERM UMR-S 1109, Centre National de Référence des Maladies Auto-immunes Systémiques Rares (RESO), Strasbourg, France
- ⁵Department of Rheumatology, Hanyang University Hospital for Rheumatic Diseases, Hanyang University Institute for Rheumatology Research and Hanyang Institute of Bioscience and Biotechnology, Seoul, South Korea
- ⁶Department of Nephrology and Renal Transplantation Unit, "Laiko" General Hospital, Medical School, National and Kapodistrian University of Athens, Athens, Greece
- ⁷Centre for Epidemiology Versus Arthritis, University of Manchester, Manchester, UK; National Institute for Health Research Manchester Biomedical Research Centre, Manchester University Hospitals NHS Foundation Trust, Manchester Academic Health Science Centre, Manchester, UK
- ⁸Department of Autoimmune Diseases, Hospital Clinic, Barcelona, Spain
- ⁹Rheumatology Unit, Department of Medicine, University of Padova, Padova, Italy
- ¹⁰Department of Rheumatology and Clinical Immunology, Charité Universitätsmedizin Berlin; Deutsches Rheumaforschungszentrum, Berlin, Germany
- ¹¹Division of Rheumatology, Northwell Health, Great Neck, New York City, New York, USA
- ¹²Lupus Program, Centre for Prognosis Studies in the Rheumatic Disease, Schroeder Arthritis Institute, Krembil Research Institute, Toronto Western Hospital, University of Toronto, Toronto, Ontario, Canada
- ¹³Service de Rhumatologie, Cliniques Universitaires Saint-Luc and Institut de Recherche Expérimentale et Clinique, Université catholique de Louvain, Brussels, Belgium
- ¹⁴Department of Rheumatology, Centro Hospitalar e Universitario de Coimbra, Coimbra, Portugal; School of Health Sciences, Universidade da Beira Interior, Covilhã, Portugal
- ¹⁵Department of Medicine, University of Cambridge, Cambridge, UK
- ¹⁶Cyprus League Against Rheumatism, Aglantzia, Cyprus
- ¹⁷Department of Rheumatology and Immunology, Faculty of Medicine, University of Szeged, Hungary
- ¹⁸Department of Medicine, Tuen Mun Hospital, Hong Kong, China
- ¹⁹Centre for Inflammatory Diseases, School of Clinical Sciences, Monash University, Melbourne, Victoria, Australia
- ²⁰Department of Biomedical Sciences Humanitas University, Nephrology and Dialysis Division, IRCCS Humanitas Research Hospital, Milan, Italy
- ²¹Rheumatology Unit, Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy
- ²²Department of Rheumatology & Hiller Research Unit Rheumatology, UKD, Heinrich-Heine University, Düsseldorf, Germany
- ²³Vasculitis Service, Rheumatology Department, Norfolk and Norwich University Hospitals NHS Foundation Trust, Norwich, UK
- ²⁴Hospital of the Hospitaller Order of Saint John of God, Budapest, Hungary
- ²⁵Department of Rheumatology and Clinical Immunology, Department of Internal Medicine and Oncology, Semmelweis University, Budapest, Hungary
- ²⁶Heart and Vascular Center, Semmelweis University, Budapest, Hungary
- ²⁷Section of Rheumatology, Department of Medicine, University of Santo Tomas, Manila, Philippines
- ²⁸Division of Rheumatology, Department of Medicine Solna, Karolinska Institutet, Stockholm, Sweden
- ²⁹Department of Gastroenterology, Dermatology and Rheumatology, Karolinska University Hospital, Stockholm, Sweden
- ³⁰Department of Rheumatology, Faculty of Medicine and Health, Örebro University, Örebro, Sweden
- ³¹Rheumatology Department, Complejo Hospitalario Universitario de Vigo, IRIDIS (Investigation in Rheumatology and Immune-Mediated Diseases) - VIGO Group, Galicia Sur Health Research Institute, Vigo, Spain
- ³²Division of Rheumatology, Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA
- ³³Grupo Oroño, Centro Regional de Enfermedades Autoinmunes y Reumáticas (GO-CREAR), Rosario, Argentina
- ³⁴Division of Rheumatology, Department of Medicine 3, Medical University of Vienna, Vienna, Austria
- ³⁵First Department of Internal Medicine, School of Medicine, University of Occupational and Environmental Health, Kitakyushu, Japan

- ³⁶Rheumatology Unit, First Department of Propaedeutic Internal Medicine, "Laiko" General Hospital, Medical School, National and Kapodistrian University of Athens, Joint Academic Rheumatology Program, Athens, Greece
- ³⁷Centre of Expertise for Lupus-, Vasculitis- and Complement-mediated Systemic autoimmune diseases, Department of Internal Medicine – section Nephrology, Leiden University Medical Center, Leiden, The Netherlands
- ³⁸Rheumatology and Clinical Immunology, Department of Clinical and Experimental Sciences, ASST Spedali Civili and University of Brescia, Brescia, Italy
- ³⁹Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, Leeds, UK
- ⁴⁰Department of Rheumatology and Clinical Immunology, Amsterdam University Medical Centers, Amsterdam, The Netherlands
- ⁴¹Department of Rheumatology, King's College Hospital, London, UK
- ⁴²Rheumatology, Clinical Immunology and Allergy, University Hospital of Heraklion, Greece, University Hospital of Heraklion, Heraklion, Greece
- ⁴³Laboratory of Autoimmunity and Inflammation, Biomedical Research Foundation of the Academy of Athens, Athens, Greece
- ⁴⁴Joint Academic Rheumatology Program, Medical School, National and Kapodistrian University of Athens, Greece, Medical School, University of Cyprus, Nicosia, Cyprus
- Twitter** Chetan B Mukhtyar @cmukhtyar, Edward M Vital @edvital, Chris Wincup @chrswincup, George Bertias @george_bertias and Dimitrios T Boumpas @none

Acknowledgements The committee wishes to acknowledge the support of the EULAR Standing Committee on Clinical Affairs. The committee also expresses its sincere appreciation and gratitude to the EULAR Secretariat, especially Simona Lupatin, executive assistant, and to Dora Togia for the outstanding organisation and coordination.

Contributors AF and MK reviewed the literature and AF drafted the manuscript. GB and CBM supervised the methodology. All authors edited the manuscript. DTB supervised the project.

Funding European League against Rheumatism (project number QoC015).

Competing interests AF reports honoraria and/or consulting fees from Lilly, Boehringer, Novartis, Abbvie, Astra Zeneca, GSK, MSD, Pfizer, UCB, Amgen, Aenorasis, support for attending meetings from UCB. MK reports honoraria and/or consulting fees from GSK, participation in advisory boards from GSK, Astra Zeneca, Amgen. JA reports honoraria and/or consulting fees from Novartis, Astra Zeneca, support for attending meetings from Novartis, Astra-Zeneca, participation in advisory boards from Roche, Astra-Zeneca (all paid to Lupus Europe). MA reports honoraria and/or consulting fees from AbbVie, Astra Zeneca, GSK, Otsuka, Roche, support for attending meetings from Pfizer. LA reports grants from GSK, honoraria and/or consulting fees from Alexion, Alpine, Amgen, Astra Zeneca, AbbVie, Biogen, BMS, Boehringer-Ingelheim, Chugai, GSK, Grifols, Idorsia, Janssen, Kezar, Lilly, Medac, Novartis, Ono pharmaceuticals, Pfizer, Roche, UCB, support for attending meetings from Novartis, Astra-Zeneca. INB reports grants from GSK, Janssen, honoraria and/or consulting fees from Astra Zeneca, GSK, Eli Lilly, UCB, MSD, participation in advisory boards from Aurinia, Astra Zeneca, ILTOO. RC reports honoraria, consulting fees and/or participation in advisory boards from AstraZeneca, Celgene, GSK, Janssen, Lilly, Pfizer, UCB, Rubió, Werfen. AD reports honoraria and consulting fees from Otsuka, Astra Zeneca, GSK, Lilly. TD reports consulting fees from Roche, Novartis, GSK, Astra Zeneca, participation in advisory boards from GNE, Roche, Novartis. DDG reports grants from Abbvie, Amgen, BMS, Eli Lilly, Janssen, Novartis, Pfizer, UCB, consulting fees from Abbvie, Amgen, Astra Zeneca, BMS, Eli Lilly, Galapagos, Gilead, Janssen, Novartis, Pfizer, UCB. FAH reports grants from GSK, Roche. DJ reports grants from GSK, Roche, CSL Vifor, honoraria and consulting fees from Astra Zeneca, Chemocentryx, GSK, Novartis, Takeda, CSL Vifor, participation in advisory boards from Chinook, GSK, stocks in Aurinia. LK reports honoraria fees from Janssen, Takeda, support for attending meetings from Abbvie, Astra Zeneca, CSL Behring, participation in advisory boards from Swixx Pharma, Roche, Astra Zeneca. CCM reports honoraria fees from GSK, support for attending meetings from Novartis. EFM reports grants from AbbVie, Amgen, AstraZeneca, Biogen, Bristol Myers Squibb, Eli Lilly and Company, EMD Serono, Genentech, GlaxoSmithKline, Janssen, and UCB, honoraria and/or consulting fees from AbbVie, Astra Zeneca, Biogen, Bristol Myers Squibb, Eli Lilly and Company, EMD Serono, Galapagos, Genentech, Gilead, GlaxoSmithKline, IGM, Janssen, Novartis, Servier, Wolf, Zenas, support for attending meetings from Astra Zeneca, Roche. GM reports honoraria fees from GSK, Roche, Otsuka, participation in advisory boards from GSK, Otsuka. MM reports honoraria and/or consulting fees from Astra Zeneca, Eli Lilly, GSK, UCB, Janssen, participation in advisory boards from Idorsia. JM reports grants from GSK, honoraria and/or consulting fees from GSK, Astra Zeneca, Otsuka, BMS, participation in advisory boards from GSK, Astra Zeneca, BMS. GN reports honoraria and/or consulting fees from Sobi, GSK, Miltenyi Biotech, Swixx Biopharma, AbbVie, Lilly, Richter, Amgen, Roche, Kedrion, support for attending meetings from Roche, AbbVie, Sobi, Biotech. SVN reports honoraria and/or consulting fees from Astra Zeneca, Boehringer Ingelheim, Idorsia, Astellas, Novartis, GSK, Roche, support for attending meetings from Pfizer, participation in advisory boards from Biogen. IP reports grants from Aurinia, BMS, GSK, Otsuka, Roche, honoraria and/or consulting fees from Astra

Zeneca, GSK, Janssen, Novartis, Otsuka, Roche, participation in advisory boards from Astra Zeneca, GSK, Novartis, Otsuka, medical writing from Astra Zeneca, GSK. JMPR reports grants from GSK, Pfizer, honoraria fees from GSK, Astra Zeneca, Lilly, support for attending meetings from GSK, Astra Zeneca, participation in advisory boards from GSK, Otsuka, Gebro, Astra Zeneca, Boehringer-Ingelheim, MSD. BAPE reports grants from Janssen, support for attending meetings from Astra Zeneca, participation in advisory boards from Astra Zeneca, GSK. MS reports grants from GSK, Astra Zeneca, honoraria and/or consulting fees from Astra Zeneca, GSK, Otsuka, UCB, participation in advisory boards from GSK. JSS reports grants from AbbVie, Astra Zeneca, Lilly, Galapagos, royalties or licenses from Elsevier, honoraria and/or consulting fees from AbbVie, Galapagos/Gilead, Novartis, BMS, Samsung, Sanofi, Chugai, R-Pharma, Lilly, MSD, Janssen, participation in advisory boards from Astra Zeneca. ES reports stocks in Astra Zeneca. YKOT reports grants from Mitsubishi-Tanabe, Eisai, Chugai, Taisho, consulting fees from Lilly, Astra Zeneca, AbbVie, Gilead, Chugai, Boehringer-Ingelheim, GSK, Eisai, Taisho, BMS, Pfizer, Taiho. MGT reports grants from Genesis, GSK, MSD, UCB, consulting fees from GSK, Lilly and UCB. CW reports grants from Versus Arthritis, British Society for Rheumatology, Lupus UK, honoraria fees from UCB, support for attending meetings from AbbVie. AT reports honoraria fees from UCB, GSK, participation in advisory boards from UCB, Galapagos. GB reports grants from GSK, Astra Zeneca, Pfizer, honoraria and/or consulting fees from Lilly, Aenorasis, Novartis, AstraZeneca, GSK, SOBI, Pfizer, participation in advisory boards from Novartis. DTB reports unrestricted investigational grants from GSK, honoraria and/or consulting fees from GSK, Astra-Zeneca, Pfizer. The remaining authors declare no conflict of interest.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

ORCID iDs

Antonis Fanouriakis <http://orcid.org/0000-0003-2696-031X>
 Myrto Kostopoulou <http://orcid.org/0000-0002-8832-7475>
 Martin Aringer <http://orcid.org/0000-0003-4471-8375>
 Laurent Arnaud <http://orcid.org/0000-0002-8077-8394>
 Sang-Cheol Bae <http://orcid.org/0000-0003-4658-1093>
 Andrea Doria <http://orcid.org/0000-0003-0548-4983>
 Thomas Dörner <http://orcid.org/0000-0002-6478-7725>
 Richard A Furie <http://orcid.org/0000-0001-6712-1585>
 Dafna D Gladman <http://orcid.org/0000-0002-9074-0592>
 Frederic A Houssiau <http://orcid.org/0000-0003-1451-083X>
 Luis Sousa Inês <http://orcid.org/0000-0003-3172-3570>
 David Jayne <http://orcid.org/0000-0002-1712-0637>
 Chi Chiu Mok <http://orcid.org/0000-0003-3696-1228>
 Eric F Morand <http://orcid.org/0000-0002-9507-3338>
 Johanna Mucke <http://orcid.org/0000-0001-8915-7837>
 Chetan B Mukhtyar <http://orcid.org/0000-0002-9771-6667>
 György Nagy <http://orcid.org/0000-0003-1198-3228>
 Ioannis Parodis <http://orcid.org/0000-0002-4875-5395>
 Michelle Petri <http://orcid.org/0000-0003-1441-5373>
 Elisabet Svenungsson <http://orcid.org/0000-0003-3396-3244>
 Yoshiya Tanaka <http://orcid.org/0000-0002-0807-7139>
 Maria G Tektonidou <http://orcid.org/0000-0003-2238-0975>
 YK Onno Teng <http://orcid.org/0000-0001-9920-2195>
 Angela Tincani <http://orcid.org/0000-0003-4355-9333>
 Edward M Vital <http://orcid.org/0000-0003-1637-4755>
 Ronald F van Vollenhoven <http://orcid.org/0000-0001-6438-8663>
 Chris Wincup <http://orcid.org/0000-0002-8742-8311>
 George Bertias <http://orcid.org/0000-0001-5299-1406>
 Dimitrios T Boumpas <http://orcid.org/0000-0002-9812-4671>

REFERENCES

- Bertias G, Ioannidis JPA, Boletis J, et al. EULAR recommendations for the management of systemic lupus erythematosus. report of a task force of the EULAR standing committee for international clinical studies including therapeutics. *Ann Rheum Dis* 2008;67:195–205.
- Bertias GK, Ioannidis JPA, Aringer M, et al. EULAR recommendations for the management of systemic lupus erythematosus with neuropsychiatric manifestations:

- report of a task force of the EULAR standing committee for clinical affairs. *Ann Rheum Dis* 2010;69:2074–82.
- 3 Bertsias GK, Tektonidou M, Amoura Z, et al. Joint European league against rheumatism and European renal association-European dialysis and transplant association (EULAR/ERA-EDTA) recommendations for the management of adult and paediatric lupus nephritis. *Ann Rheum Dis* 2012;71:1771–82.
 - 4 Andreoli L, Bertsias GK, Agmon-Levin N, et al. EULAR recommendations for women's health and the management of family planning, assisted reproduction, pregnancy and menopause in patients with systemic lupus erythematosus and/or antiphospholipid syndrome. *Ann Rheum Dis* 2017;76:476–85.
 - 5 Mosca M, Tani C, Aringer M, et al. European league against rheumatism recommendations for monitoring patients with systemic lupus erythematosus in clinical practice and in observational studies. *Ann Rheum Dis* 2010;69:1269–74.
 - 6 Parodis I, Houssiau FA. From sequential to combination and personalised therapy in lupus nephritis: moving towards a paradigm shift. *Ann Rheum Dis* 2022;81:15–9.
 - 7 Kostopoulou M, Fanouriakis A, Bertsias G, et al. Treatment of lupus: more options after a long wait. *Ann Rheum Dis* 2022;81:753–6.
 - 8 Smolen JS, Landewé RBM, Bergstra SA, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2022 update. *Ann Rheum Dis* 2023;82:3–18.
 - 9 Ramiro S, Nikiforou E, Sepiano A, et al. ASAS-EULAR recommendations for the management of axial spondyloarthritis: 2022 update. *Ann Rheum Dis* 2023;82:19–34.
 - 10 van der Heijde D, Aletaha D, Carmona L, et al. Update of the EULAR standardised operating procedures for EULAR-endorsed recommendations. *Ann Rheum Dis* 2015;74:8–13.
 - 11 Brouwers MC, Kho ME, Browman GP, et al. AGREE II: advancing guideline development, reporting and evaluation in health care. *CMAJ* 2010;182:E839–42.
 - 12 Holwick J, Chalmers I, Galsziou P, et al. Explanation of the 2011 Oxford centre for evidence-based medicine (OCEBM) levels of evidence (background document). In: *Oxford Centre for Evidence-Based Medicine*. 2011.
 - 13 Kostopoulou M, Fanouriakis A, Cheema K, et al. Management of lupus nephritis: a systematic literature review informing the 2019 update of the joint EULAR and European renal association-European dialysis and transplant association (EULAR/ERA-EDTA) recommendations. *RMD Open* 2020;6:e001263.
 - 14 Chasset F, Francès C, Barete S, et al. Influence of smoking on the efficacy of antimalarials in cutaneous lupus: a meta-analysis of the literature. *J Am Acad Dermatol* 2015;72:634–9.
 - 15 Parodis I, Sjöwall C, Jönsen A, et al. Smoking and pre-existing organ damage reduce the efficacy of Belimumab in systemic lupus erythematosus. *Autoimmunity Reviews* 2017;16:343–51.
 - 16 Parodis I, Girard-Guyonvarc'h C, Arnaud L, et al. EULAR recommendations for the non-pharmacological management of systemic lupus erythematosus and systemic sclerosis. *Ann Rheum Dis* 2023;ard-2023-224416.
 - 17 Isenberg D, Appel GB, Contreras G, et al. Influence of race/Ethnicity on response to lupus nephritis treatment: the ALMS study. *Rheumatology (Oxford)* 2010;49:128–40.
 - 18 Aringer M, Costenbader K, Daikh D, et al. European league against rheumatism/ American college of rheumatology classification criteria for systemic lupus erythematosus. *Ann Rheum Dis* 2019;78:1151–9.
 - 19 Kapsala NN, Nikolopoulos DS, Flouda SP, et al. From first symptoms to diagnosis of systemic lupus erythematosus: mapping the journey of patients in an observational study. *Clin Exp Rheumatol* 2023;41:74–81.
 - 20 Kerder A, Richter JG, Fischer-Betz R, et al. Delayed diagnosis adversely affects outcome in systemic lupus erythematosus: cross sectional analysis of the Lula cohort. *Lupus* 2021;30:431–8.
 - 21 van Vollenhoven RF, Bertsias G, Doria A, et al. DORIS definition of remission in SLE: final recommendations from an international task force. *Lupus Sci Med* 2021;8:e000538.
 - 22 Franklin K, Lau CS, Navarra SV, et al. Definition and initial validation of a lupus low disease activity state (LLDAS). *Ann Rheum Dis* 2016;75:1615–21.
 - 23 Costedoat-Chalumeau N, Pouchot J, Guettrot-Imbert G, et al. Adherence to treatment in systemic lupus erythematosus patients. *Best Pract Res Clin Rheumatol* 2013;27:329–40.
 - 24 Melles RB, Marmor MF. The risk of toxic retinopathy in patients on long-term hydroxychloroquine therapy. *JAMA Ophthalmol* 2014;132:1453–60.
 - 25 Jorge AM, Mancini C, Zhou B, et al. Hydroxychloroquine dose per ophthalmology guidelines and the risk of systemic lupus erythematosus flares. *JAMA* 2022;328:1458–60.
 - 26 Garg S, Unnithan R, Hansen KE, et al. Clinical significance of monitoring hydroxychloroquine levels in patients with systemic lupus erythematosus: a systematic review and meta-analysis. *Arthritis Care Res (Hoboken)* 2021;73:707–16.
 - 27 Marmor MF, Kellner U, Lai YYY, et al. Recommendations on screening for chloroquine and hydroxychloroquine retinopathy (2016 revision). *Ophthalmology* 2016;123:1386–94.
 - 28 Stahn C, Buttgerit F. Genomic and nongenomic effects of glucocorticoids. *Nat Rev Rheumatol* 2008;4:525–33.
 - 29 Ruiz-Irastorza G, Ugarte A, Saint-Pastou Terrier C, et al. Repeated pulses of methylprednisolone with reduced doses of prednisone improve the outcome of class III, IV and V lupus nephritis: an observational comparative study of the lupus-cruces and lupus-Bordeaux cohorts. *Autoimmun Rev* 2017;16:826–32.
 - 30 Kallas R, Li J, Petri M. Predictors of osteonecrosis in systemic lupus erythematosus: a prospective cohort study. *Arthritis Care Res (Hoboken)* 2022;74:1122–32.
 - 31 Tselios K, Gladman DD, Al-Sheikh H, et al. Medium versus high initial prednisone dose for remission induction in lupus nephritis: a propensity score-matched analysis. *Arthritis Care Res (Hoboken)* 2022;74:1451–8.
 - 32 Ruiz-Arruza I, Barbosa C, Ugarte A, et al. Comparison of high versus low-medium prednisone doses for the treatment of systemic lupus erythematosus patients with high activity at diagnosis. *Autoimmun Rev* 2015;14:875–9.
 - 33 Furie RA, Morand EF, Bruce IN, et al. Type I interferon inhibitor anifrolumab in active systemic lupus erythematosus (TULIP-1): a randomised, controlled, phase 3 trial. *Lancet Rheumatol* 2019;1:e208–19.
 - 34 Morand EF, Furie R, Tanaka Y, et al. Trial of anifrolumab in active systemic lupus erythematosus. *N Engl J Med* 2020;382:211–21.
 - 35 Singh JA, Shah NP, Mudano AS. Belimumab for systemic lupus erythematosus. *Cochrane Database Syst Rev* 2021;2:CD010668.
 - 36 Tam L-S, Li EK, Wong C-K, et al. Double-blind, randomized, placebo-controlled pilot study of leflunomide in systemic lupus erythematosus. *Lupus* 2004;13:601–4.
 - 37 Alshaiki F, Obaid E, Almuallim A, et al. Outcomes of rituximab therapy in refractory lupus: a meta-analysis. *Eur J Rheumatol* 2018;5:118–26.
 - 38 Jónsdóttir T, Gunnarsson I, Risselada A, et al. Treatment of refractory SLE with rituximab plus cyclophosphamide: clinical effects, serological changes, and predictors of response. *Ann Rheum Dis* 2008;67:330–4.
 - 39 Li EK, Tam L-S, Zhu TY, et al. Is combination rituximab with cyclophosphamide better than rituximab alone in the treatment of lupus nephritis. *Rheumatology (Oxford)* 2009;48:892–8.
 - 40 Kraaij T, Kamerling SWA, de Rooij ENM, et al. The NET-effect of combining rituximab with belimumab in severe systemic lupus erythematosus. *J Autoimmun* 2018;91:45–54.
 - 41 Atisha-Fregoso Y, Malkiel S, Harris KM, et al. Phase II randomized trial of Rituximab plus cyclophosphamide followed by belimumab for the treatment of lupus nephritis. *Arthritis Rheumatol* 2021;73:121–31.
 - 42 Kronbichler A, Brezina B, Quintana LF, et al. Efficacy of plasma exchange and immunoadsorption in systemic lupus erythematosus and antiphospholipid syndrome: a systematic review. *Autoimmun Rev* 2016;15:38–49.
 - 43 Leone A, Radin M, Almarzooqi AM, et al. Autologous hematopoietic stem cell transplantation in systemic lupus erythematosus and antiphospholipid syndrome: a systematic review. *Autoimmun Rev* 2017;16:469–77.
 - 44 Mackensen A, Müller F, Mougiakakos D, et al. Anti-CD19 CAR T cell therapy for refractory systemic lupus erythematosus. *Nat Med* 2022;28:2124–32.
 - 45 Chasset F, Arnaud L, Jachiet M, et al. Changing antimalarial agents after inefficacy or intolerance in patients with cutaneous lupus erythematosus: a multicenter observational study. *J Am Acad Dermatol* 2018;78:107–14.
 - 46 Ugarte A, Porta S, Ríos R, et al. Combined mepacrine-hydroxychloroquine treatment in patients with systemic lupus erythematosus and refractory cutaneous and articular activity. *Lupus* 2018;27:1718–22.
 - 47 Chasset F, Bouaziz J-D, Costedoat-Chalumeau N, et al. Efficacy and comparison of antimalarials in cutaneous lupus erythematosus subtypes: a systematic review and meta-analysis. *Br J Dermatol* 2017;177:188–96.
 - 48 Keyes E, Jobanputra A, Feng R, et al. Comparative responsiveness of cutaneous lupus erythematosus patients to methotrexate and mycophenolate mofetil: a cohort study. *J Am Acad Dermatol* 2022;87:447–8.
 - 49 Morand EF, Furie RA, Bruce IN, et al. Efficacy of anifrolumab across organ domains in patients with moderate-to-severe systemic lupus erythematosus: a post-hoc analysis of pooled data from the TULIP-1 and TULIP-2 trials. *Lancet Rheumatol* 2022;4:e282–92.
 - 50 Kneeland R, Montes D, Endo J, et al. Improvement in cutaneous lupus erythematosus after twenty weeks of Belimumab use: a systematic review and meta-analysis. *Arthritis Care Res (Hoboken)* 2023;75:1838–48.
 - 51 Chasset F, Tounsi T, Cesbron E, et al. Efficacy and tolerance profile of thalidomide in cutaneous lupus erythematosus: a systematic review and meta-analysis. *J Am Acad Dermatol* 2018;78:342–50.
 - 52 Aitmehdi R, Arnaud L, Francès C, et al. Long-term efficacy and safety outcomes of lenalidomide for cutaneous lupus erythematosus: a multicenter retrospective observational study of 40 patients. *J Am Acad Dermatol* 2021;84:1171–4.
 - 53 Govoni M, Hanly JG. The management of neuropsychiatric lupus in the 21st century: still so many unmet needs. *Rheumatology (Oxford)* 2020;59:v52–62.
 - 54 Bortoluzzi A, Scirè CA, Bombardieri S, et al. Development and validation of a new algorithm for attribution of neuropsychiatric events in systemic lupus erythematosus. *Rheumatology (Oxford)* 2015;54:891–8.
 - 55 Hanly JG, Su L, Farewell V, et al. Prospective study of neuropsychiatric events in systemic lupus erythematosus. *J Rheumatol* 2009;36:1449–59.
 - 56 Narváez J, Ríos-Rodríguez V, de la Fuente D, et al. Rituximab therapy in refractory neuropsychiatric lupus: current clinical evidence. *Semin Arthritis Rheum* 2011;41:364–72.

- 57 Serris A, Amoura Z, Canoui-Poitine F, *et al.* Efficacy and safety of rituximab for systemic lupus erythematosus-associated immune cytopenias: a multicenter retrospective cohort study of 71 adults. *Am J Hematol* 2018;93:424–9.
- 58 Moysidou G-S, Garantziotis P, Nikolopoulos D, *et al.* Relapses are common in severe hematologic systemic lupus erythematosus and may be prevented by early institution of immunosuppressive agents: a real-life single-center study. *Lupus* 2023;32:225–30.
- 59 Bradbury CA, Pell J, Hill Q, *et al.* Mycophenolate mofetil for first-line treatment of immune thrombocytopenia. *N Engl J Med* 2021;385:885–95.
- 60 Roussotte M, Gerfaud-Valentin M, Hot A, *et al.* Immune thrombocytopenia with clinical significance in systemic lupus erythematosus: a retrospective cohort study of 90 patients. *Rheumatology (Oxford)* 2022;61:3627–39.
- 61 Catalá-López F, Corrales I, de la Fuente-Honrubia C, *et al.* Risk of thromboembolism with thrombopoietin receptor agonists in adult patients with thrombocytopenia: systematic review and meta-analysis of randomized controlled trials. *Med Clin (Barc)* 2015;145:511–9.
- 62 Houssiau FA, Vasconcelos C, D'Cruz D, *et al.* Immunosuppressive therapy in lupus nephritis: the Euro-lupus nephritis trial, a randomized trial of low-dose versus high-dose intravenous cyclophosphamide. *Arthritis Rheum* 2002;46:2121–31.
- 63 Anders H-J, Saxena R, Zhao M-H, *et al.* Lupus nephritis. *Nat Rev Dis Primers* 2020;6:7.
- 64 Furie R, Rovin BH, Houssiau F, *et al.* Two-year, randomized, controlled trial of belimumab in lupus nephritis. *N Engl J Med* 2020;383:1117–28.
- 65 Rovin BH, Teng YKO, Ginzler EM, *et al.* Efficacy and safety of voclosporin versus placebo for lupus nephritis (AURORA 1): a double-blind, randomised, multicentre, placebo-controlled, phase 3 trial. *Lancet* 2021;397:2070–80.
- 66 Rovin BH, Furie R, Teng YKO, *et al.* A secondary analysis of the belimumab international study in lupus nephritis trial examined effects of belimumab on kidney outcomes and preservation of kidney function in patients with lupus nephritis. *Kidney Int* 2022;101:403–13.
- 67 Furie R, Rovin BH, Houssiau F, *et al.* Safety and efficacy of belimumab in patients with lupus nephritis: open-label extension of BLISS-LN study. *Clin J Am Soc Nephrol* 2022;17:1620–30.
- 68 Luís MSF, Bultink IEM, da Silva JAP, *et al.* Early predictors of renal outcome in patients with proliferative lupus nephritis: a 36-month cohort study. *Rheumatology (Oxford)* 2021;60:5134–41.
- 69 Kapsia E, Marinaki S, Michelakis I, *et al.* Predictors of early response, flares, and long-term adverse renal outcomes in proliferative lupus nephritis: a 100-month median follow-up of an inception cohort. *J Clin Med* 2022;11:5017.
- 70 Zhou T, Zhang X, Lin W, *et al.* Multitarget therapy: an effective and safe therapeutic regimen for lupus nephritis. *J Pharm Pharm Sci* 2019;22:365–75.
- 71 Arriens C, Teng YKO, Ginzler EM, *et al.* Update on the efficacy and safety profile of voclosporin: an integrated analysis of clinical trials in lupus nephritis. *Arthritis Care Res (Hoboken)* 2023;75:1399–408.
- 72 Fanouriakis A, Kostopoulou M, Cheema K, *et al.* Update of the joint European league against rheumatism and European renal association–European dialysis and transplant association (EULAR/ERA–EDTA) recommendations for the management of lupus nephritis. *Ann Rheum Dis* 2020;79:713–23.
- 73 Tai S, Anumolu N, Putman M. Challenging wisely: how to move beyond ‘1 mg/kg then taper’ *Rheumatology (Oxford)* 2022;62:3–6.
- 74 Sciascia S, Yazdany J, Dall'Era M, *et al.* Anticoagulation in patients with concomitant lupus nephritis and thrombotic microangiopathy: a multicentre cohort study. *Ann Rheum Dis* 2019;78:1004–6.
- 75 Wright RD, Bannerman F, Beresford MW, *et al.* A systematic review of the role of eculizumab in systemic lupus erythematosus-associated thrombotic microangiopathy. *BMC Nephrol* 2020;21:245.
- 76 Moroni G, Gatto M, Tamborini F, *et al.* Lack of EULAR/ERA–EDTA response at 1 year predicts poor long-term renal outcome in patients with lupus nephritis. *Ann Rheum Dis* 2020;79:1077–83.
- 77 Jourde-Chiche N, Costedoat-Chalumeau N, Baumstarck K, *et al.* Weaning of maintenance immunosuppressive therapy in lupus nephritis (WIN-lupus): results of a multicentre randomised controlled trial. *Ann Rheum Dis* 2022;81:1420–7.
- 78 Liu Z, Zhang H, Liu Z, *et al.* Multitarget therapy for induction treatment of lupus nephritis: a randomized trial. *Ann Intern Med* 2015;162:18–26.
- 79 Mok CC, Ho LY, Ying SKY, *et al.* Long-term outcome of a randomised controlled trial comparing tacrolimus with mycophenolate mofetil as induction therapy for active lupus nephritis. *Ann Rheum Dis* 2020;79:1070–6.
- 80 Saxena A, Ginzler EM, Gibson K, *et al.* Safety and efficacy of long-term Voclosporin treatment for lupus nephritis in the phase 3 AURORA 2 clinical trial. *Arthritis Rheumatol* 2023.
- 81 Walsh M, Solomons N, Lisk L, *et al.* Mycophenolate mofetil or intravenous cyclophosphamide for lupus nephritis with poor kidney function: a subgroup analysis of the Aspreva lupus management study. *Am J Kidney Dis* 2013;61:710–5.
- 82 Ji L, Xie W, Zhang Z. Low-dose glucocorticoids should be withdrawn or continued in systemic lupus erythematosus? A systematic review and meta-analysis on risk of flare and damage accrual. *Rheumatology (Oxford)* 2021;60:5517–26.
- 83 Mathian A, Pha M, Haroche J, *et al.* Withdrawal of low-dose prednisone in SLE patients with a clinically quiescent disease for more than 1 year: a randomised clinical trial. *Ann Rheum Dis* 2020;79:339–46.
- 84 Tselios K, Gladman DD, Su J, *et al.* Gradual glucocorticosteroid withdrawal is safe in clinically quiescent systemic lupus erythematosus. *ACR Open Rheumatol* 2021;3:550–7.
- 85 Fasano S, Coscia MA, Pierro L, *et al.* Which patients with systemic lupus erythematosus in remission can withdraw low dose steroids? results from a single inception cohort study. *Lupus* 2021;30:991–7.
- 86 Zen M, Fuzzi E, Loreda Martinez M, *et al.* Immunosuppressive therapy withdrawal after remission achievement in patients with lupus nephritis. *Rheumatology* 2022;61:688–95.
- 87 Moroni G, Longhi S, Giglio E, *et al.* What happens after complete withdrawal of therapy in patients with lupus nephritis. *Clin Exp Rheumatol* 2013;31:S75–81.
- 88 De Rosa M, Azzato F, Tobli JE, *et al.* A prospective observational cohort study highlights kidney biopsy findings of lupus nephritis patients in remission who flare following withdrawal of maintenance therapy. *Kidney Int* 2018;94:788–94.
- 89 Malvar A, Alberton V, Lococo B, *et al.* Kidney biopsy-based management of maintenance immunosuppression is safe and may ameliorate flare rate in lupus nephritis. *Kidney Int* 2020;97:156–62.
- 90 Jorge A, McCormick N, Lu N, *et al.* Hydroxychloroquine and mortality among patients with systemic lupus erythematosus in the general population. *Arthritis Care Res (Hoboken)* 2021;73:1219–23.
- 91 Papachristos DA, Gladman DD, Su J, *et al.* Outcomes following antimalarial withdrawal in patients with quiescent systemic lupus erythematosus. *Semin Arthritis Rheum* 2022;55:152046.
- 92 Almeida-Brasil CC, Hanly JG, Urowitz M, *et al.* Flares after hydroxychloroquine reduction or discontinuation: results from the systemic lupus international collaborating clinics (SLICC) inception cohort. *Ann Rheum Dis* 2022;81:370–8.
- 93 Almeida-Brasil CC, Pineau CA, Vinet E, *et al.* Predictors of unsuccessful hydroxychloroquine tapering and discontinuation: can we personalize decision-making in systemic lupus erythematosus treatment. *Arthritis Care Res (Hoboken)* 2022;74:1070–8.
- 94 Ji L, Xie W, Fasano S, *et al.* Risk factors of flare in patients with systemic lupus erythematosus after glucocorticoids withdrawal. A systematic review and meta-analysis. *Lupus Sci Med* 2022;9:e000603.
- 95 Tektonidou MG, Andreoli L, Limper M, *et al.* EULAR recommendations for the management of antiphospholipid syndrome in adults. *Ann Rheum Dis* 2019;78:1296–304.
- 96 Khairani CD, Bejjani A, Piazza G, *et al.* Direct oral anticoagulants vs vitamin K antagonists in patients with antiphospholipid syndromes: meta-analysis of randomized trials. *J Am Coll Cardiol* 2023;81:16–30.
- 97 Kravvariti E, Koutsogianni A, Samoli E, *et al.* The effect of hydroxychloroquine on thrombosis prevention and antiphospholipid antibody levels in primary antiphospholipid syndrome: a pilot open label randomized prospective study. *Autoimmun Rev* 2020;19:102491.
- 98 Tektonidou MG, Laskari K, Panagiotakos DB, *et al.* Risk factors for thrombosis and primary thrombosis prevention in patients with systemic lupus erythematosus with or without antiphospholipid antibodies. *Arthritis Rheum* 2009;61:29–36.
- 99 Petri M, Konig MF, Li J, *et al.* Association of higher hydroxychloroquine blood levels with reduced thrombosis risk in systemic lupus erythematosus. *Arthritis Rheumatol* 2021;73:997–1004.
- 100 López-Benjumbe B, Rodríguez-Pintó I, Amigo CP, *et al.* Eculizumab use in catastrophic antiphospholipid syndrome (CAPS): descriptive analysis from the “CAPS registry” *Autoimmun Rev* 2022;21:103055.
- 101 Furer V, Rondaan C, Heijstek MW, *et al.* 2019 update of EULAR recommendations for vaccination in adult patients with autoimmune inflammatory rheumatic diseases. *Ann Rheum Dis* 2020;79:39–52.
- 102 Drosos GC, Vedder D, Houben E, *et al.* EULAR recommendations for cardiovascular risk management in rheumatic and musculoskeletal diseases, including systemic lupus erythematosus and antiphospholipid syndrome. *Ann Rheum Dis* 2022;81:768–79.
- 103 Kwan A, Rayes HA, Lazova T, *et al.* Herpes zoster in SLE: prevalence, incidence and risk factors. *Lupus Sci Med* 2022;9:e000574.
- 104 Mok CC, Chan KH, Ho LY, *et al.* Safety and immune response of a live-attenuated herpes Zoster vaccine in patients with systemic lupus erythematosus: a randomised placebo-controlled trial. *Ann Rheum Dis* 2019;78:1663–8.
- 105 Leung J, Anderson TC, Dooling K, *et al.* Recombinant zoster vaccine uptake and risk of flares among older adults with immune-mediated inflammatory diseases in the US. *Arthritis Rheumatol* 2022;74:1833–41.
- 106 Tan SYS, Yee AM, Sim JLL, *et al.* COVID-19 vaccination in systemic lupus erythematosus: a systematic review of its effectiveness, immunogenicity, flares and acceptance. *Rheumatology* 2023;62:1757–72.
- 107 Fragoulis GE, Nikiforou E, Dey M, *et al.* EULAR recommendations for screening and prophylaxis of chronic and opportunistic infections in adults with autoimmune inflammatory rheumatic diseases. *Ann Rheum Dis* 2023;82:742–53.
- 108 Tselios K, Gladman DD, Su J, *et al.* Impact of the new American college of cardiology/American heart association definition of hypertension on atherosclerotic vascular events in systemic lupus erythematosus. *Ann Rheum Dis* 2020;79:612–7.

- 109 Hakroush S, Tampe D, Kluge IA, *et al.* Comparative analysis of SGLT-2 expression in renal vasculitis and lupus nephritis. *Ann Rheum Dis* 2022;81:1048–50.
- 110 Wang H, Li T, Sun F, *et al.* Safety and efficacy of the SGLT2 inhibitor dapagliflozin in patients with systemic lupus erythematosus: a phase I/II trial. *RMD Open* 2022;8:e002686.
- 111 Gatto M, Saccon F, Zen M, *et al.* Early disease and low baseline damage as predictors of response to belimumab in patients with systemic lupus erythematosus in a real-life setting. *Arthritis Rheumatol* 2020;72:1314–24.
- 112 Nikoloudaki M, Nikolopoulos D, Koutsoviti S, *et al.* Clinical response trajectories and drug persistence in systemic lupus erythematosus patients on Belimumab treatment: a real-life, multicentre observational study. *Front Immunol* 2023;13:1074044.
- 113 Bruce IN, Golam S, Steenkamp J, *et al.* Indirect treatment comparison of anifrolumab efficacy versus belimumab in adults with systemic lupus erythematosus. *J Comp Eff Res* 2022;11:765–77.
- 114 Neupane B, Shukla P, Slim M, *et al.* Belimumab versus anifrolumab in adults with systemic lupus erythematosus: an indirect comparison of clinical response at 52 weeks. *Lupus Sci Med* 2023;10:e000907.
- 115 Vital EM, Merrill JT, Morand EF, *et al.* Anifrolumab efficacy and safety by type I interferon gene signature and clinical subgroups in patients with SLE: post hoc analysis of pooled data from two phase III trials. *Ann Rheum Dis* 2022;81:951–61.
- 116 van Vollenhoven RF, Petri MA, Cervera R, *et al.* Belimumab in the treatment of systemic lupus erythematosus: high disease activity predictors of response. *Ann Rheum Dis* 2012;71:1343–9.
- 117 Pisetsky DS, Clowse MEB, Criscione-Schreiber LG, *et al.* A novel system to categorize the symptoms of systemic lupus erythematosus. *Arthritis Care Res (Hoboken)* 2019;71:735–41.
- 118 Geenen R, Overman CL, Christensen R, *et al.* EULAR recommendations for the health professional's approach to pain management in inflammatory arthritis and osteoarthritis. *Ann Rheum Dis* 2018;77:797–807.
- 119 Chavatzka K, Kostopoulou M, Nikolopoulos D, *et al.* Quality indicators for systemic lupus erythematosus based on the 2019 EULAR recommendations: development and initial validation in a cohort of 220 patients. *Ann Rheum Dis* 2021;80:1175–82.
- 120 Taheri N, Mageau A, Chauveheid M-P, *et al.* Impact of adherence to EULAR quality indicators on the quality of life of patients with systemic lupus erythematosus. *Eur J Intern Med* 2023;109:68–72.

Supplementary Table 1. Recommended doses of drugs used to treat systemic lupus erythematosus

Drug	Recommended dose	Dose adjustment needed in CKD
Glucocorticoids	<i>Mild-Moderate disease:</i> If needed, start with ≤ 20 mg/day with gradual tapering <i>Severe/Organ-threatening disease:</i> Consider IV MP pulses 250-1000 mg/day for 1-3 days - Continue with PO 0.3-0.5 mg/Kg/day with tapering* <i>All circumstances:</i> Keep maintenance prednisone dose at ≤ 5 mg/day	No
Hydroxychloroquine	Target dose 5 mg/Kg/day (up to 400 mg/day) In patients in long-standing remission, consider tapering to 200 mg/day	Yes
Methotrexate	10-25 mg/week in 1-2 doses (given in one day)	Yes
Azathioprine	2-3 mg/Kg/day in 1-2 doses In patients in remission, consider tapering to < 2 mg/day	Yes
Mycophenolate mofetil (MMF)/Mycophenolic acid (MPA)[§]	<i>Severe/Organ-threatening disease or "Initial" therapy in LN:</i> MMF 2-3 g/day in 2 doses – MPA: 1.44-2.16 gr/day in 2 doses <i>"Subsequent therapy" in LN:</i> MMF 1-2 g/day in 2 doses – MPA 720-1440 mg/day in 2 doses	Yes
Leflunomide	10-20 mg/day in 1 dose	Yes
Cyclophosphamide	<i>"Initial" therapy in LN:</i> IV 500 mg on weeks 0, 2, 4, 6, 8 and 10 (Low-dose - Euro-Lupus regimen) <i>Organ- or life-threatening disease:</i> IV 0.75-1 g/m ² BSA/month for 6 months (High-dose - NIH regimen)	Yes
Cyclosporine A	1-3 mg/Kg/day or up to 400 mg/day in 2 doses	Avoid overall
Tacrolimus	0.05 to 0.1 mg/Kg/day or 2-4 mg/day in 2 doses - Titrate to target blood concentration 4-6 ng/ml 12 hours after dose	Yes
Voclosporin	23.7 mg two times per day	No
Intravenous immunoglobulin	2 g/Kg total, given in 2-5 days	No
Anifrolumab	IV 300 mg every 4 weeks	

Belimumab	IV: 10 mg/Kg on weeks 0, 2, 4, then every 4 weeks SC: 200 mg weekly	No
Rituximab	1000 mg on days 1 and 15 - <i>re-administration every 6 months or “on-demand”</i>	No
Other biologic agents (off-label use usually for refractory joint and/or skin disease), e.g., • Tocilizumab • Abatacept • JAK inhibitors[#] • TNF inhibitors (rarely)[§]	Depending on agent	Depending on agent

* Recommended initial PO doses are general indication. In selected cases of organ- or life-threatening disease, higher initial doses, up to 0.7-0.8 mg/kg/day, may be given.

[§] Mycophenolate mofetil (MMF) is a prodrug of mycophenolic acid (MPA), administered orally either as MMF or as enteric-coated mycophenolate sodium (MPS). A 720 mg dose of MPS is roughly equivalent to a 1 g dose of MMF. MPS tends to be associated with more frequent gastrointestinal intolerance.

[#] Based on available data in patients with rheumatoid arthritis, risk factors for thrombosis and malignancy should be taken into account prior to use of JAK inhibitors in patients with SLE. With current knowledge, they should be avoided in patients with antiphospholipid syndrome

[§] TNF inhibitors are rarely used in SLE, due to their potential to cause drug-induced lupus. If used, regular monitoring for the appearance of anti-ds DNA antibodies and/or kidney involvement is needed.

IV: Intravenous; MP: Methylprednisolone; PO: Per os; LN: Lupus nephritis; BSA: Body surface area; NIH: National Institutes of Health; SC: Subcutaneous

Systematic Literature Review - Report

Systematic Literature Review (SLR) report for the 2023 update of the EULAR recommendations for the management of SLE

Table of Contents

Research questions and PICOs	2
PICO 1 - Therapeutic interventions	2
PICO 1a - Active SLE.....	2
PICO 1b - SLE with active mucocutaneous involvement.....	3
PICO 1c - SLE with active musculoskeletal involvement	4
PICO 1d - SLE with active neuropsychiatric involvement.....	5
PICO 1e - SLE with active haematological involvement	6
PICO 1f - SLE with active kidney involvement.....	7
PICO 2 - Low disease activity and remission.....	9
PICO 3 - SLE and antiphospholipid syndrome	9
PICO 4 - Tapering and/or withdrawal of treatment	10
PICO 5 - Vaccination against herpes zoster and SARS-CoV2 viruses	11
Inclusion and exclusion criteria:.....	11
Search strategy.....	12
Medline Search string for PICO 1 (PICOs 1a–1f).....	12
CENTRAL search string for PICO 1	13
Medline search string for PICO 2.....	16
CENTRAL search string for PICO 2	16
Medline search string for PICO 3.....	17
CENTRAL search string for PICO 3	18
Medline search for PICO 4.....	21
CENTRAL search string for PICO 4	21
Medline search string for PICO 5.....	22
CENTRAL search string for PICO 5	22
Flowchart.....	23
Summary fact sheets for all included studies.....	24
Risk of bias assessment	24
Risk of bias assessment of cohort studies and case-control studies using NOS	25
Risk of bias assessment for RCTs and quasi-RCTs using RoB2.....	32
Risk of bias assessment for meta-analyses	34
References	36

Systematic Literature Review - Report

Research questions and PICOs

For this update the research questions focused on five different domains: 1) the benefit/harm of SLE treatments (including lupus nephritis, neuropsychiatric, mucocutaneous, musculoskeletal and haematological lupus), 2) the benefits from the attainment of remission/low disease activity, 3) the risk/benefit from treatment tapering/withdrawal, 4) the management of SLE with aPL/APS and 5) the safety/toxicity of immunizations against zoster and SARS-Cov2.

Given the diversity of SLE populations, interventions, and outcomes different PICOs were developed for each individual question. As a first step a draft of the PICOs was circulated among the Task Force members who were encouraged to propose additional treatments or outcomes. The final version of PICOs was used as basis for the formulation of the respective search queries. Points-to consider for special areas/topics of interest were also included after each research question. The research questions with the respective PICOs, and points to consider are listed below.

PICO 1 – Therapeutic interventions

PICO 1a. In patients with **active SLE**, what is the evidence for the benefits and harms of therapeutic interventions including antimalarials, glucocorticoids, immunosuppressive, biological/targeted agents, plasma exchange/immunoadsorption?

Detailed PICO – see also ‘Points to consider’

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
• Patients with active SLE	• Sun protection • NSAIDs • Glucocorticoids • Hydroxychloroquine, antimalarials • Immunosuppressive agents • Cytotoxic agents • Methotrexate • Leflunomide • Azathioprine • Cyclophosphamide • Mycophenolate • Ciclosporin • Tacrolimus • Biological agents • Belimumab • Anifrolumab • Rituximab • Obinutuzumab • Ofatumumab • Ocrelizumab • Atacicept • Etanercept • Adalimumab • Abatacept • Adalimumab • Tocilizumab • Secukinumab • Ustekinumab	• Standard of care • Azathioprine • Placebo • None	• Disease activity improvement/worsening (SLEDAI, BILAG): global and specific domains • Cutaneous LE Disease Area and Severity Index • Tender joint count • Swollen joint count • Physician Global Assessment • Glucocorticoid sparing • Response (SRI-4, BICLA) • Disease control • Low disease activity (LLDAS) • Remission (various definitions including steroid-free remission) • Relapse, flare, time-to-flare • Treatment failure • Organ damage (including cataract, cognitive dysfunction, osteoporotic fracture, osteonecrosis, stroke, cardiovascular disease/MACEs, malignancy, diabetes) • Infection • Hospitalizations • Death • Adverse events/toxicity (including retinopathy) • Thrombosis

Systematic Literature Review - Report

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
	• Anakinra • JAK inhibitors (tofacitinib, baricitinib, upadacitinib, deucravacitinib) • Proteasome inhibitors (e.g., bortezomib) • Ibrdomide • Litlemab • Low-dose IL-2 • Daratumumab • CD19 CAR-T cells • Plasmapheresis • Plasma exchange • Immunoadsorption • Intravenous immunoglobulin		

Points to consider (for the SLR and/or data extraction):

- Stratification according to: patient age, ancestry/race, disease duration, prior treatments, selected biomarkers (serum complements, anti-dsDNA, IFN-signature)
- Glucocorticoids: capture dosage details such as the use of pulse methylprednisolone, initial dose, average dosage, tapering scheme
- Evidence on the efficacy of treatments in relapsing and refractory disease
- Collect data on global disease activity indices and activity from individual domains (e.g., serositis)
- Collect data on relevant safety outcomes: retinopathy, infections (including HZV, opportunistic), MACEs, hospitalizations, death

PICO 1b. In patients with **SLE and active mucocutaneous involvement**, what is the evidence for the benefits and harms of therapeutic interventions including sun protection, topical agents, antimalarials, glucocorticoids, immunosuppressive, biological/targeted agents?

Detailed PICO – see also ‘Points to consider’

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
• SLE patients with active mucocutaneous involvement	• Sun protection • Topical agents (glucocorticoids, calcineurin inhibitors) • Glucocorticoids • Hydroxychloroquine, antimalarials • Immunosuppressive agents • Cytotoxic agents • Methotrexate • Leflunomide • Azathioprine • Cyclophosphamide • Mycophenolate • Cyclosporin • Tacrolimus • Retinoids • Dapsone • Thalidomide • Lenalidomide • Biological agents	• Standard of care • Placebo • None	• Disease activity improvement/worsening (SLEDAI, BILAG): mucocutaneous-specific domains • Cutaneous LE Disease Area and Severity Index • Physician Global Assessment • Glucocorticoid sparing • Response (SRI-4, BICLA) • Disease control • Low disease activity (LLDAS) • Remission (various definitions including steroid-free remission) • Relapse, flare, time-to-flare • Treatment failure • Organ damage (including cataract, cognitive dysfunction, osteoporotic fracture, osteonecrosis, stroke, cardiovascular disease/MACEs, malignancy, diabetes) • Infection • Hospitalizations

Systematic Literature Review - Report

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
	• Belimumab • Anifrolumab • Rituximab • Obinutuzumab • Ofatumumab • Ocrelizumab • Atacicept • Etanercept • Adalimumab • Abatacept • Tocilizumab • Secukinumab • Ustekinumab • Anakinra • JAK inhibitors (tofacitinib, baricitinib, upadacitinib, deucravacitinib) • Proteasome inhibitors (e.g., bortezomib) • Ibrdomide • Litifilimab • Low-dose IL-2 • Daratumumab • CD19 CAR-T cells • Intravenous immunoglobulin		• Death • Toxicity (including retinopathy) • Thrombosis

Points to consider (for the SLR and/or data extraction):

- Stratification according to subtype: ACLE, SCLE, DLE and other forms of CCLE; patient age, ancestry/race, disease duration, prior treatments, selected biomarkers (serum complements, anti-dsDNA, IFN-signature)
- Glucocorticoids: capture dosage details such as the use of pulse methylprednisolone, initial/cumulative dose, tapering scheme
- Evidence on the efficacy of treatments in relapsing and refractory disease

PICO 1c. In patients with **SLE and active musculoskeletal involvement**, what is the evidence for the benefits and harms of therapeutic interventions including antimalarials, glucocorticoids, immunosuppressive and biological/targeted agents?

Detailed PICO – see also ‘Points to consider’

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
• SLE patients with active musculoskeletal involvement	• NSAIDs • Glucocorticoids • Hydroxychloroquine, antimalarials • Immunosuppressive agents • Cytotoxic agents • Methotrexate • Leflunomide • Azathioprine • Cyclophosphamide • Mycophenolate • Cyclosporin • Tacrolimus • Biological agents	• Standard of care • Placebo • None	• Disease activity improvement/worsening (SLEDAI, BILAG): musculoskeletal-specific domains • Tender joint count • Swollen joint count • Physician Global Assessment • Glucocorticoid sparing • Response (SRI-4, BICLA) • Disease control • Low disease activity (LLDAS) • Remission (various definitions including steroid-free remission) • Relapse, flare, time-to-flare • Treatment failure

Systematic Literature Review - Report

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
	• Belimumab • Anifrolumab • Rituximab • Obinutuzumab • Ofatumumab • Ocrelizumab • Atacicept • Etanercept • Adalimumab • Abatacept • Tocilizumab • Secukinumab • Ustekinumab • Anakinra • JAK inhibitors (tofacitinib, baricitinib, upadacitinib, deucravacitinib) • Proteasome inhibitors (e.g., bortezomib) • Ibrdomide • Litifilimab • Low-dose IL-2 • Daratumumab • CD19 CAR-T cells		• Organ damage (including cataract, cognitive dysfunction, osteoporotic fracture, osteonecrosis, stroke, cardiovascular disease/MACEs, malignancy, diabetes) • Infection • Hospitalizations • Death • Toxicity (including retinopathy) • Thrombosis

Points to consider (for the SLR and/or data extraction):

- Stratification according to arthritis phenotype (e.g., RA-like), patient age, ancestry/race, disease duration, prior treatments, selected biomarkers (serum complements, anti-dsDNA, IFN-signature)
- Glucocorticoids: capture dosage details such as the use of pulse methylprednisolone, initial dose, average dosage, tapering scheme
- Evidence on the efficacy of treatments in relapsing and refractory disease

PICO 1d. In patients with **SLE and active neuropsychiatric involvement**, what is the evidence for the benefits and harms of therapeutic interventions including antimalarials, glucocorticoids, immunosuppressive, biological/targeted agents, plasma exchange/immunoadsorption?

Detailed PICO – see also ‘Points to consider’

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
• SLE patients with active neuropsychiatric involvement	• Glucocorticoids • Hydroxychloroquine, antimalarials • Immunosuppressive agents • Cytotoxic agents • Methotrexate • Leflunomide • Azathioprine • Cyclophosphamide • Mycophenolate • Ciclosporin • Tacrolimus • Biological agents • Belimumab • Anifrolumab • Rituximab • Obinutuzumab	• Standard of care • Placebo • None	• Disease activity improvement/worsening (SLEDAI, BILAG): neuropsychiatric-specific domains • Neurological deficit (e.g., EDSS) • Neuropsychological tests • Psychiatric scales • Physician Global Assessment • Glucocorticoid sparing • Response (SRI-4, BICLA) • Disease control • Low disease activity (LLDAS) • Remission (various definitions including steroid-free remission) • Relapse, flare, time-to-flare • Treatment failure • Organ damage (including cataract, cognitive dysfunction, osteoporotic)

Systematic Literature Review - Report

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
	• Ofatumumab • Ocrelizumab • Atacicept • Etanercept • Adalimumab • Abatacept • Tocilizumab • Secukinumab • Ustekinumab • Anakinra • JAK inhibitors (tofacitinib, baricitinib, upadacitinib, deucravacitinib) • Proteasome inhibitors • Iberdomide • Litifilimab • Low-dose IL-2 • Daratumumab • CD19 CAR-T cells • Plasmapheresis • Plasma exchange • Immunoadsorption • Intravenous immunoglobulin		• fracture, osteonecrosis, stroke, cardiovascular disease/MACEs, malignancy, diabetes) • Infection • Hospitalizations • Death • Toxicity (including retinopathy) • Thrombosis

Points to consider (for the SLR and/or data extraction):

- Neuropsychiatric lupus as a single entity and according to individual manifestations (ACR nomenclature; 19 syndromes)
- Stratification according to: patient age, ancestry/race, disease duration, prior treatments, selected biomarkers (serum complements, anti-dsDNA, IFN-signature)
- Glucocorticoids: capture dosage details such as the use of pulse methylprednisolone, initial dose, average dosage, tapering scheme
- Evidence on the efficacy of treatments in relapsing and refractory disease
- Relevant safety outcomes: infections (including HZV, opportunistic), hospitalizations, death

PICO 1e. In patients with **SLE and active haematological involvement**, what is the evidence for the benefits and harms of therapeutic interventions including antimalarials, glucocorticoids, immunosuppressive, biological/targeted agents, plasma exchange/immunoadsorption?

Detailed PICO – see also ‘Points to consider’

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
• SLE patients with active haematological involvement	• Glucocorticoids • Hydroxychloroquine, antimalarials • Immunosuppressive agents • Cytotoxic agents • Methotrexate • Leflunomide • Azathioprine • Cyclophosphamide • Mycophenolate • Ciclosporin	• Standard of care • Placebo • None	• Disease activity improvement/worsening (SLEDAI, BILAG): haematological-specific domains • Complete blood count • Physician Global Assessment • Glucocorticoids sparing • Response (SRI-4, BICLA) • Disease control • Low disease activity (LLDAS) • Remission (various definitions including steroid-free remission)

Systematic Literature Review - Report

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
	• Tacrolimus • Biological agents • Belimumab • Anifrolumab • Rituximab • Obinutuzumab • Ofatumumab • Ocrelizumab • Atacicept • Etanercept • Adalimumab • Abatacept • Tocilizumab • Secukinumab • Ustekinumab • Anakinra • JAK inhibitors (tofacitinib, baricitinib, upadacitinib, deucravacitinib) • Proteasome inhibitors (e.g., bortezomib) • Iberdomide • Litifilimab • Low-dose IL-2 • Daratumumab • CD19 CAR-T cells • Plasmapheresis • Plasma exchange • Immunoabsorption • Intravenous immunoglobulin • Thrombopoietin-receptor agonists (romiplostim, eltrombopag)		• Relapse, flare, time-to-flare • Treatment failure • Organ damage (including cataract, cognitive dysfunction, osteoporotic fracture, osteonecrosis, stroke, cardiovascular disease/MACEs, malignancy, diabetes) • Infection • Hospitalizations • Death • Toxicity (including retinopathy) • Cardiovascular disease • Thrombosis

Points to consider (for the SLR and/or data extraction):

- Stratification according to patient age, ancestry/race, disease duration, prior treatments, selected biomarkers (serum complements, anti-dsDNA, IFN-signature)
- Glucocorticoids: capture dosage details such as the use of pulse methylprednisolone, initial dose, average dosage, tapering scheme
- Evidence on the efficacy of treatments in relapsing and refractory disease
- Relevant safety outcomes: infections (including HZV, opportunistic), hospitalizations, death

PICO 1f. In patients with **SLE and active kidney involvement**, what is the evidence for the benefits and harms of therapeutic interventions including antimalarials, glucocorticoids, immunosuppressive, biological/targeted agents, plasma exchange/immunoabsorption?

Detailed PICO – see also ‘Points to consider’

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
• SLE patients with active kidney involvement	• Glucocorticoids • Hydroxychloroquine, antimalarials • Immunosuppressive agents	• Standard of care • Mycophenolate • Azathioprine	• Disease activity improvement/worsening (SLEDAI, BILAG): renal-specific domains • Proteinuria improvement/worsening

Systematic Literature Review - Report

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
	•Cytotoxic agents •Methotrexate •Leflunomide •Azathioprine •Cyclophosphamide •Mycophenolate •Ciclosporin •Tacrolimus •Voclosporin •Biological agents •Belimumab •Anifrolumab •Rituximab •Obinutuzumab •Ofatumumab •Ocrelizumab •Atacicept •Telaticept •Dapagliflozin •Etanercept •Adalimumab •Abatacept •Tocilizumab •Secukinumab •Ustekinumab •Anakinra •JAK inhibitors (tofacitinib, baricitinib, upadacitinib, deucravacitinib) •Proteasome inhibitors •Iberdomide •Litifilimab •Low-dose IL-2 •Daratumumab •CD19 CAR-T cells •Plasmapheresis •Plasma exchange •Immunoadsorption •Intravenous immunoglobulin •RAAS inhibitors •SGLT2 inhibitors (Dapagliflozin)	•Cyclophosphamide •Ciclosporin •Tacrolimus •Placebo •None	•Kidney function (serum creatinine, eGFR) improvement/worsening •Chronic kidney disease •End-stage kidney disease •Histological improvement/worsening (change in activity/chronicity indices) •Physician Global Assessment •Glucocorticoid dose/tapering •Renal response (e.g., PEER, EULAR-defined endpoints) •Renal remission (complete renal response) •Relapse, flare, time-to-flare •Treatment failure •Organ damage (including cataract, cognitive dysfunction, osteoporotic fracture, osteonecrosis, stroke, cardiovascular disease/MACEs, malignancy, diabetes) •Infection •Hospitalizations •Death •Toxicity (including retinopathy) •Thrombosis

Points to consider (for the SLR and/or data extraction):

- Stratification according to kidney histology: proliferative, mixed proliferative and membranous, pure membranous (class V) lupus nephritis; presence of thrombotic microangiopathy (or other features of APS nephropathy); presence of crescents; activity and chronicity index; presence of IF/TA – limitations of current approaches to histologic classification, use of activity and chronicity scores
- Stratification according to: patient age, ancestry/race, disease duration, prior treatments, selected biomarkers (serum complements, anti-dsDNA, IFN-signature)
- Glucocorticoids: dosage details such as the use of pulse methylprednisolone, initial dose, average dosage, tapering scheme
- Evidence on the efficacy of treatments in relapsing and refractory disease
- Relevant safety outcomes: infections (including HZV, opportunistic), hospitalizations, death

Systematic Literature Review - Report

PICO 2. In patients with **SLE**, what is the evidence that attainment of low disease activity and remission are associated with improved patient and disease outcomes?

Detailed PICO – see also ‘Points to consider’

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
• SLE patients	• Low disease activity • Lupus Low Disease Activity State (LLDAS) • Remission • Inactive disease • Disease quiescence • Duration of LLDAS/remission	• Active disease • Not in low disease activity or remission or disease quiescence • None	• Relapse, flare, time-to-flare • Organ damage (including cataract, cognitive dysfunction, osteoporotic fracture, osteonecrosis, stroke, cardiovascular disease/MACEs, malignancy, diabetes) • Kidney function (serum creatinine, eGFR) improvement/worsening • Chronic kidney disease • End-stage kidney disease • Toxicity • Infection • Hospitalizations • Death

Points to consider (for the SLR and/or data extraction):

- Evidence on the prognostic value of various existing definitions and their modification, treated as binary variables (attainment or not) or (percentage of) time spent under the state
- Stratification according to: general SLE, lupus nephritis
- Stratification according to: patient age, ancestry/race, disease duration, selected biomarkers (serological activity, serum complements, anti-dsDNA, IFN-signature)

PICO 3. In patients with **SLE and antiphospholipid syndrome (including thrombotic microangiopathy)**, what is the evidence for the benefits and harms of therapeutic interventions including antiplatelets, anticoagulants, antimalarials, glucocorticoids, immunosuppressive, biological/targeted agents, plasma exchange/immunoadsorption?

Detailed PICO – see also ‘Points to consider’

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
• SLE patients with antiphospholipid syndrome	• Glucocorticoids • Hydroxychloroquine, antimalarials • Immunosuppressive agents • Cytotoxic agents • Methotrexate • Leflunomide • Azathioprine • Cyclophosphamide • Mycophenolate • Ciclosporin • Tacrolimus • Voclosporin • Biological agents • Belimumab	• Standard of care • Placebo • None	• Organ damage (including cataract, cognitive dysfunction, osteoporotic fracture, osteonecrosis, stroke, cardiovascular disease/MACEs, malignancy, diabetes) • Hospitalizations • Death • Toxicity (including bleeding) • Cardiovascular disease • Pregnancy/foetal loss • Live birth • Premature birth • Stillbirth • (Pre-)eclampsia • Vascular thrombosis (venous, arterial)

Systematic Literature Review - Report

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
	• Anifrolumab • Rituximab • Obinutuzumab • Ofatumumab • Ocrelizumab • Atacicept • Complement inhibitors (e.g., eculizumab) • Plasmapheresis • Plasma exchange • Immunoabsorption • Intravenous immunoglobulin • Aspirin • Heparin • Warfarin • Apixaban • Rivaroxaban • Eculizumab		

Points to consider (for the SLR and/or data extraction):

- Stratification according to: APS phenotype (obstetric APS, thrombotic APS, catastrophic APS), patient age, ancestry/race, selected biomarkers (serum complements, anti-dsDNA, IFN-signature)
- Evidence on the efficacy of treatments in relapsing and refractory disease

PICO 4. In patients with **SLE and quiescent disease**, what is the evidence for the benefits and harms of tapering and/or withdrawal of treatment including antimalarials, glucocorticoids, immunosuppressive, biological/targeted agents?

Detailed PICO – see also ‘Points to consider’

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
• SLE patients with quiescent disease (low disease activity or remission)	• Treatment withdrawal, discontinuation, tapering (including glucocorticoids, hydroxychloroquine, antimalarials, immunosuppressive agents, biological agents) • Duration of treatment	• Standard of care • Placebo • None	• Disease activity (SLEDAI, BILAG): global and specific domains • Physician Global Assessment • Glucocorticoid exposure • Disease control • Disease worsening • Treatment re-initiation • Low disease activity (LLDAS) • Remission (including steroid-free remission) • Relapse, flare, time-to-flare • Organ damage (including cataract, cognitive dysfunction, osteoporotic fracture, osteonecrosis, stroke, cardiovascular disease/MACEs, malignancy, diabetes) • Hospitalizations • Death • Toxicity (including bleeding)

Points to consider (for the SLR and/or data extraction):

Systematic Literature Review - Report

- Data stratification according to: patient age, ancestry/race, selected biomarkers (serum complements, anti-dsDNA, IFN-signature), disease duration, type of disease (SLE, lupus nephritis), tapering/withdrawal of glucocorticoids versus other therapeutic agents

PICO 5. In patients with SLE, what is the evidence for the benefits and harms of vaccination against infectious pathogens including [herpes zoster](#) and [SARS-CoV2](#) viruses?

Detailed PICO – see also ‘Points to consider’

Population(s)	Intervention(s)-exposure(s)	Comparison	Outcome(s)
• SLE patients	• Vaccination against zoster • Vaccination against SARS-CoV2	• No vaccination • None	• Serological response (protective antibodies) • Herpes zoster infection • SARS-CoV2 infection • COVID-19 • Need for hospitalization (e.g., need for oxygen supply, ICU) • Death • Disease activity (SLEDAI, BILAG): global and specific domains • Physician Global Assessment • Glucocorticoid exposure • Disease control • Disease worsening • Relapse, flare, time-to-flare • Toxicity

[Points to consider](#) (for the SLR and/or data extraction):

- Stratification according to: patient age, ancestry/race, disease status (active, inactive), type of disease (SLE, lupus nephritis), concomitant treatments (dose of glucocorticoids, immunosuppressives, biologics), major comorbidities (diabetes mellitus, cardiovascular disease, chronic respiratory disorders)
- Stratification according to type of vaccine (e.g., attenuated, recombinant, mRNA), number of booster vaccinations

[Inclusion and exclusion criteria:](#)

Inclusion criteria:

SLE adult population

Studies reporting data regarding efficacy/safety of treatments/withdrawal of treatments.

Studies reporting data regarding efficacy/safety of immunization against herpes or SARS-CoV2

Studies reporting outcomes associated with attainment of low disease activity or remission.

Eligible trial designs:

- Meta-analyses
- RCTs, quasi-RCTs
- Cohort studies (prospective and retrospective)
- Case-control studies
- Cross sectional-studies

Systematic Literature Review - Report

Exclusion criteria:

Reviews
 Case series
 Case reports
 Conference abstracts
 Animal studies
 Non-English language
 Trials with population <20
 Trials on paediatric populations

Search strategy

In line with the EULAR standardised operating procedures, the SLR included two databases (MEDLINE and the Cochrane Library - CENTRAL database) and one additional journal not indexed in PubMed (Lancet Rheumatology). Eligible studies had to be published between December 2017 and December 2022. The search queries for MEDLINE and CENTRAL were as follows:

Medline Search string for PICO 1 (PICOs 1a–1f)

("SLE"[Title] OR "lupus"[Title]) AND ("glucocorticoid*" [All Fields] OR "glucocorticoids"[MeSH Terms] OR "steroid*" [All Fields] OR "steroids"[MeSH Terms] OR "corticosteroid*" [All Fields] OR "anti inflammatory agents, non steroidal"[MeSH Terms] OR "non-steroidal anti-inflammatory agents"[Title] OR "nsaid"[Title] OR "nsaids"[Title] OR "nsaid s"[Title] OR ("hydroxychloroquine"[MeSH Terms] OR "hydroxychloroquine"[All Fields]) OR "antimalarial*" [All Fields] OR ("quinacrine"[MeSH Terms] OR "quinacrine"[All Fields]) OR ("methotrexate"[MeSH Terms] OR "methotrexate"[All Fields] OR "methotrexate s"[All Fields] OR "methotrexates"[All Fields]) OR ("leflunomid"[All Fields] OR "leflunomide"[MeSH Terms] OR "leflunomide"[All Fields] OR "leflunomide s"[All Fields]) OR ("calcineurin"[MeSH Terms] OR "calcineurin"[All Fields] OR "calcineurin s"[All Fields] OR "calcineurine"[All Fields] OR "calcineurins"[All Fields]) OR ("cyclosporine"[MeSH Terms] OR "cyclosporine"[All Fields] OR "ciclosporin"[All Fields] OR "ciclosporine"[All Fields] OR "cyclosporin"[All Fields] OR "cyclosporine s"[All Fields] OR "cyclosporins"[MeSH Terms] OR "cyclosporins"[All Fields] OR "cyclosporines"[All Fields]) OR ("tacrolimus"[MeSH Terms] OR "tacrolimus"[All Fields]) OR ("voclosporin"[Supplementary Concept] OR "voclosporin"[All Fields]) OR ("azathioprin"[All Fields] OR "azathioprine"[MeSH Terms] OR "azathioprine"[All Fields]) OR ("mycophenolate"[All Fields] OR "mycophenolates"[All Fields] OR "mycophenolic"[All Fields]) OR ("mycophenolate"[All Fields] OR "mycophenolates"[All Fields] OR "mycophenolic"[All Fields]) OR ("cyclophosphamide"[MeSH Terms] OR "cyclophosphamide"[All Fields] OR "cyclophosphamid"[All Fields] OR "cyclophosphamide s"[All Fields] OR "cyclophosphamides"[All Fields]) OR ("rituximab"[MeSH Terms] OR "rituximab"[All Fields] OR "rituximab s"[All Fields]) OR ("belimumab"[Supplementary Concept] OR "belimumab"[All Fields]) OR ("abatacept"[MeSH Terms] OR "abatacept"[All Fields]) OR "biologic*" [All Fields] OR "intravenous immunoglobulin"[All Fields] OR "plasma exchange"[All Fields] OR ("plasmapheresis"[MeSH Terms] OR "plasmapheresis"[All Fields] OR "plasmaphereses"[All Fields]) OR ("immunoadsorption"[All Fields] OR "immunoadsorptions"[All Fields]) OR ("anifrolumab"[Supplementary Concept] OR "anifrolumab"[All Fields]) OR ("obinutuzumab"[Supplementary Concept] OR "obinutuzumab"[All Fields]) OR ("ofatumumab"[Supplementary Concept] OR "ofatumumab"[All Fields]) OR ("ocrelizumab"[Supplementary Concept] OR "ocrelizumab"[All Fields]) OR ("taci receptor igg fc fragment fusion protein"[Supplementary Concept] OR "taci receptor igg fc fragment fusion protein"[All Fields] OR "atacept"[All Fields]) OR ("etanercept"[MeSH Terms] OR "etanercept"[All Fields]) OR ("adalimumab"[MeSH Terms] OR "adalimumab"[All Fields]) OR ("tocilizumab"[Supplementary Concept] OR "tocilizumab"[All Fields]) OR ("secukinumab"[Supplementary Concept] OR "secukinumab"[All Fields]) OR ("ustekinumab"[MeSH Terms] OR "ustekinumab"[All Fields]) OR ("interleukin 1 receptor antagonist protein"[MeSH Terms] OR "interleukin 1 receptor antagonist protein"[All Fields] OR "anakinra"[All Fields]) OR

Systematic Literature Review - Report

("tofacitinib"[Supplementary Concept] OR "tofacitinib"[All Fields] OR "tofacitinib s"[All Fields]) OR ("baricitinib"[Supplementary Concept] OR "baricitinib"[All Fields]) OR ("upadacitinib"[Supplementary Concept] OR "upadacitinib"[All Fields]) OR ("deucravacitinib"[Supplementary Concept] OR "deucravacitinib"[All Fields]) OR ("proteasome inhibitors"[MeSH Terms] OR "proteasome inhibitors"[All Fields]) OR ("bortezomib"[MeSH Terms] OR "bortezomib"[All Fields]) OR ("iberdomide"[Supplementary Concept] OR "iberdomide"[All Fields]) OR "Litifilimab"[All Fields] OR ("interleukin 2"[MeSH Terms] OR "interleukin 2"[All Fields] OR "IL-2"[All Fields]) OR ("daratumumab"[Supplementary Concept] OR "daratumumab"[All Fields]) OR "CAR-T cells"[All Fields] OR ("receptors"[All Fields] AND "thrombopoietin"[All Fields]) OR ("receptors, thrombopoietin"[MeSH Terms] OR "thrombopoietin receptors"[All Fields]) OR ("romiplostim"[Supplementary Concept] OR "romiplostim"[All Fields]) OR ("eltrombopag"[Supplementary Concept] OR "eltrombopag"[All Fields]) OR ("sodium glucose transporter 2 inhibitors"[MeSH Terms] OR "sodium glucose transporter 2 inhibitors"[All Fields] OR ("sglt2"[All Fields] AND "inhibitor"[All Fields])) OR ("dapagliflozin"[Supplementary Concept] OR "dapagliflozin"[All Fields] OR "dapagliflozin s"[All Fields]) OR ("renin"[MeSH Terms] OR "renin"[All Fields]) AND ("angiotensin s"[All Fields] OR "angiotensin"[All Fields] OR "angiotensins"[MeSH Terms] OR "angiotensins"[All Fields] OR "angiotensin"[All Fields]) AND "inhibitors"[All Fields])

Hits: 3,755

CENTRAL search string for PICO 1:

<https://www.cochranelibrary.com/advanced-search/search-manager?search=7138193>

ID	Search
#1	MeSH descriptor: [Lupus Erythematosus, Systemic] explode all trees
#2	("systemic lupus erythematosus"):ti,ab,kw (Word variations have been searched)
#3	(lupus):ti,ab,kw (Word variations have been searched)
#4	("glucocorticoid") (Word variations have been searched)
#5	MeSH descriptor: [Glucocorticoids] explode all trees
#6	MeSH descriptor: [Steroids] explode all trees
#7	(steroid) (Word variations have been searched)
#8	(corticosteroid) (Word variations have been searched)
#9	MeSH descriptor: [Anti-Inflammatory Agents, Non-Steroidal] explode all trees
#10	("non-steroidal anti-inflammatory agents"):ti (Word variations have been searched)
#11	(nsaids):ti (Word variations have been searched)
#12	MeSH descriptor: [Hydroxychloroquine] explode all trees
#13	("hydroxychloroquine") (Word variations have been searched)
#14	("antimalarial") (Word variations have been searched)
#15	MeSH descriptor: [Quinacrine] explode all trees
#16	("quinacrine") (Word variations have been searched)

Systematic Literature Review - Report

- #17 MeSH descriptor: [Methotrexate] explode all trees
- #18 ("methotrexate") (Word variations have been searched)
- #19 MeSH descriptor: [Leflunomide] explode all trees
- #20 ("leflunomide") (Word variations have been searched)
- #21 MeSH descriptor: [Calcineurin] explode all trees
- #22 ("calcineurin") (Word variations have been searched)
- #23 MeSH descriptor: [Cyclosporine] explode all trees
- #24 ("ciclosporin") (Word variations have been searched)
- #25 MeSH descriptor: [Tacrolimus] explode all trees
- #26 ("tacrolimus") (Word variations have been searched)
- #27 (voclosporin) (Word variations have been searched)
- #28 MeSH descriptor: [Azathioprine] explode all trees
- #29 ("azathioprin") (Word variations have been searched)
- #30 ("azathioprine") (Word variations have been searched)
- #31 MeSH descriptor: [Mycophenolic Acid] explode all trees
- #32 ("mycophenolate") (Word variations have been searched)
- #33 ("mycophenolic") (Word variations have been searched)
- #34 MeSH descriptor: [Cyclophosphamide] explode all trees
- #35 ("cyclophosphamide") (Word variations have been searched)
- #36 MeSH descriptor: [Rituximab] explode all trees
- #37 ("rituximab") (Word variations have been searched)
- #38 (belimumab) (Word variations have been searched)
- #39 MeSH descriptor: [Abatacept] explode all trees
- #40 ("abatacept") (Word variations have been searched)
- #41 ("biologic") (Word variations have been searched)
- #42 ("intravenous immunoglobulin") (Word variations have been searched)
- #43 ("plasma exchange") (Word variations have been searched)
- #44 MeSH descriptor: [Plasmapheresis] explode all trees
- #45 ("plasmapheresis") (Word variations have been searched)
- #46 (anifrolumab) (Word variations have been searched)
- #47 (obinutuzumab) (Word variations have been searched)
- #48 (ofatumumab) (Word variations have been searched)

Systematic Literature Review - Report

- #49 (ocrelizumab) (Word variations have been searched)
- #50 (atacept) (Word variations have been searched)
- #51 MeSH descriptor: [Etanercept] explode all trees
- #52 ("etanercept") (Word variations have been searched)
- #53 MeSH descriptor: [Adalimumab] explode all trees
- #54 ("adalimumab") (Word variations have been searched)
- #55 (tocilizumab) (Word variations have been searched)
- #56 (secukinumab) (Word variations have been searched)
- #57 (ustekinumab) (Word variations have been searched)
- #58 MeSH descriptor: [Ustekinumab] explode all trees
- #59 MeSH descriptor: [Interleukin 1 Receptor Antagonist Protein] explode all trees
- #60 (interleukin 1 receptor antagonist) (Word variations have been searched)
- #61 (anakinra) (Word variations have been searched)
- #62 (tofacitinib) (Word variations have been searched)
- #63 (baricitinib) (Word variations have been searched)
- #64 (upadacitinib) (Word variations have been searched)
- #65 (deucravacitinib) (Word variations have been searched)
- #66 MeSH descriptor: [Proteasome Inhibitors] explode all trees
- #67 ("protease inhibitor") (Word variations have been searched)
- #68 MeSH descriptor: [Bortezomib] explode all trees
- #69 ("bortezomib") (Word variations have been searched)
- #70 (iberdomide) (Word variations have been searched)
- #71 (litifilimab) (Word variations have been searched)
- #72 MeSH descriptor: [Interleukin-2] explode all trees
- #73 (interleukin 2) (Word variations have been searched)
- #74 ("IL 2") (Word variations have been searched)
- #75 (daratumumab) (Word variations have been searched)
- #76 (CAR-T cells) (Word variations have been searched)
- #77 MeSH descriptor: [Receptors, Thrombopoietin] explode all trees
- #78 (romiplostim) (Word variations have been searched)
- #79 (eltrombopag) (Word variations have been searched)
- #80 MeSH descriptor: [Sodium-Glucose Transporter 2 Inhibitors] explode all trees

Systematic Literature Review - Report

#81 (sodium glucose transporter 2 inhibitors) (Word variations have been searched)
 #82 (sglt2) (Word variations have been searched)
 #83 (dapagliflozin) (Word variations have been searched)
 #84 MeSH descriptor: [Renin] explode all trees
 #85 (renin) (Word variations have been searched)
 #86 MeSH descriptor: [Angiotensins] explode all trees
 #87 ("angiotensin") (Word variations have been searched)
 #88 (inhibitors) (Word variations have been searched)
 #89 #1 OR #2 OR #3
 #90 #84 OR #85
 #91 #86 OR #87
 #92 #90 AND #91 AND #88
 #93 #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #13 OR #14 OR #15
 OR #16 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23 OR #24 OR #25 OR #26 OR
 #27 OR #28 OR #29 OR #30 OR #31 OR #32 OR #33 OR #34 OR #35 OR #36 OR #37 OR #38
 OR #39 OR #40 OR #41 OR #42 OR #43 OR #44 OR #45 OR #46 OR #47 OR #48 OR #49 OR
 #50 OR #51 OR #52 OR #53 OR #54 OR #55 OR #56 OR #57 OR #58 OR #59 OR #60 OR #61
 OR #62 OR #63 OR #64 OR #65 OR #66 OR #67 OR #68 OR #69 OR #70 OR #71 OR #72 OR
 #73 OR #74 OR #75 OR #76 OR #77 OR #78 OR #79 OR #80 OR #81 OR #82 OR #83 OR #92
 #94 #89 AND #93
Hits: 2347

Medline search string for PICO 2:

((("SLE"[Title] OR "lupus"[Title]) AND ("remission"[All Fields] OR "remissions"[All Fields] OR "low disease activity"[All Fields] OR "LLDAS"[All Fields] OR "inactive disease"[All Fields] OR "quiescent disease"[All Fields] OR "disease quiescence"[All Fields] OR "treat to target"[All Fields])))

Hits: 929

CENTRAL search string for PICO 2:

<https://www.cochranelibrary.com/advanced-search/search-manager?search=7138194>

ID	Search
#1	MeSH descriptor: [Lupus Erythematosus, Systemic] explode all trees
#2	("systemic lupus erythematosus"):ti,ab,kw (Word variations have been searched)
#3	(lupus):ti,ab,kw (Word variations have been searched)
#4	(remission) (Word variations have been searched)
#5	(low disease activity) (Word variations have been searched)

Systematic Literature Review - Report

- #6 (LLDAS) (Word variations have been searched)
- #7 (inactive disease) (Word variations have been searched)
- #8 (quiescent disease) (Word variations have been searched)
- #9 (disease quiescence) (Word variations have been searched)
- #10 (treat to target) (Word variations have been searched)
- #11 #1 OR #2 OR #3
- #12 #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10
- #13 #11 AND #12

Hits: 911

Medline search string for PICO 3:

((("SLE"[Title] OR "lupus"[Title]) AND ("anti b2"[All Fields] OR "anti beta"[All Fields] OR "anti beta2"[All Fields] OR "anti cardiolipin"[All Fields] OR "anticardiolipin"[All Fields] OR "lupus anticoagulant"[All Fields] OR "LAC"[All Fields] OR "aPL"[All Fields] OR "antiphospholipid"[All Fields] OR ("syndrom"[All Fields] OR "syndromal"[All Fields] OR "syndromally"[All Fields] OR "syndrome"[MeSH Terms] OR "syndrome"[All Fields] OR "syndromes"[All Fields] OR "syndrome s"[All Fields] OR "syndromic"[All Fields] OR "syndroms"[All Fields]) OR ("arch plast surg"[Journal] OR "adv psychol study"[Journal] OR "acta pharmacol sin"[Journal] OR "aps"[All Fields])) AND ("manage"[All Fields] OR "managed"[All Fields] OR "management s"[All Fields] OR "managements"[All Fields] OR "manager"[All Fields] OR "manager s"[All Fields] OR "managers"[All Fields] OR "manages"[All Fields] OR "managing"[All Fields] OR "managment"[All Fields] OR "organization and administration"[MeSH Terms] OR ("organization"[All Fields] AND "administration"[All Fields]) OR "organization and administration"[All Fields] OR "management"[All Fields] OR "disease management"[MeSH Terms] OR ("disease"[All Fields] AND "management"[All Fields]) OR "disease management"[All Fields] OR ("therapeutics"[MeSH Terms] OR "therapeutics"[All Fields] OR "therapies"[All Fields] OR "therapy"[MeSH Subheading] OR "therapy"[All Fields] OR "therapy s"[All Fields] OR "therapys"[All Fields]) OR ("therapeutical"[All Fields] OR "therapeutically"[All Fields] OR "therapeuticals"[All Fields] OR "therapeutics"[MeSH Terms] OR "therapeutics"[All Fields] OR "therapeutic"[All Fields] OR ("therapeutics"[MeSH Terms] OR "therapeutics"[All Fields] OR "treatments"[All Fields] OR "therapy"[MeSH Subheading] OR "therapy"[All Fields] OR "treatment"[All Fields] OR "treatment s"[All Fields]) OR "anticoagul"[All Fields] OR "antiplatelet"[All Fields] OR "antiplatelet"[All Fields] OR ("aspirin"[MeSH Terms] OR "aspirin"[All Fields] OR "aspirins"[All Fields] OR "aspirin s"[All Fields] OR "aspirine"[All Fields]) OR ("heparin"[MeSH Terms] OR "heparin"[All Fields] OR "heparine"[All Fields] OR "heparins"[All Fields] OR "heparin s"[All Fields] OR "heparinate"[All Fields] OR "heparinated"[All Fields] OR "heparines"[All Fields] OR "heparinic"[All Fields] OR "heparinisation"[All Fields] OR "heparinised"[All Fields] OR "heparinization"[All Fields] OR "heparinize"[All Fields] OR "heparinized"[All Fields] OR "heparinizing"[All Fields]) OR ("warfarin"[MeSH Terms] OR "warfarin"[All Fields] OR "warfarin s"[All Fields] OR "warfarinization"[All Fields] OR "warfarinized"[All Fields] OR "warfarins"[All Fields]) OR ("apixaban"[Supplementary Concept] OR "apixaban"[All Fields] OR "apixaban s"[All Fields]) OR ("rivaroxaban"[MeSH Terms] OR "rivaroxaban"[All Fields]) OR "glucocorticoid"[All Fields] OR "glucocorticoids"[MeSH Terms] OR "steroid"[All Fields] OR "steroids"[MeSH Terms] OR "corticosteroid"[All Fields] OR "anti inflammatory agents, non steroidal"[MeSH Terms] OR "non-steroidal anti-inflammatory agents"[Title] OR "nsaid"[Title] OR "nsaids"[Title] OR "nsaid s"[Title] OR ("hydroxychloroquine"[MeSH Terms] OR "hydroxychloroquine"[All Fields]) OR "antimalarial"[All Fields] OR ("quinacrine"[MeSH Terms] OR "quinacrine"[All Fields]) OR ("methotrexate"[MeSH Terms] OR "methotrexate"[All Fields] OR "methotrexate s"[All Fields] OR "methotrexates"[All Fields]) OR ("leflunomid"[All Fields] OR "leflunomide"[MeSH Terms] OR "leflunomide"[All Fields] OR "leflunomide s"[All Fields]) OR ("calcineurin"[MeSH Terms] OR "calcineurin"[All Fields] OR "calcineurin s"[All Fields] OR "calcineurine"[All Fields] OR "calcineurins"[All Fields]) OR ("cyclosporine"[MeSH Terms] OR "cyclosporine"[All Fields] OR "ciclosporin"[All Fields] OR "ciclosporine"[All Fields] OR "cyclosporin"[All Fields] OR "cyclosporine s"[All Fields] OR "cyclosporins"[MeSH Terms] OR "cyclosporins"[All Fields] OR "cyclosporines"[All Fields]) OR ("tacrolimus"[MeSH Terms] OR "tacrolimus"[All Fields]) OR ("voclosporin"[Supplementary Concept] OR "voclosporin"[All Fields]) OR ("azathioprin"[All Fields] OR "azathioprine"[MeSH Terms] OR "azathioprine"[All Fields]) OR ("mycophenolate"[All Fields] OR

Systematic Literature Review - Report

"mycophenolates"[All Fields] OR "mycophenolic"[All Fields] OR ("mycophenolate"[All Fields] OR "mycophenolates"[All Fields] OR "mycophenolic"[All Fields] OR ("cyclophosphamide"[MeSH Terms] OR "cyclophosphamide"[All Fields] OR "cyclophosphamid"[All Fields] OR "cyclophosphamide s"[All Fields] OR "cyclophosphamides"[All Fields] OR ("rituximab"[MeSH Terms] OR "rituximab"[All Fields] OR "rituximab s"[All Fields] OR ("belimumab"[Supplementary Concept] OR "belimumab"[All Fields] OR ("abatacept"[MeSH Terms] OR "abatacept"[All Fields] OR "biologic*[All Fields] OR "intravenous immunoglobulin"[All Fields] OR "plasma exchange"[All Fields] OR ("plasmapheresis"[MeSH Terms] OR "plasmapheresis"[All Fields] OR "plasmaphereses"[All Fields] OR ("immunoadsorption"[All Fields] OR "immunoadsorptions"[All Fields] OR ("anifrolumab"[Supplementary Concept] OR "anifrolumab"[All Fields] OR ("obinutuzumab"[Supplementary Concept] OR "obinutuzumab"[All Fields] OR ("ofatumumab"[Supplementary Concept] OR "ofatumumab"[All Fields] OR ("ocrelizumab"[Supplementary Concept] OR "ocrelizumab"[All Fields] OR ("taci receptor igg fc fragment fusion protein"[Supplementary Concept] OR "taci receptor igg fc fragment fusion protein"[All Fields] OR "atacept"[All Fields] OR "complement inactivating agents"[MeSH Terms] OR ("complement"[All Fields] AND "inactivating"[All Fields] AND "agents"[All Fields] OR "complement inactivating agents"[All Fields] OR ("complement"[All Fields] AND "inhibitor"[All Fields] OR "complement inhibitor"[All Fields] OR ("thrombo*[All Fields] OR "pregnan*[All Fields] OR ("blood vessels"[MeSH Terms] OR ("blood"[All Fields] AND "vessels"[All Fields] OR "blood vessels"[All Fields] OR "vascular"[All Fields] OR "neovascularization, pathologic"[MeSH Terms] OR ("neovascularization"[All Fields] AND "pathologic"[All Fields] OR "pathologic neovascularization"[All Fields] OR "vascularisation"[All Fields] OR "vascularization"[All Fields] OR "vascularisations"[All Fields] OR "vascularise"[All Fields] OR "vascularised"[All Fields] OR "vascularities"[All Fields] OR "vascularitis"[All Fields] OR "vascularity"[All Fields] OR "vascularizations"[All Fields] OR "vascularize"[All Fields] OR "vascularized"[All Fields] OR "vascularizes"[All Fields] OR "vascularizing"[All Fields] OR "vasculars"[All Fields] OR "obstetric*[All Fields]))))

Hits: 1359

CENTRAL search string for PICO 3:

<https://www.cochranelibrary.com/advanced-search/search-manager?search=7138190>

ID	Search
#1	MeSH descriptor: [Lupus Erythematosus, Systemic] explode all trees
#2	("systemic lupus erythematosus"):ti,ab,kw (Word variations have been searched)
#3	(lupus):ti,ab,kw (Word variations have been searched)
#4	(anti b2) (Word variations have been searched)
#5	(anti beta) (Word variations have been searched)
#6	("anti-cardiolipin") (Word variations have been searched)
#7	(anti cardiolipin) (Word variations have been searched)
#8	("lupus anticoagulant") (Word variations have been searched)
#9	(LAC) (Word variations have been searched)
#10	(aPL) (Word variations have been searched)
#11	(antiphospholipid) (Word variations have been searched)
#12	(management) (Word variations have been searched)

Systematic Literature Review - Report

- #13 (therapeutics) (Word variations have been searched)
- #14 MeSH descriptor: [Therapeutics] explode all trees
- #15 MeSH descriptor: [Disease Management] explode all trees
- #16 (treatment) (Word variations have been searched)
- #17 ("anticoagulant") (Word variations have been searched)
- #18 ("antiplatelet") (Word variations have been searched)
- #19 ("aspirin") (Word variations have been searched)
- #20 MeSH descriptor: [Aspirin] explode all trees
- #21 ("heparin") (Word variations have been searched)
- #22 MeSH descriptor: [Heparin] explode all trees
- #23 MeSH descriptor: [Warfarin] explode all trees
- #24 ("Warfarin") (Word variations have been searched)
- #25 ("warfarin") (Word variations have been searched)
- #26 (apixaban) (Word variations have been searched)
- #27 MeSH descriptor: [Rivaroxaban] explode all trees
- #28 ("rivaroxaban") (Word variations have been searched)
- #29 ("glucocorticoid") (Word variations have been searched)
- #30 MeSH descriptor: [Glucocorticoids] explode all trees
- #31 MeSH descriptor: [Steroids] explode all trees
- #32 (steroid) (Word variations have been searched)
- #33 (corticosteroid) (Word variations have been searched)
- #34 MeSH descriptor: [Anti-Inflammatory Agents, Non-Steroidal] explode all trees
- #35 ("non-steroidal anti-inflammatory agents"):ti (Word variations have been searched)
- #36 (nsaids):ti (Word variations have been searched)
- #37 MeSH descriptor: [Hydroxychloroquine] explode all trees
- #38 ("hydroxychloroquine") (Word variations have been searched)
- #39 ("antimalarial") (Word variations have been searched)
- #40 MeSH descriptor: [Quinacrine] explode all trees
- #41 ("quinacrine") (Word variations have been searched)
- #42 MeSH descriptor: [Methotrexate] explode all trees
- #43 ("methotrexate") (Word variations have been searched)
- #44 MeSH descriptor: [Leflunomide] explode all trees

Systematic Literature Review - Report

- #45 ("leflunomide") (Word variations have been searched)
- #46 MeSH descriptor: [Calcineurin] explode all trees
- #47 ("calcineurin") (Word variations have been searched)
- #48 MeSH descriptor: [Cyclosporine] explode all trees
- #49 ("ciclosporin") (Word variations have been searched)
- #50 MeSH descriptor: [Tacrolimus] explode all trees
- #51 ("tacrolimus") (Word variations have been searched)
- #52 (voclosporin) (Word variations have been searched)
- #53 MeSH descriptor: [Azathioprine] explode all trees
- #54 ("azathioprin") (Word variations have been searched)
- #55 ("azathioprine") (Word variations have been searched)
- #56 MeSH descriptor: [Mycophenolic Acid] explode all trees
- #57 ("mycophenolate") (Word variations have been searched)
- #58 ("mycophenolic") (Word variations have been searched)
- #59 MeSH descriptor: [Cyclophosphamide] explode all trees
- #60 ("cyclophosphamide") (Word variations have been searched)
- #61 MeSH descriptor: [Rituximab] explode all trees
- #62 ("rituximab") (Word variations have been searched)
- #63 (belimumab) (Word variations have been searched)
- #64 MeSH descriptor: [Abatacept] explode all trees
- #65 ("abatacept") (Word variations have been searched)
- #66 ("biologic") (Word variations have been searched)
- #67 ("intravenous immunoglobulin") (Word variations have been searched)
- #68 ("plasma exchange") (Word variations have been searched)
- #69 MeSH descriptor: [Plasmapheresis] explode all trees
- #70 ("plasmapheresis") (Word variations have been searched)
- #71 (anifrolumab) (Word variations have been searched)
- #72 (obinutuzumab) (Word variations have been searched)
- #73 (ofatumumab) (Word variations have been searched)
- #74 (ocrelizumab) (Word variations have been searched)
- #75 (atacept) (Word variations have been searched)
- #76 MeSH descriptor: [Complement Inactivating Agents] explode all trees

Systematic Literature Review - Report

- #77 (complement inactivating factors) (Word variations have been searched)
- #78 (complement inhibitor) (Word variations have been searched)
- #79 ("thrombose") (Word variations have been searched)
- #80 ("thrombosis") (Word variations have been searched)
- #81 ("pregnancy") (Word variations have been searched)
- #82 MeSH descriptor: [Blood Vessels] explode all trees
- #83 (vascular) (Word variations have been searched)
- #84 (obstetric) (Word variations have been searched)
- #85 #1 OR #2 OR #3
- #86 #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11
- #87 #12 OR #13 OR #14 OR #15 OR #16 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23 OR #24 OR #25 OR #26 OR #27 OR #28 OR #29 OR #30 OR #31 OR #32 OR #33 OR #34 OR #35 OR #36 OR #37 OR #38 OR #39 OR #40 OR #41 OR #42 OR #43 OR #44 OR #45 OR #46 OR #47 OR #48 OR #49 OR #50 OR #51 OR #52 OR #53 OR #54 OR #55 OR #56 OR #57 OR #58 OR #59 OR #60 OR #61 OR #62 OR #63 OR #64 OR #65 OR #66 OR #67 OR #68 OR #69 OR #70 OR #71 OR #72 OR #73 OR #74 OR #75 OR #76 OR #77 OR #78 OR #79 OR #80 OR #81 OR #82 OR #83 OR #84
- #88 #85 AND #86 AND #87

Hits: 249

Medline search for PICO 4:

((("SLE"[Title] OR "lupus"[Title]) AND ("therapeutics"[MeSH Terms] OR "therapeutics"[All Fields] OR "treatments"[All Fields] OR "therapy"[MeSH Subheading] OR "therapy"[All Fields] OR "treatment"[All Fields] OR "treatment s"[All Fields] OR ("therapeutics"[MeSH Terms] OR "therapeutics"[All Fields] OR "therapies"[All Fields] OR "therapy"[MeSH Subheading] OR "therapy"[All Fields] OR "therapy s"[All Fields] OR "therapies"[All Fields]) OR ("manage"[All Fields] OR "managed"[All Fields] OR "management s"[All Fields] OR "managements"[All Fields] OR "manager"[All Fields] OR "manager s"[All Fields] OR "managers"[All Fields] OR "manages"[All Fields] OR "managing"[All Fields] OR "management"[All Fields] OR "organization and administration"[MeSH Terms] OR ("organization"[All Fields] AND "administration"[All Fields]) OR "organization and administration"[All Fields] OR "management"[All Fields] OR "disease management"[MeSH Terms] OR ("disease"[All Fields] AND "management"[All Fields]) OR "disease management"[All Fields])) AND ("stop*"[All Fields] OR "withdraw*"[All Fields] OR "discontin*"[All Fields] OR "taper*"[All Fields] OR ("duration"[All Fields] OR "durations"[All Fields])))

Hits: 829

CENTRAL search string for PICO 4:

<https://www.cochranelibrary.com/advanced-search/search-manager?search=7138188>

ID Search

Systematic Literature Review - Report

- #1 MeSH descriptor: [Lupus Erythematosus, Systemic] explode all trees
- #2 ("systemic lupus erythematosus"):ti,ab,kw (Word variations have been searched)
- #3 (lupus):ti,ab,kw (Word variations have been searched)
- #4 MeSH descriptor: [Therapeutics] explode all trees
- #5 (therapeutics) (Word variations have been searched)
- #6 (stop) (Word variations have been searched)
- #7 ("withdrawal") (Word variations have been searched)
- #8 ("discontinuation") (Word variations have been searched)
- #9 (taper) (Word variations have been searched)
- #10 ("duration") (Word variations have been searched)
- #11 #1 OR #2 OR #3
- #12 #4 OR #5
- #13 #6 OR #7 OR #8 OR #9 OR #10
- #14 #11 AND #12 AND #13

Hits: 375

Medline search string for PICO 5:

((("SLE"[Title] OR "lupus"[Title]) AND (((("vaccination"[MeSH Terms] OR "vaccination"[All Fields] OR "vaccinable"[All Fields] OR "vaccinal"[All Fields] OR "vaccinate"[All Fields] OR "vaccinated"[All Fields] OR "vaccinates"[All Fields] OR "vaccinating"[All Fields] OR "vaccinations"[All Fields] OR "vaccination s"[All Fields] OR "vaccines"[MeSH Terms] OR "vaccines"[All Fields] OR "vaccine"[All Fields] OR "vaccins"[All Fields]) AND ("herpes zoster"[MeSH Terms] OR ("herpes"[All Fields] AND "zoster"[All Fields]) OR "herpes zoster"[All Fields])) OR ("sars cov 2"[MeSH Terms] OR "sars cov 2"[All Fields] OR "covid"[All Fields] OR "covid 19"[MeSH Terms] OR "covid 19"[All Fields]))))

Hits: 333

CENTRAL search string for PICO 5:

<https://www.cochranelibrary.com/advanced-search/search-manager?search=7138094>

- ID Search
- #1 MeSH descriptor: [Lupus Erythematosus, Systemic] explode all trees
- #2 ("systemic lupus erythematosus"):ti,ab,kw (Word variations have been searched)
- #3 (lupus):ti,ab,kw (Word variations have been searched)

Systematic Literature Review - Report


- #4 MeSH descriptor: [Vaccines] explode all trees
- #5 (vaccine) (Word variations have been searched)
- #6 (vaccination) (Word variations have been searched)
- #7 MeSH descriptor: [Herpes Zoster] explode all trees
- #8 ("herpes zoster virus") (Word variations have been searched)
- #9 ("herpes virus") (Word variations have been searched)
- #10 (zoster) (Word variations have been searched)
- #11 MeSH descriptor: [COVID-19] explode all trees
- #12 MeSH descriptor: [COVID-19 Vaccines] explode all trees
- #13 ("SARS CoV") (Word variations have been searched)
- #14 (covid 19) (Word variations have been searched)
- #15 (covid) (Word variations have been searched)
- #16 #1 OR #2 OR #3
- #17 #4 OR #5 OR #6
- #18 #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #13 OR #14 OR #15
- #19 #17 AND #18
- #20 #16 AND #19

Hits: 31

Flowchart

Prisma flowchart with the use of the package DiagrammeR R [1]

Systematic Literature Review - Report



Summary fact sheets for all included studies

All relevant data are presented in a separate [Excel](#) file.

Risk of bias assessment

Systematic Literature Review - Report

A risk of bias assessment was performed for all eligible studies using the appropriate tools based on their design. The Newcastle-Ottawa scale (NOS) was used to assess cohort and case-control studies, RoB2 was used for RCTs and quasi RCTs and AMSTAR2 was used to assess meta-analyses.

Risk of bias assessment of cohort studies and case-control studies using NOS

The NOS scale is a risk of bias tool for the assessment of cohorts and case control studies based on their performance in three grouping items namely the selection of population, the comparability and the outcomes/exposures of the respective study [2]. Each cohort or case-control study is graded with a maximum of one star for each numbered item within the Selection and Outcome categories while Comparability can be graded with a maximum of two stars. For cohort studies, the number of stars and their distribution determines whether the study is of good, fair, or poor quality according to AHRQ (Agency for Healthcare Research and Quality) standards:

Good quality: 3 or 4 stars in selection domain AND 1 or 2 stars in comparability domain AND 2 or 3 stars in outcome/exposure domain

Fair quality: 2 stars in selection domain AND 1 or 2 stars in comparability domain AND 2 or 3 stars in outcome/exposure domain

Poor quality: 0 or 1 star in selection domain OR 0 stars in comparability domain OR 0 or 1 stars in outcome/exposure domain

	Risk domain								
Newcastle Ottawa scale	Selection				Comparability	Outcomes			
Cohort study	Representativeness of the exposed cohort	Selection of the non-exposed cohort	Ascertainment of exposure	Demonstration that outcome of interest was not present at start of study	Comparability of cohorts on the basis of the design or analysis controlled for confounders	Assessment of outcome	Was follow-up long enough for outcomes to occur	Adequacy of follow-up of cohorts	Quality of study
Rathoon, Indian J Nephrol, 2022	0	0	0	1	0	0	1	0	Poor
Nikoloudaki, Front Immunol, 2023	1	0	1	1	0	1	0	1	Poor
Zhang, Front Immunol 2022	1	1	1	0	1	1	0	1	Good
Floris RMD Open, 2022	0	1	1	1	1	1	1	1	Good
Aloub, Open Access Rheumatol, 2022	0	0	1	0	0	0	0	0	Poor
Hurst, AM J Med, 2022	1	1	1	1	0	1	1	1	Poor
Hunnicutt, Lupus Sci Med, 2022	0	0	1	1	0	1	1	0	Poor
Carter, Arthritis Rheumatol, 2022	1	0	1	1	0	1	0	0	Poor
Kagawa, Acta Med Okayama, 2022	0	0	1	0	0	1	1	1	Poor
Enfrein, RMD Open, 2022	0	0	1	1	0	1	1	0	Poor
Kao, J Ocul Phrmacol Ther, 2022	0	0	0	0	0	1	1	1	Poor
Ko, Semin Arthritis Rheum, 2022	1	1	0	1	2	1	1	1	Good
Dobrowolski, Rheumatology, 2022	1	1	1	1	1	1	0	1	Good
Connelly, Arthritis Rheumatol, 2022	0	0	1	1	2	1	1	1	Good

Systematic Literature Review - Report

	Risk domain								
Newcastle Ottawa scale	Selection				Comparability	Outcomes			
Cohort study	Representativeness of the exposed cohort	Selection of the non-exposed cohort	Ascertainment of exposure	Demonstration that outcome of interest was not present at start of study	Comparability of cohorts on the basis of the design or analysis controlled for confounders	Assessment of outcome	Was follow-up long enough for outcomes to occur	Adequacy of follow-up of cohorts	Quality of study
Mok, Vaccine, 2022	1	1	0	0	1	1	0	1	Fair
Hoque, Arthritis Rheumatol, 2022	1	1	0	1	2	1	1	1	Good
Wang, Lupus Sci Med, 2022	1	1	0	1	1	0	0	1	Poor
Chen, J Int Med Res, 2022	0	1	0	1	1	0	0	0	Poor
Ugarte-Gil, Ann Rheum Dis, 2022	1	1	1	1	1	1	1	1	Good
Nakai, Clin Rheumatol, 2022	0	0	1	1	0	1	1	0	Poor
Li, Pak J Med Sci, 2022	0	0	0	1	0	0	1	0	Poor
Nakai, Lupus Sci Med, 2022	0	0	1	1	1	1	0	0	Poor
Kapsia, Front Med, 2022	0	0	1	0	0	1	1	0	Poor
Khattab, Lupus, 2022	1	1	1	1	1	1	0	1	Good
Hussenbocus, Clin Rheumatol, 2022	1	1	0	1	0	1	0	1	Poor
Miyazaki, Rheumatology, 2022	1	1	1	1	2	1	1	1	Good
Almeida-Brasil, Ann Rheum Dis, 2022	1	1	0	1	2	1	0	1	Good
Ohkubo, Mod Rheumatol, 2022	0	0	1	1	1	1	1	0	Fair
Ayano, Mod Rheumatol, 2022	0	0	1	0	1	1	0	0	Poor
Yuki, Arthritis Care Res, 2022	1	0	1	0	2	1	1	0	Fair
Keyes, J Am Acad Dermatol, 2022	0	0	0	1	1	0	1	0	Poor
Simard, Lupus Sci Med, 2022	1	1	1	1	2	1	1	1	Good
Liao, J Clin Rheumatol, 2022	0	0	1	1	1	1	0	0	Poor
Izmirly, Arthritis Rheumatol, 2022	1	0	1	1	0	1	1	1	Poor
Sonigo, J Am Acad dermatol, 2021	0	0	1	0	0	1	1	0	Poor
Ruiz-Irastorza, Autoimmun Rev, 2021	0	0	1	1	1	1	0	1	Fair
Chen, Lupus, 2021	1	1	0	1	0	1	1	1	Poor
Tselios, ACR Open Rheumatol, 2021	1	1	1	1	1	1	1	1	Good
Olivieri, Joint Bone Spine, 2021	1	0	1	1	0	1	0	0	Poor
Piranavan, Clin Immunol, 2021	0	0	0	1	0	0	0	1	Poor
Abdelbaky, Egypt J Intern Med, 2021	1	0	1	0	0	1	0	0	Poor
Yoshida, Lupus, 2021	0	0	0	1	0	1	1	0	Poor
Fasano, Clin Exp Rheumatol, 2021	0	1	1	1	2	1	1	1	Good
Ugarte, Rheumatology, 2021	1	1	1	1	2	1	1	1	Good
Hill, Lupus Sci Med, 2021	1	0	0	1	2	2	1	1	Fair
Chen, Ther Adv Musculoskelet Dis, 2021	1	1	1	1	2	1	0	1	Good
Lobbess, Rheumatology, 2022	0	0	0	1	0	1	1	0	Poor
Zen, Rheumatology, 2022	1	0	1	1	2	1	1	0	Good

Systematic Literature Review - Report

	Risk domain								
Newcastle Ottawa scale	Selection				Comparability	Outcomes			
Cohort study	Representativeness of the exposed cohort	Selection of the non-exposed cohort	Ascertainment of exposure	Demonstration that outcome of interest was not present at start of study	Comparability of cohorts on the basis of the design or analysis controlled for confounders	Assessment of outcome	Was follow-up long enough for outcomes to occur	Adequacy of follow-up of cohorts	Quality of study
Roccatello, Kidney Int Rep, 2021	0	1	1	1	1	1	0	1	Good
Tselios, Arthritis Care Res, 2022	1	1	1	1	1	1	1	1	Good
Wang, Arthritis Care Res, 2021	1	1	1	1	2	1	0	1	Good
Abe, Biomed Res Int, 2021	1	0	0	1	0	1	0	0	Poor
Hoque, Arthritis Care Res, 2021	1	1	0	1	2	1	0	1	Good
Petri, Arthritis Rheumatol, 2021	1	0	1	1	2	1	0	1	Good
Choi, Rheumatology, 2021	0	1	0	1	2	1	0	1	Fair
Zickert, Rheumatology, 2021	0	0	1	1	0	1	0	1	Poor
Birt, Lupus Sci Med, 2020	1	0	1	0	0	0	1	1	Poor
Almeida-Brasil, Arthritis Care Res, 2022	1	1	0	1	2	0	1	1	Good
Haugaard, J Am Acad Dermatol, 2021	1	1	0	1	1	1	1	1	Good
Reátegui-Sokolova, RMD Open, 2021	0	0	0	1	2	1	0	5	Poor
Ceccarelli, Isr Med Assoc J, 2020	0	0	1	1	0	1	1	0	Poor
Collins, Rheumatol Ther, 2020	1	0	1	0	0	0	0	1	Poor
Sogayise, Int J Nephrol, 2020	1	1	1	1	0	1	1	0	Poor
Jin, Rheumatology, 2021	1	1	0	1	2	1	0	1	Good
Gupta, Arthritis Care Res, 2021	1	1	1	1	2	1	0	1	Good
Urowitz, Lupus Sci Med, 2020	1	1	1	1	2	1	1	1	Good
Sakai, Lupus, 2020	1	1	0	1	2	1	0	0	Fair
Nikfar, Int J Clin Pract, 2021	1	1	1	1	2	1	1	0	Good
Jakez-Ocampo, Clin Rheumatol, 2020	1	1	0	1	1	1	1	0	Good
Kang, Rheumatology, 2021	1	1	1	0	2	1	1	1	Good
Kandane-Rathnayake, Lancet Rheumatol, 2022	1	1	1	1	2	1	0	1	Good
Golder, Lancet Rheumatol, 2019	1	1	1	1	2	1	0	1	Good
28528869 Silva-Fernández et al	1	1	1	1	1	1	1	0	Good
28566017 Li et al	0	0	1	1	0	1	0	1	Poor
28704598 Ruiz-Arruza et al	0	0	1	0	0	0	1	0	Poor
28753077 Sheikholeslami et al	1	0	1	1	0	1	0	0	Poor
28856466 Sun et al	0	0	1	1	0	1	0	0	Poor
28862513 Emamikia et al	1	1	1	1	0	1	1	1	Good
28901731 Kasitanon et al	0	0	1	1	1	1	0	0	Fair
28935492 Iaccarino et al	1	0	1	1	1	1	1	0	Good
28970217 Zen et al	1	1	1	1	1	1	1	0	Good
29061479 Chasset et al	0	0	1	1	0	0	1	0	Poor
29087260 Mok et al	1	1	1	1	1	1	1	1	Good

Systematic Literature Review - Report

	Risk domain								
Newcastle Ottawa scale	Selection				Comparability	Outcomes			
Cohort study	Representativeness of the exposed cohort	Selection of the non-exposed cohort	Ascertainment of exposure	Demonstration that outcome of interest was not present at start of study	Comparability of cohorts on the basis of the design or analysis controlled for confounders	Assessment of outcome	Was follow-up long enough for outcomes to occur	Adequacy of follow-up of cohorts	Quality of study
29142034 Chen et al	1	0	1	1	1	1	0	0	Poor
29142038 Pakchotanon et al	1	0	0	1	2	1	1	0	Fair
29157178 Lee et al	1	0	1	1	2	1	0	0	Poor
29186572 Cunha et al	1	0	1	1	0	1	1	0	Poor
29216396 McCarthy et al	1	0	1	1	0	1	0	0	Poor
29222972 Deguchi et al	1	0	1	1	2	1	1	0	Good
29247540 Serris et al	1	0	1	1	0	1	1	0	Poor
29320974 Ganapati et al	1	1	1	1	0	1	0	0	Poor
29308726 Iwata et al	1	0	1	1	0	1	1	1	Poor
29409143 Furie et al	1	0	1	1	0	1	1	1	Poor
29420200 Morand et al	1	1	1	1	2	1	1	0	Good
29448881 Choi et al	0	0	1	1	0	1	1	0	Poor
29449503 Yue et al	0	0	1	1	2	1	1	0	Fair
29531772 Tani et al	1	0	1	1	2	1	1	0	Good
29515299 Sahay et al	1	1	1	1	0	1	0	0	Poor
29509932 Yap et al	1	1	1	1	0	1	1	0	Poor
29496892 Davidson et al	1	1	1	1	2	1	0	0	Poor
29460699 Furie et al	1	1	1	1	1	1	0	1	Fair
29561474 Goswami et al	1	0	1	1	2	1	0	0	Poor
29555348 Fanouriakis et al	1	0	1	1	0	1	1	1	Poor
29611341 Joo et al	1	1	1	1	0	1	1	0	Poor
29631512 Liu et al	1	0	1	1	2	1	1	0	Good
29635998 Ugarte et al	1	0	1	1	0	1	1	0	Poor
29657872 Soyuöz et al	0	0	1	1	0	1	0	0	Poor
29720229 Hanaoka et al	0	0	1	1	0	1	0	0	Poor
29792370 Tanaka et al	1	0	1	1	0	1	1	0	Poor
29806142 Petri et al	1	1	1	1	2	1	1	0	Good
29807477 Doria et al	1	0	1	1	0	1	0	1	Poor
29854814 Su et al	0	0	1	1	0	1	0	0	Poor
29855561 Burt et al	0	0	1	1	0	1	1	1	Poor
29931367 Hsu et al	1	1	1	1	1	1	1	0	Fair
29950160 Kwon et al	1	1	1	0	2	0	1	0	Poor
29954281 Spinelli et al	0	0	1	1	1	1	1	1	Fair
29987550 Monzavi et al	0	0	1	1	0	1	0	0	Poor
29998829 Park et al	1	0	1	1	0	1	0	0	Poor
30008461 Garnier et al	0	0	1	1	0	1	0	0	Poor
30055090 Tselios et al	1	1	1	1	0	1	1	0	Poor
30194649 Fasano et al	1	1	1	1	2	1	1	0	Good

Systematic Literature Review - Report

	Risk domain								
Newcastle Ottawa scale	Selection				Comparability	Outcomes			
Cohort study	Representativeness of the exposed cohort	Selection of the non-exposed cohort	Ascertainment of exposure	Demonstration that outcome of interest was not present at start of study	Comparability of cohorts on the basis of the design or analysis controlled for confounders	Assessment of outcome	Was follow-up long enough for outcomes to occur	Adequacy of follow-up of cohorts	Quality of study
30203113 Karasawa et al	0	0	1	1	0	1	1	0	Poor
30284580 Alsuwaida et al	0	0	1	1	0	1	0	0	Poor
30338639 Miyagawa et al	0	0	1	1	0	1	0	0	Poor
30451641 Gonzalez-Echavarri et al	0	0	1	1	2	1	0	0	Poor
30487482 Hossain et al	0	0	1	1	0	1	0	0	Poor
30538815 Tani et al	0	0	0	1	0	1	0	0	Poor
30523554 Goswami et al	1	0	1	1	2	1	1	0	Good
30552172 Sciascia et al	1	0	1	1	2	1	0	0	Poor
30557058 Okabayashi et al	1	0	1	1	2	1	0	0	Poor
30588322 Merrill et al	1	1	1	1	1	1	1	0	Good
30588323 van Vollenhoven et al	1	1	1	1	0	1	0	0	Poor
30626831 Hanaoka et al	0	0	1	1	0	1	0	0	Poor
30678605 Alarcon et al	1	1	1	1	0	1	0	0	Poor
30700214 Ichinose et al	1	1	1	1	0	1	0	0	Poor
30719729 Ootake et al	0	0	1	1	0	1	1	0	Poor
30755141 Martin-Iglesias et al	1	0	1	1	0	1	1	0	Poor
30771238 Wallace et al	1	0	1	1	0	1	1	1	Poor
30778862 Kawazoe et al	0	0	1	1	0	1	0	0	Poor
30852830 von Kempis et al	1	0	1	1	0	1	0	0	Poor
30937637 Sumethkul et al	0	0	1	1	0	1	0	0	Poor
30941559 Rebelo et al	1	0	1	1	0	1	1	0	Poor
30979713 Huang et al	1	0	1	1	0	1	1	0	Poor
31031386 Sharma et al	0	0	1	1	0	1	0	0	Poor
31074727 Tseng et al	1	0	1	1	0	1	1	0	Poor
31102498 Cassia et al	1	0	1	1	0	1	1	0	Poor
31122136 Geraldino-Pardilla et al	1	0	1	1	2	1	0	0	Poor
31175481 Hanaoka et al	1	0	1	1	0	1	1	0	Poor
31195632 Yang et al	1	0	1	1	2	1	1	0	Good
31199180 Tanaka et al	0	0	1	1	0	1	1	1	Poor
31264525 Anjo et al	0	0	1	1	0	1	1	0	Poor
31275608 Tani et al	0	0	1	1	2	1	1	0	Poor
31293110 Jung et al	1	0	1	1	0	1	0	0	Poor
31302695 van Vollenhoven et al	1	0	1	1	0	1	1	1	Poor
31464233 Al Hamzi et al	1	0	1	1	0	1	1	0	Good
31551028 Reategui-Sokolova et al	1	1	1	1	2	1	1	0	Good
31583978 Won et al	0	0	1	1	2	1	1	0	Poor
31600023 Floris et al	0	0	1	1	2	1	1	0	Poor

Systematic Literature Review - Report

	Risk domain								
Newcastle Ottawa scale	Selection				Comparability	Outcomes			
Cohort study	Representativeness of the exposed cohort	Selection of the non-exposed cohort	Ascertainment of exposure	Demonstration that outcome of interest was not present at start of study	Comparability of cohorts on the basis of the design or analysis controlled for confounders	Assessment of outcome	Was follow-up long enough for outcomes to occur	Adequacy of follow-up of cohorts	Quality of study
31653191 Nieto-Aristizabal et al	0	0	1	1	0	1	0	0	Poor
31769212 van Vollenhoven et al	1	1	1	1	0	1	0	1	Poor
31777844 Aouhab et al	0	0	1	1	0	1	0	0	Poor
31793379 Lee et al	1	0	1	1	2	1	1	0	Good
32020727 Miyagawa et al	1	1	1	1	0	1	0	0	Poor
32192398 Pedrosa et al	0	0	1	1	0	1	0	0	Poor
32238515 Takeuchi et al	1	0	1	1	0	1	1	0	Poor
32275125 Gatto et al	1	0	1	1	2	1	1	0	good
32321345 Sun et al	1	0	1	1	2	1	1	0	Good
32321721 Saccon et al	1	0	1	1	0	1	1	0	Poor
32434863 Vázquez-Otero et al	1	0	1	1	0	1	0	0	Poor
32437258 Prasad et al	1	1	1	1	0	1	1	0	Poor
32448782 Mok et al	1	1	1	1	2	1	1	1	Good
32452167 Padiyar et al	0	0	1	1	0	1	0	0	Poor
32462476 Argolini et al	0	0	1	1	2	1	1	1	Fair
32493152 Saleh et al	1	1	1	1	2	1	0	0	Poor
32522920 Wakiya et al	0	0	1	1	0	1	0	0	Poor
32791930 Babini et al	1	0	1	1	0	1	1	0	Poor
32813314 Bernatsky et al	1	1	1	1	2	1	1	0	Good
28857717 Pakchotanon et al	1	1	1	1	2	1	1	0	Good
28888363 Medina-Rosas et al	1	1	1	1	2	1	1	0	Good
29423203 Lay The et al	0	0	1	1	2	1	1	0	Fair
29478901 Wang et al	1	0	1	1	0	1	1	1	Poor
30045812 De Rosa et al	1	0	1	1	2	1	1	1	Good
30406967 Hanaoka et al	0	0	1	1	0	1	0	0	Poor
30755146 Ichinose et al	0	0	1	1	2	1	0	0	Poor
30821926 Sharma et al	0	0	1	1	1	1	1	0	Fair
31642908 Zen et al	0	0	1	1	2	1	1	0	Fair
31685314 Malvar et al	1	0	1	1	0	1	1	1	Poor
28659045 Watanabe et al	1	1	1	1	2	1	1	0	Good
29130759 Mecacci et al	0	0	1	1	0	1	0	0	Poor
29723256 Hanaoka et al	0	0	1	1	0	1	1	1	Poor
30837214 Gebhart et al	0	0	1	1	2	1	1	0	Fair
31905492 Dogan et al	0	0	1	1	0	1	0	0	poor
29667100 The et al	0	0	1	1	2	1	1	0	Fair
34121836 Abdelbaky et al	0	0	1	1	0	1	0	0	Poor

Systematic Literature Review - Report

Newcastle-Ottawa scale	Risk domain								Total number of stars
	Selection				Comparability	Exposure			
	Adequacy of case definition	Representativeness of cases	Selection of controls	Definition of controls	Comparability of the cases and controls on the basis of design or analysis	Ascertainment of exposure	Same method of ascertainment for cases and controls	Non-response rate	
Case control study									
Su, Front Immunol, 2022	1	0	1	1	1	1	1	0	5/9
Sada, Lupus Sci Med, 2022	1	0	0	1	1	0	1	0	4/9
Jorge, JAMA 2022	1	1	1	1	2	1	1	1	9/9
Damara, Cureus, 2022	1	0	0	0	0	1	1	0	3/9
Mancuso, Clin Exp Rheumatol, 2022	1	1	1	0	1	1	1	0	6/9
Rua-Figeroa, Semin Arthritis Rheum, 2022	0	0	1	1	0	0	1	0	3/9
Kwan, Lupus Sci Med, 2022	1	1	1	1	1	0	1	0	6/9
Jorge, Arthritis Care Res, 2022	1	1	1	1	2	1	1	1	9/9
Long, Lupus, 2021	0	1	1	1	0	0	1	1	5/9
Lo, PLOS One, 2021	1	1	1	1	2	0	1	1	8/9
Garelick, Rheumatology, 2021	0	0	1	1	1	1	1	0	5/9
Wang, Lupus, 2020	1	1	1	1	1	1	1	0	7/9
Papachristos, Semin Arthritis Rheum, 2022	1	1	1	1	2	1	1	0	8/9
29765616 Davidson et al	0	0	0	1	1	1	1	0	4/9
30103646 Yang et al	1	1	1	1	2	1	1	1	9/9
30367020 Gadakchi et al	1	0	1	0	1	1	1	0	5/9
31066646 Dall’Era	1	0	0	0	0	1	1	0	3/9
31474597 Mukwikwi et al	1	1	1	1	1	1	1	1	8/9
32407570 Jorge et al	1	1	1	1	1	1	1	0	7/9
32442312 Lenfant et al	1	1	0	1	0	1	1	0	5/9
32586407 Guo et al	1	1	1	1	1	1	1	0	7/9
32653901 Bultink et al	1	1	1	1	1	1	1	0	7/9
32807233 Fernandez-Ruiz et al	1	0	0	1	2	1	1	0	6/9
28857715 Ugarte-Gil et al	1	1	1	1	2	1	1	0	8/9

Systematic Literature Review - Report

Risk of bias assessment for RCTs and quasi-RCTs using RoB2

RoB2 is a Cochrane risk-of-bias tool for randomized trials [3]. Risk of bias is assessed in 5 different domains including bias arising from the randomization process, bias due to deviations from intended interventions, bias due to missing outcome data, bias in measurement of the outcome, and bias in selection of the reported result. The tool uses algorithms to determine the individual risk of bias for each domain. The domain-level ratings determine the overall risk of bias of a study. In brief, a trial is of low overall risk of bias if all domains are of low risk of bias, a study is considered to raise some concerns if there are concerns in at least one domain but no high risk of bias in any domain and, a study is of high risk of bias if at least one domain is of high risk of bias or multiple domains raise some concerns.

RoB2	Risk domain					
RCT	Risk of bias arising from the randomization process	Risk of bias due to deviations from the intended interventions	Missing outcome data	Risk of bias in measurement of the outcome	Risk of bias in selection of the reported result	Overall risk of bias
Zhang, Drugs R&D, 2022	High	Some concerns	Low	Low	Low	High
Morand, Arthritis Rheumatol, 2022	Low	Low	Low	Low	Low	Low
Wang, RMD Open, 2022	High	Low	Low	Some concerns	Low	High
Zheng, Mod Rheumatol, 2022	Low	Low	Low	Low	Low	Low
Wallace, Lupus, 2022	Low	Low	Low	Low	Low	Low
Furie, N Engl J Med, 2022	Low	Low	Low	Low	Low	Low
Yu, Am J Kidney Dis, 2022	Low	Low	Low	Low	Low	Low
Arriens, Arthritis Rheumatol, 2022	Low	Low	Low	Low	Low	Low
Van Vollenhoven, Ann Rheum Dis, 2022	Low	Some concerns	Some concerns	Low	Some concerns	Some concerns
Fu, Ann Rheum Dis, 2022	Some concerns	Low	Some concerns	Low	Low	Some concerns
Jourde-Chiche, Ann Rheum Dis, 2022						Some concerns
Lipsky, Ann Rheum Dis, 2022	Low	Low	Low	Low	Low	Low
Zhang, RMD Open, 2022	Low	Some concerns	Some concerns	Low	Low	Some concerns
Zheng, JAMA Netw Open, 2022	Low	Low	Low	Low	Low	Low
Vital, Ann Rheum Dis, 2022	Low	Low	Low	Low	Low	Low
Zhang, Front Med, 2022	Some concerns	Low	Some concerns	Low	Low	Some concerns
Merrill, N Engl J Med, 2022	Low	Low	Low	Low	Low	Low

Systematic Literature Review - Report

RoB2	Risk domain					
RCT	Risk of bias arising from the randomization process	Risk of bias due to deviations from the intended interventions	Missing outcome data	Risk of bias in measurement of the outcome	Risk of bias in selection of the reported result	Overall risk of bias
Ye, Am J Transl Res, 2022	Some concerns	Low	Some concerns	High	High	High
Furie, Lupus Sci Med, 2022	Low	Some concerns	Some concerns	Low	Low	Some concerns
Jayne, Ann Rheum Dis, 2022	Low	Some concerns	Some concerns	Low	Some concerns	Some concerns
Jiang, Lupus Sci Med, 2022	Low	Low	Low	Some concerns	Low	Some concerns
Bandhan, Int J Rheum Dis, 2022	High	Some concerns	Low	Low	Low	High
Furie, Ann Rheum Dis, 2022	Low	Low	Low	Low	Low	Low
Rovin, Kidney Int, 2022	Low	Low	Low	Low	Low	Low
Tanaka, RMD Open, 2022	Low	Low	Some concerns	Low	Low	Low
Rovin, Lancet, 2021	Low	Low	Low	Low	Low	Low
Ginzler, Arthritis Rheumatol, 2022	Low	Low	Low	Low	Low	Low
Hasni, Nat Communicat, 2021	Some concerns	Low	Low	Low	Some concerns	Some concerns
Isenberg, Arthritis Rheumatol, 2021	Low	Low	Low	Low	Low	Low
Furie, Rheumatology, 2021	Low	Low	Low	Low	Low	Low
Wallace, Rheumatology, 2021	Low	Low	High	Low	Low	High
Maslen, Lupus Sci Med, 2021	Low	Low	Low	Low	Low	Low
Tummala, Lupus Sci Med, 2021	Low	Low	Low	Low	Low	Low
Barua, Dermatol Ther, 2021	Some concerns	Low	Low	Low	Low	Some concerns
Chatham, Arthritis Rheumatol, 2021	Low	Low	High	Low	Low	High
Furie, N Engl J Med, 2020	Low	Low	Low	Low	Low	Low
Bruce, Lancet Rheumatol, 2021	Low	Low	Some concerns	Low	Low	Some concerns
Morand, Lancet Rheumatol, 2022	Low	Low	Low	Low	Low	Low
Sheikh, Lancet Rheumatol, 2021	Low	Low	Low	Low	Low	Low
29073347 Merrill et al	Some concerns	Some concerns	Low risk	Some concerns	Low risk	Some concerns
29105558 Kamanamool et al	Some concerns	Some concerns	Low risk	Some concerns	Low risk	Some concerns

Systematic Literature Review - Report

RoB2	Risk domain					
RCT	Risk of bias arising from the randomization process	Risk of bias due to deviations from the intended interventions	Missing outcome data	Risk of bias in measurement of the outcome	Risk of bias in selection of the reported result	Overall risk of bias
29295825 Zhang et al	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk of bias
29450636 Mehra et al	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk of bias
29671280 Doria et al	Some concerns	Some concerns	Some concerns	Some concerns	Low risk	High risk of bias
29996800 Sedhain et al	High risk	High risk	Some concerns	Low risk	Low risk	High risk of bias
30043749 Wallace et al	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk of bias
30249507 van Vollenhoven et al	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
30420324 Rovin et al	Low risk	Low risk	Some concerns	Low risk	Low risk	Some concerns
30426311 Zhang et al	Low risk	Some concerns	Some concerns	Low risk	Low risk	Some concerns
30488367 An et al	High risk	High risk	Low risk	Low risk	Some concerns	High risk of bias
31537547 He et al	Some concerns	Low risk	Low risk	Low risk	Low risk	Some concerns
31571750 Bharati et al	High risk	High risk	Low risk	Some concerns	Some concerns	High risk of bias
31851795 Morand et al	Some concerns	Low risk	Low risk	Low risk	Low risk	Some concerns
31852672 Mathian et al	High risk	High risk	Low risk	Low risk	Low risk	High risk of bias
32755035 Atisha-Fregoso et al	Some concerns	Some concerns	Low risk	Some concerns	Low risk	Some concerns
31530556 Mok et al	Some concerns	Low risk	Low risk		Low risk	Some concerns

Risk of bias assessment for meta-analyses

The AMSTAR2 (A MeaSurement Tool to Assess systematic Reviews) tool was used to assess the risk of bias of meta-analyses of RCTs and quasi-RCTs [4]. Meta-analyses of cohort studies and network meta-analyses were not considered for evaluation. Each eligible study was assessed using a checklist of sixteen items (https://amstar.ca/Amstar_Checklist.php) including seven critical domains (registration of a predefined protocol, adequacy of literature search, justification for excluding individual studies, risk of bias from individual studies, appropriateness of meta-analytical methods, consideration of risk of bias when interpreting the results of the review, and assessment of presence of publication bias). Based on the ratings a study is of high, moderate, low, or critically low quality.

Systematic Literature Review - Report

Meta-analysis

Oon, Semin Arthritis Rheum, 2018

Tunnicliffe, Cochrane Database Syst Rev, 2018

Alshaiki, Eur J Rheumatol, 2018

Deng, Turk J Med Sci, 2018

Thong, Lupus, 2019

Zhong, Drug Des Devel Ther, 2019

Zhou, Drug Des Devel Ther, 2019

Liu, Clin Rheumatol, 2019

Zhou, J Pharm Pharm Sci, 2019

Yang, Clin Rheumatol, 2020

Chasset, J Am Acad Dermatol, 2018

Gu, Arch Osteoporos, 2019

Kneeland, Arthritis Care Res, 2022

Liu, Front Immunol, 2022

Lee, Lupus, 2022

Wu, Front Immunol, 2022

Chen, J Clin Rheumatol, 2022

Chiang, Lupus, 2022

Teng, Int J Rheum Dis, 2022

Xie, Lupus Sci Med, 2021

Lee, Z Rheumatol, 2021

Zhang, Medicine, 2020

Koh, Lupus, 2020

Jiang, Medicine, 2020

Ji, Lupus Sci Med, 2022

Quality of study based on AMSTAR2

Critically Low

High

Critically Low

Critically Low

Critically Low

Critically Low

Critically Low

Critically Low

Critically Low

Critically Low

Critically Low

Critically Low

Critically Low

Critically Low

Critically Low

Critically Low

Critically Low

Critically Low

Critically Low

Low

Critically Low

Low

Critically Low

Critically Low

Critically Low

Systematic Literature Review - Report

References

1. Haddaway N, Page M, Pritchard C, McGuinness L. PRISMA2020: An R package and Shiny app for producing PRISMA 2020-compliant flow diagrams, with interactivity for optimised digital transparency and Open Synthesis. *Campbell Systematic Reviews*. 2022 03/27; 18.
2. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol*. 2010 Sep; 25(9):603-605.
3. Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ*. 2019; 366:l4898.
4. Shea BJ, Reeves BC, Wells G, Thuku M, Hamel C, Moran J, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ*. 2017; 358:j4008.