

Table of Contents

T. Quistgaard

Diffusion of Water Isotopes

Continued Analysis

Outlook

- 1 **Readying the Signals**
 - Diffusion of Water Isotopes
 - Data
 - Volcanic Horizons
 - Back Diffusion
- 2 **Continued Analysis**
 - Peak Detection
 - Random Gaps/Missing Data
 - Linear Timescale
- 3 **Outlook**

- Fick's 2nd law:

$$\frac{\partial \delta}{\partial t} = D(t) \frac{\partial^2 \delta}{\partial z^2} - \dot{\epsilon}_z(t) z \frac{\partial \delta}{\partial z} \quad (1)$$

with steady state solution

and

Diffusion in Firn

- Fick's 2nd law:

$$\frac{\partial \delta}{\partial t} = D(t) \frac{\partial^2 \delta}{\partial z^2} - \dot{\epsilon}_z(t) z \frac{\partial \delta}{\partial z} \quad (1)$$

with steady state solution

$$\delta_{\text{meas}}(z) = S(z)[\delta_{\text{init}}(z) * \mathcal{G}(z)] \quad (2)$$

and

Diffusion in Firn

T. Quistgaard

Readying the
SignalsDiffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued
Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook

- Fick's 2nd law:

$$\frac{\partial \delta}{\partial t} = D(t) \frac{\partial^2 \delta}{\partial z^2} - \dot{\epsilon}_z(t) z \frac{\partial \delta}{\partial z} \quad (1)$$

with steady state solution

$$\delta_{\text{meas}}(z) = S(z) [\delta_{\text{init}}(z) * \mathcal{G}(z)] \quad (2)$$

where $\delta_{\text{meas}}(z)$ is the measured signal, $\delta_{\text{init}}(z)$ is the initial isotopic signal

$$\mathcal{G}(z) = \frac{1}{\sigma(z)\sqrt{2\pi}} e^{-\frac{z^2}{2\sigma(z)^2}}, \quad \text{a Gaussian filter,} \quad (3)$$

and

$$S(z) = e^{\int_0^z \dot{\epsilon}_z(z') dz'}, \quad \text{the thinning function} \quad (4)$$

Diffusion in Firn

- Fick's 2nd law:

$$\frac{\partial \delta}{\partial t} = D(t) \frac{\partial^2 \delta}{\partial z^2} - \dot{\epsilon}_z(t) z \frac{\partial \delta}{\partial z} \quad (1)$$

with steady state solution

$$\delta_{\text{meas}}(z) = S(z) [\delta_{\text{init}}(z) * \mathcal{G}(z)] \quad (2)$$

where $\delta_{\text{meas}}(z)$ is the measured signal, $\delta_{\text{init}}(z)$ is the initial isotopic signal

$$\mathcal{G}(z) = \frac{1}{\sigma(z)\sqrt{2\pi}} e^{-\frac{z^2}{2\sigma(z)^2}}, \quad \text{a Gaussian filter,} \quad (3)$$

and

$$S(z) = e^{\int_0^z \dot{\epsilon}_z(z') dz'}, \quad \text{the thinning function} \quad (4)$$

Diffusion in Firn

T. Quistgaard

Diffusion of Water Isotopes

Continued Analysis

Outlook

- Fick's 2nd law:

$$\frac{\partial \delta}{\partial t} = D(t) \frac{\partial^2 \delta}{\partial z^2} - \dot{\epsilon}_z(t) z \frac{\partial \delta}{\partial z} \quad (1)$$

with steady state solution

$$\delta_{\text{meas}}(z) = S(z)[\delta_{\text{init}}(z) * \mathcal{G}(z)] \quad (2)$$

$$\mathcal{G}(z) = \frac{1}{\sigma(z)\sqrt{2\pi}} e^{-\frac{z^2}{2\sigma(z)^2}}, \quad \text{a Gaussian filter,} \quad (3)$$

and

$$\mathcal{S}(z) = e^{\int_0^z \epsilon_z(z') dz'}, \quad \text{the thinning function} \quad (4)$$

Diffusion in Firn

T. Quistgaard

Diffusion of Water Isotopes

Continued Analysis

Outlook

- Fick's 2nd law:

$$\frac{\partial \delta}{\partial t} = D(t) \frac{\partial^2 \delta}{\partial z^2} - \dot{\epsilon}_z(t) z \frac{\partial \delta}{\partial z} \quad (1)$$

with steady state solution

$$\delta_{\text{meas}}(z) = S(z)[\delta_{\text{init}}(z) * \mathcal{G}(z)] \quad (2)$$

$$\mathcal{G}(z) = \frac{1}{\sigma(z)\sqrt{2\pi}} e^{-\frac{z^2}{2\sigma(z)^2}}, \quad \text{a Gaussian filter,} \quad (3)$$

and

$$\mathcal{S}(z) = e^{\int_0^z \epsilon_z(z') dz'}, \quad \text{the thinning function} \quad (4)$$

Table of Contents

1 Readying the Signals

Diffusion of Water Isotopes

Data

Volcanic Horizons

Back Diffusion

② Continued Analysis

Peak Detection

Random Gaps/Missing Data

Linear Timescale

3 Outlook

Data

Laki to
Tambora

Example Data: Site A

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook




Figure: Example data from Alphabet Core drilled at site A near Crête.

Data

Laki to
Tambora

Unevenly Sampled Data: Spline Interpolation

T. Quistgaard

Readying the
Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued
Analysis

Peak Detection

Random

Gaps/Missing Data

Linear Timescale

Outlook




Figure: Example data from Alphabet Core drilled at site A near Crête. Shows zoom in of data from Laki to Tambora along with spline interpolated data.

Data

Laki to
Tambora

Site A

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook




Figure: Site A, raw and cubic spline interpolated data.

Data

Laki to
Tambora

Site G

T. Quistgaard

Reading the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook




Figure: Site G, raw and cubic spline interpolated data.

Data

Laki to Tambora


T. Quistgaard

Data


Continued Analysis

Outlook

Site A: Density and Diffusion Profiles



(a) Density-depth profiles based on analytical Herron-Langway model. Black is empirical data, blue is purely analytical fit and orange is fudged analytical fit



(b) Modeled diffusion length profile based on empirically computed density profile. Black dashed lines indicate ice depth corresponding to date Laki and Tambora eruptions.

Data

Laki to
Tambora

Site G: Density and Diffusion Profiles

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion


Continued Analysis

Peak Detection


Random
Gaps/Missing Data

Linear Timescale

Outlook



(a) Density-depth profiles based on analytical Herron-Langway model. Black is empirical data, blue is purely analytical fit and orange is fudged analytical fit



(b) Modelled diffusion length profile based on empirically computed density profile. Black dashed lines indicate ice depth corresponding to date Laki and Tambora eruptions.

Table of Contents

1 Readying the Signals

Diffusion of Water Isotopes

Data

Volcanic Horizons

Back Diffusion

② Continued Analysis

Peak Detection

Random Gaps/Missing Data

Linear Timescale

③ Outlook

Laki and Tambora

T. Quistgaard

Readying the
Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued
Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook

- **Electrical Conductivity Measurements (ECM)**
- **(Dielectric Profiling (DEP))**




Table of Contents

1 Readying the Signals

Diffusion of Water Isotopes

Data

Volcanic Horizons

Back Diffusion

② Continued Analysis


Peak Detection

Random Gaps/Missing Data


Linear Timescale

3 Outlook

Spectral Analysis with DCT




Spectral Analysis with DCT



$$P_{\text{tot}} = P_{\text{signal}} + |\hat{\eta}|^2$$

Spectral Analysis with DCT



$$P_{\text{tot}} = P_{\text{signal}} + |\hat{\eta}|^2$$

$$|\hat{\eta}|^2 = \frac{\sigma_\eta^2 \Delta}{|1 - a_1 e^{-ik\Delta}|^2}$$

Spectral Analysis with DCT

T. Quistgaard

Readying the
Signals

Diffusion of Water

Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued
Analysis


Peak Detection

Random

Gaps/Missing Data

Linear Timescale

Outlook



$$P_{\text{tot}} = P_{\text{signal}} + |\hat{\eta}|^2$$

$$|\hat{\eta}|^2 = \frac{\sigma_{\eta}^2 \Delta}{|1 - a_1 e^{-ik\Delta}|^2}$$

$$P_{\text{signal}} = P_0 e^{-k^2 \sigma^2}$$

Spectral Analysis with DCT

T. Quistgaard

Readying the
Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion


Continued
Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook



$$P_{\text{tot}} = P_{\text{signal}} + |\hat{\eta}|^2$$

$$|\hat{\eta}|^2 = \frac{\sigma_{\eta}^2 \Delta}{|1 - a_1 e^{-ik\Delta}|^2}$$

$$P_{\text{signal}} = P_0 e^{-k^2 \sigma^2}$$

Diffusion Lengths and Transfer Functions

$$\hat{\delta}_{\text{meas}} = \hat{\delta}_{\text{init}} \cdot \hat{M} \Leftrightarrow \hat{\delta}_{\text{init}} = \hat{\delta}_{\text{meas}} \cdot \hat{M}^{-1} \quad (5)$$

yielding a restoration filter as

Diffusion Lengths and Transfer Functions

$$\hat{\delta}_{\text{meas}} = \hat{\delta}_{\text{init}} \cdot \hat{M} \Leftrightarrow \hat{\delta}_{\text{init}} = \hat{\delta}_{\text{meas}} \cdot \hat{M}^{-1} \quad (5)$$

Add an optimal Wiener filter to enhance signal and minimize noise:

$$\hat{F} = \frac{P_{\text{signal}}}{P_{\text{signal}} + |\hat{\eta}|^2} \quad (6)$$

yielding a restoration filter as

$$\hat{\delta}_{\text{init}} = \hat{\delta}_{\text{meas}} \cdot \hat{F} \cdot \hat{M}^{-1} = \hat{\delta}_{\text{meas}} \cdot \hat{R} \quad (7)$$

$$\hat{\delta}_{\text{init}} = \hat{\delta}_{\text{meas}} \cdot \hat{F} \cdot \hat{M}^{-1} = \hat{\delta}_{\text{meas}} \cdot \hat{R} \quad (7)$$

Diffusion Lengths and Transfer Functions

yielding a restoration filter as

$$\hat{\delta}_{\text{init}} = \hat{\delta}_{\text{meas}} \cdot \hat{F} \cdot \hat{M}^{-1} = \hat{\delta}_{\text{meas}} \cdot \hat{R} \quad (7)$$




Figure: Frequency filters: The optimal filter found from the PSD (blue), the transfer function (orange), the inverse of the transfer function (green) and the combined signal restoration filter (red).

- 1 **Readying the Signals**
 - Diffusion of Water Isotopes
 - Data
 - Volcanic Horizons
 - Back Diffusion
- 2 **Continued Analysis**
 - Peak Detection
 - Random Gaps/Missing Data
 - Linear Timescale
- 3 **Outlook**

Peak Detection

Laki to
Tambora

Peak Detection

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook

- SciPy.signal.find_peaks
- $N = 32$ years btw Tambora and Laki Eruptions
- Best diffusion length estimate algorithm
- Interpolations and resampling

Peak Detection

Laki to
Tambora

Peak Detection

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook

- SciPy.signal.find_peaks
- $N = 32$ years btw Tambora and Laki Eruptions
- Best diffusion length estimate algorithm
- Interpolations and resampling

Peak Detection

Peak Detection

Laki to
Tambora

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook

- SciPy.signal.find_peaks
- $N = 32$ years btw Tambora and Laki Eruptions
- Best diffusion length estimate algorithm
- Interpolations and resampling

Peak Detection

Laki to
Tambora

Peak Detection

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook

- SciPy.signal.find_peaks
- $N = 32$ years btw Tambora and Laki Eruptions
- Best diffusion length estimate algorithm
- Interpolations and resampling

Peak Detection

Laki to
Tambora

Diffusion Length Estimate Algorithm

T. Quistgaard

Readying the Signals

Diffusion of Water

Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random

Gaps/Missing Data

Linear Timescale

Outlook




Figure: Flowchart of method for diffusion length computation, preliminary analysis steps.

Peak Detection

Laki to
Tambora

Diffusion Length Estimate Algorithm

T. Quistgaard

Readying the
Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued
Analysis

Peak Detection

Random

Gaps/Missing Data

Linear Timescale

Outlook




Figure: Flowchart of method for diffusion length computation, decision chart.

Peak Detection

Laki to
Tambora

Diffusion Length V. Peaks - No Limit

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook




Figure: Diffusion length used in back diffusion versus counted number of peaks in data series for all cores.

Peak Detection

Laki to
Tambora

Diffusion Length V. Peaks - No Limit

T. Quistgaard

Readying the Signals

Diffusion of Water

Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random

Gaps/Missing Data

Linear Timescale

Outlook




Figure: Zoom-in around $N = 32$ peaks and corresponding diffusion length used in back diffusion.

Peak Detection

Laki to
Tambora

Cubic Spline Interpolation: Before Deconvolution

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random

Gaps/Missing Data

Linear Timescale

Outlook




Figure: Cubic spline resampling of raw data.

Peak Detection

Laki to
Tambora

Resampling size V. Diffusion Length Estimate

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random

Gaps/Missing Data

Linear Timescale

Outlook







Figure: Resampling size versus diffusion length estimate to result in $N = 32$ peaks.

Peak Detection

Laki to
Tambora

Cubic Spline Interpolation: After Deconvolution

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook




Figure: Deconvoluted data with resampling of 1 and 9 cm intervals after deconvolution, but before peak



Peak Detection

Laki to
Tambora

Resampling size V. Diffusion Length Estimate

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random

Gaps/Missing Data

Linear Timescale

Outlook









Figure: Resampling size after deconvolution versus diffusion length estimate to result in $N = 32$ peaks.

Peak Detection

Laki to
Tambora

Site A: Theoretical V. Estimated Diffusion Length

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook




Figure: Data and back diffused signal, using theoretically predicted diffusion lengths and diffusion length estimated through analysis, Site A.

Peak Detection

Laki to
Tambora

Site G: Theoretical V. Estimated Diffusion Length

T. Quistgaard

Readying the Signals

Diffusion of Water

Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random

Gaps/Missing Data

Linear Timescale

Outlook




Figure: Data and back diffused signal, using theoretically predicted diffusion lengths and diffusion length estimated through analysis, Site G.

Peak Detection

Laki to
Tambora

Diffusion Lengths, All Cores

T. Quistgaard

Readying the
SignalsDiffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued
Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook

	Crete	Site A	Site B	Site E	Site G
σ_{Theo}^{min} [cm]	7.28	6.52	6.43	7.50	7.66
σ_{Theo}^{max} [cm]	7.49	6.70	6.61	7.76	7.88
σ_{est} [cm]	6.02	6.14	3.27 (N = 32) 5.85 (N = 33)	5.95	8.46

Table of Contents

T. Quistgaard

Continued Analysis

Random
Gaps/Missing Data

Outlook

1 Readying the Signals

Diffusion of Water Isotopes

Data

Volcanic Horizons

Back Diffusion

② Continued Analysis

Peak Detection

Random Gaps/Missing Data

Linear Timescale

3 Outlook

Random Gaps/Missing Data

Laki to
Tambora

Linear Interpolation, 50 and 100 cm gaps

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion



Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook



Random Gaps/Missing Data

Laki to
Tambora

Linear Interpolation, 50 and 100 cm gaps

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion



Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook



Random Gaps/Missing Data

Laki to
Tambora

Cubic Spline Interpolation

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion



Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook



Random Gaps/Missing Data

Laki to
Tambora

Cubic Spline Interpolation

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook





Table of Contents

T. Quistgaard

Continued Analysis

Linear Timescale

Outlook

1 Readying the Signals

Diffusion of Water Isotopes

Data

Volcanic Horizons

Back Diffusion

② Continued Analysis

Peak Detection

Random Gaps/Missing Data

Linear Timescale

3 Outlook




Figure: Data series on nonlinear and linearized timescales, Crete.

Linear Timescale

Laki to
Tambora

Site A

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook




Figure: Data series on nonlinear and linearized timescales, Site A.

Linear Timescale

Laki to
Tambora

Site G

T. Quistgaard

Readying the Signals

Diffusion of Water

Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random

Gaps/Missing Data

Linear Timescale

Outlook




Figure: Data series on nonlinear and linearized timescales, Site G.

Further Work

T. Quistgaard

Readying the Signals

Diffusion of Water
Isotopes

Data

Volcanic Horizons

Back Diffusion

Continued Analysis

Peak Detection

Random
Gaps/Missing Data

Linear Timescale

Outlook

- Peaks and troughs
- Accumulation seasonality
- ECM data back diffusion
- Missing data reconstruction
- Peak/cycle detection through standardization and classification

Actual Total Diffusion

Total diffusion in ice and firn

$$\sigma_{\text{tot}}(z)^2 = [S(z)\sigma_{\text{firn}}(z)]^2 + \sigma_{\text{ice}}(z)^2 \quad (8)$$

Giving an actual measured diffusion length at z_i of

$$\sigma(z_i)^2 = \sigma_{\text{firn}}(z_i)^2 S(z_i) + \sigma_{\text{ice}}(z_i)^2 + \sigma_{\text{dis}}(z_i)^2 \quad (9)$$

with

$$\sigma_{\text{dis}}(z_i)^2 = \frac{2\Delta(z_i)^2}{\pi^2} \ln\left(\frac{\pi}{2}\right) \quad (10)$$

Actual Total Diffusion

Total diffusion in ice and firn

$$\sigma_{\text{tot}}(z)^2 = [S(z)\sigma_{\text{firn}}(z)]^2 + \sigma_{\text{ice}}(z)^2 \quad (8)$$

Giving an actual measured diffusion length at z_i of

$$\sigma(z_i)^2 = \sigma_{\text{firn}}(z_i)^2 S(z_i) + \sigma_{\text{ice}}(z_i)^2 + \sigma_{\text{dis}}(z_i)^2 \quad (9)$$

with

$$\sigma_{\text{dis}}(z_i)^2 = \frac{2\Delta(z_i)^2}{\pi^2} \ln\left(\frac{\pi}{2}\right) \quad (10)$$

Laki and Tambora

- **Electrical Conductivity Measurements (ECM)**
- **Dielectric Profiling (DEP)**

