

# A sample presentation

*With ramblings from a madman*

Théo N. Dionne

APS March Meeting  
Denver Colorado  
December 31, 2025



# Table of contents

## 1 A section

- ➔ A subsection

- ➔ Another subsection

The quick brown fox jumps over the lazy dog [1].

The quick brown fox jumps over the lazy dog [1].

$$i\hbar\partial_t |\Psi\rangle = \hat{H} |\Psi\rangle \tag{1}$$

# What

# What

➡ One.

# What

- ➡ One.
- ➡ Two.

# What

- ➡ One.
- ➡ Two.
- ➡ Three.



# The state of the world

## Theorem 1.1: Birb

Birbs exist.

# Colorbox test

A very important result.

# A test of what the box environment looks like

This is the fundamental theorem of calculus:

$$\int_a^b f(t) \, dt = F(b) - F(a) \quad (2)$$

# A test of what the box environment looks like

This is the fundamental theorem of calculus:

$$\int_a^b f(t) \, dt = F(b) - F(a) \quad (2)$$

Notice:

➡ thing to say;

# A test of what the box environment looks like

This is the fundamental theorem of calculus:

$$\int_a^b f(t) \, dt = F(b) - F(a) \quad (2)$$

Notice:

- ➡ thing to say;
- ➡ another thing to say;

# A test of what the box environment looks like

This is the fundamental theorem of calculus:

$$\int_a^b f(t) \, dt = F(b) - F(a) \quad (2)$$

Notice:

- ➡ thing to say;
- ➡ another thing to say;
- ➡ consider  $F = m\ddot{x}$ .

# A simple derivation

Extremization of the action entails:

$$\boxed{0 = \delta\mathcal{S}} = \int_{t_i}^{t_f} dt \delta\mathcal{L}$$

# A simple derivation

Extremization of the action entails:

$$\boxed{0 = \delta\mathcal{S}} = \int_{t_i}^{t_f} dt \delta\mathcal{L} \\ = \int_{t_i}^{t_f} dt \left[ \frac{\delta\mathcal{L}}{\delta q} \delta q + \frac{\delta\mathcal{L}}{\delta \dot{q}} \delta \dot{q} \right]$$



# A simple derivation

Extremization of the action entails:

$$\begin{aligned}\boxed{0 = \delta\mathcal{S}} &= \int_{t_i}^{t_f} dt \delta\mathcal{L} \\ &= \int_{t_i}^{t_f} dt \left[ \frac{\delta\mathcal{L}}{\delta q} \delta q + \frac{\delta\mathcal{L}}{\delta \dot{q}} \delta \dot{q} \right] \\ &= \int_{t_i}^{t_f} dt \left[ \frac{\delta\mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta\mathcal{L}}{\delta \dot{q}} \right] \delta q + \underbrace{\int_{t_i}^{t_f} dt \frac{d}{dt} \left( \frac{\delta\mathcal{L}}{\delta \dot{q}} \delta q \right)}_{\text{vanishes (B.C.s)}}\end{aligned}$$

# A simple derivation

Extremization of the action entails:

$$\begin{aligned}\boxed{0 = \delta\mathcal{S}} &= \int_{t_i}^{t_f} dt \delta\mathcal{L} \\ &= \int_{t_i}^{t_f} dt \left[ \frac{\delta\mathcal{L}}{\delta q} \delta q + \frac{\delta\mathcal{L}}{\delta \dot{q}} \delta \dot{q} \right] \\ &= \int_{t_i}^{t_f} dt \left[ \frac{\delta\mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta\mathcal{L}}{\delta \dot{q}} \right] \delta q + \underbrace{\int_{t_i}^{t_f} dt \frac{d}{dt} \left( \frac{\delta\mathcal{L}}{\delta \dot{q}} \delta q \right)}_{\text{vanishes (B.C.s)}} \\ &\Rightarrow \boxed{\frac{\delta\mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta\mathcal{L}}{\delta \dot{q}} = 0}\end{aligned}$$

# A simple derivation

Extremization of the action entails:

$$\begin{aligned}\boxed{0 = \delta\mathcal{S}} &= \int_{t_i}^{t_f} dt \delta\mathcal{L} \\ &= \int_{t_i}^{t_f} dt \left[ \frac{\delta\mathcal{L}}{\delta q} \delta q + \frac{\delta\mathcal{L}}{\delta \dot{q}} \delta \dot{q} \right] \\ &= \int_{t_i}^{t_f} dt \left[ \frac{\delta\mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta\mathcal{L}}{\delta \dot{q}} \right] \delta q + \underbrace{\int_{t_i}^{t_f} dt \frac{d}{dt} \left( \frac{\delta\mathcal{L}}{\delta \dot{q}} \delta q \right)}_{\text{vanishes (B.C.s)}} \quad (3) \\ &\Rightarrow \boxed{\frac{\delta\mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta\mathcal{L}}{\delta \dot{q}} = 0}\end{aligned}$$

The converse is equally true, hence:

# A simple derivation

Extremization of the action entails:

$$\begin{aligned}\boxed{0 = \delta \mathcal{S}} &= \int_{t_i}^{t_f} dt \delta \mathcal{L} \\ &= \int_{t_i}^{t_f} dt \left[ \frac{\delta \mathcal{L}}{\delta q} \delta q + \frac{\delta \mathcal{L}}{\delta \dot{q}} \delta \dot{q} \right] \\ &= \int_{t_i}^{t_f} dt \left[ \frac{\delta \mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta \mathcal{L}}{\delta \dot{q}} \right] \delta q + \underbrace{\int_{t_i}^{t_f} dt \frac{d}{dt} \left( \frac{\delta \mathcal{L}}{\delta \dot{q}} \delta q \right)}_{\text{vanishes (B.C.s)}} \quad (3) \\ &\Rightarrow \boxed{\frac{\delta \mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta \mathcal{L}}{\delta \dot{q}} = 0}\end{aligned}$$

The converse is equally true, hence:

$$\boxed{\delta \mathcal{S} = 0 \text{ \& B.C.s} \iff \frac{\delta \mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta \mathcal{L}}{\delta \dot{q}} = 0} \quad (4)$$

# Questions (2 minutes)

**Thank You!**

# Bibliography I



J. Hubbard et Brian Hilton Flowers.

Electron correlations in narrow energy bands.

*Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* **276**(1365), 238–257 (1963).

[doi:10.1098/rspa.1963.0204](https://doi.org/10.1098/rspa.1963.0204).



J. Hubbard et Brian Hilton Flowers.

Electron correlations in narrow energy bands. ii. the degenerate band case.

*Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* **277**(1369), 237–259 (1964).

[doi:10.1098/rspa.1964.0019](https://doi.org/10.1098/rspa.1964.0019).

# Bibliography II



J. Hubbard et Brian Hilton Flowers.

Electron correlations in narrow energy bands iii. an improved solution.

*Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* **281**(1386), 401–419 (1964).

[doi:10.1098/rspa.1964.0190](https://doi.org/10.1098/rspa.1964.0190).