A sample presentation

With ramblings from a madman

Théo N. Dionne

APS March Meeting
Denver Colorado
5 septembre 2026





Table of contents

1 A section
- A subsection
- Another subsection

Title

The quick brown fox jumps over the lazy dog [1].

Title

The quick brown fox jumps over the lazy dog [1].

$$\boxed{i\hbar\partial_t |\Psi\rangle = \hat{\mathrm{H}} |\Psi\rangle} \tag{1}$$

What

What

- One.

What

- One.
- Two.

What

- One.
- Two.
- Three.

The state of the world

Théorème 1.1 : Birb

Birbs exist.

Colorbox test

A very important result.

A test of what the box environment looks like

This is the fundamental theorem of calculus :

$$\int_a^b f(t)\,\mathrm{d}t = F(b) - F(a) \tag{2}$$

A test of what the box environment looks like

This is the fundamental theorem of calculus :

$$\int_a^b f(t)\,\mathrm{d}t = F(b) - F(a) \tag{2}$$

Notice :

- thing to say ;

A test of what the box environment looks like

This is the fundamental theorem of calculus :

$$\int_a^b f(t)\,\mathrm{d}t = F(b) - F(a) \tag{2}$$

Notice :

- thing to say ;
- another thing to say ;

A test of what the box environment looks like

This is the fundamental theorem of calculus :

$$\int_a^b f(t)\,\mathrm{d}t = F(b) - F(a) \tag{2}$$

Notice :

- thing to say ;
- another thing to say ;
- consider $\mathbf{F} = m\ddot{\mathbf{x}}$.

A simple derivation

Extremization of the action entails :

$$\boxed{0 = \delta \mathcal{S}} = \int_{t_i}^{t_f} \mathrm{d}t \, \delta \mathcal{L}$$

A simple derivation

Extremization of the action entails :

$$\boxed{0 = \delta \mathcal{S}} = \int_{t_i}^{t_f} \mathrm{d}t \, \delta \mathcal{L}$$

$$= \int_{t_i}^{t_f} \mathrm{d}t \left[\frac{\delta \mathcal{L}}{\delta q} \delta q + \frac{\delta \mathcal{L}}{\delta \dot{q}} \delta \dot{q} \right]$$

A simple derivation

Extremization of the action entails :

$$\boxed{0 = \delta \mathcal{S}} = \int_{t_i}^{t_f} \mathrm{d}t \, \delta \mathcal{L}$$

$$= \int_{t_i}^{t_f} \mathrm{d}t \left[\frac{\delta \mathcal{L}}{\delta q} \delta q + \frac{\delta \mathcal{L}}{\delta \dot{q}} \delta \dot{q} \right]$$

$$= \int_{t_i}^{t_f} \mathrm{d}t \left[\frac{\delta \mathcal{L}}{\delta q} - \frac{\mathrm{d}}{\mathrm{d}t} \frac{\delta \mathcal{L}}{\delta \dot{q}} \right] \delta q + \underbrace{\int_{t_i}^{t_f} \mathrm{d}t \frac{\mathrm{d}}{\mathrm{d}t} \left(\frac{\delta \mathcal{L}}{\delta \dot{q}} \delta q \right)}_{\text{vanishes (B.C.s)}}$$

A simple derivation

Extremization of the action entails :

$$\boxed{0 = \delta \mathcal{S}} = \int_{t_i}^{t_f} \mathrm{d}t \, \delta \mathcal{L}$$

$$= \int_{t_i}^{t_f} \mathrm{d}t \left[\frac{\delta \mathcal{L}}{\delta q} \delta q + \frac{\delta \mathcal{L}}{\delta \dot{q}} \delta \dot{q} \right]$$

$$= \int_{t_i}^{t_f} \mathrm{d}t \left[\frac{\delta \mathcal{L}}{\delta q} - \frac{\mathrm{d}}{\mathrm{d}t} \frac{\delta \mathcal{L}}{\delta \dot{q}} \right] \delta q + \underbrace{\int_{t_i}^{t_f} \mathrm{d}t \frac{\mathrm{d}}{\mathrm{d}t} \left(\frac{\delta \mathcal{L}}{\delta \dot{q}} \delta q \right)}_{\text{vanishes (B.C.s)}}$$

$$\Rightarrow \boxed{\frac{\delta \mathcal{L}}{\delta q} - \frac{\mathrm{d}}{\mathrm{d}t} \frac{\delta \mathcal{L}}{\delta \dot{q}} = 0}$$

A simple derivation

Extremization of the action entails :

$$\boxed{0 = \delta \mathcal{S}} = \int_{t_i}^{t_f} \mathrm{d}t\, \delta \mathcal{L}$$

$$= \int_{t_i}^{t_f} \mathrm{d}t \left[\frac{\delta \mathcal{L}}{\delta q} \delta q + \frac{\delta \mathcal{L}}{\delta \dot{q}} \delta \dot{q} \right]$$

$$= \int_{t_i}^{t_f} \mathrm{d}t \left[\frac{\delta \mathcal{L}}{\delta q} - \frac{\mathrm{d}}{\mathrm{d}t} \frac{\delta \mathcal{L}}{\delta \dot{q}} \right] \delta q + \underbrace{\int_{t_i}^{t_f} \mathrm{d}t \frac{\mathrm{d}}{\mathrm{d}t} \left(\frac{\delta \mathcal{L}}{\delta \dot{q}} \delta q \right)}_{\text{vanishes (B.C.s)}}$$

$$\Rightarrow \boxed{\frac{\delta \mathcal{L}}{\delta q} - \frac{\mathrm{d}}{\mathrm{d}t} \frac{\delta \mathcal{L}}{\delta \dot{q}} = 0} \tag{3}$$

The converse is equally true, hence :

A simple derivation

Extremization of the action entails :

$$\boxed{0 = \delta\mathcal{S}} = \int_{t_i}^{t_f} \mathrm{d}t\, \delta\mathcal{L}$$

$$= \int_{t_i}^{t_f} \mathrm{d}t \left[\frac{\delta\mathcal{L}}{\delta q}\delta q + \frac{\delta\mathcal{L}}{\delta\dot{q}}\delta\dot{q}\right]$$

$$= \int_{t_i}^{t_f} \mathrm{d}t \left[\frac{\delta\mathcal{L}}{\delta q} - \frac{\mathrm{d}}{\mathrm{d}t}\frac{\delta\mathcal{L}}{\delta\dot{q}}\right]\delta q + \underbrace{\int_{t_i}^{t_f} \mathrm{d}t \frac{\mathrm{d}}{\mathrm{d}t}\left(\frac{\delta\mathcal{L}}{\delta\dot{q}}\delta q\right)}_{\text{vanishes (B.C.s)}}$$

$$\Rightarrow \boxed{\frac{\delta\mathcal{L}}{\delta q} - \frac{\mathrm{d}}{\mathrm{d}t}\frac{\delta\mathcal{L}}{\delta\dot{q}} = 0} \tag{3}$$

The converse is equally true, hence :

$$\boxed{\delta\mathcal{S} = 0 \;\&\; \text{B.C.s} \Longleftrightarrow \frac{\delta\mathcal{L}}{\delta q} - \frac{\mathrm{d}}{\mathrm{d}t}\frac{\delta\mathcal{L}}{\delta\dot{q}} = 0} \tag{4}$$

Testing columns

This is a Woodcock. Its features are :



Testing columns

This is a Woodcock. Its features are :

- A beak



Testing columns

This is a Woodcock. Its features are :

- A beak
- Wings



Testing columns

This is a Woodcock. Its features are :

- A beak
- Wings
- A brain (barely)



Testing columns

This is a Woodcock. Its features are :

- A beak
- Wings
- A brain (barely)
- Camouflage



Questions (2 minutes)

Thank You!

Bibliography I

1. Hubbard, J. & Flowers, B. H. Electron correlations in narrow energy bands. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* **276**, 238-257. eprint : https://royalsocietypublishing.org/doi/pdf/10.1098/rspa.1963.0204. https://royalsocietypublishing.org/doi/abs/10.1098/rspa.1963.0204 (1963) (cf. p. 3, 4).

2. Hubbard, J. & Flowers, B. H. Electron correlations in narrow energy bands. II. The degenerate band case. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* **277**, 237-259. eprint : https://royalsocietypublishing.org/doi/pdf/10.1098/rspa.1964.0019. https://royalsocietypublishing.org/doi/abs/10.1098/rspa.1964.0019 (1964).

Bibliography II

3. Hubbard, J. & Flowers, B. H. Electron correlations in narrow energy bands III. An improved solution. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* **281**, 401-419. eprint : https://royalsocietypublishing.org/doi/pdf/10.1098/rspa.1964.0190. https://royalsocietypublishing.org/doi/abs/10.1098/rspa.1964.0190 (1964).