

A sample presentation

With ramblings from a madman

Théo N. Dionne

APS March Meeting
Denver Colorado
3 mars 2026



Table of contents

- 1 A section
 - A subsection
 - Another subsection

The quick brown fox jumps over the lazy dog [1].

The quick brown fox jumps over the lazy dog [1].

$$i\hbar\partial_t |\Psi\rangle = \hat{H} |\Psi\rangle \quad (1)$$

What

What

➡ One.

What

- ➡ One.
- ➡ Two.

What

- ➡ One.
- ➡ Two.
- ➡ Three.

Théorème 1.1 : Birb

Birbs exist.

Colorbox test

A very important result.

A test of what the box environment looks like

This is the fundamental theorem of calculus :

$$\int_a^b f(t) dt = F(b) - F(a) \quad (2)$$

A test of what the box environment looks like

This is the fundamental theorem of calculus :

$$\int_a^b f(t) dt = F(b) - F(a) \quad (2)$$

Notice :

➡ thing to say;

A test of what the box environment looks like

This is the fundamental theorem of calculus :

$$\int_a^b f(t) dt = F(b) - F(a) \quad (2)$$

Notice :

- ➡ thing to say;
- ➡ another thing to say;

A test of what the box environment looks like

This is the fundamental theorem of calculus :

$$\int_a^b f(t) dt = F(b) - F(a) \quad (2)$$

Notice :

- ➡ thing to say;
- ➡ another thing to say;
- ➡ consider $F = m\ddot{x}$.

A simple derivation

Extremization of the action entails :

$$\boxed{0 = \delta \mathcal{S}} = \int_{t_i}^{t_f} dt \delta \mathcal{L}$$

A simple derivation

Extremization of the action entails :

$$\boxed{0 = \delta \mathcal{S}} = \int_{t_i}^{t_f} dt \delta \mathcal{L} \\ = \int_{t_i}^{t_f} dt \left[\frac{\delta \mathcal{L}}{\delta q} \delta q + \frac{\delta \mathcal{L}}{\delta \dot{q}} \delta \dot{q} \right]$$

A simple derivation

Extremization of the action entails :

$$\begin{aligned}\boxed{0 = \delta \mathcal{S}} &= \int_{t_i}^{t_f} dt \delta \mathcal{L} \\ &= \int_{t_i}^{t_f} dt \left[\frac{\delta \mathcal{L}}{\delta q} \delta q + \frac{\delta \mathcal{L}}{\delta \dot{q}} \delta \dot{q} \right] \\ &= \int_{t_i}^{t_f} dt \left[\frac{\delta \mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta \mathcal{L}}{\delta \dot{q}} \right] \delta q + \underbrace{\int_{t_i}^{t_f} dt \frac{d}{dt} \left(\frac{\delta \mathcal{L}}{\delta \dot{q}} \delta q \right)}_{\text{vanishes (B.C.s)}}\end{aligned}$$

A simple derivation

Extremization of the action entails :

$$\begin{aligned}\boxed{0 = \delta \mathcal{S}} &= \int_{t_i}^{t_f} dt \delta \mathcal{L} \\&= \int_{t_i}^{t_f} dt \left[\frac{\delta \mathcal{L}}{\delta q} \delta q + \frac{\delta \mathcal{L}}{\delta \dot{q}} \delta \dot{q} \right] \\&= \int_{t_i}^{t_f} dt \left[\frac{\delta \mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta \mathcal{L}}{\delta \dot{q}} \right] \delta q + \underbrace{\int_{t_i}^{t_f} dt \frac{d}{dt} \left(\frac{\delta \mathcal{L}}{\delta \dot{q}} \delta q \right)}_{\text{vanishes (B.C.s)}} \\&\Rightarrow \boxed{\frac{\delta \mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta \mathcal{L}}{\delta \dot{q}} = 0}\end{aligned}$$

A simple derivation

Extremization of the action entails :

$$\begin{aligned}\boxed{0 = \delta \mathcal{S}} &= \int_{t_i}^{t_f} dt \delta \mathcal{L} \\&= \int_{t_i}^{t_f} dt \left[\frac{\delta \mathcal{L}}{\delta q} \delta q + \frac{\delta \mathcal{L}}{\delta \dot{q}} \delta \dot{q} \right] \\&= \int_{t_i}^{t_f} dt \left[\frac{\delta \mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta \mathcal{L}}{\delta \dot{q}} \right] \delta q + \underbrace{\int_{t_i}^{t_f} dt \frac{d}{dt} \left(\frac{\delta \mathcal{L}}{\delta \dot{q}} \delta q \right)}_{\text{vanishes (B.C.s)}} \\&\Rightarrow \boxed{\frac{\delta \mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta \mathcal{L}}{\delta \dot{q}} = 0}\end{aligned}\tag{3}$$

The converse is equally true, hence :

A simple derivation

Extremization of the action entails :

$$\begin{aligned}\boxed{0 = \delta \mathcal{S}} &= \int_{t_i}^{t_f} dt \delta \mathcal{L} \\&= \int_{t_i}^{t_f} dt \left[\frac{\delta \mathcal{L}}{\delta q} \delta q + \frac{\delta \mathcal{L}}{\delta \dot{q}} \delta \dot{q} \right] \\&= \int_{t_i}^{t_f} dt \left[\frac{\delta \mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta \mathcal{L}}{\delta \dot{q}} \right] \delta q + \underbrace{\int_{t_i}^{t_f} dt \frac{d}{dt} \left(\frac{\delta \mathcal{L}}{\delta \dot{q}} \delta q \right)}_{\text{vanishes (B.C.s)}} \\&\Rightarrow \boxed{\frac{\delta \mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta \mathcal{L}}{\delta \dot{q}} = 0}\end{aligned}\tag{3}$$

The converse is equally true, hence :

$$\boxed{\delta \mathcal{S} = 0 \text{ \& B.C.s} \iff \frac{\delta \mathcal{L}}{\delta q} - \frac{d}{dt} \frac{\delta \mathcal{L}}{\delta \dot{q}} = 0}\tag{4}$$

Testing columns

This is a Woodcock. Its features are :



Testing columns

This is a Woodcock. Its features are :

➡ A beak



Testing columns

This is a Woodcock. Its features are :

- ➡ A beak
- ➡ Wings



Testing columns

This is a Woodcock. Its features are :

- ➡ A beak
- ➡ Wings
- ➡ A brain (barely)



Testing columns

This is a Woodcock. Its features are :

- ➡ A beak
- ➡ Wings
- ➡ A brain (barely)
- ➡ Camouflage



Questions (2 minutes)

Thank You!

Bibliography I



J. Hubbard et Brian Hilton Flowers.

Electron correlations in narrow energy bands.

Proceedings of the Royal Society of London. Series A.

Mathematical and Physical Sciences **276**(1365), 238–257
(1963).

doi:10.1098/rspa.1963.0204.



J. Hubbard et Brian Hilton Flowers.

Electron correlations in narrow energy bands. ii. the
degenerate band case.

Proceedings of the Royal Society of London. Series A.

Mathematical and Physical Sciences **277**(1369), 237–259
(1964).

doi:10.1098/rspa.1964.0019.



J. Hubbard et Brian Hilton Flowers.

Electron correlations in narrow energy bands iii. an improved solution.

Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences **281**(1386), 401–419 (1964).

doi:10.1098/rspa.1964.0190.