


Intensity Mapping for Extended Ly α Emission around $z \sim 2-7$ LAEs

Shotaro Kikuchi-hara
(D3, ICRR, University of Tokyo)


Y. Harikane, M. Ouchi, Y. Ono, T. Shibuya, R. Itoh, R. Kakuma, A. K. Inoue, R. Momose, K. Shimasaku,
H. Kusakabe, Y. Sugahara, S. Kikuta, N. Kashikawa, S. Saito, H. Zhang, & C.-H. Lee

Galaxy-IGM Workshop 2021

Why Ly α Halos?



- Gas surrounding galaxy = Circumgalactic medium (CGM)
 - CGM fuel star formation in galaxy
 - Hydrogen gas in CGM can be traced by Ly α → “Ly α halo” (LAH)
 - Surface brightness (SB) radial profiles around Ly α emitters (LAEs)?



Observation & Sample

- Data: narrow-band (NB) images from Subaru/HSC intensive surveys

HSC-SSP

NB816 ($\lambda=8177\text{\AA}$), NB921 ($\lambda=9215\text{\AA}$) → $z=5.7$ & 6.6 LAEs

Subaru Strategic Program (Aihara+19)

CHORUS

NB387 ($\lambda=3863\text{\AA}$), NB527 ($\lambda=5260\text{\AA}$) → $z=2.2$ & 3.3 LAEs

Cosmic Hydrogen Reionization Unveiled with Subaru (Inoue+20)

- UltraDeep (UD) layer images ($5\sigma \sim 26$ mag) → advantageous to detect very diffuse emission
- COSMOS+SXDS fields ($\sim 4 \text{ deg}^2$) → large LAE sample

- Sample: from Ono+21 (as a part of




- Selected from S18A images
- $N=369$ ($z=2.2$), 793 ($z=3.3$), 603 ($z=5.7$), 80 ($z=6.6$)
→ Total 1,845 LAEs

Added >1,000 LAEs at $z=2.2\&3.3$ to
Kakuma+21




©https://hsc.mtk.nao.ac.jp/ssp/instrument_jp/

Method: Intensity Mapping



Comaschi&Ferrara16, Li+16, Kovetz+17

Result: SB Radial Profiles



* SB are normalized such that luminosities in $r < 1''$ become similar
Our SB are averaged over COSMOS & SXDS fields at $z=5.7$ & 6.6

- No clear detection at $z=2.2$
- Newly detected ($S/N=4.1$) at $z=3.3$
- Potential detection ($S/N \sim 2$) at $z=5.7$ & 6.6

- Our SB agree w/ previous results at $\sim R_{\text{vir}}$ when compared under similar luminosity


Ma12: Matsuda+12 (stacking)
 Mo14: Momose+14 (stacking)
 Mo16: Momose+16 (stacking)
 Le17: Leclercq+17 (individual detection)
 Wi18: Wisotzki+18 (stacking)
 Wu20: Wu+20 (stacking)
 Ka21: Kakuma+21 (intensity mapping)

What Power Ly α Emission?

- Possible physical *processes* (*how* Ly α photons are produced):




→ Alternatively, we discuss based on Ly α emission *origins* (*where* Ly α photons are produced)



©Momose-san's review talk from IGM-Galaxy Workshop 2020
See also Umehata-san's review talk in this morning

What Power Ly α Emission? (1): Central Galaxy



- Star formation in central galaxy
 - Ly α photons via recombination
 - transfer outward while being scattered

Results

- Zheng+11 reproduces SB_{obs} ($z=5.7$)
- All model predictions lie below SB_{obs} ($z=3.3$)

Interpretations


- Contribution decreases toward low-z?
- Discrepancy among models?

Scattered Ly α from central galaxy is
candidate of ELE powering source

c.f.

- IGM scattering → plateau-like SB profile (Jeeson-Danieli+12)
- Ly α line profile asymmetry (Chen+21)
- Halo luminosity vs mass relation (Kusakabe+19)

What Power Ly α Emission? (2): CGM



- Star formation in central galaxy
 - ionizing photons
 - fluorescent emission in CGM

Results


- Predicted SB drop at R_{vir}
 - CGM contribution alone cannot reproduce SB_{obs}

Interpretations

CGM fluorescence *unlikely* to power ELE

c.f. Cooling radiation contribution is not dominant
(Rosdahl&Blaizot+12, Byrohl+20)

What Power Ly α Emission? (3): Satellite Galaxies



- Star formation in satellite galaxies
→ Ly α photons via recombination

Results


- Predicted SB \ll SB_{obs} (except at z=2.2)

Interpretations

Satellite galaxies *only marginally* contribute to ELE

c.f. SB(Ly α) is more extended than SB(UV)
(e.g., Momose+14, +16, Wu+20)

What Power Ly α Emission? (4): Other Halos



※ SB are normalized such that luminosities in $r < 1''$ become similar

- Ly α emission from other halos

Results

- Model predictions agree w/ SB_{obs} ($z=3.3$)
- Zheng+11 overpredicts SB_{obs} ($z=5.7$)

Interpretations

- Masking bright sources $\rightarrow SB_{obs} < SB_{model}$ (Kakuma+21)
 \rightarrow Then why $SB_{model} = SB_{obs}$ at $z=3.3$?

Other halos *may contribute* to ELE

c.f. MXDF (Bacon+21):
population of faint galaxies can reproduce ELE

Summary

- Intensity mapping to 1,845 LAEs ($z=2.2\text{-}6.6$) + Subaru/HSC NB images (HSC-SSP & CHORUS)
 - Very diffuse “extended Ly α emission” (ELE) beyond R_{vir} identified around $z=3.3\text{-}6.6$ LAEs
 - ELE can be reproduced by ...
 - Scattered Ly α photons produced in central galaxy
 - Contribution from other halos
- This is first systematic investigation of ELE across $z\sim 2\text{-}7$
 - Systematic comparison w/ models is necessary to tackle physical origins

Origins differ among LAEs, populations (LAE/LBG/QSO), radial scales ($<R_{\text{vir}}$ or $>R_{\text{vir}}$), redshifts, etc. Thus systematic comparison is crucial.