GLOW LAB
PROTOCOLS

GLOW FUEL

NAD+ 100 mg + MOTS-C 10 mg + 5-Amino-1MQ 10 mg
120 mg total per bottle

Final volume: 30 mL
Concentration: 4 mg/mL combined
(NAD+ 3.3333 · MOTS-C 0.3333 · 5-Amino-1MQ 0.3333 mg/mL)
Calculated amount: 600 mcg (0.6 mg) per 0.15 mL spray
(NAD+ 500 + MOTS-C 50 + 5-Amino-1MQ 50 mcg)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

GLOW FUEL

NAD+ 100 mg + MOTS-C 10 mg + 5-Amino-1MQ 10 mg
120 mg total per bottle

Final volume: 30 mL
Concentration: 4 mg/mL combined
(NAD+ 3.3333 · MOTS-C 0.3333 · 5-Amino-1MQ 0.3333 mg/mL)
Calculated amount: 600 mcg (0.6 mg) per 0.15 mL spray
(NAD+ 500 + MOTS-C 50 + 5-Amino-1MQ 50 mcg)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

GLOW FUEL

NAD+ 100 mg + MOTS-C 10 mg + 5-Amino-1MQ 10 mg
120 mg total per bottle

Final volume: 30 mL
Concentration: 4 mg/mL combined
(NAD+ 3.3333 · MOTS-C 0.3333 · 5-Amino-1MQ 0.3333 mg/mL)
Calculated amount: 600 mcg (0.6 mg) per 0.15 mL spray
(NAD+ 500 + MOTS-C 50 + 5-Amino-1MQ 50 mcg)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

GLOW FUEL

NAD+ 100 mg + MOTS-C 10 mg + 5-Amino-1MQ 10 mg
120 mg total per bottle

Final volume: 30 mL
Concentration: 4 mg/mL combined
(NAD+ 3.3333 · MOTS-C 0.3333 · 5-Amino-1MQ 0.3333 mg/mL)
Calculated amount: 600 mcg (0.6 mg) per 0.15 mL spray
(NAD+ 500 + MOTS-C 50 + 5-Amino-1MQ 50 mcg)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

GLOW FUEL

NAD+ 100 mg + MOTS-C 10 mg + 5-Amino-1MQ 10 mg
120 mg total per bottle

Final volume: 30 mL
Concentration: 4 mg/mL combined
(NAD+ 3.3333 · MOTS-C 0.3333 · 5-Amino-1MQ 0.3333 mg/mL)
Calculated amount: 600 mcg (0.6 mg) per 0.15 mL spray
(NAD+ 500 + MOTS-C 50 + 5-Amino-1MQ 50 mcg)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

GLOW FUEL

NAD+ 100 mg + MOTS-C 10 mg + 5-Amino-1MQ 10 mg
120 mg total per bottle

Final volume: 30 mL
Concentration: 4 mg/mL combined
(NAD+ 3.3333 · MOTS-C 0.3333 · 5-Amino-1MQ 0.3333 mg/mL)
Calculated amount: 600 mcg (0.6 mg) per 0.15 mL spray
(NAD+ 500 + MOTS-C 50 + 5-Amino-1MQ 50 mcg)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

GLOW FUEL

NAD+ 100 mg + MOTS-C 10 mg + 5-Amino-1MQ 10 mg
120 mg total per bottle

Final volume: 30 mL
Concentration: 4 mg/mL combined
(NAD+ 3.3333 · MOTS-C 0.3333 · 5-Amino-1MQ 0.3333 mg/mL)
Calculated amount: 600 mcg (0.6 mg) per 0.15 mL spray
(NAD+ 500 + MOTS-C 50 + 5-Amino-1MQ 50 mcg)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

GLOW FUEL

NAD+ 100 mg + MOTS-C 10 mg + 5-Amino-1MQ 10 mg
120 mg total per bottle

Final volume: 30 mL
Concentration: 4 mg/mL combined
(NAD+ 3.3333 · MOTS-C 0.3333 · 5-Amino-1MQ 0.3333 mg/mL)
Calculated amount: 600 mcg (0.6 mg) per 0.15 mL spray
(NAD+ 500 + MOTS-C 50 + 5-Amino-1MQ 50 mcg)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

GLOW FUEL

NAD+ 100 mg + MOTS-C 10 mg + 5-Amino-1MQ 10 mg
120 mg total per bottle

Final volume: 30 mL
Concentration: 4 mg/mL combined
(NAD+ 3.3333 · MOTS-C 0.3333 · 5-Amino-1MQ 0.3333 mg/mL)
Calculated amount: 600 mcg (0.6 mg) per 0.15 mL spray
(NAD+ 500 + MOTS-C 50 + 5-Amino-1MQ 50 mcg)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

GLOW FUEL

NAD+ 100 mg + MOTS-C 10 mg + 5-Amino-1MQ 10 mg
120 mg total per bottle

Final volume: 30 mL
Concentration: 4 mg/mL combined
(NAD+ 3.3333 · MOTS-C 0.3333 · 5-Amino-1MQ 0.3333 mg/mL)
Calculated amount: 600 mcg (0.6 mg) per 0.15 mL spray
(NAD+ 500 + MOTS-C 50 + 5-Amino-1MQ 50 mcg)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

GLOW FUEL

NAD+ 100 mg + MOTS-C 10 mg + 5-Amino-1MQ 10 mg
120 mg total per bottle

Final volume: 30 mL
Concentration: 4 mg/mL combined
(NAD+ 3.3333 · MOTS-C 0.3333 · 5-Amino-1MQ 0.3333 mg/mL)
Calculated amount: 600 mcg (0.6 mg) per 0.15 mL spray
(NAD+ 500 + MOTS-C 50 + 5-Amino-1MQ 50 mcg)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

GLOW FUEL

NAD+ 100 mg + MOTS-C 10 mg + 5-Amino-1MQ 10 mg
120 mg total per bottle

Final volume: 30 mL
Concentration: 4 mg/mL combined
(NAD+ 3.3333 · MOTS-C 0.3333 · 5-Amino-1MQ 0.3333 mg/mL)
Calculated amount: 600 mcg (0.6 mg) per 0.15 mL spray
(NAD+ 500 + MOTS-C 50 + 5-Amino-1MQ 50 mcg)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption