GLOW LAB
PROTOCOLS

WOLVERINE

BPC-157 10 mg + TB-500 10 mg
20 mg total per bottle

Final volume: 30 mL
Concentration: 0.3333 mg/mL each; 0.6667 mg/mL combined
Calculated amount: 100 mcg per 0.15 mL spray
(50 mcg BPC-157 + 50 mcg TB-500)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

WOLVERINE

BPC-157 10 mg + TB-500 10 mg
20 mg total per bottle

Final volume: 30 mL
Concentration: 0.3333 mg/mL each; 0.6667 mg/mL combined
Calculated amount: 100 mcg per 0.15 mL spray
(50 mcg BPC-157 + 50 mcg TB-500)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

WOLVERINE

BPC-157 10 mg + TB-500 10 mg
20 mg total per bottle

Final volume: 30 mL
Concentration: 0.3333 mg/mL each; 0.6667 mg/mL combined
Calculated amount: 100 mcg per 0.15 mL spray
(50 mcg BPC-157 + 50 mcg TB-500)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

WOLVERINE

BPC-157 10 mg + TB-500 10 mg
20 mg total per bottle

Final volume: 30 mL
Concentration: 0.3333 mg/mL each; 0.6667 mg/mL combined
Calculated amount: 100 mcg per 0.15 mL spray
(50 mcg BPC-157 + 50 mcg TB-500)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

WOLVERINE

BPC-157 10 mg + TB-500 10 mg
20 mg total per bottle

Final volume: 30 mL
Concentration: 0.3333 mg/mL each; 0.6667 mg/mL combined
Calculated amount: 100 mcg per 0.15 mL spray
(50 mcg BPC-157 + 50 mcg TB-500)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

WOLVERINE

BPC-157 10 mg + TB-500 10 mg
20 mg total per bottle

Final volume: 30 mL
Concentration: 0.3333 mg/mL each; 0.6667 mg/mL combined
Calculated amount: 100 mcg per 0.15 mL spray
(50 mcg BPC-157 + 50 mcg TB-500)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

WOLVERINE

BPC-157 10 mg + TB-500 10 mg
20 mg total per bottle

Final volume: 30 mL
Concentration: 0.3333 mg/mL each; 0.6667 mg/mL combined
Calculated amount: 100 mcg per 0.15 mL spray
(50 mcg BPC-157 + 50 mcg TB-500)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

WOLVERINE

BPC-157 10 mg + TB-500 10 mg
20 mg total per bottle

Final volume: 30 mL
Concentration: 0.3333 mg/mL each; 0.6667 mg/mL combined
Calculated amount: 100 mcg per 0.15 mL spray
(50 mcg BPC-157 + 50 mcg TB-500)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

WOLVERINE

BPC-157 10 mg + TB-500 10 mg
20 mg total per bottle

Final volume: 30 mL
Concentration: 0.3333 mg/mL each; 0.6667 mg/mL combined
Calculated amount: 100 mcg per 0.15 mL spray
(50 mcg BPC-157 + 50 mcg TB-500)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

WOLVERINE

BPC-157 10 mg + TB-500 10 mg
20 mg total per bottle

Final volume: 30 mL
Concentration: 0.3333 mg/mL each; 0.6667 mg/mL combined
Calculated amount: 100 mcg per 0.15 mL spray
(50 mcg BPC-157 + 50 mcg TB-500)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

WOLVERINE

BPC-157 10 mg + TB-500 10 mg
20 mg total per bottle

Final volume: 30 mL
Concentration: 0.3333 mg/mL each; 0.6667 mg/mL combined
Calculated amount: 100 mcg per 0.15 mL spray
(50 mcg BPC-157 + 50 mcg TB-500)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

WOLVERINE

BPC-157 10 mg + TB-500 10 mg
20 mg total per bottle

Final volume: 30 mL
Concentration: 0.3333 mg/mL each; 0.6667 mg/mL combined
Calculated amount: 100 mcg per 0.15 mL spray
(50 mcg BPC-157 + 50 mcg TB-500)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption

GLOW LAB
PROTOCOLS

WOLVERINE

BPC-157 10 mg + TB-500 10 mg
20 mg total per bottle

Final volume: 30 mL
Concentration: 0.3333 mg/mL each; 0.6667 mg/mL combined
Calculated amount: 100 mcg per 0.15 mL spray
(50 mcg BPC-157 + 50 mcg TB-500)
Approximately 200 sprays before priming and residual loss
Calculated values assume 0.15 mL pump output.

RESEARCH USE ONLY

Not for human consumption