

DSC 462 HW#2, Kefu Zhu

Question 1

(a) $S_x = \{2, 3, 4, 5, 6, 7\}$

(b)(c)

Game Process:

(1) The 1st number (6 possible choices)

(2) The 2nd number

- The 2nd number is the same as the 1st number (1 possible choices) $\rightarrow X = 2, P(X = 2) = \frac{6 \cdot 1}{6^2}$
- The 2nd number is not the same as the 1st number (5 possible choices)

(3) The 3rd number

- The 3rd number is the same as one of the first 2 numbers (2 possible choices)
 $\rightarrow X = 3, P(X = 3) = \frac{(6 \cdot 5) \cdot 2}{6^3}$
- The 3rd number is not the same as any of the first 2 numbers (4 possible choices)

...

Therefore, we have the following calculation

$$P(X = 2) = \frac{(6) \cdot 1}{6^2} = \frac{1}{6}$$

$$P(X = 3) = \frac{(6 \cdot 5) \cdot 2}{6^3} = \frac{5}{18}$$

$$P(X = 4) = \frac{(6 \cdot 5 \cdot 4) \cdot 3}{6^4} = \frac{5}{18}$$

$$P(X = 5) = \frac{(6 \cdot 5 \cdot 4 \cdot 3) \cdot 4}{6^5} = \frac{5}{27}$$

$$P(X = 6) = \frac{(6 \cdot 5 \cdot 4 \cdot 3 \cdot 2) \cdot 5}{6^6} = \frac{25}{324}$$

$$P(X = 7) = \frac{(6!) \cdot 6}{6^7} = \frac{5}{324}$$

Sanity check: $P(X = 2) + P(X = 3) + P(X = 4) + P(X = 5) + P(X = 6) + P(X = 7)$
 $= \frac{1}{6} + \frac{5}{18} + \frac{5}{18} + \frac{5}{27} + \frac{25}{324} + \frac{5}{324} = 1$

Since we already have $P(X = i)$ for every $i \in S_x$, we can easily calculate every $P(X > i)$

$$P(X > 2) = 1 - P(X \leq 2) = 1 - P(X = 2) = 1 - \frac{1}{6} \approx 0.83$$

$$P(X > 3) = 1 - P(X = 2) - P(X = 3) = 1 - \frac{1}{6} - \frac{5}{18} \approx 0.56$$

$$P(X > 4) = 1 - P(X = 2) - P(X = 3) - P(X = 4) = 1 - \frac{1}{6} - \frac{5}{18} - \frac{5}{18} \approx 0.28$$

$$P(X > 5) = P(X = 6) + P(X = 7) = \frac{25}{324} + \frac{5}{324} \approx 0.09$$

$$P(X > 6) = P(X = 7) \approx 0.02$$

$$P(X > 7) = 0$$

Question 2

(a)

$$f(x) = \begin{cases} -c(x-2), & x \in (0, 2) \\ c(x-2), & x \in [2, 4) \\ 0, & \text{otherwise} \end{cases}$$

$$\sum p(x) = 1 = \int_0^2 -c(x-2)dx + \int_2^4 c(x-2)dx$$

$$= -c\left(\frac{x^2}{2} - 2x \Big|_0^2\right) + c\left(\frac{x^2}{2} - 2x \Big|_2^4\right)$$

$$= -c(2 - 4) + c[(8 - 8) - (2 - 4)]$$


$$= 4c \Rightarrow c = \frac{1}{4}$$

(b)

$$\therefore f(x) = \begin{cases} -\frac{1}{4}x + \frac{1}{2}, & x \in (0, 2) \\ \frac{1}{4}x - \frac{1}{2}, & x \in [2, 4) \\ 0, & \text{otherwise} \end{cases}$$

$$\therefore F(x) = \begin{cases} 0, & x \in (-\infty, 0] \\ \int_0^x -\frac{1}{4}x + \frac{1}{2}dx = \frac{x}{2} - \frac{x^2}{8}, & x \in (0, 2) \\ (\int_0^2 -\frac{1}{4}x + \frac{1}{2}dx) + (\int_2^x \frac{1}{4}x - \frac{1}{2}dx) = 1 + \frac{x^2}{8} - \frac{x}{2}, & x \in [2, 4) \\ 1, & x \in [4, +\infty) \end{cases}$$

```
# F(x) for x in (0,2)
f1 = function(x) {x/2 - x^2/8}
# F(x) for x in (2,4)
f2 = function(x) {1 + x^2/8 - x/2}
# Plot F(x) in (0,2)
plot(f1, from = 0, to = 2, xlim = c(-6,6), ylim = c(0,1), xlab = 'X', ylab = 'F(X)')
# Plot F(x) in (2,4)
plot(f2, from = 2, to = 4, add = TRUE)
# Add F(x) = 0 from -infinity to 0
segments(-6,0,0,0)
# Add F(x) = 1 from 4 to +infinity
segments(4,1,6,1)
```



Question 3

$$\text{var}(S_N) = \sum_{i=1}^n \sigma_i^2 + 2 \sum_{i < j} \sigma_{ij}$$

$$\therefore \sigma_i^2 = E[X_i^2] - E[X_i]^2 = \frac{1}{N} - \left(\frac{1}{N}\right)^2 = \frac{1}{N} \cdot \left(1 - \frac{1}{N}\right)$$

$$\because \sigma_{ij} = E[X_i X_j] - E[X_i]E[X_j] = 1 \cdot \frac{1}{N} \frac{1}{N-1} - \frac{1}{N} \frac{1}{N} = \frac{1}{N^2(N-1)}$$

$$\therefore \text{var}(S_N) = \sum_{i=1}^n \frac{1}{N} \cdot (1 - \frac{1}{N}) + 2 \sum_{i,j=1, i < j}^n \frac{1}{N^2(N-1)} = 1 + 2 \cdot (\frac{N(N-1)}{2}) \cdot \frac{1}{N^2(N-1)} = 1$$

Question 4

(a)

Define:

1. X = the test score of an individual
2. N = the number of candidates who passed the test

$$P(X \geq 625) = P(Z \geq \frac{625-500}{75}) \approx 0.04779035$$

```
# Calculate the P(Z >= (625-500)/75)
1 - pnorm(625, mean = 500, sd = 75) # 0.04779035
```

Define:

1. p = the probability that an individual passes the test = 0.04779035
2. q = the probability that an individual does not pass the test = $1 - p = 0.9522097$

$$P(N \geq 3) = 1 - P(N = 0) - P(N = 1) - P(N = 2)$$

$$= 1 - \binom{35}{0} \cdot q^{35} - \binom{35}{1} \cdot p \cdot q^{34} - \binom{35}{2} \cdot p^2 \cdot q^{33}$$

$$= 1 - 1 \cdot (0.9522097)^{35} - 35 \cdot 0.04779035 \cdot (0.9522097)^{34} - 595 \cdot (0.04779035)^2 \cdot (0.9522097)^{33}$$

$$\approx 0.2333842$$

(b)

$$P(N \geq 3) = 0.8 = 1 - \binom{35}{0} \cdot q^{35} - \binom{35}{1} \cdot p \cdot q^{34} - \binom{35}{2} \cdot p^2 \cdot q^{33}$$

$$\Rightarrow \binom{35}{0} \cdot q^{35} + \binom{35}{1} \cdot p \cdot q^{34} + \binom{35}{2} \cdot p^2 \cdot q^{33} - 0.2 = 0$$

```
# Define function
f = function(x) {
  choose(35,0)*(1-x)^35 +
  choose(35,1)*x*(1-x)^34 +
  choose(35,2)*(x^2)*(1-x)^33 - 0.2}
# Calculate the value of x
uniroot(f,c(0,1))
```

```
# Result
$root
[1] 0.1183283

$f.root
[1] -2.792853e-05

$iter
[1] 9

$init.it
[1] NA

$estim.prec
[1] 6.103516e-05
```

Therefore, $p \approx 0.1183283$. Based on the Z -score table, $Z \approx 1.18$

```
# Check the Z-score in R
qnorm(0.1183283, 0, 1, lower.tail = FALSE) # 1.183385
```

$$\therefore Z \approx 1.18 = \frac{X-500}{75} \rightarrow X \approx 589$$

Question 5

$$F(M) = P(X_1 \leq M, X_2 \leq M, \dots, X_n \leq M)$$

$$\because X_1, \dots, X_n \text{ are i.i.d}$$

$$\therefore F_x(M) = P(X_1 \leq M)P(X_2 \leq M) \cdots P(X_n \leq M) = F(M)^k = \left(\frac{M-0}{c-0}\right)^n = \left(\frac{M}{c}\right)^n$$

$$\Rightarrow f(x) = \frac{d}{dM} F_x(M) = \frac{nM^{n-1}}{c^n}$$

Then, $E(M) = \int_0^c M \cdot f(M) dM$

$$= \int_0^c M \cdot \frac{n}{c^n} \cdot M^{n-1} dM$$

$$= \frac{n}{c^n \cdot (n+1)} \cdot M^{n+1} \Big|_0^c$$

$$= \frac{n}{n+1} \cdot c$$