

UNIVERSAL ANALYTIC GRÖBNER BASES AND TROPICAL GEOMETRY

Tristan Vaccon¹ Thibaut Verron²

International Symposium on Symbolic and Algebraic Computation, 27 July 2023

1. Université de Limoges, XLIM, Limoges, France
2. Johannes Kepler University, Institute for Algebra, Linz, Austria




Algebraic
geometry



Analytic
geometry

TROPICAL ANALYTIC GEOMETRY



Rigid
geometry


Tate

Algebraic
geometry



GAGA

Analytic
geometry



Rigid
geometry

Tate

Algebraic
geometry







GAGA

Analytic
geometry

Tropical
geometry

TROPICAL ANALYTIC GEOMETRY





TROPICAL VARIETIES

K field with valuation (e.g. $\mathbb{Q}((t)), \mathbb{Q}_p, \dots$)

$I \subseteq K[X]$ ideal

Definition with valuation:

$$\text{trop}(I) = \text{clo}\{\text{val}(\mathbf{x}) : \mathbf{x} \in V(I)\}$$



Newton-Puiseux, Hensel, etc.

Fund. th. of tropical geometry



Definition with initial forms:

System of weights: $\mathbf{w} = (w_0, \dots, w_n) \in \mathbb{R}^n$

\mathbf{w} -valuation: $\text{val}_{\mathbf{w}}(aX_1^{\alpha_1} \dots X_n^{\alpha_n}) = w_0 \text{val}(a) + w_1 \alpha_1 + \dots + w_n \alpha_n$

$\text{init}_{\mathbf{w}}(f)$ = sum of terms with minimal \mathbf{w} -valuation

$\text{init}_{\mathbf{w}}(I) = \langle \text{init}_{\mathbf{w}}(f) : f \in I \rangle$

$\text{trop}(I) = \{\mathbf{w} \text{ such that } \text{init}_{\mathbf{w}}(I) \text{ does not contain a monomial}\}$

$$\begin{aligned} x &= \alpha t^1 + \dots, y = \beta t^1 + \dots \\ f(x, y) &= \alpha^2 t^2 + \beta^2 t^4 - t^4 + \dots \\ &= \alpha^2 t^2 + \dots \neq 0 \\ \text{init}_{(1,1)}(I) &= \langle x^2 \rangle \end{aligned}$$

$(1, 1) \bullet$

$$\begin{aligned} x &= \alpha t^2 + \dots, y = \beta t^2 + \dots \\ f(x, y) &= \alpha^2 t^4 + \beta^2 t^6 - t^4 + \dots \\ &= (\alpha^2 - 1)t^4 + \dots \\ &= 0 \implies \alpha^2 - 1 = 0 \\ \text{init}_{(2,2)}(I) &= \langle x^2 - t^4 \rangle \end{aligned}$$

$(2, 2) \bullet$

$(2, 1)$

$$\begin{aligned} f &= x^2 + t^2 y^2 - t^4 \in \mathbb{Q}((t))[x, y] \\ I &= \langle f \rangle \end{aligned}$$


CONNECTION WITH THE GRÖBNER FAN

Gröbner fan:

- Partition of \mathbb{R}^n according to

$$\mathbf{a} \sim \mathbf{b} \iff \text{init}_{\mathbf{a}}(I) = \text{init}_{\mathbf{b}}(I)$$

- Finite union of rational cones
- Maximal dim. = term orders
- Lower dim. = boundaries = collisions
- Contains the tropical variety



Tropical variety of I (without valuations)

1. Compute a universal GB of I
2. Compute its Newton polytope
3. Compute the Gröbner fan of I
4. For each non-maximal cone \mathcal{C} , pick a $\mathbf{w} \in \mathcal{C}$ and test if $\text{init}_{\mathbf{w}}(I)$ contains a monomial

In practice: traverse only the necessary parts of the Gröbner fan

[Bogard Jensen Speyer Sturmfels Thomas 2006]

$$I \subseteq \mathbb{Q}[X]$$

[Bogard Jensen
Speyer Sturmfels
Thomas 2006]

Gröbner
fan

Tropical
variety

$$I \subseteq \mathbb{Q}(t)[X]$$


$$J \subseteq \mathbb{Q}[t, X]$$

[Bogard Jensen
Speyer Sturmfels
Thomas 2006]


Gröbner
fan

Tropical
variety


COMPUTING TROPICAL VARIETIES WITH VALUATION



COMPUTING TROPICAL VARIETIES WITH VALUATION



COMPUTING TROPICAL VARIETIES WITH VALUATION



Definition: convergent series with coefficients in a valued field or ring ($\mathbb{Q}(t)$, $\mathbb{Q}((t))$, $\mathbb{Q}_p \dots$)

$$K\{\mathbf{X}; \mathbf{r}\} = \left\{ \sum_{\alpha \in \mathbb{N}^n} a_{\alpha} \mathbf{X}^{\alpha} \text{ with } \text{val}(a_{\alpha}) - \alpha \cdot \mathbf{r} \xrightarrow{|\alpha| \rightarrow \infty} \infty \right\}$$

$\mathbf{r} = (r_1, \dots, r_n)$: convergence radii

- If $\mathbf{r} = (0, \dots, 0)$, equivalent: $a_{\alpha} \rightarrow 0$
- If $\mathbf{r} \in \mathbb{Z}^n$, equivalent to change of variable $K\{\mathbf{X}; \mathbf{r}\} = K\{(X_i / p^{r_i}); (0)\}$
- $K[\mathbf{X}] = K\{\mathbf{X}; \infty\}$ (everywhere convergent)


Term ordering:

$$a\mathbf{X}^{\alpha} <_{\mathbf{r}} b\mathbf{X}^{\beta} \iff \begin{cases} \text{val}(a) - \mathbf{r} \cdot \alpha > \text{val}(b) - \mathbf{r} \cdot \beta \\ \text{or they are equal and } \mathbf{X}^{\alpha} < \mathbf{X}^{\beta} \end{cases}$$


- Every Tate series has a leading term
- Every Tate series ideal has a finite Gröbner basis
- Different \mathbf{r} give different leading terms and Gröbner bases

OVERCONVERGENCE

Fact: If $r \leq s$, then $K\{X; s\} \subseteq K\{X; r\}$:




$$K\{X; (4, 3)\} \subseteq K\{X; (1, 2)\}$$





$$K\{X; (\infty, \infty)\} = K[X] \subseteq K\{X; (1, 2)\}$$

OVERCONVERGENCE

Fact: If $\mathbf{r} \leq \mathbf{s}$, then $K\{\mathbf{X}; \mathbf{s}\} \subseteq K\{\mathbf{X}; \mathbf{r}\}$:



$$K\{\mathbf{X}; (4, 3)\} \subseteq K\{\mathbf{X}; (1, 2)\}$$



$$K\{\mathbf{X}; (\infty, \infty)\} = K[\mathbf{X}] \subseteq K\{\mathbf{X}; (1, 2)\}$$

Theorem (Caruso, Vaccon, V. 2022)

Let $\mathbf{r} \geq \mathbf{s}$, $I \subset K\{\mathbf{X}; \mathbf{r}\}$ and $I_{\mathbf{s}} = K\{\mathbf{X}; \mathbf{s}\} / I$ (completion of the ideal).

Then $I_{\mathbf{s}}$ admits a Gröbner basis comprised only of elements of $K\{\mathbf{X}; \mathbf{r}\}$.

In particular, the completion of a polynomial ideal has a polynomial basis.

Key component: Mora's reduction algorithm

Input: $G \subset K\{\mathbf{X}; \mathbf{r}\}$, $f \in K\{\mathbf{X}; \mathbf{r}\}$

Output: $h, u \in K\{\mathbf{X}; \mathbf{r}\}$, such that:

- uf reduces to h and is irreducible modulo G (in $K\{\mathbf{X}; \mathbf{s}\}$)
- $\text{LT}_{\mathbf{s}}(u) = 1$, or equivalently, u is invertible in $K\{\mathbf{X}; \mathbf{s}\}$

$$\text{LT}_{r, \leq}(f) = a_0 \mathbf{X}^{\alpha_0} + a_1 \mathbf{X}^{\alpha_1} + \dots + a_k \mathbf{X}^{\alpha_k} + a_{k+1} \mathbf{X}^{\alpha_{k+1}} + \dots \text{ with } \text{val}(a_i) - \alpha_i \cdot \mathbf{r} \xrightarrow{|\alpha_i| \rightarrow \infty} \infty$$

minimal $\text{val}(a_i) - \alpha_i \cdot \mathbf{r}$

$$\text{LT}_{\mathbf{r}, \leq}(f) = a_0 \mathbf{X}^{\alpha_0} + a_1 \mathbf{X}^{\alpha_1} + \dots + a_k \mathbf{X}^{\alpha_k} + a_{k+1} \mathbf{X}^{\alpha_{k+1}} + \dots \text{ with } \text{val}(a_i) - \alpha_i \cdot \mathbf{r} \xrightarrow{|\alpha_i| \rightarrow \infty} \infty$$

$$\text{minimal } \text{val}(a_i) - \alpha_i \cdot \mathbf{r} = \text{init}_{(-1, \mathbf{r})}(f)$$

Theorem (Vaccon, V. 2023)

Let $I \subseteq K[\mathbf{X}]$, and $\mathbf{w} \in \mathbb{Q}^{n+1}$ a system of weights.

Let $\mathbf{r} = (-w_1/w_0, \dots, -w_n/w_0)$ and $I_{\mathbf{r}} = I K[\mathbf{X}; \mathbf{r}]$ the corresponding completion, then

$$\text{init}_{\mathbf{w}}(I) = \text{init}_{\mathbf{w}}(I_{\mathbf{r}}) \cap K[\mathbf{X}].$$

This is a local result, which translates globally as:

$$V_{\text{trop}}(I) = \bigcup_{\mathbf{s} \in \mathbb{Q}^n} \text{trop}(I_{\mathbf{s}}).$$

Theorem (Vaccon, V. 2023)

Let G be a Gröbner basis of $I_{\mathbf{r}}$, then

$$\text{init}_{\mathbf{w}}(I_{\mathbf{r}}) = \langle \text{init}_{\mathbf{w}}(g) : g \in G \rangle.$$

Theorem (Caruso, Vaccon, V. 2022; Vaccon, V. 2023)

Let $I \subseteq K[\mathbf{X}]$ be a homogeneous ideal.

There exists a finite subset $G \subseteq I$ s.t. for all $\mathbf{r} \in \mathbb{Q}^n$, G is a Gröbner basis of $I_{\mathbf{r}} = I K\{\mathbf{X}; \mathbf{r}\}$.

Furthermore:

- this is independent of the order used for breaking ties
- G can be computed

As a consequence, a homogeneous polynomial ideal always has a finite Gröbner fan.

Why homogeneous?

- Key question: how does $\text{init}_{\mathbf{a}}(I) = \text{init}_{\mathbf{b}}(I)$ relate to $\text{init}_{\mathbf{a}}(G) = \text{init}_{\mathbf{b}}(G)$?
- Usually, this is answered by taking **reduced** Gröbner bases
- We can have reduced GB (with the usual algorithm),
or overconvergent GB (using Mora's algorithm)... **but not both in general**
- For homogeneous ideals, reduced overconvergent bases exist

Tropical variety with valuation

- I. $I \subseteq K[X]$ homogeneous ideal.
- O. the tropical variety of I , given as a union of rational cones
 1. compute a universal analytic Gröbner basis G of I
 2. get all the maximal dimensional cones in the Gröbner fan
 3. compute the rest of the cones
 4. for each non-maximal cone \mathcal{C}
 - 4.1 pick a $\mathbf{w} = (-1, \mathbf{r}) \in \mathcal{C}$
 - 4.2 Then $\text{init}_{\mathbf{w}}(G)$ is a Gröbner basis of $\text{init}_{\mathbf{w}}(I_{\mathbf{r}})$
 - 4.3 test if $\text{init}_{\mathbf{w}}(I)$ contains a monomial
 - 4.4 conclude whether $\mathbf{w} \in \text{trop}(I)$ and therefore $\mathcal{C} \subseteq \text{trop}(I)$.

Tropical variety with valuation

- I. $I \subseteq K[X]$ homogeneous ideal.
- O. the tropical variety of I , given as a union of rational cones
 1. compute a universal analytic Gröbner basis G of I
 2. get all the maximal dimensional cones in the Gröbner fan
 3. compute the rest of the cones
 4. for each non-maximal cone \mathcal{C}
 - 4.1 pick a $\mathbf{w} = (-1, \mathbf{r}) \in \mathcal{C}$
 - 4.2 Then $\text{init}_{\mathbf{w}}(G)$ is a Gröbner basis of $\text{init}_{\mathbf{w}}(I_{\mathbf{r}})$
 - 4.3 test if $\text{init}_{\mathbf{w}}(I)$ contains a monomial
 - 4.4 conclude whether $\mathbf{w} \in \text{trop}(I)$ and therefore $\mathcal{C} \subseteq \text{trop}(I)$.

Mora's reduction
UAGB algo.

Tropical variety with valuation

- I. $I \subseteq K[X]$ homogeneous ideal.
- O. the tropical variety of I , given as a union of rational cones
 1. compute a universal analytic Gröbner basis G of I
 2. get all the maximal dimensional cones in the Gröbner fan
 3. compute the rest of the cones
 4. for each non-maximal cone \mathcal{C}
 - 4.1 pick a $\mathbf{w} = (-1, \mathbf{r}) \in \mathcal{C}$
 - 4.2 Then $\text{init}_{\mathbf{w}}(G)$ is a Gröbner basis of $\text{init}_{\mathbf{w}}(I_{\mathbf{r}})$
 - 4.3 test if $\text{init}_{\mathbf{w}}(I)$ contains a monomial
 - 4.4 conclude whether $\mathbf{w} \in \text{trop}(I)$ and therefore $\mathcal{C} \subseteq \text{trop}(I)$.

Mora's reduction
UAGB algo.

Newton polytope

Discrete geometry

Tropical variety with valuation

- I. $I \subseteq K[X]$ homogeneous ideal.
- O. the tropical variety of I , given as a union of rational cones
 1. compute a universal analytic Gröbner basis G of I
 2. get all the maximal dimensional cones in the Gröbner fan
 3. compute the rest of the cones
 4. for each non-maximal cone \mathcal{C}
 - 4.1 pick a $\mathbf{w} = (-1, \mathbf{r}) \in \mathcal{C}$
 - 4.2 Then $\text{init}_{\mathbf{w}}(G)$ is a Gröbner basis of $\text{init}_{\mathbf{w}}(I_{\mathbf{r}})$
 - 4.3 test if $\text{init}_{\mathbf{w}}(I)$ contains a monomial
 - 4.4 conclude whether $\mathbf{w} \in \text{trop}(I)$ and therefore $\mathcal{C} \subseteq \text{trop}(I)$.

Mora's reduction
UAGB algo.

Newton polytope
Discrete geometry

Th. on Tate GB
Saturation

Tropical variety with valuation

- I. $I \subseteq K[X]$ homogeneous ideal.
0. the tropical variety of I , given as a union of rational cones
 1. compute a universal analytic Gröbner basis G of I
 2. get all the maximal dimensional cones in the Gröbner fan
 3. compute the rest of the cones
 4. for each non-maximal cone \mathcal{C}
 - 4.1 pick a $\mathbf{w} = (-1, \mathbf{r}) \in \mathcal{C}$
 - 4.2 Then $\text{init}_{\mathbf{w}}(G)$ is a Gröbner basis of $\text{init}_{\mathbf{w}}(I_{\mathbf{r}})$
 - 4.3 test if $\text{init}_{\mathbf{w}}(I)$ contains a monomial
 - 4.4 conclude whether $\mathbf{w} \in \text{trop}(I)$ and therefore $\mathcal{C} \subseteq \text{trop}(I)$.

Mora's reduction
UAGB algo.

Newton polytope
Discrete geometry


Th. on Tate GB
Saturation

- Proof of concept showing that the main algorithmic ingredients are in place:
universal Gröbner basis, Gröbner fan, connection to the tropical variety
- Next task: transposing the advanced traversal techniques used in the classical setting

So far, we have shown that for (poly)disks:

- We can compute overconvergent Gröbner bases
- We can compute a universal analytic Gröbner basis

What about more general convergence conditions?




Questions:

- Local: can we compute overconvergent Gröbner bases?
- Global: Can we compute a universal analytic Gröbner basis?

So far, we have shown that for (poly)disks:

- We can compute overconvergent Gröbner bases
- We can compute a universal analytic Gröbner basis

Example: upper polyhedral domains:



Questions:


- Local: can we compute overconvergent Gröbner bases?
- Global: Can we compute a universal analytic Gröbner basis?

Th. (Vaccon V. 2023) YES

So far, we have shown that for (poly)disks:

- We can compute overconvergent Gröbner bases
- We can compute a universal analytic Gröbner basis

Example: annuli (with Laurent terms):




Questions:

- Local: can we compute overconvergent Gröbner bases?
- Global: Can we compute a universal analytic Gröbner basis?

So far, we have shown that for (poly)disks:

- We can compute overconvergent Gröbner bases
- We can compute a universal analytic Gröbner basis

Example: annuli (with Laurent terms):



Questions:

- Local: can we compute overconvergent Gröbner bases?
- Global: Can we compute a universal analytic Gröbner basis?

Thank you for your attention!