

Incorporating neutron star physics into gravitational wave inference with neural priors

Thibeau Wouters, Peter T. H. Pang, Tim Dietrich, Chris Van
Den Broeck



Utrecht
University

Nikhef

Table of Contents

① Introduction

② Methods

③ Results

④ Conclusion

Contents

① Introduction

② Methods

③ Results

④ Conclusion

Motivation

- Bayesian inference depends on choice of **priors**:

$$\mathcal{P}(\theta_{\text{GW}}|d) \propto \frac{\mathcal{L}(d|\theta_{\text{GW}})\pi(\theta_{\text{GW}})}{\mathcal{Z}}$$


- By default, we use **agnostic priors**, but what if we *do* have non-trivial prior information?
- Case study: neutron stars (NSs) and information from
 - Population models
 - Equation of state (EOS) constraints

Neural priors

Flexible way to encode NS physics into GW inference

Key idea

Train normalizing flow (NF) on samples informed by populations and EOS
→ **neural prior**



Contents

① Introduction

② Methods

③ Results

④ Conclusion

NS population models

Three fiducial population models for NS masses:

① Uniform [1–3]:

- Only use EOS constraints for maximum mass (M_{TOV})
- NS mass $\sim U[1 \text{ M}_\odot, M_{\text{TOV}}]$

② Gaussian [4]:

- NS mass $\sim \mathcal{N}(1.33 \text{ M}_\odot, (0.09 \text{ M}_\odot)^2)$

③ Double Gaussian [5, 6]:

- Weighted mixture of two Gaussians
- $0.65 \times \mathcal{N}(1.34 \text{ M}_\odot, (0.07 \text{ M}_\odot)^2) + 0.35 \times \mathcal{N}(1.80 \text{ M}_\odot, (0.21 \text{ M}_\odot)^2)$


For NSBH systems:

- BH mass m_1^{src} : from $[M_{\text{TOV}}, 5 \text{ M}_\odot]$
- NS mass m_2^{src} : above models


EOS constraints

- We use three EOS constraints [7]:
 - ① **Heavy pulsars:** must support $2 M_{\odot}$ NSs
 - ② **Chiral EFT (χ_{EFT}):** nuclear theory predictions (softer EOS)
 - ③ **NICER:** mass-radius observations of NSs (stiffer EOS)
- Posterior samples obtained with JESTER [8] 

Chiral EFT




NICER



Normalizing flows

Normalizing flows [9, 10]

- Neural density estimators: trainable bijections
- Often used in PE, e.g., DINGO [11, 12], NESSAI [13, 14]
- Generate samples, evaluate density: can be used as priors [15]



Construction of neural priors


Steps to generate training data:

- ① Draw EOS posterior curve: determines $M_{\text{TOV}}, \Lambda(m)$
- ② Draw masses from population model
- ③ Compute $\Lambda_i = \Lambda(m_i)$ for NSs (NSBH: $\Lambda_1 = 0$)

Implementation:

- Created with GLASFLOW [16, 17]
- CouplingNSF architecture (neural spline flows [18])
- Use as a JointPrior in BILBY (NFPrior)
 - Sample & logpdf: evaluate NF
 - Rescale: unit hypercube \rightarrow multivariate Gaussian $\xrightarrow{\text{NF}}$ data space

All neural priors



Contents

① Introduction

② Methods

③ Results

GW170817

GW190425

GW230529

④ Conclusion

Setup

Analyze GW170817, GW190425, GW230529 with:

- 4096 live points, multibanding likelihood
- IMRPhenomXP_NRTidalv3
- Neural priors for m_i, Λ_i (standard priors for other parameters)

Two contributions:

- ① Narrower constraints with neural priors
- ② Model selection with Bayes factors

Jeffreys' scale for Bayes factors (\log_{10} scale)

$\log_{10}(\mathcal{B}_1^2)$	Interpretation	Color
$[0, \frac{1}{2}]$	Barely worth mentioning	Light orange
$[\frac{1}{2}, 1]$	Substantial	Orange
$[1, \frac{3}{2}]$	Strong	Red-orange
$[\frac{3}{2}, 2]$	Very strong	Dark red
> 2	Decisive	Red

Contents

① Introduction

② Methods

③ Results

GW170817

GW190425

GW230529


④ Conclusion

GW170817: Source classification

Showing \log_{10} Bayes factors relative to model with highest evidence

Source	Population	EOS	GW170817
BNS	Uniform	PSRs	-1.83
		PSRs+ χ EFT	-0.80
		PSRs+NICER	-1.58
	Gaussian	PSRs	-0.68
		PSRs+ χ EFT	ref.
		PSRs+NICER	-0.76
	Double Gaussian	PSRs	-1.36
		PSRs+ χ EFT	-0.59
		PSRs+NICER	-0.92
NSBH	Uniform	PSRs	-224.65
		PSRs+ χ EFT	-224.66
		PSRs+NICER	-224.66
	Gaussian	PSRs	-224.67
		PSRs+ χ EFT	-224.66
		PSRs+NICER	-224.66
	Double Gaussian	PSRs	-224.67
		PSRs+ χ EFT	-224.68
		PSRs+NICER	-224.67

GW170817: Parameter constraints (Gaussian)



GW170817: Discussion

Source classification:

- Decisive evidence for BNS over NSBH
- Prefer Gaussian population model
- Slight preference for softer EOS (PSRs+ χ_{EFT})

Parameter constraints:

- More equal mass ratio: $q \geq 0.9$
- Constrained tidal deformability $\tilde{\Lambda}$
- Higher luminosity distance compared to agnostic prior
- Matches multimessenger analyses of GW170817 [19–23]

Contents

① Introduction

② Methods

③ Results

GW170817

GW190425

GW230529


④ Conclusion

GW190425: Source classification

Showing \log_{10} Bayes factors relative to best model

Source	Population	EOS	GW190425
BNS	Uniform	PSRs	-0.07
		PSRs+ χ EFT	-0.11
		PSRs+NICER	ref.
	Gaussian	PSRs	-6.89
		PSRs+ χ EFT	-8.47
		PSRs+NICER	-5.45
	Double Gaussian	PSRs	-0.55
		PSRs+ χ EFT	-0.79
		PSRs+NICER	-0.57
NSBH	Uniform	PSRs	-1.52
		PSRs+ χ EFT	-1.35
		PSRs+NICER	-1.63
	Gaussian	PSRs	-0.82
		PSRs+ χ EFT	-1.11
		PSRs+NICER	-1.43
	Double Gaussian	PSRs	-4.11
		PSRs+ χ EFT	-3.83
		PSRs+NICER	-24.31

GW190425: Parameter constraints (Uniform)



GW190425: Discussion

Source classification:

- Prefer BNS over NSBH (but less conclusive than GW170817)
- Most consistent with uniform population
- Masses are outliers compared to galactic binaries

Parameter constraints:

- Less equal masses: $q \leq 0.9$
- $\tilde{\Lambda} \approx 200$ (more prior-dominated due to lower SNR)
- Higher luminosity distance: 182_{-49}^{+41} Mpc vs. 157_{-65}^{+64} Mpc (90% credibility)

Contents

① Introduction

② Methods

③ Results

GW170817

GW190425

GW230529


④ Conclusion

GW230529: Source classification

Showing \log_{10} Bayes factors relative to best model

Source	Population	EOS	GW230529
BNS	Uniform	PSRs	-13.14
		PSRs+ χ EFT	-13.12
		PSRs+NICER	-12.92
	Gaussian	PSRs	-18.82
		PSRs+ χ EFT	-18.83
		PSRs+NICER	-18.81
	Double Gaussian	PSRs	-13.75
		PSRs+ χ EFT	-13.77
		PSRs+NICER	-13.71
NSBH	Uniform	PSRs	-0.08
		PSRs+ χ EFT	-0.02
		PSRs+NICER	-0.25
	Gaussian	PSRs	-0.05
		PSRs+ χ EFT	-0.20
		PSRs+NICER	ref.
	Double Gaussian	PSRs	-0.14
		PSRs+ χ EFT	-0.13
		PSRs+NICER	-0.05

GW230529: Parameter constraints (Gaussian)



GW230529: Discussion

Source classification:

- Decisive evidence for NSBH over BNS (agrees with LVK [24])
- No evidence between hypotheses (low SNR)

Parameter constraints:

- Mass ratio more constrained: $q \leq 0.4$
 - As a result, improved spin constraints (χ_{1z} closer to zero)
- Tidal deformability posteriors dominated by priors
- Luminosity distance: 235^{+59}_{-58} Mpc vs. 201^{+84}_{-97} Mpc (90% credibility)
- Less tidal information: NRTidalv3 tapers at (90% credibility)
 - 641^{+318}_{-158} Hz (agnostic priors)
 - 858^{+113}_{-108} Hz (neural prior, reference model)

Contents

① Introduction

② Methods

③ Results

④ Conclusion

Conclusion

- PE often uses agnostic priors, missing valuable information
- **Neural priors**: flexible way to encode non-trivial prior information
- Case study: neutron star physics from
 - Population models
 - EOS constraints
- Two highlights:
 - ① **Bayesian source classification** (BNS vs. NSBH)
 - ② **Informed parameter constraints** (narrower posteriors)
- Implemented in BILBY
- Data-driven approach: easy to extend/generalize



Thanks for listening!



References I

- [1] R. Abbott et al. "Observation of Gravitational Waves from Two Neutron Star–Black Hole Coalescences". In: *Astrophys. J. Lett.* 915.1 (2021), p. L5. DOI: [10.3847/2041-8213/ac082e](https://doi.org/10.3847/2041-8213/ac082e). arXiv: [2106.15163 \[astro-ph.HE\]](https://arxiv.org/abs/2106.15163).
- [2] Philippe Landry and Jocelyn S. Read. "The Mass Distribution of Neutron Stars in Gravitational-wave Binaries". In: *Astrophys. J. Lett.* 921.2 (2021), p. L25. DOI: [10.3847/2041-8213/ac2f3e](https://doi.org/10.3847/2041-8213/ac2f3e). arXiv: [2107.04559 \[astro-ph.HE\]](https://arxiv.org/abs/2107.04559).
- [3] Jacob Golomb et al. "Interplay of astrophysics and nuclear physics in determining the properties of neutron stars". In: *Phys. Rev. D* 111.2 (2025), p. 023029. DOI: [10.1103/PhysRevD.111.023029](https://doi.org/10.1103/PhysRevD.111.023029). arXiv: [2410.14597 \[astro-ph.HE\]](https://arxiv.org/abs/2410.14597).
- [4] Feryal Özel and Paulo Freire. "Masses, Radii, and the Equation of State of Neutron Stars". In: *Ann. Rev. Astron. Astrophys.* 54 (2016), pp. 401–440. DOI: [10.1146/annurev-astro-081915-023322](https://doi.org/10.1146/annurev-astro-081915-023322). arXiv: [1603.02698 \[astro-ph.HE\]](https://arxiv.org/abs/1603.02698).
- [5] Justin Alsing, Hector O. Silva, and Emanuele Berti. "Evidence for a maximum mass cut-off in the neutron star mass distribution and constraints on the equation of state". In: *Mon. Not. Roy. Astron. Soc.* 478.1 (2018), pp. 1377–1391. DOI: [10.1093/mnras/sty1065](https://doi.org/10.1093/mnras/sty1065). arXiv: [1709.07889 \[astro-ph.HE\]](https://arxiv.org/abs/1709.07889).

References II

- [6] Dong-Sheng Shao et al. "Maximum mass cutoff in the neutron star mass distribution and the prospect of forming supramassive objects in the double neutron star mergers". In: *Phys. Rev. D* 102.6 (2020), p. 063006. DOI: [10.1103/PhysRevD.102.063006](https://doi.org/10.1103/PhysRevD.102.063006). arXiv: [2009.04275 \[astro-ph.HE\]](https://arxiv.org/abs/2009.04275).
- [7] Hauke Koehn et al. "From existing and new nuclear and astrophysical constraints to stringent limits on the equation of state of neutron-rich dense matter". In: *Phys. Rev. X* 15.2 (2025), p. 021014. DOI: [10.1103/PhysRevX.15.021014](https://doi.org/10.1103/PhysRevX.15.021014). arXiv: [2402.04172 \[astro-ph.HE\]](https://arxiv.org/abs/2402.04172).
- [8] Thibeau Wouters et al. "Leveraging differentiable programming in the inverse problem of neutron stars". In: (Apr. 2025). arXiv: [2504.15893 \[astro-ph.HE\]](https://arxiv.org/abs/2504.15893).
- [9] Ivan Kobyzev, Simon J. D. Prince, and Marcus A. Brubaker. "Normalizing Flows: An Introduction and Review of Current Methods". In: *IEEE Trans. Pattern Anal. Machine Intell.* 43.11 (2021), pp. 3964–3979. DOI: [10.1109/tpami.2020.2992934](https://doi.org/10.1109/tpami.2020.2992934). arXiv: [1908.09257 \[stat.ML\]](https://arxiv.org/abs/1908.09257).
- [10] George Papamakarios et al. "Normalizing Flows for Probabilistic Modeling and Inference". In: *J. Machine Learning Res.* 22.1 (2021), pp. 2617–2680. DOI: [10.5555/3546258.3546315](https://doi.org/10.5555/3546258.3546315). arXiv: [1912.02762 \[stat.ML\]](https://arxiv.org/abs/1912.02762).

References III




- [11] Maximilian Dax et al. "Real-Time Gravitational Wave Science with Neural Posterior Estimation". In: *Phys. Rev. Lett.* 127.24 (2021), p. 241103. DOI: [10.1103/PhysRevLett.127.241103](https://doi.org/10.1103/PhysRevLett.127.241103). arXiv: [2106.12594 \[gr-qc\]](https://arxiv.org/abs/2106.12594).
- [12] Maximilian Dax et al. "Neural Importance Sampling for Rapid and Reliable Gravitational-Wave Inference". In: *Phys. Rev. Lett.* 130.17 (2023), p. 171403. DOI: [10.1103/PhysRevLett.130.171403](https://doi.org/10.1103/PhysRevLett.130.171403). arXiv: [2210.05686 \[gr-qc\]](https://arxiv.org/abs/2210.05686).
- [13] Michael J. Williams, John Veitch, and Chris Messenger. "Nested sampling with normalizing flows for gravitational-wave inference". In: *Phys. Rev. D* 103.10 (2021), p. 103006. DOI: [10.1103/PhysRevD.103.103006](https://doi.org/10.1103/PhysRevD.103.103006). arXiv: [2102.11056 \[gr-qc\]](https://arxiv.org/abs/2102.11056).
- [14] Michael J. Williams, John Veitch, and Chris Messenger. "Importance nested sampling with normalising flows". In: *Mach. Learn. Sci. Tech.* 4.3 (2023), p. 035011. DOI: [10.1088/2632-2153/acd5aa](https://doi.org/10.1088/2632-2153/acd5aa). arXiv: [2302.08526 \[astro-ph.IM\]](https://arxiv.org/abs/2302.08526).
- [15] Justin Alsing and Will Handley. "Nested sampling with any prior you like". In: *Mon. Not. Roy. Astron. Soc.* 505.1 (2021), pp. L95–L99. DOI: [10.1093/mnrasl/slab057](https://doi.org/10.1093/mnrasl/slab057). arXiv: [2102.12478 \[astro-ph.IM\]](https://arxiv.org/abs/2102.12478).
- [16] Michael J. Williams et al. *uofgravity/glasflow: v0.4.1*. Oct. 2024. DOI: [10.5281/zenodo.13914483](https://doi.org/10.5281/zenodo.13914483). URL: <https://doi.org/10.5281/zenodo.13914483>.
- [17] Conor Durkan et al. *nflows: normalizing flows in PyTorch*. Version v0.14. Nov. 2020. DOI: [10.5281/zenodo.4296287](https://doi.org/10.5281/zenodo.4296287). URL: <https://doi.org/10.5281/zenodo.4296287>.

References IV

- [18] Conor Durkan et al. "Neural Spline Flows". In: June 2019. arXiv: [1906.04032 \[stat.ML\]](https://arxiv.org/abs/1906.04032).
- [19] David Radice and Liang Dai. "Multimessenger Parameter Estimation of GW170817". In: *Eur. Phys. J. A* 55.4 (2019), p. 50. doi: [10.1140/epja/i2019-12716-4](https://doi.org/10.1140/epja/i2019-12716-4). arXiv: [1810.12917 \[astro-ph.HE\]](https://arxiv.org/abs/1810.12917).
- [20] Michael W. Coughlin et al. "Multimessenger Bayesian parameter inference of a binary neutron star merger". In: *Mon. Not. Roy. Astron. Soc.* 489.1 (2019), pp. L91–L96. doi: [10.1093/mnrasl/slz133](https://doi.org/10.1093/mnrasl/slz133). arXiv: [1812.04803 \[astro-ph.HE\]](https://arxiv.org/abs/1812.04803).
- [21] B. P. Abbott et al. "Properties of the binary neutron star merger GW170817". In: *Phys. Rev. X* 9.1 (2019), p. 011001. doi: [10.1103/PhysRevX.9.011001](https://doi.org/10.1103/PhysRevX.9.011001). arXiv: [1805.11579 \[gr-qc\]](https://arxiv.org/abs/1805.11579).
- [22] Peter T. H. Pang et al. "An updated nuclear-physics and multi-messenger astrophysics framework for binary neutron star mergers". In: *Nature Commun.* 14.1 (2023), p. 8352. doi: [10.1038/s41467-023-43932-6](https://doi.org/10.1038/s41467-023-43932-6). arXiv: [2205.08513 \[astro-ph.HE\]](https://arxiv.org/abs/2205.08513).
- [23] Matteo Breschi et al. "Bayesian inference of multimessenger astrophysical data: Joint and coherent inference of gravitational waves and kilonovae". In: *Astron. Astrophys.* 689 (2024), A51. doi: [10.1051/0004-6361/202449173](https://doi.org/10.1051/0004-6361/202449173). arXiv: [2401.03750 \[astro-ph.HE\]](https://arxiv.org/abs/2401.03750).
- [24] A. G. Abac et al. "Observation of Gravitational Waves from the Coalescence of a 2.5–4.5 M_⊙ Compact Object and a Neutron Star". In: *Astrophys. J. Lett.* 970.2 (2024), p. L34. doi: [10.3847/2041-8213/ad5beb](https://doi.org/10.3847/2041-8213/ad5beb). arXiv: [2404.04248 \[astro-ph.HE\]](https://arxiv.org/abs/2404.04248).




Likelihood distributions

- Likelihood distributions, obtained from final posterior samples



Information gain

- “Prior” = Λ_i computed from EOSs conditioned on the posterior source-frame masses m_i^{src}
- “Posterior” = Λ_i from posterior samples
- KL divergence between prior and posterior in brackets



More posteriors

Event	Pop	EOS	$m_1^{\text{src}} [M_{\odot}]$	$m_2^{\text{src}} [M_{\odot}]$	q	Λ_1	Λ_2	$\bar{\Lambda}$	$\delta\bar{\Lambda}$	$d_L [\text{Mpc}]$	
GW170817 (BNS)	Uninformed			$1.47^{+0.12}_{-0.11}$	$1.26^{+0.10}_{-0.10}$	$0.86^{+0.14}_{-0.12}$	286^{+473}_{-286}	448^{+603}_{-448}	364^{+403}_{-234}	0^{+158}_{-170}	38^{+8}_{-11}
	U	PSRs	$1.51^{+0.09}_{-0.06}$	$1.23^{+0.05}_{-0.07}$	$0.81^{+0.06}_{-0.09}$	413^{+110}_{-133}	1477^{+636}_{-334}	795^{+108}_{-71}	119^{+68}_{-43}	41^{+6}_{-8}	
		PSRs+ χ_{EFT}	$1.52^{+0.08}_{-0.07}$	$1.23^{+0.06}_{-0.06}$	$0.81^{+0.08}_{-0.07}$	215^{+87}_{-111}	712^{+219}_{-175}	390^{+98}_{-93}	58^{+19}_{-25}	40^{+6}_{-8}	
		PSRs+NICER	$1.57^{+0.05}_{-0.07}$	$1.19^{+0.05}_{-0.03}$	$0.75^{+0.06}_{-0.05}$	264^{+86}_{-52}	1399^{+367}_{-342}	620^{+99}_{-82}	116^{+37}_{-38}	40^{+6}_{-8}	
	G	PSRs	$1.41^{+0.04}_{-0.03}$	$1.32^{+0.03}_{-0.03}$	$0.93^{+0.04}_{-0.05}$	578^{+131}_{-150}	874^{+229}_{-174}	713^{+143}_{-122}	35^{+28}_{-23}	39^{+7}_{-9}	
		PSRs+ χ_{EFT}	$1.42^{+0.05}_{-0.04}$	$1.31^{+0.03}_{-0.04}$	$0.92^{+0.05}_{-0.06}$	298^{+130}_{-99}	495^{+173}_{-118}	385^{+141}_{-85}	21^{+19}_{-13}	40^{+6}_{-9}	
		PSRs+NICER	$1.39^{+0.02}_{-0.02}$	$1.34^{+0.02}_{-0.02}$	$0.96^{+0.03}_{-0.03}$	641^{+186}_{-128}	826^{+277}_{-178}	729^{+219}_{-133}	21^{+22}_{-17}	38^{+7}_{-8}	
	DG	PSRs	$1.40^{+0.03}_{-0.02}$	$1.33^{+0.02}_{-0.02}$	$0.95^{+0.02}_{-0.03}$	687^{+140}_{-156}	960^{+183}_{-231}	814^{+143}_{-185}	29^{+21}_{-13}	40^{+6}_{-7}	
		PSRs+ χ_{EFT}	$1.39^{+0.02}_{-0.02}$	$1.34^{+0.02}_{-0.02}$	$0.96^{+0.03}_{-0.03}$	382^{+148}_{-111}	507^{+153}_{-152}	444^{+136}_{-131}	11^{+12}_{-9}	39^{+7}_{-7}	
		PSRs+NICER	$1.39^{+0.02}_{-0.02}$	$1.33^{+0.02}_{-0.02}$	$0.96^{+0.02}_{-0.03}$	703^{+184}_{-153}	934^{+200}_{-200}	806^{+157}_{-184}	26^{+14}_{-17}	38^{+7}_{-8}	
GW190425 (BNS)	Uninformed			$1.75^{+0.13}_{-0.10}$	$1.56^{+0.10}_{-0.10}$	$0.89^{+0.11}_{-0.11}$	292^{+541}_{-292}	415^{+694}_{-415}	374^{+436}_{-333}	0^{+187}_{-173}	157^{+64}_{-65}
	U	PSRs	$1.81^{+0.10}_{-0.09}$	$1.50^{+0.07}_{-0.08}$	$0.83^{+0.08}_{-0.09}$	109^{+56}_{-57}	428^{+209}_{-200}	230^{+73}_{-71}	39^{+24}_{-23}	173^{+54}_{-47}	
		PSRs+ χ_{EFT}	$1.83^{+0.12}_{-0.10}$	$1.48^{+0.07}_{-0.10}$	$0.81^{+0.09}_{-0.09}$	58^{+40}_{-28}	259^{+174}_{-107}	132^{+44}_{-40}	23^{+21}_{-12}	187^{+48}_{-49}	
		PSRs+NICER	$1.82^{+0.10}_{-0.12}$	$1.48^{+0.10}_{-0.07}$	$0.81^{+0.11}_{-0.09}$	108^{+81}_{-56}	449^{+132}_{-150}	232^{+57}_{-51}	40^{+15}_{-23}	182^{+41}_{-49}	
	DG	PSRs	$1.93^{+0.08}_{-0.08}$	$1.41^{+0.05}_{-0.05}$	$0.73^{+0.06}_{-0.05}$	67^{+37}_{-38}	636^{+217}_{-206}	238^{+65}_{-61}	63^{+21}_{-21}	186^{+44}_{-44}	
		PSRs+ χ_{EFT}	$1.94^{+0.07}_{-0.06}$	$1.40^{+0.04}_{-0.05}$	$0.72^{+0.04}_{-0.05}$	51^{+23}_{-24}	408^{+125}_{-100}	156^{+37}_{-45}	37^{+10}_{-9}	190^{+41}_{-47}	
		PSRs+NICER	$1.90^{+0.08}_{-0.07}$	$1.42^{+0.06}_{-0.05}$	$0.75^{+0.06}_{-0.06}$	96^{+41}_{-36}	690^{+188}_{-164}	283^{+45}_{-51}	66^{+18}_{-17}	186^{+41}_{-48}	
GW230529 (NSBH)	Uninformed			$3.65^{+0.62}_{-0.85}$	$1.43^{+0.26}_{-0.24}$	$0.39^{+0.17}_{-0.13}$	—	2791^{+2113}_{-1950}	182^{+243}_{-182}	91^{+115}_{-91}	201^{+84}_{-97}
	U	PSRs	$3.73^{+0.28}_{-0.32}$	$1.38^{+0.10}_{-0.09}$	$0.37^{+0.06}_{-0.05}$	—	698^{+344}_{-303}	41^{+8}_{-8}	21^{+4}_{-4}	238^{+61}_{-63}	
		PSRs+ χ_{EFT}	$3.52^{+0.46}_{-0.45}$	$1.46^{+0.18}_{-0.14}$	$0.41^{+0.12}_{-0.08}$	—	322^{+206}_{-202}	26^{+8}_{-7}	13^{+4}_{-4}	220^{+70}_{-64}	
		PSRs+NICER	$3.27^{+0.27}_{-0.57}$	$1.54^{+0.25}_{-0.12}$	$0.47^{+0.17}_{-0.08}$	—	368^{+213}_{-291}	40^{+13}_{-14}	19^{+6}_{-7}	237^{+65}_{-67}	
	G	PSRs	$3.86^{+0.28}_{-0.26}$	$1.34^{+0.08}_{-0.08}$	$0.35^{+0.04}_{-0.05}$	—	933^{+497}_{-451}	46^{+14}_{-16}	23^{+7}_{-8}	236^{+59}_{-58}	
		PSRs+ χ_{EFT}	$3.83^{+0.25}_{-0.25}$	$1.35^{+0.07}_{-0.07}$	$0.35^{+0.04}_{-0.04}$	—	582^{+261}_{-198}	30^{+7}_{-8}	15^{+3}_{-4}	242^{+59}_{-57}	
		PSRs+NICER	$3.89^{+0.22}_{-0.24}$	$1.33^{+0.07}_{-0.06}$	$0.34^{+0.04}_{-0.03}$	—	1000^{+321}_{-311}	49^{+9}_{-9}	25^{+4}_{-5}	235^{+59}_{-58}	
	DG	PSRs	$3.78^{+0.23}_{-0.24}$	$1.36^{+0.07}_{-0.06}$	$0.36^{+0.04}_{-0.04}$	—	890^{+341}_{-333}	49^{+12}_{-13}	25^{+6}_{-7}	237^{+63}_{-64}	
		PSRs+ χ_{EFT}	$3.85^{+0.18}_{-0.21}$	$1.34^{+0.05}_{-0.06}$	$0.35^{+0.03}_{-0.03}$	—	516^{+194}_{-205}	26^{+8}_{-8}	13^{+4}_{-4}	237^{+66}_{-71}	
		PSRs+NICER	$3.86^{+0.15}_{-0.16}$	$1.34^{+0.04}_{-0.04}$	$0.35^{+0.02}_{-0.02}$	—	1026^{+254}_{-306}	52^{+11}_{-13}	26^{+6}_{-6}	229^{+65}_{-68}	