

Analyzing GW231109_235456 in the ET era and incorporating neutron star physics into future GW inference

arXiv:2510.22290 & arXiv:2511.22987

Thibeau Wouters, Anna Puecher, Peter T. H. Pang, Tim Dietrich, Chris Van Den Broeck



Utrecht
University

Nikhef

Table of Contents


- ① Part 1: Analyzing GW231109_235456 in the ET era
- ② Part 2: Neural priors for GW inference
- ③ Conclusion

Structure of this talk

Data analysis of neutron stars forms a **loop**:

- ① **Constraining** the EOS with neutron star observations
- ② **Applying** EOS knowledge in neutron star data analysis (e.g., GW)

How can we efficiently perform this loop?



Contents

- ① Part 1: Analyzing GW231109_235456 in the ET era
- ② Part 2: Neural priors for GW inference
- ③ Conclusion

GWTC-4.0 and GW231109_235456

- GWTC-4.0 released recently [1]:
 - Over 200 gravitational wave events analyzed in total

GWTC-4.0 and GW231109_235456

- GWTC-4.0 released recently [1]:
 - Over 200 gravitational wave events analyzed in total
- **No confident binary neutron star detections**

GWTC-4.0 and GW231109_235456

- GWTC-4.0 released recently [1]:
 - Over 200 gravitational wave events analyzed in total
- **No confident binary neutron star detections**
- However: sub-threshold candidate **GW231109_235456** identified [2]
 - $\text{SNR} \sim 9.7$ (distance: ~ 165 Mpc)
 - Fainter, but mass closer to GW170817 than GW190425


GWTC-4.0 and GW231109_235456

- GWTC-4.0 released recently [1]:
 - Over 200 gravitational wave events analyzed in total
- **No confident binary neutron star detections**
- However: sub-threshold candidate **GW231109_235456** identified [2]
 - $\text{SNR} \sim 9.7$ (distance: ~ 165 Mpc)
 - Fainter, but mass closer to GW170817 than GW190425

What can we learn about the EOS from such a merger?


GW231109_235456: component masses

Component masses compared to other low-mass GW events [3–5]



Parameter estimation on GW231109_235456

- IMRPhenomXP_NRTidalv3
- Standard priors for $m_i, \Lambda_i \leq 5000$, spins below 0.05 or 0.4




Constraining EOS from GW231109_235456

Put information on jester here


Constraining EOS from GW231109_235456

Constraints on radius of $1.4 M_{\odot}$ neutron star ($R_{1.4}$):




Projection: Einstein Telescope & Cosmic Explorer

- Simulate GW231109-like event with third-generation detectors
- Einstein Telescope: SNR ~ 134 , with Cosmic Explorer: SNR ~ 294
- Recovery improved



Projection: radius constraints

Recover radius with accuracy of 300-400 meters (ET+CE vs ET)



Conclusion (part 1)

- GW231109_235456: sub-threshold BNS candidate from O4a
- SNR matters for EOS inference
 - Current detectors: poor constraints
 - ET and CE: precise radius measurements ($\sim 300\text{-}400$ m)
- JESTER can handle individual high SNR signals: efficiently go from NS observations to EOS constraints

Contents


- ① Part 1: Analyzing GW231109_235456 in the ET era
- ② Part 2: Neural priors for GW inference
- ③ Conclusion

Structure of this talk

Data analysis of neutron stars forms a **loop**:

- ① **Constraining** the EOS with neutron star observations
- ② **Applying** EOS knowledge in neutron star data analysis (e.g., GW)

How can we efficiently perform this loop?



Neural priors: motivation

- Bayesian inference depends on choice of **priors**:

$$\mathcal{P}(\theta_{\text{GW}}|d) \propto \frac{\mathcal{L}(d|\theta_{\text{GW}})\pi(\theta_{\text{GW}})}{\mathcal{Z}}$$


- By default, we use **agnostic priors**, but what if we *do* have non-trivial prior information?
- Case study: neutron stars (NSs) and information from
 - Population models
 - Equation of state (EOS) constraints

Neural priors

Flexible way to encode NS physics into GW inference

Neural priors: key idea

Train normalizing flow (NF) on samples informed by populations and EOS
→ **neural prior**



NS population models

Three fiducial population models for NS masses:

① **Uniform** [6–8]:

- Only use EOS constraints for maximum mass (M_{TOV})
- NS mass $\sim U[1 \text{ M}_\odot, M_{\text{TOV}}]$

② **Gaussian** [9]:

- NS mass $\sim \mathcal{N}(1.33 \text{ M}_\odot, (0.09 \text{ M}_\odot)^2)$

③ **Double Gaussian** [10, 11]:

- Weighted mixture of two Gaussians
- $0.65 \times \mathcal{N}(1.34 \text{ M}_\odot, (0.07 \text{ M}_\odot)^2) + 0.35 \times \mathcal{N}(1.80 \text{ M}_\odot, (0.21 \text{ M}_\odot)^2)$


For NSBH systems:

- BH mass m_1^{src} : from $[M_{\text{TOV}}, 5 \text{ M}_\odot]$
- NS mass m_2^{src} : above models


EOS constraints

- We use three EOS constraints [12]:
 - ① **Heavy pulsars:** must support $2 M_{\odot}$ NSs
 - ② **Chiral EFT (χ_{EFT}):** nuclear theory predictions (softer EOS)
 - ③ **NICER:** mass-radius observations of NSs (stiffer EOS)
- Posterior samples obtained with JESTER [13] 

Chiral EFT




NICER



Normalizing flows

Normalizing flows [14, 15]

- Neural density estimators: trainable bijections
- Often used in PE, e.g., DINGO [16, 17], NESSAI [18, 19]
- Generate samples, evaluate density: can be used as priors [20]



Construction of neural priors


Steps to generate training data:

- ① Draw EOS posterior curve: determines $M_{\text{TOV}}, \Lambda(m)$
- ② Draw masses from population model
- ③ Compute $\Lambda_i = \Lambda(m_i)$ for NSs (NSBH: $\Lambda_1 = 0$)

Implementation:

- Created with GLASFLOW [21, 22]
- CouplingNSF architecture (neural spline flows [23])
- Use as a JointPrior in BILBY (NFPrior)
 - Sample & logpdf: evaluate NF
 - Rescale: unit hypercube \rightarrow multivariate Gaussian $\xrightarrow{\text{NF}}$ data space

All neural priors



Setup

Analyze GW170817, GW190425, GW230529 with:

- 4096 live points, multibanding likelihood
- IMRPhenomXP_NRTidalv3
- Neural priors for m_i, Λ_i (standard priors for other parameters)

Two contributions:

- ① Narrower constraints with neural priors
- ② Model selection with Bayes factors

Jeffreys' scale for Bayes factors (\log_{10} scale)


| $\log_{10}(\mathcal{B}_1^2)$ | Interpretation | Color |
|------------------------------|-------------------------|---------------|
| $[0, \frac{1}{2}]$ | Barely worth mentioning | Light orange |
| $[\frac{1}{2}, 1]$ | Substantial | Orange |
| $[1, \frac{3}{2}]$ | Strong | Red-orange |
| $[\frac{3}{2}, 2]$ | Very strong | Dark red |
| > 2 | Decisive | Very dark red |

GW170817: Source classification

Showing \log_{10} Bayes factors relative to model with highest evidence

| Source | Population | EOS | GW170817 |
|--------|-----------------|------------------|----------|
| BNS | Uniform | PSRs | -1.83 |
| | | PSRs+ χ EFT | -0.80 |
| | | PSRs+NICER | -1.58 |
| | Gaussian | PSRs | -0.68 |
| | | PSRs+ χ EFT | ref. |
| | | PSRs+NICER | -0.76 |
| | Double Gaussian | PSRs | -1.36 |
| | | PSRs+ χ EFT | -0.59 |
| | | PSRs+NICER | -0.92 |
| NSBH | Uniform | PSRs | -224.65 |
| | | PSRs+ χ EFT | -224.66 |
| | | PSRs+NICER | -224.66 |
| | Gaussian | PSRs | -224.67 |
| | | PSRs+ χ EFT | -224.66 |
| | | PSRs+NICER | -224.66 |
| | Double Gaussian | PSRs | -224.67 |
| | | PSRs+ χ EFT | -224.68 |
| | | PSRs+NICER | -224.67 |

GW170817: Parameter constraints (Gaussian pop.)



GW170817: Discussion

Source classification:

- Decisive evidence for BNS over NSBH
- Prefer Gaussian population model
- Slight preference for softer EOS (PSRs+ χ_{EFT})

Parameter constraints:


- More equal mass ratio: $q \geq 0.9$
- Constrained tidal deformability $\tilde{\Lambda}$
- Higher luminosity distance compared to agnostic prior
- Matches multimessenger analyses of GW170817 [24–28]

GW190425: Source classification

Showing \log_{10} Bayes factors relative to best model

| Source | Population | EOS | GW190425 |
|--------|-----------------|------------------|----------|
| BNS | Uniform | PSRs | -0.07 |
| | | PSRs+ χ EFT | -0.11 |
| | | PSRs+NICER | ref. |
| | Gaussian | PSRs | -6.89 |
| | | PSRs+ χ EFT | -8.47 |
| | | PSRs+NICER | -5.45 |
| | Double Gaussian | PSRs | -0.55 |
| | | PSRs+ χ EFT | -0.79 |
| | | PSRs+NICER | -0.57 |
| NSBH | Uniform | PSRs | -1.52 |
| | | PSRs+ χ EFT | -1.35 |
| | | PSRs+NICER | -1.63 |
| | Gaussian | PSRs | -0.82 |
| | | PSRs+ χ EFT | -1.11 |
| | | PSRs+NICER | -1.43 |
| | Double Gaussian | PSRs | -4.11 |
| | | PSRs+ χ EFT | -3.83 |
| | | PSRs+NICER | -24.31 |

GW190425: Parameter constraints (Uniform pop.)



GW190425: Discussion

Source classification:

- Prefer BNS over NSBH (but less conclusive than GW170817)
- Most consistent with uniform population
- Masses are outliers compared to galactic binaries

Parameter constraints:


- Less equal masses: $q \leq 0.9$
- $\tilde{\Lambda} \approx 200$ (more prior-dominated due to lower SNR)
- Higher luminosity distance: 182_{-49}^{+41} Mpc vs. 157_{-65}^{+64} Mpc (90% credibility)

GW230529: Source classification

Showing \log_{10} Bayes factors relative to best model

| Source | Population | EOS | GW230529 |
|--------|-----------------|------------------|----------|
| BNS | Uniform | PSRs | -13.14 |
| | | PSRs+ χ EFT | -13.12 |
| | | PSRs+NICER | -12.92 |
| | Gaussian | PSRs | -18.82 |
| | | PSRs+ χ EFT | -18.83 |
| | | PSRs+NICER | -18.81 |
| | Double Gaussian | PSRs | -13.75 |
| | | PSRs+ χ EFT | -13.77 |
| | | PSRs+NICER | -13.71 |
| NSBH | Uniform | PSRs | -0.08 |
| | | PSRs+ χ EFT | -0.02 |
| | | PSRs+NICER | -0.25 |
| | Gaussian | PSRs | -0.05 |
| | | PSRs+ χ EFT | -0.20 |
| | | PSRs+NICER | ref. |
| | Double Gaussian | PSRs | -0.14 |
| | | PSRs+ χ EFT | -0.13 |
| | | PSRs+NICER | -0.05 |

GW230529: Parameter constraints (Gaussian pop.)



Source classification:

- Decisive evidence for NSBH over BNS (agrees with LVK [5])
- No evidence between hypotheses (low SNR)

Parameter constraints:

- Mass ratio more constrained: $q \leq 0.4$
 - As a result, improved spin constraints (χ_{1z} closer to zero)
- Tidal deformability posteriors dominated by priors
- Luminosity distance: 235^{+59}_{-58} Mpc vs. 201^{+84}_{-97} Mpc (90% credibility)
- Less tidal information: NRTidalv3 tapers at (90% credibility)
 - 641^{+318}_{-158} Hz (agnostic priors)
 - 858^{+113}_{-108} Hz (neural prior, reference model)

Conclusion (part 2)

- Flexible way to encode non-trivial prior information
- Two highlights:
 - ① Bayesian model selection
 - ② Informed parameter constraints
- Implemented in BILBY
- Data-driven approach: easy to extend/generalize


Contents

- ① Part 1: Analyzing GW231109_235456 in the ET era
- ② Part 2: Neural priors for GW inference
- ③ Conclusion

Conclusion

Methods for closing the neutron star data analysis loop:

- ① JESTER: from nuclear physics and multimessenger astronomy to EOS constraints
- ② Neural priors: incorporate NS physics into GW inference



References I

- [1] A. G. Abac et al. "GWTC-4.0: Updating the Gravitational-Wave Transient Catalog with Observations from the First Part of the Fourth LIGO-Virgo-KAGRA Observing Run". In: (Aug. 2025). arXiv: [2508.18082 \[gr-qc\]](#).
- [2] Wanting Niu et al. "GW231109_235456: A Sub-threshold Binary Neutron Star Merger in the LIGO-Virgo-KAGRA O4a Observing Run?" In: (Sept. 2025). arXiv: [2509.09741 \[astro-ph.HE\]](#).
- [3] B. P. Abbott et al. "GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral". In: *Phys. Rev. Lett.* 119.16 (2017), p. 161101. DOI: [10.1103/PhysRevLett.119.161101](#). arXiv: [1710.05832 \[gr-qc\]](#).
- [4] B. P. Abbott et al. "GW190425: Observation of a Compact Binary Coalescence with Total Mass $\sim 3.4 M_{\odot}$ ". In: *Astrophys. J. Lett.* 892.1 (2020), p. L3. DOI: [10.3847/2041-8213/ab75f5](#). arXiv: [2001.01761 \[astro-ph.HE\]](#).
- [5] A. G. Abac et al. "Observation of Gravitational Waves from the Coalescence of a 2.5–4.5 M_{\odot} Compact Object and a Neutron Star". In: *Astrophys. J. Lett.* 970.2 (2024), p. L34. DOI: [10.3847/2041-8213/ad5beb](#). arXiv: [2404.04248 \[astro-ph.HE\]](#).
- [6] R. Abbott et al. "Observation of Gravitational Waves from Two Neutron Star–Black Hole Coalescences". In: *Astrophys. J. Lett.* 915.1 (2021), p. L5. DOI: [10.3847/2041-8213/ac082e](#). arXiv: [2106.15163 \[astro-ph.HE\]](#).

References II

- [7] Philippe Landry and Jocelyn S. Read. "The Mass Distribution of Neutron Stars in Gravitational-wave Binaries". In: *Astrophys. J. Lett.* 921.2 (2021), p. L25. DOI: [10.3847/2041-8213/ac2f3e](https://doi.org/10.3847/2041-8213/ac2f3e). arXiv: [2107.04559 \[astro-ph.HE\]](https://arxiv.org/abs/2107.04559).
- [8] Jacob Golomb et al. "Interplay of astrophysics and nuclear physics in determining the properties of neutron stars". In: *Phys. Rev. D* 111.2 (2025), p. 023029. DOI: [10.1103/PhysRevD.111.023029](https://doi.org/10.1103/PhysRevD.111.023029). arXiv: [2410.14597 \[astro-ph.HE\]](https://arxiv.org/abs/2410.14597).
- [9] Feryal Özal and Paulo Freire. "Masses, Radii, and the Equation of State of Neutron Stars". In: *Ann. Rev. Astron. Astrophys.* 54 (2016), pp. 401–440. DOI: [10.1146/annurev-astro-081915-023322](https://doi.org/10.1146/annurev-astro-081915-023322). arXiv: [1603.02698 \[astro-ph.HE\]](https://arxiv.org/abs/1603.02698).
- [10] Justin Alsing, Hector O. Silva, and Emanuele Berti. "Evidence for a maximum mass cut-off in the neutron star mass distribution and constraints on the equation of state". In: *Mon. Not. Roy. Astron. Soc.* 478.1 (2018), pp. 1377–1391. DOI: [10.1093/mnras/sty1065](https://doi.org/10.1093/mnras/sty1065). arXiv: [1709.07889 \[astro-ph.HE\]](https://arxiv.org/abs/1709.07889).
- [11] Dong-Sheng Shao et al. "Maximum mass cutoff in the neutron star mass distribution and the prospect of forming supramassive objects in the double neutron star mergers". In: *Phys. Rev. D* 102.6 (2020), p. 063006. DOI: [10.1103/PhysRevD.102.063006](https://doi.org/10.1103/PhysRevD.102.063006). arXiv: [2009.04275 \[astro-ph.HE\]](https://arxiv.org/abs/2009.04275).

References III

- [12] Hauke Koehn et al. "From existing and new nuclear and astrophysical constraints to stringent limits on the equation of state of neutron-rich dense matter". In: *Phys. Rev. X* 15.2 (2025), p. 021014. DOI: [10.1103/PhysRevX.15.021014](https://doi.org/10.1103/PhysRevX.15.021014). arXiv: [2402.04172 \[astro-ph.HE\]](https://arxiv.org/abs/2402.04172).
- [13] Thibeau Wouters et al. "Leveraging differentiable programming in the inverse problem of neutron stars". In: (Apr. 2025). arXiv: [2504.15893 \[astro-ph.HE\]](https://arxiv.org/abs/2504.15893).
- [14] Ivan Kobyzev, Simon J. D. Prince, and Marcus A. Brubaker. "Normalizing Flows: An Introduction and Review of Current Methods". In: *IEEE Trans. Pattern Anal. Machine Intell.* 43.11 (2021), pp. 3964–3979. DOI: [10.1109/tpami.2020.2992934](https://doi.org/10.1109/tpami.2020.2992934). arXiv: [1908.09257 \[stat.ML\]](https://arxiv.org/abs/1908.09257).
- [15] George Papamakarios et al. "Normalizing Flows for Probabilistic Modeling and Inference". In: *J. Machine Learning Res.* 22.1 (2021), pp. 2617–2680. DOI: [10.5555/3546258.3546315](https://doi.org/10.5555/3546258.3546315). arXiv: [1912.02762 \[stat.ML\]](https://arxiv.org/abs/1912.02762).
- [16] Maximilian Dax et al. "Real-Time Gravitational Wave Science with Neural Posterior Estimation". In: *Phys. Rev. Lett.* 127.24 (2021), p. 241103. DOI: [10.1103/PhysRevLett.127.241103](https://doi.org/10.1103/PhysRevLett.127.241103). arXiv: [2106.12594 \[gr-qc\]](https://arxiv.org/abs/2106.12594).
- [17] Maximilian Dax et al. "Neural Importance Sampling for Rapid and Reliable Gravitational-Wave Inference". In: *Phys. Rev. Lett.* 130.17 (2023), p. 171403. DOI: [10.1103/PhysRevLett.130.171403](https://doi.org/10.1103/PhysRevLett.130.171403). arXiv: [2210.05686 \[gr-qc\]](https://arxiv.org/abs/2210.05686).

References IV

- [18] Michael J. Williams, John Veitch, and Chris Messenger. “Nested sampling with normalizing flows for gravitational-wave inference”. In: *Phys. Rev. D* 103.10 (2021), p. 103006. DOI: [10.1103/PhysRevD.103.103006](https://doi.org/10.1103/PhysRevD.103.103006). arXiv: [2102.11056 \[gr-qc\]](https://arxiv.org/abs/2102.11056).
- [19] Michael J. Williams, John Veitch, and Chris Messenger. “Importance nested sampling with normalising flows”. In: *Mach. Learn. Sci. Tech.* 4.3 (2023), p. 035011. DOI: [10.1088/2632-2153/acd5aa](https://doi.org/10.1088/2632-2153/acd5aa). arXiv: [2302.08526 \[astro-ph.IM\]](https://arxiv.org/abs/2302.08526).
- [20] Justin Alsing and Will Handley. “Nested sampling with any prior you like”. In: *Mon. Not. Roy. Astron. Soc.* 505.1 (2021), pp. L95–L99. DOI: [10.1093/mnrasl/slab057](https://doi.org/10.1093/mnrasl/slab057). arXiv: [2102.12478 \[astro-ph.IM\]](https://arxiv.org/abs/2102.12478).
- [21] Michael J. Williams et al. *uofgravity/glasflow: v0.4.1*. Oct. 2024. DOI: [10.5281/zenodo.13914483](https://doi.org/10.5281/zenodo.13914483). URL: <https://doi.org/10.5281/zenodo.13914483>.
- [22] Conor Durkan et al. *nflows: normalizing flows in PyTorch*. Version v0.14. Nov. 2020. DOI: [10.5281/zenodo.4296287](https://doi.org/10.5281/zenodo.4296287). URL: <https://doi.org/10.5281/zenodo.4296287>.
- [23] Conor Durkan et al. “Neural Spline Flows”. In: June 2019. arXiv: [1906.04032 \[stat.ML\]](https://arxiv.org/abs/1906.04032).
- [24] David Radice and Liang Dai. “Multimessenger Parameter Estimation of GW170817”. In: *Eur. Phys. J. A* 55.4 (2019), p. 50. DOI: [10.1140/epja/i2019-12716-4](https://doi.org/10.1140/epja/i2019-12716-4). arXiv: [1810.12917 \[astro-ph.HE\]](https://arxiv.org/abs/1810.12917).

References V

- [25] Michael W. Coughlin et al. "Multimessenger Bayesian parameter inference of a binary neutron star merger". In: *Mon. Not. Roy. Astron. Soc.* 489.1 (2019), pp. L91–L96. DOI: [10.1093/mnrasl/slz133](https://doi.org/10.1093/mnrasl/slz133). arXiv: [1812.04803](https://arxiv.org/abs/1812.04803) [astro-ph.HE].
- [26] B. P. Abbott et al. "Properties of the binary neutron star merger GW170817". In: *Phys. Rev. X* 9.1 (2019), p. 011001. DOI: [10.1103/PhysRevX.9.011001](https://doi.org/10.1103/PhysRevX.9.011001). arXiv: [1805.11579](https://arxiv.org/abs/1805.11579) [gr-qc].
- [27] Peter T. H. Pang et al. "An updated nuclear-physics and multi-messenger astrophysics framework for binary neutron star mergers". In: *Nature Commun.* 14.1 (2023), p. 8352. DOI: [10.1038/s41467-023-43932-6](https://doi.org/10.1038/s41467-023-43932-6). arXiv: [2205.08513](https://arxiv.org/abs/2205.08513) [astro-ph.HE].
- [28] Matteo Breschi et al. "Bayesian inference of multimessenger astrophysical data: Joint and coherent inference of gravitational waves and kilonovae". In: *Astron. Astrophys.* 689 (2024), A51. DOI: [10.1051/0004-6361/202449173](https://doi.org/10.1051/0004-6361/202449173). arXiv: [2401.03750](https://arxiv.org/abs/2401.03750) [astro-ph.HE].

Posterior distributions for ET/ET+CE injections

