

Fast parameter estimation of gravitational waves from binary neutron stars with JAX and normalizing flows

Thibau Wouters, Peter T. H. Pang, Tim Dietrich, Chris Van Den Broeck

t.r.i.wouters@uu.nl



[arXiv:2404.11397](https://arxiv.org/abs/2404.11397)

LVK Barcelona, September 2024



Utrecht
University



Table of Contents

① Introduction

② Methods

③ Results


④ Conclusion

Parameter estimation

Parameter estimation (PE): get **posterior** of GW parameters θ

$$p(\theta|d) = \frac{p(d|\theta)p(\theta)}{p(d)} = \frac{\text{likelihood} \times \text{prior}}{\text{evidence}}$$

Problem: Markov Chain Monte Carlo (MCMC): computationally expensive for binary neutron stars (BNS)



Overview

JIM: fast parameter estimation of GW signals with JAX

- MCMC sampler: FLOWMC
- Waveforms: RIPPLE




Table of Contents

① Introduction

② Methods

③ Results

④ Conclusion

JAX = NUMPY + composable transformations:


- ① Automatic differentiation
- ② Just-in-time (JIT) compilation
- ③ GPU acceleration
- ④ Parallelization



Potential for massive acceleration, such as in parameter estimation!

Normalizing flows

- Generative machine learning model
- Learn mapping between **latent** and **parameter** space
- Enable approximate sampling from complicated distributions
- Training data: MCMC samples



FLOWMC: normalizing-flow (NF) enhanced MCMC sampling

- 1 Gradient-based sampler (local sampler)
- 2 Train NF with samples from local sampler
- 3 Sample normalizing flow (global sampler)




Table of Contents

① Introduction

② Methods

③ Results

④ Conclusion

Results

- Waveforms: TaylorF2 (TF2), IMRPhenomD_NRTidalv2 (NRTv2)
- JIM wall time: (i) computing reference parameters for relative binning, (ii) training NF, (iii) sampling

Event	Waveform	JIM (1 GPU)	pBILBY (480 cores)	RB-BILBY (24 cores)	ROQ-BILBY (24 cores)
GW170817	TF2	(4.85 + 15.33) min	9.64 h	3.8 h	–
	NRTv2	(5.38 + 25.59) min	10.99 h	4.11 h	1.65 h
GW190425	TF2	(2.63 + 18.30) min	8.18 h	2.81 h	–
	NRTv2	(3.26 + 21.20) min	4.91 h	2.42 h	0.97 h
Injection	TF2	24.76 min	–	–	–
	NRTv2	18.02 min	–	–	–

(pBILBY = PARALLEL BILBY, RB = relative binning, ROQ = reduced order quadrature)

Environmental impact

JIM is **more environmentally friendly** than existing pipelines

- Energy consumption for all 204 runs of paper
- Convert to number of trees to capture the emitted CO_2 in a year.

Method	Trees
JIM	0.55
pBILBY	67.68
RB-BILBY	1.32
ROQ-BILBY	sampling 0.52
	precompute [†] 0.44

[†]Estimated cost to build ROQ bases.

Table of Contents

① Introduction

② Methods


③ Results

④ Conclusion

Conclusion

JIM: a fast and environmentally friendly PE pipeline for GW signals. Our contribution:

- TaylorF2 and IMRPhenomD_NRTidalv2 in RIPPLE
- Parameter estimation of BNS in 15 – 30 minutes sampling time without pretraining



References

- [1] Kaze WK Wong, Maximiliano Isi, and Thomas DP Edwards. “Fast gravitational wave parameter estimation without compromises”. In: *arXiv preprint arXiv:2302.05333* (2023). Available at: <https://github.com/kazewong/jim>.
- [2] Thomas DP Edwards et al. “RIPPLE: Differentiable and Hardware-Accelerated Waveforms for Gravitational Wave Data Analysis”. In: *arXiv preprint arXiv:2302.05329* (2023). Available at: <https://github.com/tedwards2412/ripple/>.
- [3] Marylou Gabrié, Grant M Rotskoff, and Eric Vanden-Eijnden. “Efficient bayesian sampling using normalizing flows to assist markov chain monte carlo methods”. In: *arXiv preprint arXiv:2107.08001* (2021).
- [4] Kaze WK Wong, Marylou Gabrié, and Daniel Foreman-Mackey. “FLOWMC: Normalizing-flow enhanced sampling package for probabilistic inference in Jax”. In: *arXiv preprint arXiv:2211.06397* (2022). Available at: <https://github.com/kazewong/flowMC>.
- [5] Roy Frostig, Matthew James Johnson, and Chris Leary. “Compiling machine learning programs via high-level tracing”. In: *Systems for Machine Learning 4.9* (2018). Available at: <https://github.com/google/jax>.
- [6] Peter T. H. Pang et al. “An updated nuclear-physics and multi-messenger astrophysics framework for binary neutron star mergers”. In: *Nature Commun.* 14.1 (2023). Available at <https://github.com/nuclear-multimessenger-astronomy/nmma>, p. 8352. DOI: 10.1038/s41467-023-43932-6. arXiv: 2205.08513 [astro-ph.HE].

APPENDIX

Normalizing flow details

- Rational-quadratic neural spline flows
- 10 layers, 8 bins
- 128 neurons in hidden layers
- Adam optimizer, learning rate decayed (polynomial schedule)
- Deep learning library: EQUINOX


Loss function: KL divergence on sampled data

$$\mathcal{L}(T) = -\frac{1}{n} \sum_{i=1}^n \log \hat{\rho}(x_i)$$

Stopping criterion

We stop training the NF if we achieve a mean Metropolis-Hastings acceptance rate of 10% (20%) for real events (injections).


Example: GW170817, TaylorF2 with 20%:



Validation – Mismatch waveforms

Cross-check against LALSUITE: mismatch histogram based on 10 000 waveforms, from uniform samples with following ranges:

Parameter	Range
Component masses	$[0.5M_{\odot}, 3M_{\odot}]$
Component aligned spins	$[-0.05, 0.05]$
Dimensionless tidal deformabilities	$[0, 5000]$
Inclination angle	$[0, \pi]$



Validation – p-p plot

We demonstrate the robustness of JIM:

- 100 GW events with HLV at design sensitivity and $T = 128$ s,
- NRTv2: reference waveform relative binning without taper,
- Priors: Table 1.





Table 1: Prior ranges used in our analyses. All priors are uniform priors with the specified range.

Parameter	Injection	GW170817	GW190425
$\mathcal{M} [M_{\odot}]$	[0.88, 2.61]	[1.18, 1.21]	[1.485, 1.490]
q	[0.5, 1]	[0.125, 1]	[0.125, 1]
χ_i	$[-0.05, 0.05]$	$[-0.05, 0.05]$	$[-0.05, 0.05]$
Λ_i	[0, 5000]	[0, 5000]	[0, 5000]
d_L [Mpc]	[30, 300]	[1, 75]	[1, 500]
t_c [s]	$[-0.1, 0.1]$	$[-0.1, 0.1]$	$[-0.1, 0.1]$
ϕ_c	$[0, 2\pi]$	$[0, 2\pi]$	$[0, 2\pi]$
$\cos \iota$	$[-1, 1]$	$[-1, 1]$	$[-1, 1]$
ψ	$[0, \pi]$	$[0, \pi]$	$[0, \pi]$
α	$[0, 2\pi]$	$[0, 2\pi]$	$[0, 2\pi]$
$\sin \delta$	$[-1, 1]$	$[-1, 1]$	$[-1, 1]$

GW170817 & GW190425: Jensen-Shannon divergences

Table 2: Jensen-Shannon divergences (in bits) between the marginal posterior obtained for GW170817 and GW190425 using TaylorF2 and IMRPhenomD_NRTidalv2 with JIM and PBILBY, with the highest value of each comparison in bold. The divergences are bound between $[0, 1]$.

Parameter	GW170817		GW190425	
	TF2	NRTv2	TF2	NRTv2
\mathcal{M}	0.001725	0.000516	0.003557	0.002461
q	0.005212	0.007894	0.004837	0.002960
χ_1	0.005633	0.004301	0.002794	0.004825
χ_2	0.003030	0.002671	0.002416	0.003041
Λ_1	0.001062	0.002208	0.008556	0.000783
Λ_2	0.000559	0.002186	0.005808	0.003576
d_L	0.001544	0.01847	0.001273	0.002878
ϕ_c	0.003500	0.010714	0.003338	0.006126
$\cos \iota$	0.001615	0.012851	0.006400	0.005279
ψ	0.004048	0.011036	0.001516	0.003730
α	0.014008	0.001258	0.009822	0.012291
$\sin \delta$	0.009570	0.001761	0.008934	0.009228

GW170817 with TaylorF2

Figure 1





Figure 2



GW190425 with TaylorF2

Figure 3




Figure 4

