

OpenMalaria

ITN deployment strategies

ITN parameterisation

Settings

Outcomes

Simulation set 1

- **PYR-ITNs** (baseline coverage, every 3 years)
- **CFP-PYR-ITN** without compromise
- **CFP-PYR-ITN** with **25% coverage reduction**
- **CFP-PYR-ITN** with **50% coverage reduction**
- **CFP-PYR-ITN** every **4 years**
- **CFP-PYR-ITN** every **6 years**

Parametrisation 1

- Efficacy: entomological study in Tanzania
- Usage: coverage survey in Tanzania

Parametrisation 2A

- Efficacy: entomological study in Benin
- Usage: coverage survey in Benin

Parametrisation 2B

- Efficacy: entomological study in Benin
- Usage: attrition study in Benin

- **Human blood index:** 50% or 99%

- **Entomological inoculation rate:** 5, 10, or 100 infective bites/person/year

- **Transmission seasonality:** perennial or seasonal

- **Access to first-line treatment:** 10 or 40% receiving treatment within 14 days of symptom onset

Incidence of uncomplicated malaria cases in the **whole population** over a **12-year period**

Simulation set 2

- **PYR-ITNs** (baseline coverage, every 3 years)
- **CFP-PYR-ITN** at **reduced coverage** (for various coverage levels, every 3 years)

Incidence of uncomplicated malaria cases among **ITN users and ITN non-users** over a **3-year period**