

# Lecture 5

## Epipolar Geometry



Professor Silvio Savarese

*Computational Vision and Geometry Lab*



# Lecture 5


## Epipolar Geometry

- Why is stereo useful?
- Epipolar constraints
- Essential and fundamental matrix
- Estimating  $F$
- Examples

**Reading:** [AZ] Chapter: 4 “Estimation – 2D perspective transformations”  
Chapter: 9 “Epipolar Geometry and the Fundamental Matrix Transformation”  
Chapter: 11 “Computation of the Fundamental Matrix  $F$ ”  
[FP] Chapter: 7 “Stereopsis”  
Chapter: 8 “Structure from Motion”



# Recovering structure from a single view



From calibration rig

→ location/pose of the rig,  $K$


From points and lines at infinity  
+ orthogonal lines and planes

→ structure of the scene,  $K$

---

Knowledge about scene (point correspondences, geometry of lines & planes, etc...)

# Recovering structure from a single view



## Why is it so difficult?

Intrinsic ambiguity of the mapping from 3D to image (2D)


# Recovering structure from a single view

Intrinsic ambiguity of the mapping from 3D to image (2D)



Courtesy slide S. Lazebnik


# Two eyes help!



# Two eyes help!

$$P = l \times l'$$

[Eq. 1]




This is called **triangulation**

# Triangulation

- Find  $P^*$  that minimizes

$$d(p, M P^*) + d(p', M' P^*) \quad [\text{Eq. 2}]$$






# Multi (stereo)-view geometry

- **Camera geometry:** Given corresponding points in two images, find camera matrices, position and pose.
- **Scene geometry:** Find coordinates of 3D point from its projection into 2 or multiple images.
- **Correspondence:** Given a point  $p$  in one image, how can I find the corresponding point  $p'$  in another one?

# Epipolar geometry





- Epipolar Plane
- Baseline
- Epipolar Lines


- Epipoles  $e, e'$

= intersections of baseline with image planes  
= projections of the other camera center

# Example of epipolar lines



# Example: Parallel image planes




- Baseline intersects the image plane at infinity
- Epipoles are at infinity
- Epipolar lines are parallel to  $u$  axis



# Example: Parallel Image Planes



# Example: Forward translation



- The epipoles have same position in both images
- Epipole called FOE (focus of expansion)






# Epipolar Constraint




- Two views of the same object
- Given a point on left image, how can I find the corresponding point on right image?

# Epipolar geometry






# Epipolar Constraint



# Epipolar Constraint



$$M = K \begin{bmatrix} I & 0 \end{bmatrix}$$


$$M P = \begin{bmatrix} u \\ v \\ 1 \end{bmatrix} = p \quad \text{[Eq. 3]}$$

$$M' = K' \begin{bmatrix} R^T & -R^T T \end{bmatrix}$$

$$M' P = \begin{bmatrix} u' \\ v' \\ 1 \end{bmatrix} = p' \quad \text{[Eq. 4]}$$

# Epipolar Constraint

$$K_{canonical} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



$$M = K \begin{bmatrix} I & 0 \end{bmatrix}$$



$$M = \begin{bmatrix} I & 0 \end{bmatrix} \quad \text{[Eq. 5]}$$


$K = K'$  are known  
(canonical cameras)

$$M' = K' \begin{bmatrix} R^T & -R^T T \end{bmatrix}$$



$$M' = \begin{bmatrix} R^T & -R^T T \end{bmatrix} \quad \text{[Eq. 6]}$$

# Epipolar Constraint



$$p^T \cdot [T \times (R p')] = 0 \rightarrow p^T \cdot [T_{\times}] \cdot R p' = 0$$

[Eq. 8] [Eq. 9]


# Cross product as matrix multiplication

$$\mathbf{a} \times \mathbf{b} = \begin{bmatrix} 0 & -a_z & a_y \\ a_z & 0 & -a_x \\ -a_y & a_x & 0 \end{bmatrix} \begin{bmatrix} b_x \\ b_y \\ b_z \end{bmatrix} = [\mathbf{a}_\times] \mathbf{b}$$

$$\mathbf{a} = [a_x \ a_y \ a_z]^T$$

$$\mathbf{b} = [b_x \ b_y \ b_z]^T$$

# Epipolar Constraint



$$p^T \cdot [T \times (R p')] = 0 \rightarrow p^T \cdot [T_{\times}] \cdot R p' = 0$$

[Eq. 8] [Eq. 9]


$\mathbf{E}$  = Essential matrix

(Longuet-Higgins, 1981)

# Epipolar Constraint


$$p^T \cdot E p' = 0$$

[Eq. 10]



- $l = E p'$  is the epipolar line associated with  $p'$
- $l' = E^T p$  is the epipolar line associated with  $p$
- $E e' = 0$  and  $E^T e = 0$
- $E$  is  $3 \times 3$  matrix; 5 DOF
- $E$  is singular (rank two)

# Epipolar Constraint



$$M = K \begin{bmatrix} I & 0 \end{bmatrix}$$

$$M' = K' \begin{bmatrix} R^T & -R^T T \end{bmatrix}$$

$$p_c = K^{-1} p \quad \text{[Eq. 11]}$$

$$p'_c = K'^{-1} p' \quad \text{[Eq. 12]}$$




# Epipolar Constraint

$$p_c = K^{-1} p$$

[Eq. 11]

$$p'_c = K'^{-1} p'$$

[Eq. 12]




[Eq.9]

$$p_c^T \cdot [T_{\times}] \cdot R p'_c = 0 \rightarrow (K^{-1} p)^T \cdot [T_{\times}] \cdot R K'^{-1} p' = 0$$

$$p^T \boxed{K^{-T} \cdot [T_{\times}] \cdot R K'^{-1}} p' = 0 \rightarrow p^T \boxed{F} p' = 0 \quad \text{[Eq. 13]}$$

# Epipolar Constraint



[Eq. 13]

$$p^T F p' = 0$$

$$F = K^{-T} \cdot [T_{\times}] \cdot R K'^{-1}$$


**F = Fundamental Matrix**

(Faugeras and Luong, 1992)

[Eq. 14]



# Epipolar Constraint

$$p^T \cdot F p' = 0$$



- $l = F p'$  is the epipolar line associated with  $p'$
- $l' = F^T p$  is the epipolar line associated with  $p$
- $F e' = 0$  and  $F^T e = 0$
- $F$  is 3x3 matrix; 7 DOF
- $F$  is singular (rank two)

# Why F is useful?



- Suppose  $F$  is known
- No additional information about the scene and camera is given
- Given a point on left image, we can compute the corresponding epipolar line in the second image

# Why $F$ is useful?


- $F$  captures information about the epipolar geometry of 2 views + camera parameters
- **MORE IMPORTANTLY:**  $F$  gives constraints on how the scene changes under view point transformation (without reconstructing the scene!)
- Powerful tool in:
  - 3D reconstruction
  - Multi-view object/scene matching

# Estimating F

## The Eight-Point Algorithm

(Longuet-Higgins, 1981)



(Hartley, 1995)



$$p^T F p' = 0$$



# Estimating F



# Estimating F

**[Eq. 13]**  $\mathbf{p}^T \mathbf{F} \mathbf{p}' = 0 \quad \longrightarrow$

$$\mathbf{p} = \begin{bmatrix} u \\ v \\ 1 \end{bmatrix} \quad \mathbf{p}' = \begin{bmatrix} u' \\ v' \\ 1 \end{bmatrix}$$

$$(u, v, 1) \begin{pmatrix} F_{11} & F_{12} & F_{13} \\ F_{21} & F_{22} & F_{23} \\ F_{31} & F_{32} & F_{33} \end{pmatrix} \begin{pmatrix} u' \\ v' \\ 1 \end{pmatrix} = 0$$

$$\longrightarrow (uu', uv', u, vu', vv', v, u', v', 1) \begin{pmatrix} F_{11} \\ F_{12} \\ F_{13} \\ F_{21} \\ F_{22} \\ F_{23} \\ F_{31} \\ F_{32} \\ F_{33} \end{pmatrix} = 0$$

**[Eq. 14]**

Let's take 8 corresponding points



# Estimating F

$$\left(u_i u'_i, u_i v'_i, u_i, v_i u'_i, v_i v'_i, v_i, u'_i, v'_i, 1\right) \begin{pmatrix} F_{11} \\ F_{12} \\ F_{13} \\ F_{21} \\ F_{22} \\ F_{23} \\ F_{31} \\ F_{32} \\ F_{33} \end{pmatrix} = 0 \quad \text{[Eq. 14]}$$

# Estimating F

$$\mathbf{W} \begin{pmatrix} u_1 u'_1 & u_1 v'_1 & u_1 & v_1 u'_1 & v_1 v'_1 & v_1 & u'_1 & v'_1 & 1 \\ u_2 u'_2 & u_2 v'_2 & u_2 & v_2 u'_2 & v_2 v'_2 & v_2 & u'_2 & v'_2 & 1 \\ u_3 u'_3 & u_3 v'_3 & u_3 & v_3 u'_3 & v_3 v'_3 & v_3 & u'_3 & v'_3 & 1 \\ u_4 u'_4 & u_4 v'_4 & u_4 & v_4 u'_4 & v_4 v'_4 & v_4 & u'_4 & v'_4 & 1 \\ u_5 u'_5 & u_5 v'_5 & u_5 & v_5 u'_5 & v_5 v'_5 & v_5 & u'_5 & v'_5 & 1 \\ u_6 u'_6 & u_6 v'_6 & u_6 & v_6 u'_6 & v_6 v'_6 & v_6 & u'_6 & v'_6 & 1 \\ u_7 u'_7 & u_7 v'_7 & u_7 & v_7 u'_7 & v_7 v'_7 & v_7 & u'_7 & v'_7 & 1 \\ u_8 u'_8 & u_8 v'_8 & u_8 & v_8 u'_8 & v_8 v'_8 & v_8 & u'_8 & v'_8 & 1 \end{pmatrix} \begin{pmatrix} F_{11} \\ F_{12} \\ F_{13} \\ F_{21} \\ F_{22} \\ F_{23} \\ F_{31} \\ F_{32} \\ F_{33} \end{pmatrix} = 0 \quad [\text{Eqs. 15}] \quad \mathbf{f}$$

- Homogeneous system  $\mathbf{W} \mathbf{f} = 0$
- Rank 8  $\longrightarrow$  A non-zero solution exists (unique)
- If  $N > 8 \longrightarrow$  Lsq. solution by SVD!  $\longrightarrow \hat{\mathbf{F}}$   
 $\|\mathbf{f}\| = 1$

$\hat{F}$  satisfies:  $p^T \hat{F} p' = 0$

and estimated  $\hat{F}$  may have full rank ( $\det(\hat{F}) \neq 0$ )

**But remember:** fundamental matrix is Rank2

Find  $F$  that minimizes  $\|F - \hat{F}\| = 0$   
Frobenius norm (\*)

Subject to  $\det(F)=0$

SVD (again!) can be used to solve this problem

(\*) Sq. root of the sum of squares of all entries




Find  $F$  that minimizes  $\|F - \hat{F}\| = 0$   
 Frobenius norm (\*)

Subject to  $\det(F)=0$



$$F = U \begin{bmatrix} s_1 & 0 & 0 \\ 0 & s_2 & 0 \\ 0 & 0 & 0 \end{bmatrix} V^T$$

Where:

$$U \begin{bmatrix} s_1 & 0 & 0 \\ 0 & s_2 & 0 \\ 0 & 0 & s_3 \end{bmatrix} V^T = SVD(\hat{F})$$




Data courtesy of R. Mohr and B. Boufama.



Mean errors:  
10.0pixel  
9.1pixel

# Problems with the 8-Point Algorithm



$$\begin{array}{l} \mathbf{W} \mathbf{f} = 0, \\ \|\mathbf{f}\| = 1 \end{array} \xrightarrow{\text{Lsq solution by SVD}} \mathbf{F}$$

- Recall the structure of  $\mathbf{W}$ :
  - do we see any potential (numerical) issue?

# Problems with the 8-Point Algorithm

$$\mathbf{W}\mathbf{f} = 0$$

$$\begin{pmatrix} u_1 u'_1 & u_1 v'_1 & u_1 & v_1 u'_1 & v_1 v'_1 & v_1 & u'_1 & v'_1 & 1 \\ u_2 u'_2 & u_2 v'_2 & u_2 & v_2 u'_2 & v_2 v'_2 & v_2 & u'_2 & v'_2 & 1 \\ u_3 u'_3 & u_3 v'_3 & u_3 & v_3 u'_3 & v_3 v'_3 & v_3 & u'_3 & v'_3 & 1 \\ u_4 u'_4 & u_4 v'_4 & u_4 & v_4 u'_4 & v_4 v'_4 & v_4 & u'_4 & v'_4 & 1 \\ u_5 u'_5 & u_5 v'_5 & u_5 & v_5 u'_5 & v_5 v'_5 & v_5 & u'_5 & v'_5 & 1 \\ u_6 u'_6 & u_6 v'_6 & u_6 & v_6 u'_6 & v_6 v'_6 & v_6 & u'_6 & v'_6 & 1 \\ u_7 u'_7 & u_7 v'_7 & u_7 & v_7 u'_7 & v_7 v'_7 & v_7 & u'_7 & v'_7 & 1 \\ u_8 u'_8 & u_8 v'_8 & u_8 & v_8 u'_8 & v_8 v'_8 & v_8 & u'_8 & v'_8 & 1 \end{pmatrix} \begin{pmatrix} F_{11} \\ F_{12} \\ F_{13} \\ F_{21} \\ F_{22} \\ F_{23} \\ F_{31} \\ F_{32} \\ F_{33} \end{pmatrix} = 0$$

- Highly un-balanced (not well conditioned)
- Values of  $\mathbf{W}$  must have similar magnitude
- This creates problems during the SVD decomposition




# Normalization

IDEA: Transform image coordinates such that the matrix  **$W$**  becomes better conditioned (**pre-conditioning**)



For each image, apply a transformation  $T$  (translation and scaling) acting on image coordinates such that:

- Origin = centroid of image points
- Mean square distance of the image points from origin is  $\sim 2$  pixels

# Example of normalization




Coordinate system of the image before applying  $T$




Coordinate system of the image after applying  $T$

- Origin = centroid of image points
- Mean square distance of the image points from origin is  $\sim 2$  pixels

# Normalization



$$q_i = T \ p_i$$



$$q'_i = T' \ p'_i$$

# The Normalized Eight-Point Algorithm

0. Compute  $T$  and  $T'$  for image 1 and 2, respectively

1. Normalize coordinates in images 1 and 2:

$$q_i = T p_i \quad q'_i = T' p'_i$$



2. Use the eight-point algorithm to compute  $\hat{F}_q$  from the corresponding points  $q_i$  and  $q'_i$ .

1. Enforce the rank-2 constraint:  $\rightarrow F_q$  such that:

$$\begin{cases} q^T F_q q' = 0 \\ \det(F_q) = 0 \end{cases}$$



2. De-normalize  $F_q$ :  $F = T^T F_q T'$

Without normalization




Mean errors:  
10.0pixel  
9.1pixel

With normalization



Mean errors:  
1.0pixel  
0.9pixel

# The Fundamental Matrix Song






# Next lecture: Stereo systems





# Example: Parallel image planes



$K_1 = K_2 = \text{known}$

$x$  parallel to  $O_1O_2$

$E = ?$

Hint :

$R = I$

$T = (T, 0, 0)$

# Essential matrix for parallel images

$$\mathbf{E} = [\mathbf{T}_\times] \cdot \mathbf{R}$$


$$\mathbf{E} = \begin{bmatrix} 0 & -T_z & T_y \\ T_z & 0 & -T_x \\ -T_y & T_x & 0 \end{bmatrix} \mathbf{R} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -T \\ 0 & T & 0 \end{bmatrix}$$

[Eq. 20]

$$\mathbf{T} = [T \ 0 \ 0]$$

$$\mathbf{R} = \mathbf{I}$$


# Example: Parallel image planes



What are the directions of epipolar lines?

$$l = E p' = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -T \\ 0 & T & 0 \end{bmatrix} \begin{bmatrix} u' \\ v' \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ -T \\ T v' \end{bmatrix} \quad \text{horizontal!}$$


# Example: Parallel image planes



How are  $p$   
and  $p'$   
related?

$$p^T \cdot E p' = 0$$

# Example: Parallel image planes




How are  $p$   
and  $p'$   
related?

$$\Rightarrow \begin{pmatrix} u & v & 1 \end{pmatrix} \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -T \\ 0 & T & 0 \end{bmatrix} \begin{pmatrix} u' \\ v' \\ 1 \end{pmatrix} = 0 \Rightarrow \begin{pmatrix} u & v & 1 \end{pmatrix} \begin{pmatrix} 0 \\ -T \\ Tv' \end{pmatrix} = 0 \Rightarrow Tv = Tv' \Rightarrow v = v'$$



# Example: Parallel image planes




Rectification: making two images “parallel”


- Why it is useful?
- Epipolar constraint  $\rightarrow v = v'$
  - New views can be synthesized by linear interpolation


# Application: view morphing




S. M. Seitz and C. R. Dyer, *Proc. SIGGRAPH 96*, 1996, 21-30





# Rectification













From its reflection!





# Deep view morphing

D. Ji, J. Kwon, M. McFarland, S. Savarese, CVPR 2017




# Deep view morphing

D. Ji, J. Kwon, M. McFarland, S. Savarese, CVPR 2017



# Deep view morphing


D. Ji, J. Kwon, M. McFarland, S. Savarese, CVPR 2017



$I_1$

GT

$I_2$



n here.