

(AW	(ALONG WITH
(PRN)	(AS NEEDED)
(UD	(AS DIRECTED
(UD)	(AS DIRECTED)
*1	TAKE 1 TABLET ONCE DAILY
*1/4	1/4
*1C	TAKE 1 CAPSULE ONCE A DAY
*2	TAKE 1 TABLET TWICE DAILY
*2C	TAKE 1 CAPSULE TWICE A DAY
*3	TAKE 1 TABLET 3 TIMES A DAY
*3/4	3/4
*3C	TAKE 1 CAPSULE 3 TIMES A DAY
*3P	TAKE 1 TEASPOON 3 TIMES A DAY
*4	TAKE 1 TABLET 4 TIMES DAILY
*4C	TAKE 1 CAPSULE 4 TIMES A DAY
*AM	T1T AM
*HS	TAKE 1 TABLET AT BEDTIME
*PM	T1T PM
*SS	TSST QD
1.52T	TAKE 1½ TO 2 TABLETS
1.5T	TAKE 1½ TABLETS
1/21T	TAKE ½ TO 1 TABLET
1/2H	½ HOUR
1/2T	TAKE ½ TABLET
1/41T	TAKE ¼ TO 1 TABLET
1/4T	TAKE ¼ TABLET
115T	TAKE 1-1½ TABS
12C	TAKE 1 OR 2 CAPSULE(S)
12D	ONCE OR TWICE A DAY
12T	TAKE 1 OR 2 TABLET(S)
12TS	TAKE 1 TO 2 TEASPOONSFUL
1BID	TAKE TWICE A DAY FOR 1 DAY
1C	TAKE 1 CAPSULE
1D	ONCE A DAY
1GTT	INSTILL 1 DROP INTO
1H	1 HOUR
1P	APPLY ONE PATCH

1PU	1 PUF
1ST	FIRST
1T	TAKE 1 TABLET
1TB	TAKE 1 TABLESPOONFUL (15ML)
1THS	AND 1 TABLET AT BEDTIME
1TS	TAKE 1 TEASPOONFUL (5MLS)
2*2	TAKE 2 TABLETS TWICE DAILY
2*3	TAKE 2 TABLETS THREE TIMES DAILY
2*4	TAKE 2 TABLETS FOUR TIMES DAILY
2-3	2 TO 3
2.5T	TAKE 2½ TABLETS
2C	TAKE 2 CAPSULES
2GTT	INSTILL 2 DROPS INTO
2P	TAKE 2 PUFFS
2PBID	TAKE 2 PUFFS BID
2PQID	TAKE 2 PUFFS QID
2PTID	TAKE 2 PUFFS TID
2STAT	TAKE 2 AT ONCE, THEN
2T	TAKE 2 TABLETS
2TB	TAKE 2 TABLESPOONFULS (30MLS)
2TS	TAKE 2 TEASPOONSFUL (10MLS)
2XD	TWO TIMES DAILY
3-4	3 OR 4
3/4T	TAKE ¾ OF A TAB
34GTT	INSTILL 3 TO 4 DROPS INTO
3C	TAKE 3 CAPSULES
3D	THREE TIMES A DAY
3GTTS	INSTILL 3 DROPS INTO
3T	TAKE 3 TABLETS
3XD	THREE TIMES DAILY
4GTTS	INSTILL 4 DROPS INTO
4STAT	TAKE 4 AT ONCE, THEN
4T	TAKE 4 TABLETS
4XD	FOUR TIMES DAILY
5GTTS	INSTILL 5 DROPS INTO
5T	TAKE 5 TABLETS
6T	TAKE 6 TABLETS
6XD	SIX TIMES DAILY
7T	TAKE 7 TABLETS
8T	TAKE 8 TABLETS
AA	THE AFFECTED AREA(S)
AAA	APPLY TO AFFECTED AREA(S)

AAAS	APPLY TO AFFECTED AREAS
AAO	TAKE ALL AT ONCE
AB	BEFORE BREAKFAST
AC	BEFORE MEALS
ACETA	TAKE ½ TAB TWICE DAILY. START 1 DAY BEFORE ASCENT AND CONTINUE FOR 2 DAYS AT HIGHEST ALTITUDE
ACHS	BEFORE MEALS & AT BEDTIME
ACU	INSTILL GTT QID F7D, THEN INSTILL GTT TID F7D, THEN INSTILL GTT BID F7D, THEN INSTILL GTT 1D F7D, THEN STOP
AD	RIGHT EAR
AE	THE AFFECTED EAR(S)
AEY	THE AFFECTED EYE(S)
AFF	AFFECTED
AFT	IN THE AFTERNOON
AGI	AGITATION
AL	LEFT EAR
ALEN	TAKE 1 TABLET ONCE WEEKLY WITH A FULL GLASS OF WATER AT LEAST 30 MINUTES PRIOR TO FIRST FOOD/DRINK . AVOID LAYING DOWN FOR AT LEAST 30 MINUTES AND AFTER FIRST FOOD OF THE DAY
ALLER	ALLERGY
ALMO	TAKE 1 TABLET AT FIRST SIGN OF MIGRAINE. MAY REPEAT IN 2 HOURS TO A MAXIMUM OF 25MG IN 24 HOURS
AM	IN THE MORNING
AM,	IN THE MORNING,
AMHS	IN THE MORNING & AT BEDTIME
AMO	TAKE ONE CAPSULE THREE TIMES DAILY UNTIL FINISHED
AMOX	TAKE 2 CAPSULES TO START, THEN TAKE 1 CAPSULE THREE TIMES A DAY UNTIL FINISHED
ANA	ANAL AREA
ANX	ANXIETY
AP	APPLY
APL	APPLICATORFUL
APP	APPLICATORFUL
APS	APPLY SPARINGLY
AR	AREA
ART	ARTHRITIS
ASDE	AS DECONGESTANT
AST	ASTHMA
ASU	AFTER SUPPER
AU	EACH EAR
AUD	APPLY AS DIRECTED TO AFFECTED AREA
AUS	IN THE EAR
AW	ALONG WITH
AWHCT	(ALONG WITH HYDROCHLOROTHIAZIDE)

AWHCT12	(TAKE ALONG WITH ONE TABLET OF 12.5 MG HYDROCHLOROTHIAZIDE)
AWHCT25	(TAKE ALONG WITH ONE TABLET OF 25 MG HYDROCHLOROTHIAZIDE)
AWHCTZ12	(TAKE ALONG WITH ONE TABLET OF 12.5 MG HYDROCHLOROTHIAZIDE)
AWHCTZ25	(TAKE ALONG WITH ONE TABLET OF 25 MG HYDROCHLOROTHIAZIDE)
B	BREAKFAST
BAP	RELIEF FROM BACK PAIN
BAQ	GENTLY INSERT THE TIP OF THE SPRAY NOZZLE INTO ONE NOSTRIL, THEN FIRMLY PRESS THE PLUNGER TO GIVE THE DOSE. U UD FOR SEVERE HYPOGLYCEMIA
BC	TAKE 1 TABLET DAILY FOR
BC21	TAKE 1 TABLET ONCE A DAY FOR 21 DAYS, THEN STOP FOR 7 DAYS, THEN REPEAT
BELL1	INSTILL 1 DROP TO THE OPERATED EYE QID FOR THE FIRST WEEK, TH TID FOR THE SECOND WEEK, TH BID FOR THE THIRD WEEK TH STOP
BELL2	INSTILL 1 DROP TO THE OPERATED EYE QID F7D
BELL3	INSTILL 1 DROP TO THE OPERATED EYE TID F2W
BET	BETWEEN
BETM	BETWEEN MEALS
BEXSERO	FOR INTRAMUSCULAR INJECTION TO BE GIVEN AFTER 2 MONTHS OF AGE AS TWO-DOSE PRIMARY SERIES SPACED AT LEAST 2 MONTHS APART. BOOSTER IN SECOND YEAR OF LIFE >6 MONTHS FROM LAST DOSE.
BF	BEFORE
BG	U UD F BLOOD GLUCOSE MORNITORING **USE A NEW LANCET AND STRIP EACH TIME YOU TEST**
BID	TWICE A DAY
BID)	TWICE A DAY)
BID,	TWICE A DAY,
BKF	BREAKFAST
BM	BOWEL MOVEMENT
BMS	BOWEL MOVEMENTS
BP	BLOOD PRESSURE
BS	BLEEDING STOPS
BTID	2 OR 3 TIMES A DAY
BW	TWICE EACH WEEK
C	WITH
CA	WITH WATER
CAC	COUGH & CONGESTION
CAP	CAPSULE
CAP)	CAPSULE)
CAPS	CAPSULES
CAPS)	CAPSULES)
CC	WITH MEALS
CCP	(CONTINUING CARE PRESCRIPTION)
CDS	TO CLEAN DRY SKIN
CF	WITH FOOD
CG	COUGH

CH	CHEW
CHAM	TAKE A 0.5MG TABLET ONCE A DAY FOR 3 DAYS, THEN TAKE A 0.5 MG TABLET TWICE A DAY FOR 4 DAYS, THEN TAKE A 1MG TABLET TWICE A DAY.
CHAM1	TAKE ½ TABLET ONCE A DAY FOR 3 DAYS, THEN TAKE SS TABLET TWICE A DAY FOR 4 DAYS, THEN TAKE 1 TABLET TWICE A DAY
CHE	CHEST
CHLORO	STARTING 1 WEEK BEFORE TRAVEL TAKE 2 TABS WEEKLY, THEN WEEKLY WHILE AWAY AND FOR 4 WEEKS AFTER RETURNING HOME
CIPRODEX	INSTILL 4 DROPS INTO AFFECTED EAR(S) BID FOR 7 DAYS
CO	COLD
COJ	TAKE WITH ORANGE JUICE
COL	TAKE 2 TABS RIGHT AWAY TH T1T 1 HOUR LATER TH T1T
CON	CONGESTION
CONG	CONGESTION
CONS	CONSTIPATION
CONTRAVE	TAKE 1 TABLET EVERY MORNING FOR 1 WEEK, THEN INCREASE TO 1 TABLET TWICE DAILY FOR 1 WEEK, THEN TAKE 2 TABLETS IN THE MORNING AND 1 TABLET IN THE EVENING FOR 1 WEEK, THEN TAKE 2 TABLETS TWICE DAILY
COU	COUGH
CP	RELIEF FROM CHEST PAIN
CPL	CAPLET
CRA	CRAMPS
CS	CAPSULE(S)
D	DAY
DA	DAILY
DA.	DAILY.
DAU	(DISCARD REMAINDER)
DC	DISCONTINUE
DEC	DECREASE BY
DEP	DEPRESSION
DIA	DIARRHEA
DIAM	TAKE ½ TABLET BID STARTING 1 DAY BEFORE ASCENT AND CONTINUE FOR TWO DAYS AT HIGHEST ALTITUDE
DICL	TAKE 2 TABLETS AT BEDTIME MAY TAKE 1 TABLET IN THE MORNING AND 1 TABLET MID-AFTERNOON
DIS	DISSOLVE
DIW	DISSOLVED IN WATER
DIZ	DIZZINESS
DOU	DOUCHE
DOXYMAL	TAKE 1 TABLET DAILY STARTING 2 DAYS PRIOR TO ARRIVAL IN MALARIOUS REGION, DAILY WHILE THERE, AND DAILY FOR 28 DAYS UPON DEPARTURE FROM REGION
DP	DENTAL PROCEDURE
DPFL	DROPPERFUL
DPS	DROPS
DS	DAYS

DUKORAL	MIX CONTENTS OF ONE POUCH IN 5 OZ OF COLD WATER. POUR VIAL CONTENTS INTO MIXTURE AND TAKE ORALLY ON AN EMPTY STOMACH. REPEAT DOSE IN 1 WEEK. START AT LEAST 2 WEEKS BEFORE DEPARTURE
E	EACH
ECZ	ECZEMA
ELE	TAKE 1 TABLET AT FIRST SIGN OF MIGRAINE. MAY REPEAT IN 2 HOURS TO A MAXIMUM OF 80MG IN 24 HOURS
EN	EACH NOSTRIL
EOD	EVERY OTHER DAY
EOW	EVERY OTHER WEEK
ES	TAKE ON AN EMPTY STOMACH
EST	APPLY ONE PATCH TWICE WEEKLY
EVRA	APPLY 1 PATCH TO CLEAN SKIN ONCE WEEKLY FOR 3 WEEKS FOLLOWED BY 1 WEEK PATCH FREE
F	FOR
F10D	FOR 10 DAYS
F10D,	FOR 10 DAYS,
F12D	FOR 12 DAYS
F14D	FOR 14 DAYS
F14D,	FOR 14 DAYS,
F15D	FOR 15 DAYS
F16	DAILY ON DAYS 16 TO 25
F1D	FOR 1 DAY
F1M	FOR ONE MONTH
F1W	FOR 1 WEEK
F1W,	FOR 1 WEEK,
F1WT	FOR ONE WEEK, THEN
F20D	FOR 20 DAYS
F21	FOR 21 DAYS SKIP 7 THEN REPEAT
F21D	FOR 21 DAYS
F25D	FOR 25 DAYS
F25M	FOR 1ST 25 DAYS OF EACH MONTH
F28	FOR 28 DAYS, THEN REPEAT
F28D	FOR 28 DAYS
F2D	FOR 2 DAYS
F2W	FOR 2 WEEKS
F2W,	FOR 2 WEEKS,
F30D	FOR 30 DAYS
F30D,	FOR 30 DAYS,
F3D	FOR 3 DAYS
F3D,	FOR 3 DAYS,
F3W	FOR 3 WEEKS
F3W,	FOR 3 WEEKS,
F4D	FOR 4 DAYS

F4W	FOR 4 WEEKS
F4W,	FOR 4 WEEKS,
F5D	FOR 5 DAYS
F5D,	FOR 5 DAYS,
F5W	FOR 5 WEEKS
F6D	FOR 6 DAYS
F6W	FOR 6 WEEKS
F7D	FOR 7 DAYS
F7D,	FOR 7 DAYS,
F8D	FOR 8 DAYS
F8W	FOR 8 WEEKS
F9D	FOR 9 DAYS
FAC	TO FACE
FC	FOR COUGH
FC1	BLISTER PACK FILLING CHART START DATE:
FGTT	A FEW DROPS
FIBH	FOR INJECTION BY HEALTHCARE PROFESSIONAL
FIBP	FOR INJECTION BY PHYSICIAN
FIBPH	FOR INJECTION BY PHARMACIST
FII	FOR INTRAMUSCULAR INJECTION
FIHP	FOR INTRAMUSCULAR INJECTION BY A HEALTHCARE PROFESSIONAL
FILL CHART	PILL PACK FILLING CHART / START DATE:
FOSFO	ADD SACHET CONTENTS TO 125ML OF COLD WATER, STIR TO DISSOLVE, THEN DRINK IMMEDIATELY
FOU	FOR OFFICE USE
FP	AS NEEDED FOR PAIN
FROVA	TAKE 1 TABLET AT FIRST SIGN OF MIGRAINE. MAY REPEAT IN 4 HOURS TO A MAXIMUM OF 5MG IN 24 HOURS
FS	ONLY AS NEEDED FOR SLEEP
FSCI	FOR SUBCUTANEOUS INJECTION
FSIHP	FOR SUBCUTANEOUS INJECTION BY A HEALTHCARE PROFESSIONAL
FW	FOR WHEEZING
G	GIVE
G1.5TS	GIVE 1 & 1/2 TEASPOONSFUL 7.5M
G1/2T	GIVE 1/2 TABLET
G1/2TS	GIVE 1/2 TEASPOONFUL (2.5ML)
G1T	GIVE 1 TABLET
G1TS	GIVE 1 TEASPOONFUL (5.0ML)
G2/3TS	GIVE 2/3 (3.3ML) TEASPOONFUL
G2T	GIVE 2 TABLETS
G2TS	GIVE 2 TEASPOONSFUL (10.0ML)
G3/4TS	GIVE 3/4 TEASPOONFUL (3.75MLS)
GAR	GARGLE

GARD	FOR INTRAMUSCULAR INJECTION BY HEALTHCARE PROFESSIONAL AT 0, 2 AND 6 MONTHS
GLA	GLAUCOMA
GTT	1 DROP
GTTIN	DROPS INTO EAR CANAL
GTTN	DROPS INTO THE EAR CANAL
GTTQID	INS GTT IN THE OPERATIVE EYE QID F1W, THEN TID FOR ONE WEEK, THEN BID FOR ONE WEEK, THEN ONCE DAILY FOR ONE WEEK, THEN STOP
GTTS	DROPS
H	HOUR
HA	HEADACHE
HAV	FOR IM INJECTION BY HEALTHCARE PROFESSIONAL AT 0 AND 6-12 MONTHS
HCP	HEALTHCARE PROFESSIONAL
HRS	HOURS
HS	AT BEDTIME
HS)	AT BEDTIME)
IAEAR	IN AFFECTED EAR(S)
IAEYE	IN AFFECTED EYE(S)
IAN	IN AFFECTED NOSTRIL(S)
IBU600	TAKE 1 TABLET EVERY 6 HOURS FOR 3 DAYS, THEN TAKE AS NEEDED FOR PAIN THEREAFTER
IEN	IN EACH NOSTRIL
IH	INHALE
IM	INTRAMUSCULAR
IMHCP	FOR INTRAMUSCULAR INJECTION BY A HEALTHCARE PROFESSIONAL
IMHP	FOR INTRAMUSCULAR INJECTION BY HEALTH PROFESSIONAL
IMJ	FOR INTRAMUSCULAR INJECTION
IMLY	INTRAMUSCULARLY
INC	INCREASING TO
INF	INFECTION
INFL	INFLAMATION
INH	INHALE
INHS	INHALATIONS
INJ	INJECT
INJT	INJECTION
INS	INSTILL
INS1	INSTILL 1 DROP INTO
INS1OV	INSERT 1 OVULE
INS2	INSTIL 2 DROPS
INSA	INSERT 1 APPLICATORFUL
INSO	INSERT ONE OVULE
INSS	INSERT 1 SUPPOSITORY
INSUT	INSERT 1 TABLET UNDER TONGUE

INTRAN	INTRANASALLY
ITT	INTO THE
IUD	INTERUTERINE DEVICE FOR INSERTION BY PHYSICIAN - THIS ITEM IS NON-RETURNABLE
IXIARO	FOR IM INJECTION AT 0 AND 28 DAYS
JUBLIA	APPLY TO AFFECTED NAIL(S) ONCE DAILY AS DIRECTED
JUI	JUICE
LBM	LOOSE BOWEL MOVEMENT
LEPTO	TAKE 2 TABLETS ONCE WEEKLY. START 2 DAYS BEFORE ENTERING LEPTOSPIROSIS ENDEMIC AREA OR ENGAGING IN RISK ACTIVITY AND CONTINUE WEEKLY UNTIL YOU TAKE A FINAL DOSE AFTER YOUR LAST POSSIBLE EXPOSURE
LES	LESIONS
LOA	LEAVE OF ABSENCE
LOC	LOCALLY
LS	LOOSE STOOL
LTDZITH	ONCE DAILY FOR 1-3 DAYS (TAKE AT ONSET OF 3RD LOOSE STOOL PER 24 HOUR PERIOD - FOR TRAVELLER'S DIARRHEA)
LUN	LUNCH
MAL	TAKE 1 TABLET DAILY STARTING 2 DAYS PRIOR TO ARRIVAL IN MALARIA ENDEMIC AREA. TAKE DAILY WHILE EXPOSED AND CONTINUE FOR 7 DAYS UPON LEAVING
MALP	TABLETS DAILY STARTING 2 DAYS PRIOR TO DEPARTURE DAILY WHILE AWAY AND DAILY FOR 7 DAYS UPON RETURNING HOME
MAX	TO A MAXIMUM OF
MDU	TAKE AS DIRECTED
MED	FOR MEDICAL REVIEW
MEN	MENSTRUAL
MIN	MINUTES
MIN)	MINUTES)
MIST	SPRAY 1/2 OF SYRINGE CONTENTS INTO EACH NOSTRIL. IN AN UPRIGHT POSITION, PLACE TIP JUST INSIDE NOSTRIL AND DEPRESS PLUNGER AS RAPIDLY AS POSSIBLE UNTIL THE DOSE DIVIDER CLIP PREVENTS FURTHER MOVEMENT. REMOVE CLIP AND REPEAT IN OTHER NOSTIL.
ML	ML(S)
MON	MONTH
MOUN10	INJ 10 MG SUB ONCE WEEKLY
MOUN12.5	INJ 12.5 MG SUB ONCE WEEKLY
MOUN15	INJ 15 MG SUB ONCE WEEKLY
MOUN2.5	INJ 2.5 MG SUB ONCE WEEKLY
MOUN5	INJ 5 MG SUB ONCE WEEKLY
MOUN7.5	INJ 7.5 MG SUB ONCE WEEKLY
MREP	MAY BE REPEATED AS NEEDED
MS	MUSCLE SPASM(S)
MUS	MUSCLE
MWF	MONDAY, WEDNESDAY AND FRIDAY
N	NIGHT

NARA	TAKE 1 TABLET AT FIRST SIGN OF MIGRAINE. MAY REPEAT IN 4 HOURS TO A MAXIMUM OF 5MG IN 24 HOURS
NAS	NASAL
NASO	USE 1 SPRAY IN EACH NOSTRIL ONCE DAILY
NASY	NASALLY
NAU	NAUSEA
NAV	NAUSEA & VOMITING
NER	NERVES
NIC	CHEW 1 PIECE SLOWLY AS NEEDED
NITRO	USE 1 SPRAY SUBLINGUALLY WHEN NEEDED; REPEAT EVERY 5 MINUTES IF PAIN PERSISTS; MAY USE UP TO 3 SPRAYS IN A 15-MINUTE PERIOD. CALL 911, IF SYMPTOMS HAVE NOT RESOLVED 3-5 MINUTES AFTER 1 DOSE.
NN	NOON
NOCTE	AT NIGHT
NOS	NOSTRIL
NOSS	NOSTRILS
NS	DO NOT SWALLOW
NT	NOW, THEN
NU	IN EACH NOSTRIL
NUVA	INSERT ONE RING VAGINALLY AND LEAVE IN PLACE FOR 3 CONSECUTIVE WEEKS, THEN REMOVED FOR 1 WEEK.
O	OMIT
OAU	INTO THE AFFECTED EYE
OD	RIGHT EYE
OES	ON AN EMPTY STOMACH
OJ	ORANGE JUICE
OS	LEFT EYE
OTC	OVER THE COUNTER
OU	EACH EYE
OUS	IN THE EYE(S)
OZ	OUNCE
OZEM1	INJECT 1 MG SUB ONCE WEEKLY
OZEM2	INJ 0.25 MG SUB ONCE WEEKLY
OZEM5	INJ 0.5 MG SUB ONCE WEEKLY
OZEMPIC	INJECT 0.25 MG SUB ONCE WEEKLY F4W, THEN INJ 0.5 MG SUB ONCE WEEKLY
OZEMPIC,	INJECT 0.25 MG SUB ONCE WEEKLY F4W, THEN INJ 0.5 MG SUB ONCE WEEKLY,
OZS	OUNCES
P1T	PLACE 1 TABLET
PAC	PACKET
PAT	APPLY ONE PATCH
PB	AFTER BREAKFAST
PC	AFTER MEALS
PCHS	AFTER MEALS & AT BEDTIME
PD	PER DAY

PDUKORAL	MIX CONTENTS OF ONE POUCH IN 5 OZ OF COLD WATER THEN DISCARD 1/2 THE PREPARED MIXTURE. POUR VIAL CONTENTS INTO MIXTURE AND TAKE ORALLY ON AN EMPTY STOMACH. REPEAT FOR A TOTAL OF THREE DOSES SPACED 7 DAYS APART
PENLAC	APPLY TO THE AFFECTED NAILS ONCE DAILY.
PERI	RINSE WITH 15 MLS TWICE DAILY AFTER BREAKFAST AND BEFORE BEDTIME
PERIDEX	RINSE WITH 15 MLS TWICE DAILY AFTER BREAKFAST AND BEFORE BEDTIME
PF	PUFF
PFS	PUFFS
PHY	PHYSICIAN
PI	PIECE
PKG	PACKAGE
PL	PLACE
PM	IN THE EVENING
PM,	IN THE EVENING,
PN	PAIN
PO	ORALLY
POI	**POISON**
PR	PAIN RELIEF
PRE	INJECT THE CONTENTS OF ONE
PREDFORT	INSTILL 1 DROP IN THE OPERATED EYE FOUR TIMES DAILY FOR 1 WEEK THEN THREE TIMES DAILY FOR 1 WEEK THEN TWICE DAILY FOR 1 WEEK THEN ONCE DAILY F1W THEN STOP
PRES	PATIENT CONSENTED TO SERVICE AND WILL SELF-MONITOR. PATIENT TO FOLLOW-UP IF ANY CONCERNS. DOCTOR HAS BEEN NOTIFIED ABOUT PRESCRIBING VIA FAX.
PRN	AS NEEDED
PRN)	AS NEEDED)
PRN.	AS NEEDED.
PRNA	AS REQUIRED FOR ASTHMA
PRNAL	AS REQUIRED FOR ALLERGIES
PRNB	WHEN NEEDED FOR BREATHING
PRNBAP	AS REQUIRED FOR BACK PAIN
PRNBP	AS NEEDED FOR BACK PAIN
PRNC	AS REQUIRED FOR COUGH
PRNCP	AS NEEDED FOR CHEST PAIN
PRND	AS REQUIRED FOR DIARRHEA
PRNDI	AS REQUIRED FOR DIZZINESS
PRNE	AS REQUIRED FOR NERVES
PRNF	AS NEEDED FOR
PRNFE	AS NEEDED FOR FEVER
PRNFL	AS REQUIRED FOR FLUID
PRNH	AS REQUIRED FOR HEADACHE
PRNI	AS REQUIRED FOR ITCHING
PRNM	AS REQUIRED FOR MENSTRUAL PAIN
PRNMS	AS NEEDED FOR MUSCLE SPASM

PRNN	AS REQUIRED FOR NAUSEA
PRNNE	AS REQUIRED FOR NERVES
PRNP	WHEN NEEDED FOR PAIN
PRNR	AS REQUIRED FOR ARTHRITIS
PRNS	AS REQUIRED FOR SLEEP
PRNSP	WHEN NEEDED FOR SPASM
PRNW	AS REQUIRED FOR WHEEZING
PRNZ	AS REQUIRED FOR DIZZINESS
PS	BEFORE SUPPER
PTDAZITH	ONCE DAILY FOR 1-3 DAYS (TAKE AT ONSET OF 3RD LOOSE STOOL PER 24 HOUR PERIOD - FOR TRAVELLER'S DIARRHEA)
PUF	PUFF(S)
Q	EVERY
Q1-2H	EVERY 1 OR 2 HOURS
Q10H	EVERY 10 HOURS
Q12H	EVERY 12 HOURS
Q1H	EVERY HOUR
Q1W	ONCE WEEKLY AT THE SAME TIME
Q23H	EVERY 2 TO 3 HOURS
Q24H	EVERY 24 HOURS
Q2D	EVERY SECOND DAY
Q2H	EVERY 2 HOURS
Q34H	EVERY 3 TO 4 HOURS
Q3D	EVERY 3 DAYS
Q3H	EVERY 3 HOURS
Q45H	EVERY 4 TO 5 HOURS
Q46H	EVERY 4 TO 6 HOURS
Q4H	EVERY 4 HOURS
Q4H,	EVERY FOUR HOURS,
Q5D	5 TIMES DAILY
Q5H	EVERY 5 HOURS
Q6-8H	EVERY 6 TO 8 HOURS
Q68H	EVERY 6 TO 8 HOURS
Q6H	EVERY 6 HOURS
Q810H	EVERY 8 TO 10 HOURS
Q812H	EVERY 8 TO 12 HOURS
Q8H	EVERY 8 HOURS
QAM	EVERY MORNING
QAM,	EVERY MORNING,
QD	ONCE DAILY
QH	EVERY HOUR
QHS	AT BEDTIME
QID	FOUR TIMES A DAY

QID)	FOUR TIMES DAILY)
QID,	FOUR TIMES DAILY,
QO	EVERY OTHER
QOAM	EVERY OTHER MORNING
QOD	EVERY OTHER DAY
QPM	EACH EVENING
QSD	EVERY OTHER DAY
R	INTO THE RECTUM
RA	RASH
RABA	FOR IM INJECTION AT DAYS 0, 7, AND 21 OR 28
REC	RECTALLY
REF	REFRIGERATE
REL	RELIEF FROM
REN	PATIENT CONSENTED TO SERVICE, ESTABLISHED THERAPY AND WILL SELF MONITOR. PATIENT TO FOLLOWUP IF ANY CONCERNS.
RENEW	INTERVENTION CODE: ED REASON CODE: 91 IN PERSON / 92 PHONE SPECIAL SERVICES CODE 2 FEE 12 OR 20\$
REP	REPEAT
RIN1	RINSE FOR 1 MINUTE IN MOUTH
RIZA	TAKE 1 TABLET AT FIRST SIGN OF MIGRAINE. MAY REPEAT IN 2 HOURS TO A MAXIMUM OF 20MG IN 24 HOURS
RMW	RINSE MOUTH WITH
RT	RIGHT
S	WITHOUT
S&S	SWISH AND SPIT
SAS	SWISH AND SWALLOW
SAT	SATURDAY
SAXENDA	INJECT 0.6 MG SUB ONCE DAILY F7D, THEN 1.2 MG QD F7D, THEN 1.8 MG QD F7D, THEN 2.4 MG QD F7D, THEN 3 MG QD ONGOING
SC	SUBCUTANEOUS
SCA	SCALP
SD	AS A SINGLE DOSE
SEV	SEVERE
SG	SHAKE WELL AND GIVE
SHB	SHORTNESS OF BREATH
SHIN	FOR INTRAMUSCULAR INJECTION BY HEALTH PROFESSIONAL AT 0 AND 2-6 MONTHS
SHING	FOR INTRAMUSCULAR INJECTION BY HEALTH PROFESSIONAL AT 0 AND 2-6 MONTHS
SHINGRIX	FOR INTRAMUSCULAR INJECTION BY HEALTH PROFESSIONAL AT 0 AND 2-6 MONTHS
SIA	SEE INSTRUCTIONS ATTACHED
SIN	SINUS
SK	SKIN
SL	DISSOLVE 1 TABLET UNDER TONGUE
SL2	DISSOLVE 2 TABLETS UNDER THE TONGUE
SLE	SLEEP

SP	SPARINGLY
SP1	SPRAY ONCE IN TO EACH NOSTRIL
SP2	SPRAY TWICE INTO EACH NOSTRIL
SPA	SPARINGLY
SPIR	INHALE THE CONTENTS OF ONE CAPSULE ONCE DAILY VIA DEVICE
SPIRIVA	INHALE THE CONTENTS OF ONE CAPSULE ONCE DAILY
SPP	SUPPER
SPR	SPRAY
SPRS	SPRAYS
SR	SUPPER
SS	½
SS1	½ TO 1
SST	TAKE ½ TABLET
ST	AT ONCE THEN
STAN1	STARTING 1 DAY BEFORE SURGERY USE 1 DROP IN THE OPERATED EYE SIX TIMES DAILY AND CONTINUE FOR 1 WEEK AFTER SURGERY THEN STOP
STAN2	STARTING 1 DAY BEFORE SURGERY USE 1 DROP IN THE OPERATED EYE SIX TIMES DAILY FOR 1 WEEK THEN FOUR TIMES DAILY FOR 1 WEEK THE THREE TIMES DAILY FOR 1 WEEK THEN TWICE DAILY FOR 1 WEEK THEN STOP
STAT	NOW, THEN
STO	STOMACH
STT	AT ONCE, THEN
SUB	SUBCUTANEOUSLY
SUMA	TAKE 1 TABLET AT FIRST SIGN OF MIGRAINE. MAY REPEAT IN 2 HOURS TO A MAXIMUM OF 200MG IN 24 HOURS
SUN	SUNDAY
SUP	SUPPOSITORY
SW	.. SHAKE WELL ..
SWEG0.25	STEP 1: INJECT 0.25 MG SUB ONCE WEEKLY F4W
SWEG0.5	STEP 2: INJECT 0.5 MG SUB ONCE WEEKLY F4W
SWEG1	STEP 3: INJECT 1 MG SUB ONCE WEEKLY F4W
SWEG1.7	STEP 4: INJECT 1.7 MG SUB ONCE WEEKLY F4W
SWEG2.4	STEP 5: INJECT 2.4 MG SUB ONCE WEEKLY
SYM	SYMPTOMS
SYMBI	USE 2 INHALATIONS TWICE DAILY
SYS	SPRAYS
T	TAKE
T1	TAKE 1
T1.5	TAKE 1.5 TABS
T1C	TAKE 1 CAPSULE
T1P	TAKE 1 TEASPOONFUL
T1T	TAKE 1 TABLET
T1W	TAKE 1 TABLET ONCE WEEKLY
T2	TAKE 2 TABLETS UNTIL DONE

T2C	TAKE 2 CAPSULES
T2P	TAKE 2 PUFFS
T2S	TAKE 2 NOW THEN
T2T	TAKE 2 TABLETS
T3C	TAKE 3 CAPSULES
T3P	TAKE 3 TEASPOONSFUL
T3T	TAKE 3 TABLETS
T4C	TAKE 4 CAPSULES
T4T	TAKE 4 TABLETS
T5C	TAKE 5 CAPSULES
T5T	TAKE 5 TABLETS
T6	TAKE 6 TABLETS
T6C	TAKE 6 CAPSULES
T6T	TAKE 6 TABLETS
T7C	TAKE 7 CAPSULES
T7T	TAKE 7 TABLETS
T8C	TAKE 8 CAPSULES
T8T	TAKE 8 TABLETS
TA	THEREAFTER
TA)	THEREAFTER)
TA,	THEREAFTER,
TAB	TABLET
TAB)	TABLET)
TAB]	TABLET]
TABS	TABLETS
TABS)	TABLETS)
TABS]	TABLETS]
TAE	INTO THE AFFECTED EYE(S)
TB	TABLESPOONSFUL(S)
TBA	TO BE ADMINISTERED AS DIRECTED
TBLS	TABLESPOONFUL
TDAZITH	TAKE 4 TABLETS AS A SINGLE DOSE. IF SYMPTOMATIC AFTER 24 HOURS: CONTINUE WITH 2 TABLETS ONCE PER DAY FOR 2 MORE DOSES (FOR TRAVELLER'S DIARRHEA)
TDZITH	TAKE 4 TABLETS AS A SINGLE DOSE. IF SYMPTOMATIC AFTER 24 HOURS: CONTINUE WITH 2 TABS ONCE PER DAY FOR 2 MORE DOSES (FOR TRAVELLER'S DIARRHEA)
TEN	TENSION
TF	UNTIL FINISHED
TG	UNTIL MEDICATION IS GONE
TH	THEN
TID	THREE TIMES DAILY
TID)	THREE TIMES DAILY)
TID,	THREE TIMES DAILY,
TIL	UNTIL

TL	THIN LAYER
TOD	TODAY
TOL	AS TOLERATED
TOM	TOMORROW
TON	TONIGHT
TOP	AND APPLY CREAM TOPICALLY
TQD	THEN ONE DAILY
TQID	3 OR 4 TIMES A DAY
TR	THEN REPEAT
TRIPTAN	T1T PRN AT ONSET OF MIGRAINE. IF INADEQUATE RELIEF AFTER 2 HOURS, MAY REPEAT DOSE (MAX OF 2 TABLETS WITHIN A 24 HOUR PERIOD)
TS	TEASPOONFUL(S)
TSP	TEASPOONFUL
TSPS	TEASPOONSFUL
TSS	TEASPOONFUL
TSST	TAKE ½ TABLET
TT	TO THE
TTC	TAKE 1 OR 2 CAPSULE(S)
TTP	TAKE 1 OR 2 TEASPOONSFUL
TTT	TAKE 1 OR 2 TABLET(S)
TWIN	FOR INTRAMUSCULAR INJECTION BY HEALTHCARE PROFESSIONAL AT 0, 1 AND 6-12 MONTHS
TWIN1	FOR INTRAMUSCULAR INJECTION BY HEALTHCARE PROFESSIONAL AT 0, 7 AND 21 DAYS, WITH A BOOSTER AT 1 YEAR
TWIN6	OR INTRAMUSCULAR INJECTION BY HEALTHCARE PROFESSIONAL AT 0, 1 AND 6 MONTHS
U	USE
UC	UNTIL COMPLETED
UD	AS DIRECTED
UD)	AS DIRECTED)
UD,	AS DIRECTED,
UF	UNTIL FINISHED
UF)	UNTIL FINISHED)
ULC	ULCER
UNI	UNITS
UP10	UP TO 10 PIECES PER DAY
UTD	TAKE AS DIRECTED
VA	APPLICATORFUL INTO THE VAGINA
VAB	VAGINALLY AT BEDTIME
VAG	VAGINA
VAGIFEM	INSERT 1 TABLET VAGINALLY ONCE DAILY AT BEDTIME FOR 2 WEEKS, THEN DECREASE USE TO TWICE WEEKLY UD
VAGY	VAGINALLY
VAL	TAKE 4 TABLETS AS A SINGLE DOSE, THEN REPEAT IN 12 HOURS UD

VAL1000	T2T AT THE FIRST SIGN OF COLD SORE, REPEAT IN 12 HOURS FOR A TOTAL OF TWO DOSES PER OUTBREAK.
VAL500	TAKE 4 TABS AT THE FIRST SIGN OF COLD SORE, REPEAT IN 12 HOURS FOR A TOTAL OF TWO DOSES PER OUTBREAK.
VAR	TAKE ½ TAB ONCE DAILY FOR 3 DAYS, THEN TAKE ½ TAB TWICE DAILY FOR 4 DAYS, THEN TAKE ONE TAB TWICE DAILY
VIVOTIF	T1C ONCE PER DAY ON DAYS 1, 3, 5 AND 7. TAKE 1 HOUR BEFORE FOOD WITH A COOL GLASS OF WATER (START AT LEAST 2 WEEKS BEFORE TRAVEL) - REFRIGERATE CAPULES
VS	SUPPOSITORY INTO THE VAGINA
W	WITH
WA	WHEN AWAKE
WAT	WITH A GLASS OF WATER
WEF	WITH EXTRA FLUIDS
WEG0.25	INJECT 0.25 MG SUB ONCE WEEKLY
WEG0.5	INJECT 0.5 MG SUB ONCE WEEKLY
WEG1	INJECT 1 MG SUB ONCE WEEKLY
WEG1.7	INJECT 1.7 MG SUB ONCE WEEKLY
WEG2.4	INJECT 2.4 MG SUB ONCE WEEKLY
WF	WITH FOOD
WHE	WHEEZING
WK	WEEK
WKLY	WEEKLY
WKS	WEEKS
WM	WITH MEALS
WOJ	WITH ORANGE JUICE
WS	WITH SUPPER
WW	WITH WATER
WZ	WHEEZE
X	TIMES
YF	FOR SUBCUTANEOUS INJECTION
ZAKIM	INS GTT TO OPERATED EYE QID F1W (START AFTER SURGERY)
ZAKIP	INS GTT TO OPERATED EYE QID F1W, THEN TID F1W, THEN BID F1W, THEN 1D F1W, THEN STOP (START AFTER SURGERY)
ZAKIPRO	INS GTT QAM TO OPERATED EYE F3W INS GTT TO OPERATED EYE QID F1W, THEN TID F1W, THEN BID F1W, THEN 1D F1W, THEN STOP (START AFTER SURGERY)
ZITH	T 2 TABS STAT T 1 TAB QD UF
ZOLMI	TAKE 1 TABLET AT FIRST SIGN OF MIGRAINE. MAY REPEAT IN 2 HOURS TO A MAXIMUM OF 10MG IN 24 HOURS