

Sig Code Listing Report

Report Parameters
Language - English
Sort By Text

Sig Code Listing Report
Token Text

Printed on: 01-10-2026 20:07:31

LOCARABIC	استخدام 2 إلى 3 قطرات داخل الأذن مرتين يوميا لمدة 7 أيام
(UD	(AS DIRECTED
(UD)	(AS DIRECTED)
(PRN	(AS NEEDED
(DWI)	(DAILY WITNESSED INGESTION)
(F7D UD)	(FOR 7 DAYS AS DIRECTED)
RUP	(RETURN UNUSED PORTION TO PHARMACY FOR DISPOSAL)
POI	**POISON**
SW	.. SHAKE WELL ..
FLUVIRAL	0.5 ML SINGLE DOSE IM INJECTION INTO RIGHT / LEFT DELTOID AT ON
GTT	1 DROP
1H	1 HOUR
1-2	1 OR 2
SS	1/2
1/2H	1/2 HOUR
HH	1/2 HOUR
SS1	1/2 TO 1
SMOKE	1P CDS DA DNSM
2-3	2 OR 3
BTID	2 OR 3 TIMES A DAY
3-4	3 OR 4
TQID	3 OR 4 TIMES A DAY
TID	3 TIMES A DAY
Q5D	5 TIMES DAILY
FGTT	A FEW DROPS
ABD	ABDOMINAL DISCOMFORT
ABP	ABDOMINAL PAIN
ACN	ACNE
MONURO	ADD 1 SACHET TO 125 ML. (HALF A CUP) OF COLD WATER. STIR TO DISSOLVE AND DRINK IMMEDIATELY.
AFF	AFFECTED
P	AFTER
PB	AFTER BREAKFAST
PC	AFTER MEALS
PCHS	AFTER MEALS & AT BEDTIME
ASU	AFTER SUPPER

AGI	AGITATION
ALLER	ALLERGY
ANA	ANAL AREA
1THS	AND 1 TABLET AT BEDTIME
TOP	AND APPLY CREAM TOPICALLY
FLUS	ANNUAL INFLUENZA IMMUNIZATION.
FLUSHOT	ANNUAL INFLUENZA IMMUNIZATION.
AFV	ANNUAL INFLUENZA VACCINATION
ANX	ANXIETY
APL	APPLICATORFUL
APP	APPLICATORFUL
VA	APPLICATORFUL INTO THE VAGINA
AP	APPLY
1P	APPLY ONE PATCH
EST	APPLY ONE PATCH TWICE WEEKLY
PAT	APPLY PATCH
APS	APPLY SPARINGLY
AAA	APPLY TO AFFECTED AREA(S)
AR	AREA
ART	ARTHRITIS
ASDE	AS DECONGESTANT
UD	AS DIRECTED
UD)	AS DIRECTED)
PRN	AS NEEDED
PRNF	AS NEEDED FOR
PRNBP	AS NEEDED FOR BACK PAIN
PRNCP	AS NEEDED FOR CHEST PAIN
PRNFE	AS NEEDED FOR FEVER
PRNMS	AS NEEDED FOR MUSCLE SPASM
FP	AS NEEDED FOR PAIN
PRN)	AS NEEDED)
PRNAL	AS REQUIRED FOR ALLERGIES
PRNR	AS REQUIRED FOR ARTHRITIS
PRNA	AS REQUIRED FOR ASTHMA
PRNBAP	AS REQUIRED FOR BACK PAIN
PRNC	AS REQUIRED FOR COUGH
PRND	AS REQUIRED FOR DIARRHEA
PRNDI	AS REQUIRED FOR DIZZINESS
PRNZ	AS REQUIRED FOR DIZZINESS
PRNFL	AS REQUIRED FOR FLUID
PRNH	AS REQUIRED FOR HEADACHE
PRNI	AS REQUIRED FOR ITCHING

PRNM	AS REQUIRED FOR MENSTRUAL PAIN
PRNN	AS REQUIRED FOR NAUSEA
PRNNE	AS REQUIRED FOR NERVES
PRNE	AS REQUIRED FOR NERVES
PRNS	AS REQUIRED FOR SLEEP
PRNW	AS REQUIRED FOR WHEEZING
TOL	AS TOLERATED
AST	ASTHMA
HS	AT BEDTIME
QHS	AT BEDTIME
NOCTE	AT NIGHT
ST	AT ONCE THEN
STT	AT ONCE, THEN
BBH	BABY BE HEALTHY PROGRAM: TAKE 1 TABLET ONCE DAILY WITH A MEAL.
BF	BEFORE
AB	BEFORE BREAKFAST
AC	BEFORE MEALS
ACHS	BEFORE MEALS & AT BEDTIME
PS	BEFORE SUPPER
BET	BETWEEN
BETM	BETWEEN MEALS
IC	BETWEEN MEALS
BS	BLEEDING STOPS
BP	BLOOD PRESSURE
BM	BOWEL MOVEMENT
BKF	BREAKFAST
B	BREAKFAST
CPL	CAPLET
CAP	CAPSULE
CS	CAPSULE(S)
CAPS	CAPSULES
CHE	CHEST
CH	CHEW
NIC	CHEW 1 PIECE SLOWLY AS NEEDED
CO	COLD
CCDP	COLLABORATIVE CARE DEMONSTRATION PROJECT
CON	CONGESTION
CONG	CONGESTION
CONS	CONSTIPATION
CCP	CONTINUED CARE PRESCRIPTION (NO REFILLS REMAIN - CONTACT YOUR PHYSICIAN'S OFFICE)
COU	COUGH
CG	COUGH

CAC	COUGH & CONGESTION
CRA	CRAMPS
CR	CRUSH
DA	DAILY
QD	DAILY
F16	DAILY ON DAYS 16 TO 25
DWI	DAILY WITNESSED INGESTION
D	DAY
DS	DAYS
DEC	DECREASE BY
DP	DENTAL PROCEDURE
DEP	DEPRESSION
DR	DIAPER RASH
DIA	DIARRHEA
DISC	DISCOMFORT
DC	DISCONTINUE
DIS	DISSOLVE
SL	DISSOLVE 1 TABLET UNDER TONGUE
DIW	DISSOLVED IN WATER
DIZ	DIZZINESS
NS	DO NOT SWALLOW
DOU	DOUCHE
DPFL	DROPPERFUL
DPS	DROPS
GTTS	DROPS
GTTIN	DROPS INTO EAR CANAL
GTTN	DROPS INTO THE EAR CANAL
E	EACH
AU	EACH EAR
QPM	EACH EVENING
OU	EACH EYE
ECZ	ECZEMA
Q	EVERY
Q1-2H	EVERY 1 OR 2 HOURS
Q10H	EVERY 10 HOURS
Q12H	EVERY 12 HOURS
Q12H)	EVERY 12 HOURS)
Q2H	EVERY 2 HOURS
Q23H	EVERY 2 TO 3 HOURS
Q24H	EVERY 24 HOURS
Q3D	EVERY 3 DAYS
Q3H	EVERY 3 HOURS

Q34H	EVERY 3 TO 4 HOURS
Q4H	EVERY 4 HOURS
Q45H	EVERY 4 TO 5 HOURS
Q46H	EVERY 4 TO 6 HOURS
Q5H	EVERY 5 HOURS
Q6H	EVERY 6 HOURS
Q6-8H	EVERY 6 TO 8 HOURS
Q68H	EVERY 6 TO 8 HOURS
Q8H	EVERY 8 HOURS
Q810H	EVERY 8 TO 10 HOURS
Q812H	EVERY 8 TO 12 HOURS
Q4-8	EVERY FOUR TO EIGHT HOURS
4-6	EVERY FOUR TO SIX HOURS
Q1H	EVERY HOUR
QH	EVERY HOUR
QAM	EVERY MORNING
QO	EVERY OTHER
QOD	EVERY OTHER DAY
QSD	EVERY OTHER DAY
EOD	EVERY OTHER DAY
QOAM	EVERY OTHER MORNING
EOW	EVERY OTHER WEEK
Q2D	EVERY SECOND DAY
6-8	EVERY SIX TO EIGHT HOURS
EXP	EXPECTORATE
FE	FEET
1ST	FIRST
FO	FOOT
F	FOR
F1D	FOR 1 DAY
F1W	FOR 1 WEEK
F1W,	FOR 1 WEEK,
F10D	FOR 10 DAYS
F10D,	FOR 10 DAYS,
F12D	FOR 12 DAYS
F14D	FOR 14 DAYS
F15D	FOR 15 DAYS
F25M	FOR 1ST 25 DAYS OF EACH MONTH
F2D	FOR 2 DAYS
F2W	FOR 2 WEEKS
F20D	FOR 20 DAYS
F21D	FOR 21 DAYS

F21	FOR 21 DAYS SKIP 7 THEN REPEAT
F25D	FOR 25 DAYS
F28D	FOR 28 DAYS
F28	FOR 28 DAYS, THEN REPEAT
F3D	FOR 3 DAYS
F3D,	FOR 3 DAYS,
F3W	FOR 3 WEEKS
F30D	FOR 30 DAYS
F4D	FOR 4 DAYS
F4W	FOR 4 WEEKS
F4W:	FOR 4 WEEKS:
F5D	FOR 5 DAYS
F5D,	FOR 5 DAYS,
F5W	FOR 5 WEEKS
F6D	FOR 6 DAYS
F6D,	FOR 6 DAYS,
F6W	FOR 6 WEEKS
F7D	FOR 7 DAYS
F7D,	FOR 7 DAYS,
F8D	FOR 8 DAYS
F9D	FOR 9 DAYS
FC	FOR COUGH
FIHP	FOR INJECTION BY HEALTH CARE PRACTITIONER
FI	FOR INJECTION BY HEALTH CARE PRACTITIONER
SHINGRIX	FOR INJECTION BY HEALTH CARE PROFESSIONAL 0.5 ML (IM) FOR 1ST DOSE, THEN AGAIN IN 2 TO 6 MONTHS
FIBP	FOR INJECTION BY HEALTHCARE PROFESSIONAL
FOU	FOR OFFICE USE
F1WT	FOR ONE WEEK, THEN
FW	FOR WHEEZING
QID	FOUR TIMES A DAY
GAR	GARGLE
PERICHLOR	GARGLE WITH 15 MLS THEN SPIT, REPEAT TWICE DAILY AS DIRECTED.
G	GIVE
G.3	GIVE 0.3ML DAILY
G1.5TS	GIVE 1 & 1/2 TEASPOONSFUL 7.5M
G1T	GIVE 1 TABLET
G1TS	GIVE 1 TEASPOONFUL (5.0ML)
G1/2T	GIVE 1/2 TABLET
G1/2TS	GIVE 1/2 TEASPOONFUL (2.5ML)
G2T	GIVE 2 TABLETS
G2TS	GIVE 2 TEASPOONSFUL (10.0ML)
G2/3TS	GIVE 2/3 (3.3ML) TEASPOONFUL

G3/4TS	GIVE 3/4 TEASPOONFUL (3.75MLS)
GLA	GLAUCOMA
HA	HEADACHE
HIV	HIVES
H	HOUR
HRS	HOURS
HYP	HYPERTENSION
IFN	IF NECESSARY
GL	IN A GLASS OF
IEN	IN EACH NOSTRIL
NU	IN EACH NOSTRIL
EY	IN THE AFFECTED EYE
AFT	IN THE AFTERNOON
AUS	IN THE EAR
PM	IN THE EVENING
OUS	IN THE EYE(S)
AM	IN THE MORNING
AMHS	IN THE MORNING & AT BEDTIME
INC	INCREASING TO
INF	INFECTION
INFL	INFLAMATION
INH	INHALATIONS
IH	INHALE
INJ	INJECT
OZEMPIC25	INJECT 0.25 MG SUBCUTANEOUSLY ONCE WEEKLY AS DIRECTED
WEGOVY25	INJECT 0.25 MG SUBCUTANEOUSLY ONCE WEEKLY AS DIRECTED
OZEMPICST	INJECT 0.25 MG SUBCUTANEOUSLY ONCE WEEKLY FOR 4 WEEKS, THEN INJECT 0.5MG SUBCUTANEOUSLY ONCE WEEKLY AS DIRECTED
OZEMPIC50	INJECT 0.5 MG SUBCUTANEOUSLY ONCE WEEKLY AS DIRECTED
WEGOVY50	INJECT 0.5 MG SUBCUTANEOUSLY ONCE WEEKLY AS DIRECTED
WEGOVY1	INJECT 1 MG SUBCUTANEOUSLY ONCE WEEKLY AS DIRECTED
OZEMPIC1	INJECT 1 MG SUBCUTANEOUSLY ONCE WEEKLY AS DIRECTED
WEGOVY17	INJECT 1.7 MG SUBCUTANEOUSLY ONCE WEEKLY AS DIRECTED
WEGOVY24	INJECT 2.4 MG SUBCUTANEOUSLY ONCE WEEKLY AS DIRECTED
GARDASIL	INJECT IM AS DIRECTED (SECOND DOSE AFTER 2 MONTHS OF INITIAL AND FINAL INJECTION AFTER 6 MONTHS OF INITIAL DOSE)
TWINRIX	INJECT IM AS DIRECTED (SECOND INJECTION AFTER ONE MONTH OF INITIAL AND FINAL INJECTION AFTER 6 MONTHS OF INITIAL DOSE)
TWINRIX2	INJECT IM AS DIRECTED (SECOND INJECTION AFTER ONE WEEK OF INITIAL, THIRD INJECTION AFTER THREE WEEKS OF INITIAL AND LAST INJECTION AFTER 12 MONTHS OF INITIAL DOSE)
FRAGMIN	INJECT THE CONTENTS OF ONE PREFILLED SYRINGE ONCE DAILY
INJT	INJECTION
IR	INSERT

INSA	INSERT 1 APPLICATORFUL
INS1OV	INSERT 1 OVULE
IN1OV	INSERT 1 OVULE
NUVARING	INSERT 1 RING VAGINALLY AND KEEP IN PLACE FOR THREE WEEKS. REMOVE FOR ONE WEEK BREAK THEN REPEAT CYCLE.
INSS	INSERT 1 SUPPOSITORY
INSUT	INSERT 1 TABLET UNDER TONGUE
VAGIFEMNEW	INSERT 1 TABLET VAGINALLY ONCE DAILY AT BEDTIME FOR 2 WEEKS, THEN INSERT 1 TABLET VAGINALLY TWICE WEEKLY THEREAFTER
INSO	INSERT ONE OVULE
INST	INSERT ONE TAMPON
INS2	INSTIL 2 DROPS
INS	INSTILL
12GTT	INSTILL 1 OR 2 DROPS INTO
2GTT	INSTILL 2 DROPS INTO
3GTTS	INSTILL 3 DROPS INTO
34GTT	INSTILL 3 TO 4 DROPS INTO
4GTTS	INSTILL 4 DROPS INTO
5GTTS	INSTILL 5 DROPS INTO
INS1	INSTILL I DROP INTO
ITT	INTO THE
OAU	INTO THE AFFECTED EYE
R	INTO THE RECTUM
IM	INTRAMUSCULARLY
ITC	ITCHING
JUI	JUICE
K	KNEE
LOA	LEAVE OF ABSENCE
L	LEFT
AL	LEFT EAR
OS	LEFT EYE
LES	LESIONS
LID	LID
LOC	LOCALLY
LBM	LOOSE BOWEL MOVEMENT
LS	LOOSE STOOL
LUN	LUNCH
MREP	MAY BE REPEATED AS NEEDED
M	MEALS
MR	MEDICATION REVIEW
MEN	MENSTURAL
MIN	MINUTES
LAXADA	MIX 17 GM. OF POWDER IN A LIQUID AND DRINK.

RESTORA	MIX 17 GM. OF POWDER IN A LIQUID AND DRINK.
ML	ML(S)
MWF	MONDAY, WEDNESDAY AND FRIDAY
MON	MONTH
MUS	MUSCLE
MS	MUSCLE SPASM(S)
NAS	NASAL
NASY	NASALLY
NAU	NAUSEA
NAV	NAUSEA & VOMITING
NER	NERVES
N	NIGHT
NN	NOON
NOS	NOSTRIL
NOSS	NOSTRILS
NT	NOW, THEN
STAT	NOW, THEN
O	OMIT
OES	ON AN EMPTY STOMACH
OCNE	ONCE
1D	ONCE DAILY
12D	ONCE OR TWICE A DAY
Q1W	ONCE WEEKLY AT THE SAME TIME
FS	ONLY AS NEEDED FOR SLEEP
PO	ORALLY
OJ	ORANGE JUICE
OZ	OUNCE
OZS	OUNCES
PKG	PACKAGE
PAC	PACKET
PN	PAIN
PR	PAIN RELIEF
PD	PER DAY
PI	PIECE
PL	PLACE
P1T	PLACE 1 TABLET
FASTAB	PLACE ONE TABLET ON TONGUE AND ALLOW TO DISINTEGRATE UNTIL PARTICLES CAN BE SWALLOWED. DO NOT CRUSH OR CHEW THE TABLETS.
PF	PUFF
PUF	PUFF(S)
PFS	PUFFS
RA	RASH
REC	RECTALLY

REF	REFRIGERATE
REL	RELIEF FROM
BAP	RELIEF FROM BACK PAIN
CP	RELIEF FROM CHEST PAIN
REP	REPEAT
RT	RIGHT
AD	RIGHT EAR
OD	RIGHT EYE
RIN1	RINSE FOR 1 MINUTE IN MOUTH
RMW	RINSE MOUTH WITH
SAT	SATURDAY
SCA	SCALP
SIA	SEE INSTRUCTIONS ATTACHED
SEV	SEVERE
SG	SHAKE WELL AND GIVE
SHB	SHORTNESS OF BREATH
SIN	SINUS
SK	SKIN
SLE	SLEEP
SPA	SPARINGLY
SP	SPARINGLY
SP1	SPRAY ONCE IN TO EACH NOSTRIL
SP2	SPRAY TWICE INTO EACH NOSTRIL
SPR	SPRAY(S)
SPRS	SPRAYS
SYS	SPRAYS
CHAMPIX	START IN BLUE SECTION ; *1 F 3 DAYS, THEN *2 F4D TH START GREEN SECTION AND T 1 TAB BID UD
STO	STOMACH
STST	STORE TO STORE TRANSFER
SC	SUBCUTANEOUS
SUB	SUBCUTANEOUSLY
SQ	SUBCUTANEOUSLY
SUN	SUNDAY
SR	SUPPER
SPP	SUPPER
SUP	SUPPOSITORY
VS	SUPPOSITORY INTO THE VAGINA
SAS	SWISH AND SWALLOW
SYM	SYMPTONS
ZITH	T 2 TABS STAT T 1 TAB QD UF
TBLS	TABLESPOONFUL
TB	TABLESPOONSFUL(S)

TAB	TABLET
TABS	TABLETS
T	TAKE
T1	TAKE 1
1.5T	TAKE 1 AND 1/2 TABLETS
1C	TAKE 1 CAPSULE
T1C	TAKE 1 CAPSULE
*3C	TAKE 1 CAPSULE 3 TIMES A DAY
*4C	TAKE 1 CAPSULE 4 TIMES A DAY
*1C	TAKE 1 CAPSULE ONCE A DAY
*2C	TAKE 1 CAPSULE TWICE A DAY
12C	TAKE 1 OR 2 CAPSULES
TTC	TAKE 1 OR 2 CAPSULES
TTT	TAKE 1 OR 2 TABLETS
12T	TAKE 1 OR 2 TABLETS
TTP	TAKE 1 OR 2 TEASPOONFUL
1TB	TAKE 1 TABLESPOONFUL (15ML)
1T	TAKE 1 TABLET
T1T	TAKE 1 TABLET
*3	TAKE 1 TABLET 3 TIMES A DAY
*4	TAKE 1 TABLET 4 TIMES DAILY
*HS	TAKE 1 TABLET AT BEDTIME
BC	TAKE 1 TABLET DAILY FOR
IBUPROFEN	TAKE 1 TABLET EVERY 6 HOURS FOR PAIN FOR 3 DAYS, THEN TAKE 1 TABLET EVERY 6 HOURS WHEN NECESSARY FOR PAIN.
*AM	TAKE 1 TABLET EVERY MORNING
*1T	TAKE 1 TABLET ONCE A DAY
*1	TAKE 1 TABLET ONCE DAILY
BC21	TAKE 1 TABLET ONCE DAILY FOR 21 DAYS AS DIRECTED. (STOP FOR 7 DAYS IF INSTRUCTED BY PRESCRIBER THEN REPEAT)
*2	TAKE 1 TABLET TWICE DAILY
*2P	TAKE 1 TEASPOON 2 TIMES A DAY
*3P	TAKE 1 TEASPOON 3 TIMES A DAY
*4P	TAKE 1 TEASPOON 4 TIMES A DAY
*1P	TAKE 1 TEASPOON(5ML)ONCE DAILY
T1P	TAKE 1 TEASPOONFUL
1TS	TAKE 1 TEASPOONFUL (5MLS)
12TS	TAKE 1 TO 2 TEASPOONSFUL
1.5TS	TAKE 1&1/2 TEASPOONSFUL(7.5ML)
THT	TAKE 1/2 (ONE-HALF) OF A TABLET
SST	TAKE 1/2 TABLET
1/2T	TAKE 1/2 TABLET
1/2TS	TAKE 1/2 TEASPOONFUL (2.5MLS)

1/21T	TAKE 1/2 TO 1 TABLET
TQT	TAKE 1/4 (ONE-QUARTER) OF A TABLET
2STAT	TAKE 2 AT ONCE, THEN
2C	TAKE 2 CAPSULES
T2C	TAKE 2 CAPSULES
T2S	TAKE 2 NOW THEN
2TB	TAKE 2 TABLESPOONFULS (30MLS)
2T	TAKE 2 TABLETS
T2T	TAKE 2 TABLETS
2*2	TAKE 2 TABLETS 2 TIMES A DAY
2*3	TAKE 2 TABLETS 3 TIMES A DAY
2*4	TAKE 2 TABLETS 4 TIMES A DAY
T2	TAKE 2 TABLETS UNTIL DONE
2TS	TAKE 2 TEASPOONSFUL (10MLS)
3C	TAKE 3 CAPSULES
T3C	TAKE 3 CAPSULES
T3T	TAKE 3 TABLETS
3T	TAKE 3 TABLETS
T3P	TAKE 3 TEASPOONSFUL
4STAT	TAKE 4 AT ONCE, THEN
T4C	TAKE 4 CAPSULES
T4T	TAKE 4 TABLETS
4T	TAKE 4 TABLETS
T5C	TAKE 5 CAPSULES
T5T	TAKE 5 TABLETS
5T	TAKE 5 TABLETS
T6C	TAKE 6 CAPSULES
T6T	TAKE 6 TABLETS
T6	TAKE 6 TABLETS
6T	TAKE 6 TABLETS
T7C	TAKE 7 CAPSULES
T7T	TAKE 7 TABLETS
7T	TAKE 7 TABLETS
T8C	TAKE 8 CAPSULES
T8T	TAKE 8 TABLETS
8T	TAKE 8 TABLETS
AAO	TAKE ALL AT ONCE
UTD	TAKE AS DIRECTED
MDU	TAKE AS DIRECTED
PREDCAL	TAKE AS DIRECTED (PLEASE FOLLOW MEDICATION DOSING CALENDAR)
PP	TAKE AS DIRECTED WITH PILL PACK MEDICATIONS
PAXLO	TAKE DOSING REGIMEN TWICE DAILY FOR 5 DAYS (EACH DOSE = TWO 150 MG. TABLETS & ONE 100 MG. TABLET)

ES	TAKE ON AN EMPTY STOMACH
MALARONE	TAKE ONE TABLET ONCE A DAY STARTING ONE OR TWO DAYS PRIOR TO TRAVEL, CONTINUE THROUGHOUT AND FOR ANOTHER SEVEN DAYS AFTER RETURN.
1BID	TAKE TWICE A DAY FOR 1 DAY
DUKORAL	TAKE TWO DOSES ORALLY AT LEAST ONE WEEK APART. START WITH THE FIRST DOSE AT LEAST TWO WEEKS PRIOR TO TRAVEL.
COJ	TAKE WITH ORANGE JUICE
TSP	TEASPOONFUL
TSS	TEASPOONFUL
TS	TEASPOONFUL(S)
TSPS	TEASPOONSFUL
TEN	TENSION
FD	THE 1ST DOSE
AA	THE AFFECTED AREA(S)
AE	THE AFFECTED EAR(S)
AER	THE AFFECTED EAR(S)
AEY	THE AFFECTED EYE(S)
AFE	THE AFFECTED EYE(S)
TH	THEN
NEXTRX	THEN FILL NEXT PRESCRIPTION REFILL ON FILE (DIFFERENT RX#)
TQD	THEN ONE DAILY
TR	THEN REPEAT
TA	THEREAFTER
TL	THIN LAYER
3D	THREE TIMES A DAY
X	TIMES
MAX	TO A MAXIMUM OF
TBA	TO BE ADMINISTERED AS DIRECTED
TBI	TO BE INJECTED BY QUALIFIED HEALTH PROVIDER
CDS	TO CLEAN DRY SKIN
FAC	TO FACE
TT	TO THE
TOD	TODAY
TOM	TOMORROW
TON	TONIGHT
BID	TWICE A DAY
BW	TWICE EACH WEEK
ULC	ULCER
UNI	UNITS
TIL	UNTIL
UC	UNTIL COMPLETED
UF	UNTIL FINISHED
TF	UNTIL FINISHED

TG	UNTIL MEDICATION IS GONE
UP10	UP TO 10 PIECES PER DAY
U	USE
NITROSP	USE 1 SPRAY UNDER THE TONGUE FOR CHEST PAIN. AFTER 5 MINUTES, USE SECOND SPRAY IF NECESSARY TO ALLEVIATE CHEST PAIN. IF NOT RESOLVED, USE THIRD SPRAY AND PROCEED TO THE CLOSEST EMERGENCY DEPARTMENT OR CALL 9-1-1.
VAG	VAGINA
VAGY	VAGINALLY
PV	VAGINALLY
VAB	VAGINALLY AT BEDTIME
WK	WEEK
WKLY	WEEKLY
WKS	WEEKS
WZ	WHEEZE
WHE	WHEEZING
WA	WHEN AWAKE
WN	WHEN NECESSARY
PRN2	WHEN NECESSARY
PRN1	WHEN NEEDED
PRNB	WHEN NEEDED FOR BREATHING
PRNP	WHEN NEEDED FOR PAIN
PRNSP	WHEN NEEDED FOR SPASM
W	WITH
C	WITH
WAT	WITH A GLASS OF WATER
WEF	WITH EXTRA FLUIDS
CF	WITH FOOD
WF	WITH FOOD OR MILK
WM	WITH MEALS
CC	WITH MEALS
WOJ	WITH ORANGE JUICE
WS	WITH SUPPER
WW	WITH WATER
CA	WITH WATER
S	WITHOUT
WPF	WORK-PLACE FLU SHOT CLINIC
WFC	WORKPLACE FLU-SHOT CLINIC