Shaking hands COMS20010 2020, Video 3-3

John Lapinskas, University of Bristol



Counting the edges of this graph seems unpleasant...

but adding up the vertex degrees would be much easier.



Counting the edges of this graph seems unpleasant...

but adding up the vertex degrees would be much easier.

Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.



Counting the edges of this graph seems unpleasant...

but adding up the vertex degrees would be much easier.

Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Proof: All edges contain two vertices, and each vertex v is in d(v) edges.



Counting the edges of this graph seems unpleasant...

but adding up the vertex degrees would be much easier.

Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Proof: All edges contain two vertices, and each vertex v is in d(v) edges. Count the number of vertex-edge pairs: Let $X = \{(v, e) \in V \times E \colon v \in E\}$. Then |X| = 2|E| and $|X| = \sum_{v \in V} d(v)$, so we're done. \Box .




Counting the edges of this graph seems unpleasant...

but adding up the vertex degrees would be much easier.

Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Proof: All edges contain two vertices, and each vertex v is in d(v) edges. Count the number of vertex-edge pairs: Let $X = \{(v, e) \in V \times E \colon v \in E\}$. Then |X| = 2|E| and $|X| = \sum_{v \in V} d(v)$, so we're done.

Here,
$$\sum_{v} d(v) =$$




Counting the edges of this graph seems unpleasant...

but adding up the vertex degrees would be much easier.

Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Proof: All edges contain two vertices, and each vertex v is in d(v) edges. Count the number of vertex-edge pairs: Let $X = \{(v, e) \in V \times E \colon v \in E\}$. Then |X| = 2|E| and $|X| = \sum_{v \in V} d(v)$, so we're done.

Here,
$$\sum_{v} d(v) = 3 +$$




Counting the edges of this graph seems unpleasant...

but adding up the vertex degrees would be much easier.

Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Proof: All edges contain two vertices, and each vertex v is in d(v) edges. Count the number of vertex-edge pairs: Let $X = \{(v,e) \in V \times E \colon v \in E\}$. Then |X| = 2|E| and $|X| = \sum_{v \in V} d(v)$, so we're done.

Here,
$$\sum_{v} d(v) = 3 + 5 +$$




Counting the edges of this graph seems unpleasant...

but adding up the vertex degrees would be much easier.

Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Proof: All edges contain two vertices, and each vertex v is in d(v) edges. Count the number of vertex-edge pairs: Let $X = \{(v, e) \in V \times E \colon v \in E\}$. Then |X| = 2|E| and $|X| = \sum_{v \in V} d(v)$, so we're done.

Here,
$$\sum_{v} d(v) = 3 + 5 + 4 +$$




Counting the edges of this graph seems unpleasant...

but adding up the vertex degrees would be much easier.

Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Proof: All edges contain two vertices, and each vertex v is in d(v) edges. Count the number of vertex-edge pairs: Let $X = \{(v, e) \in V \times E \colon v \in E\}$. Then |X| = 2|E| and $|X| = \sum_{v \in V} d(v)$, so we're done.

Here,
$$\sum_{v} d(v) = 3 + 5 + 4 + 4 + 4$$




Counting the edges of this graph seems unpleasant...

but adding up the vertex degrees would be much easier.

Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Proof: All edges contain two vertices, and each vertex v is in d(v) edges. Count the number of vertex-edge pairs: Let $X = \{(v, e) \in V \times E \colon v \in E\}$. Then |X| = 2|E| and $|X| = \sum_{v \in V} d(v)$, so we're done.



Counting the edges of this graph seems unpleasant...

but adding up the vertex degrees would be much easier.

Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Proof: All edges contain two vertices, and each vertex v is in d(v) edges. Count the number of vertex-edge pairs: Let $X = \{(v, e) \in V \times E \colon v \in E\}$. Then |X| = 2|E| and $|X| = \sum_{v \in V} d(v)$, so we're done.

Here,
$$\sum_{v} d(v) = 3 + 5 + 4 + 4 + 5 + 4 + 4$$




Counting the edges of this graph seems unpleasant...

but adding up the vertex degrees would be much easier.

Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Proof: All edges contain two vertices, and each vertex v is in d(v) edges. Count the number of vertex-edge pairs: Let $X = \{(v,e) \in V \times E \colon v \in E\}$. Then |X| = 2|E| and $|X| = \sum_{v \in V} d(v)$, so we're done.

Here,
$$\sum_{v} d(v) = 3 + 5 + 4 + 4 + 5 + 4 + 5 + 4$$



Counting the edges of this graph seems unpleasant...

but adding up the vertex degrees would be much easier.

Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Proof: All edges contain two vertices, and each vertex v is in d(v) edges. Count the number of vertex-edge pairs: Let $X = \{(v, e) \in V \times E \colon v \in E\}$. Then |X| = 2|E| and $|X| = \sum_{v \in V} d(v)$, so we're done. \Box .

Here,
$$\sum_{v} d(v) = 3 + 5 + 4 + 4 + 5 + 4 + 5 + 6 =$$



Counting the edges of this graph seems unpleasant...

but adding up the vertex degrees would be much easier.

Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Proof: All edges contain two vertices, and each vertex v is in d(v) edges. Count the number of vertex-edge pairs: Let $X = \{(v, e) \in V \times E \colon v \in E\}$. Then |X| = 2|E| and $|X| = \sum_{v \in V} d(v)$, so we're done.

(This proof idea is called double-counting.)

Here, $\sum_{\nu} d(\nu) = 3 + 5 + 4 + 4 + 5 + 4 + 5 + 6 = 36$, so 18 edges total.

Handshake Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Question: How many edges does an *n*-vertex cycle have?



Handshake Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Question: How many edges does an *n*-vertex cycle have?




Answer: Every vertex has degree 2, so

$$\#(\text{edges}) = \frac{1}{2} \sum_{v} d(v) = \frac{1}{2} \cdot n \cdot 2 = n.$$

Handshake Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Question: How many edges does an *n*-vertex cycle have?



Answer: Every vertex has degree 2, so

$$\#(\text{edges}) = \frac{1}{2} \sum_{v} d(v) = \frac{1}{2} \cdot n \cdot 2 = n.$$

A graph is k-regular if every vertex has degree k (so cycles are 2-regular).

Question: Are there 3-regular graphs G = (V, E) with |V| odd?

Handshake Lemma: For any graph G = (V, E), $\sum_{v \in V} d(v) = 2|E|$.

Question: How many edges does an *n*-vertex cycle have?



Answer: Every vertex has degree 2, so

$$\#(\text{edges}) = \frac{1}{2} \sum_{v} d(v) = \frac{1}{2} \cdot n \cdot 2 = n.$$

A graph is k-regular if every vertex has degree k (so cycles are 2-regular).

Question: Are there 3-regular graphs G = (V, E) with |V| odd?

Answer: No, as then $\sum_{v \in V} d(v)$ would be 3|V| (which is odd). 2|E| is even, so this can't happen.

John Lapinskas Video 3-3 3 / 4

In **directed** graphs, can we express the number of edges in terms of inand out-degrees? Yes!

Directed Handshake Lemma:


For any digraph
$$G = (V, E)$$
, $\sum_{v \in V} d^+(v) = \sum_{v \in V} d^-(v) = |E|$.

In **directed** graphs, can we express the number of edges in terms of inand out-degrees? Yes!

Directed Handshake Lemma:

For any digraph
$$G = (V, E)$$
, $\sum_{v \in V} d^+(v) = \sum_{v \in V} d^-(v) = |E|$.

Proof: Terminology: we call the first vertex in a directed edge the **tail**, and the second vertex the **head**. (Matching the direction of the arrow!)



In **directed** graphs, can we express the number of edges in terms of inand out-degrees? Yes!

Directed Handshake Lemma:

For any digraph
$$G = (V, E)$$
, $\sum_{v \in V} d^+(v) = \sum_{v \in V} d^-(v) = |E|$.

Proof: Terminology: we call the first vertex in a directed edge the **tail**, and the second vertex the **head**. (Matching the direction of the arrow!)



Instead of counting vertex-edge pairs, we count tail-edge pairs.

In **directed** graphs, can we express the number of edges in terms of inand out-degrees? Yes!

Directed Handshake Lemma:

For any digraph
$$G = (V, E)$$
, $\sum_{v \in V} d^+(v) = \sum_{v \in V} d^-(v) = |E|$.

Proof: Terminology: we call the first vertex in a directed edge the **tail**, and the second vertex the **head**. (Matching the direction of the arrow!)



Instead of counting vertex-edge pairs, we count tail-edge pairs.

So let $X = \{(v, e) \in V \times E : e = (v, w) \text{ for some } w\}.$

Each edge has one tail, so |X| = |E|.

And each vertex v is the tail of $d^+(v)$ edges, so $|X| = \sum_{v \in V} d^+(v)$.

John Lapinskas Video 3-3 4/4

In **directed** graphs, can we express the number of edges in terms of inand out-degrees? Yes!

Directed Handshake Lemma:

For any digraph
$$G = (V, E)$$
, $\sum_{v \in V} d^+(v) = \sum_{v \in V} d^-(v) = |E|$.

Proof: Terminology: we call the first vertex in a directed edge the **tail**, and the second vertex the **head**. (Matching the direction of the arrow!)



Instead of counting vertex-edge pairs, we count tail-edge pairs.

So let $X = \{(v, e) \in V \times E : e = (v, w) \text{ for some } w\}.$

Each edge has one tail, so |X| = |E|.

And each vertex v is the tail of $d^+(v)$ edges, so $|X| = \sum_{v \in V} d^+(v)$.

Similarly, counting head-edge pairs gives $\sum_{v \in V} d^-(v) = |E|$.

John Lapinskas Video 3-3 4 / 4