

Lecture Notes: Linear Second-Order Differential Equations — Non-Homogeneous

(Course by: Professor Dave on YouTube)

Thobias K. Høivik

April 23, 2025

$$a \frac{d^2 y}{dx^2} + b \frac{dy}{dx} + cy = f(x)$$

Is non-homogeneous because instead of being equal to 0, it is equal to a function of x . We start by finding the general solution to the equivalent homogeneous case.

$$a\lambda^2 + b\lambda + c = 0$$

This is known as the complementary solution. Now we must find the **particular integral** (solution compatible with the right side $f(x)$).

Suppose we want to solve

$$y'' - 5y' + 6y = e^x$$

If we pretend that the left hand side equals 0 instead, we get a homogeneous differential equation with general solution

$$y_{CF} = Ae^{2x} + Be^{3x}$$

Trial Functions

We find a corresponding trial function from the table below and proceed.

Form of $f(x)$	Trial Solution $y_p(x)$
$P_n(x)$ (polynomial of degree n)	$A_0 + A_1x + \cdots + A_nx^n$
$e^{\alpha x}$	$Ae^{\alpha x}$
$\sin(\beta x), \cos(\beta x)$	$A \cos(\beta x) + B \sin(\beta x)$
$e^{\alpha x} \cdot P_n(x)$	$(A_0 + A_1x + \cdots + A_nx^n)e^{\alpha x}$
$e^{\alpha x} \cdot \cos(\beta x)$ or $e^{\alpha x} \cdot \sin(\beta x)$	$e^{\alpha x}(A \cos(\beta x) + B \sin(\beta x))$
$x^n e^{\alpha x} \sin(px)$ or $x^n e^{\alpha x} \cos(px)$	$(C_nx^n + C_{n-1}x^{n-1} + \cdots + C_0)(C_s \sin px + C_c \cos px)e^{\alpha x}$

Table 1: Choosing Trial Functions for $y'' + ay' + by = f(x)$

We must choose a trial function from this table based on the form of $f(x)$. We have $f(x) = e^x$ so we get the trial function

$$y_{PI}(x) = Ce^x$$

The function must be linearly independent of our complementary function, which this function is. We then substitute in our trial function for y in the original differential equation to get:

$$Ce^x - 5Ce^x + 6Ce^x = e^x \Rightarrow 2Ce^x = e^x \Rightarrow C = \frac{1}{2}$$

$$y_{PI} = \frac{1}{2}e^x$$

Our general solution then becomes

$$y = y_{CF} + y_{PI}Ae^{2x} + Be^{3x} + \frac{1}{2}e^x$$

If we find that our trial function is not linearly independent of the complementary solution we multiply the constant term by x .

Variation of Parameters

Trial functions is not the only way to find the solution to the differential equation. We begin by calculating the **Wronskian**:

$$W(x) = \det \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$$

In the case of our earlier example we have

$$W(x) = \det \begin{vmatrix} e^{2x} & e^{3x} \\ 2e^{2x} & 3e^{3x} \end{vmatrix} = 3e^{5x} - 2e^{5x} = e^{5x}$$

To find the particular integral we use the formula

$$\begin{aligned} y_{PI} &= y_2 \int \frac{y_1 \cdot f(x)}{W(x)} dx - y_1 \int \frac{y_2 \cdot f(x)}{W(x)} dx \\ y_{PI} &= e^{3x} \int \frac{e^{3x} \cdot e^x}{e^{5x}} dx - e^{2x} \int \frac{e^{3x} \cdot e^x}{e^{5x}} dx \\ &= e^{3x} \left(-\frac{1}{2} e^{-2x} \right) - e^{2x} (-e^{-x}) = \frac{1}{2} e^x \end{aligned}$$

So we see that we get the same solution as before.