# Groups, Rings and Fields

Thobias Høivik

Spring 2026

# Contents


**1 Groups** **4**

1.1 Basic Examples . . . . . 4

1.2 Basic Properties . . . . . 5

1.3 Example Problem . . . . . 5

1.4 Abelian Groups . . . . . 6

**2 Important Groups** **7**

**3 Subgroups** **9**

3.1 Structure of Cyclic Subgroups . . . . . 10

**4 Cosets and Lagrange's Theorem** **11**

4.1 Problems relating to subgroups . . . . . 13

**5 Normal subgroups** **15**

**6 Exercises for Week 1** **17**

**7 Problems for week of Jan 26** **19**

**8 External- and Internal Direct Products** **23**

**9 Problems from week of 8 Feb** **25**

**10 Problems week of 16 Feb, Quotient grps, simple grps** **29**

**11 Group Actions** **32**

**12 Rings, Fields & Integral domains** **36**

**13 Some problems pertaining to Rings, Fields, etc.** **39**

**14 Fermat's and Euler's theorems, Quotient Fields** **44**

14.1Quotient Fields . . . . . 44

**15 Polynomial Rings** **52**

15.1Degree of a Polynomial . . . . . 52

15.2Units in Polynomial Ring . . . . . 53

15.3 Evaluation Homomorphisms . . . . . 53
15.4 Roots . . . . . 53

**16 Polynomial Factorisation** **53**
16.1 Useful Irreducibility Criterion . . . . . 54

**17 Problems from week of March 22** **55**

**18 Homomorphisms of rings, Prime/maximal ideals** **58**
18.1 Prime Ideals . . . . . 59
18.2 Maximal Ideals . . . . . 59

**19 Problems problems and more problems** **60**

**20 Tackling some extension field problems early** **65**

**21 Trying an earlier exam** **67**

**22 Extension fields & Vector spaces** **72**

**23 Some problems relevant to Fraleigh's sections 29-33** **75**

**24 Some problems from Section 33** **79**

**25 Automorphisms of Field Extensions & Isomorphism Extension** **82**

**26 Exam Practice** **84**

# 1 Groups

**Definition 1.1** (Group). *A group is a set $S$ together with a binary operation $\circ$ such that the following properties hold:*

1. *Identity: There exists an element $e \in S$, satisfying*

$$e \circ a = a \circ e = a$$

   *for every $a \in S$.*

2. *Inverses: For every $a \in S$ there exists $b \in S$ such that*

$$a \circ b = b \circ a = e, \text{ the identity element}$$

   *We usually denote this element as $a^{-1}$ or $-a$, depending on context.*

3. *Associativity: For any $a, b, c \in S$, we require*

$$a \circ (b \circ c) = (a \circ b) \circ c$$

*A group is then the tupple $(S, \circ)$. We will often just write the set to refer to the group, e.g. refering to the group $(\mathbb{Z}, +)$ as just $\mathbb{Z}$.*

## 1.1 Basic Examples

**Example 1.1** (Integers under addition). *The set of integers $\mathbb{Z}$ with the operation $+$ forms a group:*

- *Identity: $0$ since $0 + n = n + 0 = n$ for all $n \in \mathbb{Z}$.*
- *Inverses: For $n \in \mathbb{Z}$, the inverse is $-n$.*
- *Associativity: Addition is associative.*

*Hence $(\mathbb{Z}, +)$ is a group.*

**Example 1.2** (Non-example: Natural numbers under addition). *The set $\mathbb{N}$ under $+$ is not a group since there is no inverse for $n > 0$.*

## 1.2 Basic Properties

**Theorem 1.1** (Uniqueness of identity). *The identity element in a group is unique.*

*Proof.* Suppose $e$ and $e'$ are both identities. Then

$$e = e \circ e' = e',$$

so the identity is unique. □

**Theorem 1.2** (Uniqueness of inverses). *Each element in a group has a unique inverse.*

*Proof.* Suppose $b$ and $c$ are inverses of $a$. Then

$$b = b \circ e = b \circ (a \circ c) = (b \circ a) \circ c = e \circ c = c.$$

□

## 1.3 Example Problem

**Problem 1.1.** *Determine whether the set*

$$G = \{1, -1, i, -i\} \subset \mathbb{C}$$

*with multiplication is a group.*

*Solution.* We check the group axioms:

1. **Closure:** Multiplying any two elements of $G$ yields another element in $G$. True.
2. **Identity:** The element $1$ acts as identity. True.
3. **Inverses:** Each element has an inverse in $G$: $1^{-1} = 1$, $(-1)^{-1} = -1$, $i^{-1} = -i$, $(-i)^{-1} = i$. True.
4. **Associativity:** Multiplication of complex numbers is associative. True.

Hence $(G, \cdot)$ is a group, in fact it is an abelian group which we shall describe below in definition 1.2. □

## 1.4 Abelian Groups

**Definition 1.2** (Abelian Group)**.** *A group* $(G, \circ)$ *is* abelian *(or commutative) if*

$$a \circ b = b \circ a \quad \forall a, b \in G.$$

**Example 1.3.** *The group* $(\mathbb{Z}, +)$ *is abelian because* $m + n = n + m$*.*

## 2 Important Groups

First let's define some often used notation.

For $g \in G$, define:

- $g^n = gg \dots g (n > 0)$
- $g^0 = e$
- $g^{-n} = (g^{-1})^n$

and recognize the following identities (which are provable by induction):

$$g^m g^n = g^{m+n}, \quad (g^m)^n = g^{mn}$$

**Definition 2.1** (Roots of Unity).

$$U_n = \{e^{2\pi i k/n} : 0 \leq k < n\}$$

*are called the n-th roots of unity.*

**Definition 2.2.** *An subset $\{g_1, \dots, g_k\} \subseteq G$ is a generating set if every $g \in G$ can be expressed as*

$$g_1^{n_1} \cdots g_k^{n_k}$$

*If a singleton $g'$ alone generates a group then it is called a generator of $G$ and $G$ is said to be cyclic.*

The $n$-th roots of unity are finite, with exactly $n$ elements. They are cyclic, with generator $e^{2i\pi/n}$.

**Definition 2.3** (Symmetric Group). *The symmetric group $S_n$ is the set of all permutations of $n$ elements under composition. In other words it is the set of all $\sigma : [n] \to [n]$ with composition as its operation.*

The symmetric group is non-abelian for $n \geq 3$. The identity corresponds to the identity permutation (doing nothing) and inverses are undoing a permutation. We have the following notation for permutations in the symmetric Group (assume $n = 3$ for this example):

$$\begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}$$

corresponds to $1 \mapsto 2, 2 \mapsto 3, 3 \mapsto 1$.

Another way to write the same permutation is:

$$(1\,2\,3)$$

# 3 Subgroups

**Definition 3.1** (Subgroup). *Let $G$ be a group under $\star$. A subgroup $H \leq G$ is a subset $H \subseteq G$ that is itself a group under $\star$.*

Having to mechanically check if $H \subseteq G$ satisfies all the required axioms can be a bit tedious so next we introduce a powerful theorem which let's us easily determine whether some subset is a subgroup or not.

**Theorem 3.1** (Subgroup Test). *A nonempty subset $H \subseteq G$ is a subgroup if and only if*

$$\forall x, y \in H,\ xy^{-1} \in H.$$

*Proof Idea.* If $H$ is a subgroup the required conditions follows.

Conversely:

- Nonempty so some $h \in H$.
- $hh^{-1} = e \in H$.
- Closure under $xy^{-1}$ will give inverses and closure.

□

We get some nice corollaries from this, as well as a nice and tidy way to prove whether or not a subset is a subgroup.

**Corollary 3.1.** *The intersection of any collection of subgroups is a subgroup.*
*Given any subset $S \subseteq G$, there is a smallest subgroup containing it.*

**Definition 3.2** (Cyclic Subgroups). *Let $G$ be a group.*
*For any $g \in G$, define:*

$$\langle g \rangle = \{g^n : n \in \mathbb{Z}\}.$$

*$\langle g \rangle$ is called the cyclic subgroup generated by $g$.*

It is very straightforward to show that the cyclic subgroup generated by some $g \in G$ is indeed a subgroup with the use of theorem 3.1.

On page 7 we say that a group is cyclic if it has a signle element which generates it, but now we can simply say that a group $G$ is cyclic if $G = \langle g \rangle$ for some $g \in G$.

**Theorem 3.2.** *Suppose a group $G$ is cyclic, i.e. $G = \langle g \rangle$ for some $g \in G$.*
*Then it is isomorphic to one of the following:*

- $(\mathbb{Z}, +)$, *or*
- $(\mathbb{Z}/n\mathbb{Z}, +)$ *for some* $n \geq 1$.

## 3.1 Structure of Cyclic Subgroups

**Definition 3.3** (Order). *The order of an element $g \in G$ is*

$$\mathit{ord}(g) = |\langle g \rangle|$$

Furthermore, if $\mathrm{ord}(g) = n < \infty$, then $g^n = e$ and $n$ is minimal, otherwise we say $g$ has infinite order.

If $\mathrm{ord}(g) = n$, then:

$$\langle g \rangle = \{e, g, g^2, \dots, g^{n-1}\}$$

Moreover:

- $g^k = e \Leftrightarrow n \mid k$
- $\langle g^k \rangle = \langle g \rangle \Leftrightarrow \gcd(k, n) = 1$

**Theorem 3.3.** *Every subgroup of a cyclic group is cyclic.*
*More precisely:*

- *Subgroups of $\mathbb{Z}$ are exactly $n\mathbb{Z}$.*
- *Subgroups of $\mathbb{Z}/n\mathbb{Z}$ correspond to the divisors of $n$.*

With all of this machinery we can approach group theory with geometric and algebraic intuition. Cyclic groups are like repeated motion, finite cyclic groups corresponding to rotations by rational angles and infinite ones corresponding to translations.

# 4 Cosets and Lagrange's Theorem

**Definition 4.1** (Left and Right Cosets). *Let $H \leq G$, and let $g \in G$. The left- and right cosets are:*

$$gH = \{gh : h \in H\} \quad Hg = \{hg : h \in H\}$$

*respectively.*

A key intuition is that a coset is like a copy of $H$ shifted by $g$. We may also recognize that the cosets are the same size as $H$ by identifying $h \mapsto gh$ to be a bijection and so forth.

The next natural question is when are two cosets equal? Like with showing when a subset is a subgroup we have a little trick for this problem.

**Theorem 4.1.** *Let $G$ be a group.*
*For $g_1, g_2 \in G$, the following are equivalent:*

$$\begin{gathered} g_1 H = g_2 H \\ \Updownarrow \\ g_2^{-1} g_1 \in H \\ \Updownarrow \\ g_1 \in g_2 H \end{gathered}$$

**Theorem 4.2.** *The set of all left cosets of $H \leq G$ forms a partition of $G$.*
*That is,*

- *Every element of $G$ is in exactly on coset.*
- *Two cosets are either the same or disjoint.*

**Definition 4.2.** *Let $G$ be a group with $H \leq G$.*
*The index of $H$ in $G$, written $[G : H]$, is the number of left cosets of $H$.*

If $G$ is finite we get $|G| = [G : H] \cdot |H|$.

**Theorem 4.3** (Lagrange's Theorem). *Let $G$ be a group with $H \leq G$. Then:*

$$|H| \mid |G|$$

*Proof.* Let $G$ be a group with $H \leq G$.

Recall that by theorem 4.2 we have that the set of left cosets of $H$, $g_1H, g_2H, \dots, g_kH$ (this set is finite since $G$ is finite) are pairwise disjoint, satisfying

$$G = \bigcup_{i=1}^{k} g_iH$$

meaning

$$|G| = \sum_{i=1}^{k} |g_iH|$$

Recall also that $|H| = |g_iH|$, $\forall i \in [k]$. Thus

$$|G| = \sum_{i=1}^{k} |H| = k|H|$$

In other words $|G|$ is $|H|$ times some integer $k$, therefore $|H|$ divides $|G|$. □

Theorem 4.3 has the following immediate consequences:

- The order of any element divides $|G|$.
- $g^{|G|} = e$ for every $g \in G$.

Beware that it does not follow from Lagrange's Theorem that there exists a subgroup with the order of a divisor of $|G|$ for every divisor of $G$. The alternating group of order $12$, $A_4$, has no subgroup of order $6$ for example.

**Proposition 4.1.** *A group of order $p$ where $p$ is prime, is cyclic.*

*Proof Sketch.* Let $G$ be a group with $|G| = p$ (prime).

Let $g \in G \setminus \{1\}$. Consider it's generated subgroup $\langle g \rangle$. By Lagrange's Theorem the order of this subgroup divides $p$ so $|\langle g \rangle|$ is $1$ or $p$, but it can't be $1$ as $g \neq 1$ so it's order is $p$, i.e. $g$ generates the entirety of $G$. □

## 4.1 Problems relating to subgroups

**Problem 4.1.** *Let $G$ be a group and $g \in G$ with ord$(g) = 12$.*

- *List all distinct subgroups of $\langle g \rangle$.*
- *For which integers $k$ does $g^k$ generate $\langle g \rangle$?*

*Solution.* Recalling that $\langle g \rangle = \{g^k : k \in \mathbb{Z}\}$, we want to find all the distinct subgroups of $\langle g \rangle$. In other words, which subsets of $\langle g \rangle$ are also groups with respect to the operation of $G$?

ord$(g) = |\langle g \rangle| = 12$ hence $\langle g \rangle = \{g^0, g^1, \dots, g^{11}\}$. We know that there are two trivial subgroups of $\langle g \rangle$, namely $\{g^0\}$ and $\langle g \rangle$. Otherwise, we know that every cyclic subgroub of a cyclic group (which $\langle g \rangle$ is) is itself cyclic, so we can find the subgroups by looking at the generated groups of each element of $\langle g \rangle$. Otherwise, we could use the fact that $\langle g \rangle$ is isomorphic to $\mathbb{Z}/12\mathbb{Z}$ together with realizing that the number of subgroups is then the same as the number of positive divisors of $12$.

Using the first approach we get:

$$\begin{aligned}
\langle g^2 \rangle &= \{g^n : n = 2k,\ k \in \mathbb{Z}\} \\
&= \{e, g^2, \dots, g^10\} = \langle g^10 \rangle \\
\langle g^3 \rangle &= \{e, g^3, g^6, g^9\} = \langle g^9 \rangle \\
\langle g^4 \rangle &= \{e, g^4, g^8\} = \langle g^8 \rangle \\
\langle g^6 \rangle &= \{e, g^6\}
\end{aligned}$$

Those integers which are coprime to $12$, like $1, 5, 7, 11$, will generate $\langle g \rangle$. Thus we have $6$ distinct subgroups, exactly the same as the number of positive divisors of $12$, as expected. □

**Problem 4.2.** *Let $G$ be any group $H = \langle g \rangle$ where ord$(g) < \infty$. Prove that if $g^m \in H$, then $\langle g^m \rangle \leq H$.*

*Proof.* Assume, for contradiction, that $g^m \in H$, but $\langle g^m \rangle$ is not a subgroup of $H$.

Let $s = m \mod \text{ord}(g)$. Then $\langle g^m \rangle = \{g^{sk} : k \in \mathbb{Z}\} \neq \emptyset$. It is clear that this is a subset of $\langle g \rangle = H$, since $\langle g \rangle$ is closed under the group operation and $g^m \in H$. Then the only way our assumption is true is if there exists some $x, y \in \langle g^m \rangle$ such that $xy^{-1} \notin \langle g^m \rangle$.

Every element $x, y \in \langle g^m \rangle$ is of the form $x = g^{sn}, y = g^{st}$ for some $n, l \in \mathbb{N}$. By assumption, $xy^{-1} = g^{s(n-t)} \notin \langle g^m \rangle$ which would necessarily mean that $n - t \notin \mathbb{Z}$, impossible. □

The above argument can be made more rigorous by considering the isomorphism between $\langle g^m \rangle$ and $\mathbb{Z}/s\mathbb{Z}$, or by proving the claim directly (I am an idiot, and this was a bad approach).

# 5 Normal subgroups

**Definition 5.1** (Normal subgroup). *A subgroup $H \leq G$ is normal, written $H \trianglelefteq G$, if:*

$$gH = Hg, \forall g \in G$$

*with the equivalent characterization:*

$$H \trianglelefteq G \Leftrightarrow gHg^{-1} = H, \forall g \in G$$

A few things to tick off immediately;

1. Every subgroup of an abelian group is normal.
2. {e} and $G$ are always normal.
3. The alternating group $A_n \trianglelefteq S_n$.
4. Subgroups of index $2$ are always normal.

If you consider the map (conjugation)

$$x \mapsto gxg^{-1},$$

which constitutes an automorphism of $G$ we can say the following.

Normal subgroups are precisely those subgroups which are fixed under every conjugation.

**Definition 5.2** (Quotient Group). *If $H \trianglelefteq G$, define:*

$$G/H := \{gH : g \in G\}$$

*with an operation we'll call multiplication by:*

$$(gH)(kH) = (gk)H$$

**Theorem 5.1.** *The quotient group with multiplication, as defined above, is a well-defined group.*

**Definition 5.3.** *We define the* ***canonical projection*** *as*

$$\pi : G \to G/H,\ \pi(g) = gH.$$

Notably, the projection $\pi$ constitutes a homomorphism with kernel $\ker \pi = H$. Notice that this claim of the kernel being $H$ itself should be immediately obvious as $hH$ for some $h \in H$ is itself $H$ as it's closed.

# 6 Exercises for Week 1

**Problem 6.1** (29, p.48). *Show that if $G$ is a finite group with identity $e$ and an even number of elements, then there is $a \neq e$ in the group such that $a * a = e$.*

*Proof.* We are being asked to show that in a finite group with even order, there exists some element other than the identity which is it's own inverse.

Recall uniqueness of inverses and that for $g \in G$, $(g^{-1})^{-1} = g$. Thus for every element there exists one and only one element which is it's inverse. $e^{-1} = e$ is covered. In $G \setminus \{e\}$ there are an odd number of elements. Each of these have exactly one unique inverse meaning that there is one element "left" over which then necessarily has itself as it's inverse. □

**Problem 6.2** (33, p.49). *Let $G$ be an abelian group and let $c^n = c * c * \cdots * c$ for $n$ factors c, where $c \in G$ and $n \in \mathbb{Z}^+$. Give a proof by mathematical induction that $(a * b)^n = a^n * b^n$ for all $a, b \in G$.*

*Proof.* Let $a, b \in G$. The base case where $n = 1$ is trivial so we proceed to our assumption that for any $k \in \mathbb{Z}^+$ we have $(a * b)^k = a^k * b^k$.

Let $n = k + 1$.

$$\begin{aligned}
(a * b)^n &= (a * b)^{k+1} \\
&= (a * b)^k * (a * b) \\
&= a^k * b^k * (a * b) && \text{by hypothesis} \\
&= a^k * (b^k * a) * b && \text{by associativity} \\
&= a^k * (a * b^k) * b && \text{by commutativity} \\
&= (a^k * a) * (b^k * b) \\
&= (a^{k+1}) * (b^{k+1}) \\
&= a^n * b^n
\end{aligned}$$

By the principle of mathematical induction we have that

$$(a * b)^n = a^n * b^n, \ \forall n \in \mathbb{Z}^+$$

□

**Problem 6.3** (35, p. 49). *Show that if $(a*b)^2 = a^2 * b^2$ for $a, b$ in a group, then $a*b = b*a$.*

*Proof.*

$$\begin{aligned}(a*b)^2 &= (a*b)*(a*b)\\ a^2*b^2 &= (a*a)*(b*b)\\ \Rightarrow (a*b)*(a*b) &= (a*a)*(b*b)\\ a*(b*a)*b &= a*(a*b)*b\end{aligned}$$

By one application each of left- and right cancellation we have

$$a*b = b*a$$

□

**Problem 6.4** (41, p. 49). *Let $G$ be a group and fix $g \in G$. Show that the map $i_g$, s.t. $i_g(x) = gxg^{-1}$ for $x \in G$. Show that $i_g$ is an automorphism on $G$.*

*Proof.* Let $i_g : G \to G$ be defined as above. We need to show that it satisfies all the properties of an isomorphism.

It is clear, through left- and right cancellation, that $i_g$ is injective.

Notice also that for any $x \in G$ we have the pre-image $g^{-1}xg$ since $i_g(g^{-1}xg) = gg^{-1}xgg^{-1} = x$, hence $i_g$ is surjective.

Lastly we verify the homomorphism property:

$$\begin{aligned}i_g(xy) &= gxyg^{-1}\\ &= gxeyg^{-1}\\ &= gxg^{-1}gyg^{-1}\\ &= i_g(x)i_g(y)\end{aligned}$$

completing the proof. □

# 7 Problems for week of Jan 26

**Problem 7.1** (Problem 44 Fraleigh's p.67). *Let $G = \langle a \rangle$ be a cyclic group isomorphic to a group $G'$. If $\phi : G \to G'$ is an isomorphism, show that for every $x \in G$, $\phi(x)$ is uniquely determined by $\phi(a)$. That is, if $\phi$ and $\psi$ constitutes isomorphisms from $G \to G'$ with $\phi(a) = \psi(a)$, then $\phi(x) = \psi(x)$ for every $x \in G$.*

*Proof.* Let $G$ be generated by $a$, isomorphic to $G'$ via $\phi$ and $\psi$, satisfying

$$\phi(a) = \psi(a)$$

Let $x \in G$. Then $x = a^n$ for some $n \in \mathbb{Z}$, as $G = \langle a \rangle$. Observe the following:

$$\begin{aligned} \phi(x) &= \phi(a^n) \\ &= \phi(a)^n \\ &= \psi(a)^n \\ &= \psi(a^n) = \psi(x) \end{aligned}$$

This completes the proof. □

**Problem 7.2** (Problem 6 Fraleigh's p.83). *Compute $|\sigma|$ for*

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 3 & 1 & 4 & 5 & 6 & 2 \end{pmatrix}$$

*Solution.* Observe that $\sigma$ is the permutation which sends

$$\begin{aligned} 1 &\to 3 \\ 2 &\to 1 \\ &\vdots \\ 6 &\to 2 \end{aligned}$$

so $\sigma^2$ is

$$\begin{aligned} 1 &\to 4 \\ 2 &\to 3 \\ &\vdots \\ 6 &\to 1 \end{aligned}$$

We want to find $n \in \mathbb{N}$ such that $\sigma^n =$ the identity permutation. It's easy to see that performing the permutation $6$ times gives $1 \to 1$ which we would require. We also see that $2 \to 2$, $3 \to 3$ and so on.

We could also write the permutation in compact form and see that $\sigma$ is a cycle of length $6$, so the order is $6$. □

**Problem 7.3** (Problem 7 Fraleigh's p.83). *Compute $|\tau|$ for*

$$\tau = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 2 & 4 & 1 & 3 & 6 & 5 \end{pmatrix}$$

*Solution.* Note that $\tau$ in compact notation is

$$\tau = (1\ 2\ 4\ 3)(5\ 6)$$

so $\tau$ is the composition of a $2$- and $4$-cycle.

$$|\tau| = \texttt{lcm}(4, 2) = 4$$

□

**Problem 7.4** (Problem 49 Fraleigh's p.86). *If $A$ is a set, then a subgroup $H \leq S_A$ is transitive on $A$ if for each $a, b \in A$ there exists $\sigma \in H$ such that $\sigma(a) = b$. Show that if $A$ is a nonempty finite set, then there exists a finite cylic subgroup $H$ of $S_A$ with $|H| = |A|$ that is transitive on $A$.*

*Proof.* Let $A = \{a_1, a_2, \dots, a_n\}$ with $S_A$ being the permutations on $A$. Let $\sigma \in S_A$ be the permutation which sends $a_i \to a_{i+1}$ (take $i$ modulo $n$). It is clear that $\sigma^k$ is the permutation which sends $a_i$ to $a_{i+k}$ and furthermore $|\sigma| = n = |A|$. It is not hard to see that $\langle \sigma \rangle$ forms a subgroup where each permutation shifts all elements of $A$.

It is clear that for any $a_i, a_j \in A$ we can find a permutation which sends one to the other. Suppose, without loss of generality, $i < j$. Then $\sigma^{j-i}(a_i) = a_j$, and the inverse of said permutation sends $a_j$ to $a_i$.

A bit inelegantly written, but I digress. □

**Problem 7.5** (Problem 1 Fraleigh's p.94). *Find the orbit of*

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 5 & 1 & 3 & 6 & 2 & 4 \end{pmatrix}$$

*Solution.* We are looking at $\sigma \in S_6$ acting on $[6]$ (I assume) so we want to find

$$\{\texttt{Orb}_\sigma(n) = \{\sigma^k(x) : k \in \mathbb{Z}\} \mid n = 1, 2, \dots, 6\}$$

Let us rewrite $\sigma$ more compactly.

$$1 \to 5 \to 2 \to 1$$

$$3 \to 3$$

$$4 \to 6 \to 4$$

so

$$\sigma = (1\ 5\ 2)(4\ 6)$$

Thus we have the following orbits:

$$\begin{aligned} \mathrm{Orb}_\sigma(1) = \mathrm{Orb}_\sigma(2) = \mathrm{Orb}_\sigma(5) &= \{1, 2, 5\} \\ \mathrm{Orb}_\sigma(4) = \mathrm{Orb}_\sigma(6) &= \{4, 6\} \\ \mathrm{Orb}_\sigma(3) &= \{3\} \end{aligned}$$

□

**Problem 7.6** (Problem 7 Fraleigh's p.94). *Compute:*

$$(1\ 4\ 5)(7\ 8)(2\ 5\ 7)$$

*Solution.*

$$\begin{aligned} (1\ 4\ 5)(7\ 8)(2\ 5\ 7) &= (1\ 4\ 5) \circ ((7\ 8)(2\ 5\ 7)) \\ &= (1\ 4\ 5)(2\ 5\ 8\ 7) \\ &= (2\ 1\ 4\ 5\ 8\ 7) \end{aligned}$$

□

**Problem 7.7** (Problem 29 Fraleigh's p.96). *Show that for every subgroup $H$ of $S_n$ for $n \geq 2$, either all permutations in $H$ are even or exactly half are.*

*Proof.* Recall that an even permutation is a permutation which can be expressed as an even number of transpositions (swaps of $2$ elements).

Let $\mathrm{sgn}(\sigma) = 1$ if $\sigma$ is even, $\mathrm{sgn}(\sigma) = -1$ otherwise for all $\sigma \in S_n$. $\mathrm{sgn} : S_n \to \{1, -1\}$ is a homomorphism. Restrict this map to $H$. Then, the image:

$$\mathrm{sgn}[H] = \begin{cases} \{+1\} \\ \{+1, -1\} \end{cases}$$

In the first case it is clear that every element of $H$ must be even.

In the other case where sgn is surjective, the kernel

$$\ker(\mathrm{sgn}[H]) = H \cap A_n,$$

the set of even permutations in $H$. By the first isomorphism theorem,

$$H/(H \cap A_n) \cong \{+1, -1\}$$

so

$$[H : H \cap A_n] = 2$$

hence

$$|H \cap A_n| = \frac{1}{2}H$$

i.e. half of $H$ is even. $\square$

# 8 External- and Internal Direct Products

**Definition 8.1.** *Let* $H, K \leq G$,

$$G' = H \times K = \{(h,k) \mid h \in H,\ \ k \in K\}$$
$$G = H \oplus K = \{hk \mid h \in H,\ \ k \in K\}$$

*The internal direct product* $H \oplus K$ *presupposes that* $H, K \trianglelefteq G$, *and* $H \cap K = \{e\}$.

We really want to be able to say that $H \times K \cong H \oplus K$.

**Example 8.1.** *Consider the dihedral group* $D_3 = \{1, x, x^2, y, xy, x^y\}$. *Let* $H = \langle x \rangle$ *and* $K = \langle y \rangle$. *It's easy to see that* $H \cap K = \{e\} = \{1\}$, *and* $H \trianglelefteq D_3$. *However* $xK = \{x, xy\}$, *but* $Kx = \{x, yx\} = \{x, x^2y\}$. *Hence*

$$D_3 \neq H \oplus K$$

**Lemma 8.1.** $H, K \trianglelefteq G$ *and* $H \cap K = \{e\}$.

1. *if* $ab = a'b'$ *where* $a, a' \in H$ *and* $b, b' \in K$ *then* $a = a'$ *and* $b = b'$.
2. *if* $a \in H$ *and* $b \in K$ *then* $ab = ba$.

**Theorem 8.1.** *Assuming* $H, K$ *normal subgroups of* $G$ *with trivial intersection, then*

$$H \times K \cong H \oplus K$$

*Proof.* Define a map

$$\varphi : H \times K \to HK, \qquad \varphi(h,k) = hk.$$

This map is well-defined since $h \in H$, $k \in K$ implies $hk \in HK$.

Define a second map

$$\psi : HK \to H \times K$$

by sending $g \in HK$ to the unique pair $(h,k)$ such that $g = hk$. Such a decomposition exists since $HK = G$, and is unique because $H \cap K = \{e\}$.

We now verify that these maps are inverse to one another. For all $(h,k) \in H \times K$,

$$(\psi \circ \varphi)(h,k) = \psi(hk) = (h,k),$$

and for all $g \in HK$,

$$(\varphi \circ \psi)(g) = g.$$

Thus $\varphi$ is bijective with inverse $\psi$.

Finally, $\varphi$ is a homomorphism, since for $(h_1, k_1), (h_2, k_2) \in H \times K$,

$$\begin{aligned}\varphi((h_1, k_1)(h_2, k_2)) &= \varphi(h_1 h_2, k_1 k_2) \\ &= h_1 h_2 k_1 k_2 \\ &= h_1 k_1 h_2 k_2 \\ &= \varphi(h_1, k_1)\varphi(h_2, k_2),\end{aligned}$$

where we use that elements of $H$ and $K$ commute.

Therefore $\varphi$ is an isomorphism, and $HK \cong H \times K$. $\square$

**Theorem 8.2.** *If $|x|, |y| < \infty$ and $(x, y) \in G \times H$ then $|(x, y)| = lcm(|x|, |y|)$.*

# 9 Problems from week of 8 Feb

**Problem 9.1** (Problem 47 Fraleigh's p.113). *Let $G$ be an abelian group. Let $H \subseteq G$ consisting of the identity and all elements of $H$ with order $2$. Show that $H \leq G$.*

*Proof.* Let $H = \{e\} \cup \{g \in G \mid |g| = 2\} \subseteq G$. Let $x, y \in H$, and for the sake of simplicity assume they are nontrivial, i.e. $x \neq e \neq y$. To show that $H$ constitutes a subgroup of $G$ it suffices to show that $xy^{-1} \in H$ (we already know it is nonempty).

$$\begin{aligned}
(xy^{-1})^2 &= xy^{-1}xy^{-1} \\
&= xxy^{-1}y^{-1} && (G \text{ abelian}) \\
&= x^2(y^2)^{-1} \\
&= e(e) \\
&= e
\end{aligned}$$

Hence $|xy^{-1}| = 2$, so $xy^{-1} \in H$. Thus $H$ is a subgroup of $G$. □

**Problem 9.2** (Problem 52 Fraleigh's p.113). *Show that a finie abelian group is not cyclic if and only if it contains a subgroup isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$ for some prime $p$.*

*Proof.* Let $G$ be a finite abelian group.

We begin by proving the right-to-left implication. Assume then that there is some $H \leq G$ with $H \cong \mathbb{Z}_p \times \mathbb{Z}_p$. Suppose, for a contradiction, that $G = \langle g \rangle$ (cyclic) for some $g \in G$. Let $\phi : H \to \mathbb{Z}_p \times \mathbb{Z}_p$ be an isomorphism.

Recall that as $G$ is cyclic, generated by $g$, we have that there is some $k \in \mathbb{Z}$, for every $h \in H$, such that $h = k \cdot g$. As $G$ is finite, $k \cdot g = (|G| - k) \cdot g$, so we can safely assume $k \geq 0$ without any loss of generality.

Since $\phi$ is bijective we know that, for every $y \in \mathbb{Z}_p \times \mathbb{Z}_p$, there is some $h \in H$ such that

$$\phi(h) = y$$

Using the homomorphism property, and the fact that $h = kg,\ k \in \mathbb{N}$, we

get that:

$$\begin{aligned} y &= \phi(h) \\ &= \phi(k \cdot g) \\ &= \phi\left(\sum_{i=1}^{k} g\right) \\ &= \sum_{i=1}^{k} \phi(g) \\ &= k \cdot \phi(g) \end{aligned}$$

As $y$ was chosen arbitrarily we now see that $\mathbb{Z}_p \times \mathbb{Z}_p = \langle \phi(g) \rangle$, i.e. cyclic. Being cyclic, together with the fact that $|\mathbb{Z}_p \times \mathbb{Z}_p| = p^2$, it is the case that there must be some element with order $p^2$, but no such element exists so $\mathbb{Z}_p \times \mathbb{Z}_p$ is not cyclic. We have arrived at a contradiction so our assumption that $G$ is cyclic must be false.

Now we prove the other direction. We now assume that we have a finite abelian group $G$ which is not cyclic. As $G$ is finite we know that $G$ is finitely generated. Thus by the fundamental theorem of finitely generated abelian groups we know that $G$ is isomorphic to a direct product of cyclic groups of the form

$$\mathbb{Z}_{(p_1)^{\gamma_1}} \times \mathbb{Z}_{(p_2)^{\gamma_2}} \times \cdots \times \mathbb{Z}_{(p_n)^{\gamma_n}} \times \mathbb{Z} \times \mathbb{Z} \times \cdots \times \mathbb{Z}$$

As $G$ is finite, the direct product does not include $\mathbb{Z}$. Furthermore as $G$ is not cyclic we know that

$$G \ncong \mathbb{Z}_{p^\gamma}$$

Hence $G$ must be the direct product of at least 2 $\mathbb{Z}_{(p_i)^{\gamma_i}}$. As $\gamma_i \in \mathbb{N} \setminus \{0\}$ and distinct from $\gamma_j$ for $j \neq i$ there must be some $\mathbb{Z}_{(p_i)^{\gamma_l}}$ with $\gamma_l \geq 2$. Notice that this group necessarily has

$$H = \{0\} \times \cdots \times \{0\} \times \mathbb{Z}_{p_i} \times \{0\} \times \cdots \times \{0\} \times \mathbb{Z}_{p_i} \times \{0\} \times \cdots \times \{0\}$$

as a subgroup, which itself is isomorphic to $\mathbb{Z}_{p_i} \times \mathbb{Z}_{p_i}$. Then $\phi^{-1}[H]$ is a subgroup of $G$, which is isomorphic to $\mathbb{Z}_{p_i} \times \mathbb{Z}_{p_i}$. $\square$

**Problem 9.3** (Problem 46 Fraleigh's p.135). *Let a group $G$ be generated by $\{a_i \mid i \in I\}$, where $I$ is some indexing set and $a_i \in G$ for all $i \in I$. Let $\phi, \mu$ be two homomorphisms from $G$ into a group $G'$, such that $\phi(a_i) = \mu(a_i)$ for every $i \in I$. Prove that $\phi = \mu$.*

*Proof.* Let $G = \{a_i \mid i \in I\}$ with homomorphisms $\phi, \mu : G \to G'$ which

coincide on $a_i$ for all $i \in I$. Let $x \in G$.

$$\begin{aligned}
\phi(x) &= \phi\left(\prod_{i\in I} a_i^{k_i}\right),\ k_i \in \mathbb{Z} \\
&= \prod_{i\in I} \phi(a_i^{k_i}) \\
&= \prod_{i\in I} \underbrace{\phi(a_i)\cdots\phi(a_i)}_{k_i \text{ times}} \\
&= \prod_{i\in I} \underbrace{\mu(a_i)\cdots\mu(a_i)}_{k_i \text{ times}} \\
&= \prod_{i\in I} \mu(a_i^{k_i}) \\
&= \mu\left(\prod_{i\in I} a_i^{k_i}\right) = \mu(x)
\end{aligned}$$

$x$ was chosen arbitrarily hence $\phi$ and $\mu$ coincide on all $x \in G$. In other words, $\phi = \mu$. □

**Problem 9.4** (52 Fraleigh's p.135). *Let $\phi : G \to G'$ be a homomorphism with kernel $H$ and let $a \in G$. Prove the set equality $\{x \in G \mid \phi(x) = \phi(a)\} = Ha$.*

*Proof.* We begin by proving that $Ha \subseteq \{x \in G \mid \phi(x) = \phi(a)\}$.

Suppose $ha \in Ha$. Then $h \in H$ so

$$\phi(ha) = \underbrace{\phi(h)}_{\in H}\phi(a) = e'\phi(a) = \phi(a)$$

Thus $ha \in \{x \in G \mid \phi(x) = \phi(a)\}$. As $x$ was arbitrarily chosen, we conclude that

$$Ha \subseteq \{x \in G \mid \phi(x) = \phi(a)\}$$

Let $x \in \{x \in G \mid \phi(x) = \phi(a)\}$. Then

$$\begin{aligned}
\phi(x) &= \phi(a) \\
\phi(x)\phi(a)^{-1} &= e' \\
\phi(xa^{-1}) &= e'
\end{aligned}$$

hence $xa^{-1} \in H$ which implies $x \in Ha$. Once again $x$ was arbitrarily chosen so

$$\{x \in G \mid \phi(x) = \phi(a)\} \subseteq Ha$$

Summing up we conclude that

$$Ha = \{x \in G \mid \phi(x) = \phi(a)\}$$

□

# 10 Problems week of 16 Feb, Quotient grps, simple grps

**Problem 10.1** (Problem 3 Fraleigh's p.142). *Find the order of the given factor group:*

$$(\mathbb{Z}_4 \times \mathbb{Z}_2)/\langle(2,1)\rangle$$

*Solution.* We first compute the subgroup generated by $(2,1)$:

$$\langle(2,1)\rangle = \{(0,0),(2,1)\},$$

since $(2,1)+(2,1)=(0,0)$.

Because $\mathbb{Z}_4 \times \mathbb{Z}_2$ is abelian, every subgroup is normal, so the quotient group is well-defined.

The elements of the quotient

$$(\mathbb{Z}_4 \times \mathbb{Z}_2)/\langle(2,1)\rangle$$

are the cosets

$$(x,y)+\langle(2,1)\rangle = \{(x,y),(x+2,y+1)\}, \quad (x,y) \in \mathbb{Z}_4 \times \mathbb{Z}_2.$$

Each coset has exactly $2$ elements. Since

$$|\mathbb{Z}_4 \times \mathbb{Z}_2| = 8,$$

the number of distinct cosets is

$$\frac{8}{2} = 4.$$

Hence the quotient group has order $4$. □

**Problem 10.2** (Problem 34 Fraleigh's p.143). *Show that if a finite group $G$ has exactly one subgroup $H$ of a given order, then $H$ is a normal subgroup of $G$.*

*Proof.* Suppose $G$ is a finite group with only one subgroup $H$ of some order $k \in \mathbb{N}$. Recall that for the conjugacy of $H$ is still a subgroup, furthermore

$$|gHg^{-1}| = |H| = k,\ g \in G$$

However $H$ is the only subgroup with order $k$ so

$$gHg^{-1} = H,\ \forall g \in G$$

which implies that $H$ is normal.

□

**Problem 10.3** (Problem 35 Fraleigh's p.143). *Show that if $H$ and $N$ are subgroups of a group $G$, and $N$ is normal in $G$, then $H \cap N$ is normal in $H$.*

*Proof.* Let $G$ be a group with $H \leq G$ and $N \trianglelefteq G$. We take it for granted that $H \cap N$ is indeed a group (this is not hard to show).

Fix some arbitrary $h \in H$. We must show that $h(H \cap N)h^{-1} \subseteq H \cap N$.

Let $x \in H \cap N$. As $x, h \in H$ we ceirtanly have $hxh^{-1} \in H$. As $x \in N$, $N$ normal, we have that for any $h \in H \subseteq G$, $hxh^{-1} \in N$.

Hence

$$hxh^{-1} \in H \cap N$$

□

**Problem 10.4** (Problem 3 Fraleigh's p.151). *Classify the groups according to the fundamental theorem of finitely generated abelian groups.*

$$(\mathbb{Z}_2 \times \mathbb{Z}_4)/\langle(1,2)\rangle$$
$$(\mathbb{Z}_4 \times \mathbb{Z}_8)/\langle(1,2)\rangle$$
$$(\mathbb{Z}_4 \times \mathbb{Z}_4 \times \mathbb{Z}_8)/\langle(1,2,4)\rangle$$

*Solution.* We begin with $(\mathbb{Z}_2 \times \mathbb{Z}_4)/\langle(1,2)\rangle$.

$$\begin{aligned}
\langle(1,2)\rangle &= \{(0,0),(1,2)\} \\
(0,0)\{(0,0),(1,2)\} &= \{(0,0),(1,2)\} = A \\
(1,0)\{(0,0),(1,2)\} &= \{(1,0),(0,2)\} = B \\
(0,1)\{(0,0),(1,2)\} &= \{(0,1),(1,3)\} = C \\
(1,1)\{(0,0),(1,2)\} &= \{(1,1),(0,3)\} = D \\
(0,2)\{(0,0),(1,2)\} &= \{(1,0),(0,2)\} = B \\
(1,2)\{(0,0),(1,2)\} &= \{(1,2),(0,0)\} = A \\
(0,3)\{(0,0),(1,2)\} &= \{(0,3),(1,1)\} = D \\
(1,3)\{(0,0),(1,2)\} &= \{(1,3),(0,1)\} = C
\end{aligned}$$

We get an abelian group with $4$ distinct elements. Crucially, we can see that the coset of $(0,1)$ has order $4$ and is thus a generator. We conclude that $(\mathbb{Z}_2 \times \mathbb{Z}_4)/\langle(1,2)\rangle \cong \mathbb{Z}_4$.

Now we look at $(\mathbb{Z}_4 \times \mathbb{Z}_8)/\langle(1,2)\rangle$.

Notice that

$$(\mathbb{Z}_4 \times \mathbb{Z}_8) = \langle a, b \mid 4a = 0,\ 8b = 0,\ ab = ba\rangle$$

so quotienting with $\langle(1,2)\rangle$ only imposes the extra relation that

$$a + 2b = 0 \Leftrightarrow a = -2b$$

then $\text{ord}(1,2) = \text{lcm}(4,4) = 4$ so $\langle(1,2)\rangle$ has $4$ elements. $|\mathbb{Z}_4 \times \mathbb{Z}_8| = 32$ so the quotient group has

$$\frac{32}{4} = 8$$

elements. Up to isomorphism there exist $3$ abelian groups of order $8$. The relation imposed by $\langle(1,2)\rangle$ changes nothing so we require $8b = 0$. Clearly

$$(\mathbb{Z}_4 \times \mathbb{Z}_8)/\langle(1,2)\rangle \cong \mathbb{Z}_8$$

Lastly we look at

$$(\mathbb{Z}_4 \times \mathbb{Z}_4 \times \mathbb{Z}_8)/\langle(1,2,4)\rangle$$

Notice that

$$\mathbb{Z}_4 \times \mathbb{Z}_4 \times \mathbb{Z}_8 = \{a, b, c \mid 4a = 0,\ 4b = 0,\ 8c = 0,\ ab = ba,\ ac = ca,\ bc = ca\}$$

with

$$\langle(1,2,4)\rangle$$

imposing $a + 2b + 4c = 0$ which then means that $a = -2b - 4c$. The order of $\mathbb{Z}_4 \times \mathbb{Z}_4 \times \mathbb{Z}_8$ is $128$. The order of the subgroup is $\text{lcm}(4,2,2)$ which is $4$. Thus the order of the quotient group is

$$\frac{128}{4} = 32 = 2^5$$

There are precisely $7$ abelian groups unique up to isomorphism. $b$ has order $4$ and $c$ has $8$ so we conclude that the quotient group is isomorphic to

$$\mathbb{Z}_4 \times \mathbb{Z}_8$$

□

**Problem 10.5** (Problem 34 Fraleigh's p.153). *Let $G$ be finite, containing a nontrivial subgroup of index $2$ in $G$. Show that $G$ is not simple.*

*Proof.* There exists a subgroup $H \leq G$ with $2$ left cosets. Since $H$ is always a left coset of $H$ there must be one more left coset $gH$ for any $g \notin H$. Since the number of right cosets is the same we have right cosets $H$ and $Hg$. There are only $2$ cosets so the non-$H$ cosets must be the same

$$gH = Hg$$

Thus left and right cosets coincide. Therefore

$$H \trianglelefteq G$$

$H$ is nontrivial, and crucially $H \neq G$ (since the index is $2$). Hence $G$ has a nontrivial proper subgroup. In other words, $G$ is not simple. □

# 11 Group Actions

**Definition 11.1.** *Let $X$ be a set, and $G$ a group. A (left) action of $G$ on $X$ is a map*

$$G \times X \to X$$
$$(g, x) \mapsto gx$$

*such that $ex = x$ and $g_1(g_2x) = (g_1g_2)x$ for all $x \in X$ and $g_1, g_2 \in G$.*

The right action is similarily defined, and in the case where $G$ is abelian the actions coincide.

**Example 11.1.** *The following constitute group actions when considering matrix-vector multiplication.*

1. *$GL_2(\mathbb{R})$ and its subgroups on $\mathbb{R}^2$.*
2. *same for $GL_n(\mathbb{R})$ on $\mathbb{R}^n$.*

*A very natural group action is $D_n$ acting on the regular polygon with $n$ sides, as $D_n$ is the symmetries of the regular $n$-gon. Furthermore $S_n$ (which is often thought of as the bijections $f : [n] \to [n]$) can act on said $[n]$.*
*Another group action which we have also seen many times is conjugation where $G$ acts on itself.*

**Definition 11.2.** *Let $G$ be a group and $X$ a set.*
*We say that a group action $G \times X \to X$ is* faithful *if for any $g \in G$, $g \neq e$ implies there is some $x \in X$ such that $gx \neq x$.*
*Alternatively it can be defined to be a group action for which any $g_1, g_2 \in G$ with $g_1 \neq g_2$ implies that there exists $x \in X$ such that $g_1x \neq g_2x$.*

**Proposition 11.1.** *Consider a group action $G \times X \to X$. It defines a group homomorphism $\phi : G \to \mathrm{Sym}(X)$ for which $\ker \phi = \{e\}$ if and only if the group action is faithful.*

**Problem 11.1** (Problem 11 Fraleigh's p.160). *Let $X$ be a $G$-set. Show that $G$ acts faithfully on $X$ if and only if no two distinct elements of $G$ have the same action on each element $X$.*
*(I.e. prove the equivalent definition posed in definition 11.2)*

*Proof.* Let $G$ be a group, $X$ a $G$-set.

Assume first that the action of $G$ on $X$ is faithful. Then $e \neq g \in G$ implies the existence of some $x \in X$ for which

$$gx \neq x$$

Suppose, for a contradiction, that there exist some distinct $g_1, g_2 \in G$ which coincide on every $x \in X$. Then

$$\begin{aligned} g_1 x &= g_2 x \\ g_2^{-1} g_1 x &= x \end{aligned}$$

In other words there exists $g = g_2^{-1} g_1 \in G$ for which every $x \in X$ is a fixed-point, contradicting the assumption that $G$ acts faithfully on $X$. Thus $G$ acting faithfully on $X$ must imply that any distinct pair of elements of $G$ have some $x \in X$ for which they do not coincide.

Assume now that $G$ acts on $X$ such that for every distinct $g_1, g_2 \in G$ for which there exists $x \in X$ such that $g_1 x \neq g_2 x$. Let $g_2 = e \neq g_1$. They are clearly distinct and furthermore

$$g_1 x \neq g_2 x = x$$

This holds for any $g_1 \in G$ so $G$ acts faithfully on $X$. $\square$

**Definition 11.3** (Orbit). *Let $G$ be a group acting on a $G$-set $X$. Let $x \in X$.*
*The orbit of $x$, often denoted $\mathit{Orb}(x)$ or $Gx$ is the set*

$$\mathit{Orb}(x) := \{gx \mid g \in G\}$$

**Definition 11.4** (Stabilizer). *Let $G$ be a group acting on a $G$-set $X$. Let $x \in X$.*
*The stabilizer of $x$, often denoted $\mathit{Stab}_G(x)$ or $G_x$ is the set*

$$\{g \in G \mid gx = x\}$$

**Theorem 11.1** (Orbit-Stabilizer). *For a finite group $G$ acting on a set $X$, the following holds (conditions are equivalent).*

$$|\mathit{Orb}(x)| = [G : \mathit{Stab}_G(x)]$$
$$|G| = |\mathit{Orb}(x)| \cdot |\mathit{Stab}_G(x)|$$

**Problem 11.2.** *Let $\{X_i \mid i \in I\}$ be a disjoint collection of sets. Let each $X_i$ be a $G$-set for the same group $G$.*
*a) Show that $\bigcup_{i \in I} X_i$ can be viewed in a natural way as a $G$-set, the **union** of the $G$-sets $X_i$.*
*b) Show that every $G$-set $X$ is the union of its orbits.*

*Proof of (a).* Let $X = \bigcup_{i \in I} X_i$. Since the union is disjoint, each $x \in X$ lies in a unique $X_i$. Define

$$g \cdot x := g \cdot_{X_i} x$$

where the right-hand side denotes the given $G$-action on $X_i$.

This is well-defined because the union is disjoint. For $x \in X_i$ we have

$$e \cdot x = x$$

since $X_i$ is a $G$-set, and

$$(gh) \cdot x = (gh) \cdot_{X_i} x = g \cdot_{X_i} (h \cdot_{X_i} x) = g \cdot (h \cdot x).$$

Thus the action axioms hold, and $X$ becomes a $G$-set. □

*Proof of (b).* For $x \in X$, define the orbit

$$G \cdot x = \{g \cdot x \mid g \in G\}.$$

Since $x = e \cdot x$, every element lies in its orbit, so

$$X = \bigcup_{x \in X} G \cdot x.$$

If $G \cdot x \cap G \cdot y \neq \emptyset$, then $g \cdot x = h \cdot y$ for some $g, h \in G$. Then

$$x = g^{-1} h \cdot y,$$

so $x \in G \cdot y$, which implies $G \cdot x = G \cdot y$. Thus distinct orbits are disjoint, and $X$ is the disjoint union of its orbits. □

**Theorem 11.2** (Burnside's Formula). *Let $G$ be a finite group and $X$ a finite $G$-set. If $r$ is the number of orbits in $X$ under $G$, then*

$$r \cdot |G| = \sum_{g \in G} |X_g|$$

*where $X_g$ are the* fixed points *of $g$, namely all $x \in X$ satisfying $gx = x$.*

**Problem 11.3** (Problem 1 Fraileigh's p.164). *Find the number of orbits in $\{1, \dots, 8\}$ under the cyclic subgroup $\langle (1\ 3\ 5\ 6) \rangle \leq S_8$.*

*Solution.* Both $[8]$ and the given subgroup of $S_8$ are finite so we use Burnside's formula. Denote $\langle \sigma \rangle := \langle 1\ 3\ 5\ 6 \rangle$. We know that $|\sigma| = 4$ so we must simply calculate the sum of sizes of fixed points of each permutation in $\langle \sigma \rangle$. Obviously $|X_e| = |X| = 8$. Next up $\sigma$ has $2, 4, 7, 8$ as fixed points so $|X_\sigma| = 4$. Next up $\sigma^2 = (1\ 5)(3\ 6)$ and has the same fixed points. $\sigma^3 = (1\ 6\ 5\ 3)$ with the same fixed points once again. Thus

$$r = 8 + 4 + 4 + 4/|\sigma| = 5$$

$\square$

# 12 Rings, Fields & Integral domains

**Definition 12.1** (Ring). *A ring $R$ is a set together with two binary operations $+$ and $\cdot$, written $(R, +, \cdot)$, satisfying*

1. *$(R, +)$ is an abelian group (identity denoted $0$),*
2. *$(R, \cdot)$ has closure and associativity,*
3. *with distributive laws, i.e.*

$$a \cdot (b + c) = ab + ac, \quad (b + c) \cdot a = ba + ca.$$

*If $R$ has a multiplicative identity $1$, we say $R$ is a ring with identity, or $R$ is a ring with unity or $R$ is unital.*
*If $ab = ba$ for all $a, b \in R$, then $R$ is a commutative ring.*

So what, informally, a ring is, is a structure where you can add, multiply and subtract, but not necessarily divide.

Rings have a bunch of immediately nice properties such as

- $0 \in R$ is unique (follows from $(R, +)$ being abelian grp),
- $-a$ is unique (follows from the same logic),
- $a \cdot 0 = 0 = 0 \cdot a$ (very nice),
- $|R| \geq 2$, then $1 \neq 0$,
- $-(a + b) = (-a) + (-b)$,
- $-(-a) = a$,
- $-(ab) = (-a)b = a(-b)$,
- $ab = (-a)(-b)$.

**Definition 12.2** (Unit). *Let $(R, +, \cdot)$ be a ring with identity, and let $u \in R$. $u$ is called a unit if $u$ has a multiplicative inverse $u^{-1}$ such that*

$$u \cdot u^{-1} = u^{-1} \cdot u = 1$$

*We denote the set of units*

$$\{u \in R \mid u \text{ unit}\} =: R^x$$

*(This set is a group under the multiplication of $R$)*

**Definition 12.3** (Zero divisor). *Let $(R, +, \cdot)$ be a ring. A zero divisor is a non-zero element $a \in R$ such that there exists non-zero $b \in R$ satisfying*

$$ab = 0 \quad \text{left zero divisor} ba = 0 \quad \text{right zero divisor}$$

*Where these coincide in commutative rings.*

**Definition 12.4** (Field). *Let $(R, +, \cdot)$ be a ring.*
*$R$ is a **field** if $(R \setminus \{0\}, \cdot)$ is an abelian group.*

**Definition 12.5.** *Let $(R, +, \cdot)$ be a ring.*
*$R$ is a **division ring** if $(R \setminus \{0\}, \cdot)$ is a group.*

Crucially, in both a divison ring and a field we know that $\exists 1 \in R$, the multiplicative identity, meaning a field/division ring is unital.

The prototypical example of a commutative unital ring is $(\mathbb{Z}, +, \cdot)$. Importantly, $\mathbb{Z}$ is not a field. So a big area of focus will be to look at larger more powerful structures.

**Definition 12.6** (Integral Domain). *Let $(R, +, \cdot)$ be a ring.*
*$R$ is an **integral domain** if it is a commutative, unital ring.*

**Proposition 12.1.** *A ring $R$ is an integral domain if and only if $R$ has no zero-divisors and $|R| \geq 2$.*

The nice property we can gleam from integral domains is that $a \cdot b = 0$ if and only if either $a$ or $b$ is zero.

Usually, when we say integral domain, or field, we will be refering to said structure when it has $2$ or more elements. The reason for this is that a ring or field with $1$ element is not worth dwelling on.

**Proposition 12.2.** *A field is an integral domain.*

*Proof sketch.* If $ab = 0$ and $a \neq 0$, then $b = a^{-1}(ab) = a^{-1}0 = 0$. □

**Theorem 12.1.** *Every finite integral domain is a field.*

*Proof.* Let $D$ be a finite integral domain.

$(D \setminus \{0\}, \cdot)$ is closed, because $D$ has no zero divisors. Furthermore it

is associative and $1 \in D \setminus \{0\}$ by definition. The last thing we need is inverses.

Let $r \in D$, $r \neq 0$. Consider $r, r^2, r^3, \dots r^k, \dots$

As $D$ is finite there must be repeted powers in this list. Let $r^i = r^j$, $i < j$ wlog. Then

$$\underbrace{r \cdot r \cdots r}_{i} \cdot 1 = \underbrace{r \cdot r \cdots r}_{j}$$

Use left cancellation repeatedly to get

$$r^{j-i} = 1$$

Let $s = r^{j-i-1}$. Then $rs = sr = 1$, so $s = r^{-1}$. $\square$

**Theorem 12.2.** *Every finite division ring is a field.*

# 13 Some problems pertaining to Rings, Fields, etc.

**Problem 13.1** (Problem 3 & 5 Fraleigh's p.174). *Compute the products*

$$(11)(-4) \in \mathbb{Z}_{15}$$

$$(2,3)(3,5) \in \mathbb{Z}_5 \times \mathbb{Z}_9$$

*Solution.*

$$\begin{aligned} 11 \cdot -4 &= -(11 \cdot 4) \\ &= -(44) \\ &\equiv 1 \mod 15 \end{aligned}$$

$$\begin{aligned} (2,3)(3,5) &= (2 \cdot 3, 3 \cdot 5) \\ &= (6, 15) \\ &\equiv (1, 6) \end{aligned}$$

□

**Problem 13.2** (Problem 11 Fraleigh's p.175). *Determine whether the following structure is a ring, then if it is a commutative ring, whether it is unital, and whether it is a field.*

$$\left\{a + b\sqrt{2} \mid a, b \in \mathbb{Z}\right\}$$

*Solution.* Let's denote this structure as $\mathbb{Z}(\sqrt{2})$.

It is quite clear that $(\mathbb{Z}(\sqrt{2}), +)$ forms an abelian group as the integers is an abelian group and the identity here is $0 + 0\sqrt{2} \in \mathbb{Z}(\sqrt{2})$. Furthermore $(a + b\sqrt{2}) + (c + d\sqrt{2}) = (a + c) + (b + d)\sqrt{2} \in \mathbb{Z}(\sqrt{2})$, so we're good.

The $\mathbb{Z} \setminus \{0\}$ with multiplication is closed, but not every element has an inverse (in fact only $1$ and $-1$ do), so we'd expect something similar here.

Suppose $a + b\sqrt{2}, c + d\sqrt{2} \in \mathbb{Z}(\sqrt{2}) \setminus \{0\}$. Then

$$\begin{aligned} (a + b\sqrt{2})(c + d\sqrt{2}) &= ac + cb\sqrt{2} + ad\sqrt{2} + 2bd \\ &= (ac + 2bd) + (ad + cb)\sqrt{2} \in \mathbb{Z}(\sqrt{2}) \setminus \{0\} \end{aligned}$$

So the multiplicative structure is closed with identity, but there does not exist inverses for every element as for instance $2 \in \mathbb{Z}(\sqrt{2})^x$, but

$2^{-1}$ does not exist. It is also not hard to see that distributive laws work fine.

Thus we conclude that $\mathbb{Z}(\sqrt{2})$ is a commutative unital ring. $\square$

**Problem 13.3** (Problem 20 Fraleigh's p.175). *Consider the matrix ring $M_2(\mathbb{Z}_2)$.*
*a. Find the order (number of elements) of the ring.*
*b. List all units in the ring.*

*Solution of (a).* Recall that

$$M_2(\mathbb{Z}_2) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \middle| \, a, b, c, d \in \mathbb{Z}_2 \right\}$$

There are $4$ entries in each matrix in the ring, all of which can take on one of two values from $\mathbb{Z}_4$. Thus there are $2^4 = 16$ elements. $\square$

**Problem 13.4** (Problem 37 Fraleigh's p.176). *Show that if $U$ is the collection of all units in a ring $(R, +, \cdot)$ with unity, then $(U, \cdot)$ is a group.*

*Proof.* We conisder $U$ the collection of the units of a ring $R$. $1 \in R$ (as $R$ has unity), $(R, \cdot)$ has associativity by definition and the set of units $U$ is precisely those $u \in R$ such that

$$u \cdot u^{-1} = u^{-1} \cdot u = 1$$

so it remains to show that $U$ is closed. Let $u, v \in U$. Then, as said, there exists inverses $u^{-1}, v^{-1} \in U$. Then

$$\begin{aligned} (uv)(v^{-1}u^{-1}) &= u(vv^{-1})u^{-1} \\ &= u(1)u^{-1} \\ &= uu^{-1} \\ &= 1 \end{aligned}$$

so $(uv)^{-1} = (v^{-1}u^{-1}) \in U$. Thus the product of two units is itself a unit. We have shown $(U, \cdot)$ to have all the properties of a group. $\square$

**Problem 13.5** (Problem 46 Fraleigh's p.176). *An element $a$ of a ring $R$ is nilpotent if $a^n = 0$ for some $n \in \mathbb{Z}^+$. Show that if $a$ and $b$ are nilpotent in some ring commutative ring $R$, then $a + b \in R$ is also nilpotent.*

*Proof.* Suppose $a, b \in R$ (commutative) such that $a^n = 0$ and $b^m = 0$ for some $n, m \in \mathbb{Z}^+$.

Let $k \in \mathbb{Z}^+$. Then

$$\begin{aligned}(a+b)^k &= \sum_{i=0}^{k} \binom{k}{i} a^{k-i} b^i \\ &= a^k + ka^{k-1}b + \cdots + kab^{k-1} + b^k\end{aligned}$$

in particular, if $k := n + m$ then we observe the following:

$$\begin{aligned}(a+b)^{n+m} &= \sum_{k=0}^{n+m} \binom{n+m}{k} a^{n+m-k} b^k \\ &= \sum_{k=m+1}^{n+m} \binom{n+m}{k} a^{n+m-k} b^k + \sum_{k=0}^{m} \binom{n+m}{k} a^{n+m-k} b^k \\ &= \underbrace{\binom{n+m}{m+1} a^{n+m-m-1} b^{m+1}}_{\cdots\, a^{n-1} \cdot 0 \cdot b = 0} + \ldots + \underbrace{b^{n+m}}_{(b^n) \cdot 0 = 0} \\ &+ \underbrace{a^{n+m}}_{0 \cdot a^m = 0} + \cdots + \underbrace{\binom{n+m}{m} a^{n+m-m} b^m}_{\cdots\, 0 \cdot 0 = 0} \\ &= 0\end{aligned}$$

since every term is $0$. □

**Problem 13.6** (Problem 1 Fraleigh's p.182). *Find all solutions of $x^3 - 2x^2 - 3x = 0$ in $\mathbb{Z}_{12}$.*

*Solution.*

$$\begin{aligned}f(x) = x^3 - 2x^2 - 3x &= 0 \\ x(x^2 - 2x - 3) &= 0 \\ x(x-3)(x+1) &= 0\end{aligned}$$

Since we are in a ring with zero-divisors we are really asking for which $x \in \mathbb{Z}_{12}$ the following holds

$$12 \mid x(x-3)(x+1)$$

We begin by solving  mod $3$ where

$$f(x) = x^2(x+1) \in \mathbb{Z}_3[x]$$

with solutions $0, 2 \in \mathbb{Z}_3$. $f(x)$ reduces to

$$f(x) = x(x-1)^2 \in \mathbb{Z}_4[x]$$

with solutions $0, 1, 2 \in \mathbb{Z}_4$. We get the solutions

$$\{0, 2, 5, 6, 8, 9\} \subset \mathbb{Z}_{12}$$

□

**Problem 13.7** (Problem 11 Fraleigh's p.182). *Let $R$ be a commutative unital ring with characteristic $4$. Compute and simplify $(a+b)^4$ for $a, b \in R$.*

*Solution.*

$$\begin{aligned}(a+b)^4 &= a^4 + \binom{4}{1}a^3b + \binom{4}{2}a^2b^2 + \binom{4}{3}ab^3 + b^4 \\ &= a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4\end{aligned}$$

As $4 \equiv 0 \mod 4$ and $6 \equiv 2 \mod 4$ we get $(a+b)^4 \equiv a^4 + 2a^2b^2 + b^4$ in integral domain $R$ with $\text{char}(R) = 4$. □

**Problem 13.8** (Problem 23 Fraleigh's p.183). *Show that a division ring has exactly two idempotent elements.*

*Proof.* Recall that we define division rings as non-trivial so $|R| \geq 2$ crucially. The candidates for idempotent elements are $0, 1 \in R$ as $0 \cdot 0 = 0$ as $0 \cdot a = 0 \; \forall a \in R$. Then since $1 \cdot a = a \; \forall a \in R$ we conclude that $1 \cdot 1 = 1$. Now we need to show that there do not exist any other elements which satisfy this property. Assume, for a contradiction, that there exists $a \in R \setminus \{0, 1\}$ such that

$$a^2 = a$$

As $R$ is a division ring we know there exists $a^{-1} \in R$ such that

$$\begin{aligned}a^2a^{-1} &= aa^{-1} \\ a(aa^{-1}) &= 1 \\ a &= 1\end{aligned}$$

contradiction our assumption that $a \neq 1$. □

**Problem 13.9** (Problem 29 Fraleigh's p.183). *Show that the characteristic of an integral domain $D$ is either $0$ or a prime $p$.*

*Proof.* Let $D$ be an integral domain, i.e. a commutative ring with unity and no zero-divisors. Assume, for a contradiction, that $\text{char}(D) = k \in$

$\mathbb{Z}^+$ (not prime). Then char$(D) = k = nm$, $n, m \in \mathbb{Z}^+$. Then we can conclude that

$$k \cdot 1 = \underbrace{1 + 1 + \cdots + 1 + 1}_{k=nm \text{ times}} = 0$$

but then

$$(n \cdot 1)(m \cdot 1) = 0$$

so $n \cdot 1 = 0$ or $m \cdot 1 = 0$ so $k$ is not the characteristic of $D$. $\square$

# 14 Fermat's and Euler's theorems, Quotient Fields

**Theorem 14.1** (Fermat's Little Theorem). *If $p$ is prime and* $\gcd(a, p) = 1$, *then*

$$a^{p-1} \equiv 1 \mod p$$

*Equivalently*

$$a^p \equiv a \mod p$$

*Proof idea.* The key fact is that

$$(\mathbb{Z}/p\mathbb{Z})^\times$$

the nonzero elements mod $p$, form a group under multiplication. This group has $p - 1$ elements (finite) and cyclic. By Lagrange's Theorem, if $G$ is finite and $g \in G$ then

$$g^{|G|} = e$$

Applying this to $G = (\mathbb{Z}/p\mathbb{Z})^\times$ we get

$$a^{p-1} \equiv 1$$

□

**Theorem 14.2.** *Euler's Theorem If* $\gcd(a, n) = 1$, *then*

$$a^{\phi(n)} \equiv 1 \mod n,$$

*where $\phi(n)$ is Euler's totient function.*

*Proof idea.* We consider

$$(\mathbb{Z}/n\mathbb{Z})^\times$$

the group of units mod $n$. It has exactly $\phi(n)$ elements. Again by Lagrange:

$$a^{\phi(n)} \equiv 1$$

□

So Fermat's Little Theorem is just Euler's Theorem for $n$ prime.

## 14.1 Quotient Fields

In $\mathbb{Z}$, you generally cannot divide by $2$. But in $\mathbb{Q}$, you can. The questions we want to answer is:

"Can we build a field from a ring?"

As it turns out the answer is yes (conditions may apply...).

Recall that an integral domain is

- a commutative ring,
- with identity,
- with no zero divisors

Suppose $D$ is an integral domain. We construct Frac$(D)$ like rational numbers.

**Definition 14.1** (Quotient Field). *The **field of fractions** or **quotient field** of an integral domain $D$ is the smallest field $K$ (often denoted* Frac$(D)$*) containing $D$ as a subring, constructed by forming formal fractions $\frac{a}{b}$ where $a, b \in K$ $(b \neq 0)$.*
*The operations on $K$ are defined as*

$$\frac{a}{b} + \frac{c}{d} = \frac{ad + bd}{bd}$$

*and*

$$\frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd}$$

*We equip $K$ with an equivalence relation $\sim$ where we say*

$$\frac{a}{b} \sim \frac{c}{d}$$

*if $ad = bc$.*

As we see we get a field Frac$(D)$ and we get an injective embedding

$$D \hookrightarrow \text{Frac}(D)$$

via

$$a \mapsto \frac{a}{1}$$

**Problem 14.1** (Problem 20.5 Fraleigh's p.189). *Use Fermat's theorem to find the remainder of $37^{49}$ when it is divided by $7$.*

*Solution.* gcd$(37, 7) = 1$ and $p = 7$ is ceirtainly prime so Fermat's Little Theorem applies.

We can rewrite

$$37^{49} = 37^{7^2} \equiv 2^{7^2}$$

From FLT we have $2^6 \equiv 1$.

$$2^{7^2} = 2^{6 \cdot 8 + 1} \equiv 1^8 \cdot 2 \equiv 2 \mod 7$$

□

**Problem 14.2** (Problem 20.12 Fraleigh's p.189). *Describe all solutions to*

$$22x \equiv 5 \mod 15$$

*Solution.* The greatest common divisor $\gcd(22, 15) = 1$ which is a divisor of $5$ so there does exist solutions (in fact, as $\gcd(22, 15) = 1$ there exist exactly one solution). We can simplify to get

$$7x \equiv 5 \mod 15$$

As the greates common divisor is $1$ we know that $7$ is a unit in $\mathbb{Z}/15\mathbb{Z}$. We want to find $7^{-1}$. Notice that $7 \cdot 13 = 91 \equiv 1$.

$$\begin{aligned} x &\equiv 13 \cdot 5 \mod 15 \\ 13 \cdot 5 &= 65 \\ 65 &\equiv 5 \mod 15 \end{aligned}$$

Thus

$$x \equiv 5 \mod 15$$

□

**Problem 14.3** (Problem 20.15 Fraleigh's p.189). *As in the task above for*

$$39x \equiv 125 \mod 9$$

*Solution.* Notice that

$$39 \equiv 3 \mod 9$$

and

$$125 \equiv -1 \equiv 8 \mod 9$$

so we can transform the problem into

$$3x \equiv 8 \mod 9$$

Notice that $\gcd(3, 9) = 3$, but $3$ is not a divisor of $8$ thus the solution DNE. □

**Problem 14.4** (Problem 20.27 Fraleigh's p.190). *Show that $1$ and $p - 1$ are the only elements of the field $\mathbb{Z}_p$ that are their own multiplicative inverse. [Hint: Consider the equation $x^2 - 1 = 0$].*

*Proof.* Let $p$ be prime and let $x \in \mathbb{Z}_p$. Notice that $x = x^{-1}$ iff

$$x^2 \equiv 1 \mod p \Leftrightarrow x^2 - 1 \equiv 0 \mod p$$

It is not hard to see that $1$ and $p-1$ are solutions to these equations, but we have to show that they are the only ones. Notice that $x^2-1$ can be rewritten so we get

$$(x-1)(x+1) \equiv 0 \mod p$$

Recall that since $\mathbb{Z}_p$ is a field (no zero divisors) it is the case that $x-1 \equiv 0$ or $x+1 \equiv 0$. I.e.

$$x \equiv 1 \mod p, \quad x \equiv -1 \mod p.$$

Therefore

$$x = \begin{cases} 1 \\ p-1 \end{cases}$$

□

**Problem 14.5** (Problem 21.1 Fraleigh's p.196). *Describe the field $F$ of quotients of the integral subdomain*

$$D = \{n + mi \mid n, m \in \mathbb{Z}\}$$

*of $\mathbb{C}$. "Describe" means give the elements of $\mathbb{C}$ that make up the field of quotients of $D$ in $\mathbb{C}$.*

*Solution.* Recall that

$$\text{Frac}(D) = \left\{ \frac{a}{b} \,\middle|\, a, b \in D \ (b \neq 0) \right\}$$

$a, b \in D$ means that $a, b$ can be written as

$$a = n + mi, \quad b = k + li, \quad n, m, k, l \in \mathbb{Z}$$

(note: $k + li \neq 0 \Rightarrow k \neq 0 \neq li$) so

$$\text{Frac}(D)$$

are elements of the form

$$\begin{aligned} \frac{n+mi}{k+li} &= \frac{n+mi}{k+li} \cdot \frac{k-li}{k-li} \\ &= \frac{(n+mi)(k-li)}{k^2+l^2} \\ &= \frac{nk + kmi - nli + ml}{k^2+l^2} \\ &= \frac{nk + ml + (km - nl)i}{k^2+l^2} \\ &= \frac{nk+ml}{k^2+l^2} + \frac{km-nl}{k^2+l^2} i \end{aligned}$$

Now we know $\text{Frac}(D) \subseteq \mathbb{Q}(i)$, but is this subset relation strict? Let $a = r/s,\ b = t/s$. Then

$$a + bi = \frac{r}{s} + \frac{t}{s}i$$

and since

$$r + ti \in \mathbb{Z}(i) = D, \text{ and } s \in \mathbb{Z} \subset \mathbb{Z}(i) = D$$

we conclude that $\mathbb{Q}(i) \subseteq D$. I.e.

$$D = \text{Frac}(\mathbb{Z}(i)) = \mathbb{Q}(i)$$

□

**Problem 14.6** (Problem 21.12 Fraleigh's p.197). *Let $R$ be a nonzero commutative ring, and let $T$ be a nonempty subset of $R$ closed under multiplication and containing neither $0$ nor divisors of $0$. Starting with $R \times T$ and otherwise exactly following the construction in this section, we can show that the ring $R$ can be enlarged to a partial ring of quotients $Q(R,T)$. Think about this for $15$ minutes or so; look back over the construction and see why things still work. In particular, show the following:*
*a. $Q(R,T)$ has unity even if $R$ does not.*
*b. In $Q(R,T)$, every nonzero element of $T$ is a unit.*

*Proof of (a).* We let $R$ be some nonzero commutative ring and take a closed subset $T$ not containing $0$ or any divisors of it and construct $Q(R,T)$ to be

$$\frac{r}{t},\ r \in R \text{ and } t \in T$$

Let $x \in Q(R,T)$ then

$$x = \frac{r}{t}$$

for some $r \in R,\ t \in T$. As $T \subseteq R$ we know that

$$\frac{t}{t} \in Q(R,T)$$

Notice that

$$\frac{t}{t} \cdot \frac{r}{t} = \frac{tr}{t^2} = \frac{r}{t} \cdot \frac{t}{t}$$

but

$$\frac{tr}{t^2} = \frac{r}{t}$$

as

$$t^2 r = trt$$

so $t/t = 1_{Q(R,T)}$. □

*Proof.* Suppose $t, s \in T$, nonzero, with

$$\frac{t}{s} \in Q(R, T)$$

We claim this arbitrary nonzero element is invertible. Consider

$$\frac{s}{t}$$

Observe that

$$\frac{t}{s}\frac{s}{t} = \frac{t}{t} = \frac{s}{t}\frac{t}{s}$$

and as we showed in (a), and following from $st = ts$, we get

$$1 = \frac{s}{t}\frac{t}{s}$$

□

**Problem 14.7** (Problem 22.2 Fraleigh's p.207). *Find the sum and product of the given polynomial in the polynomial ring:*

$$f(x) = x + 1,\ g(x) = 3x^2 + 2x + 3, \text{ in } \mathbb{Z}_2[x].$$

*Solution.*

$$\begin{aligned} f(x) + g(x) &= 3x^2 + 3x + 4 \\ &\equiv x^2 + x \mod 2 \end{aligned}$$

$$\begin{aligned} f(x)g(x) &= (x+1)(3x^2 + 2x + 3) \\ &\equiv (x+1)(x^2+1) \mod 2 \\ &= x^3 + x^2 + x + 1 \end{aligned}$$

□

**Problem 14.8** (Problem 22.5 Fraleigh's p.207). *How many polynomials are there of degree $\leq 3$ in $\mathbb{Z}_2[x]$? (Include $0$)*

*Solution.* The polynomials in $\mathbb{Z}_2$ are precisely

$$ax^3 + bx^2 + cx + d,\ \ a, b, c, d \in \mathbb{Z}_2$$

So there are $4$ independent choices of coefficients which each can take on one of two values. Thus there are

$$2^4 = 16$$

possible polynomials in $\mathbb{Z}_2$. □

**Problem 14.9** (Problem 22.12/13 Fraleigh's p.207). *Find all zeros of the indicated finite field of the given polynomial in said field.*

$$x^2 + 1, \text{ in } \mathbb{Z}_2$$

*and*

$$x^3 + 2x + 2, \text{ in } \mathbb{Z}_7$$

*Solution.*

$$x^2 + 1 = 0$$
$$x^2 = -1 \equiv 1 \mod 2$$

so the only zero is $1 \in \mathbb{Z}_2$.

Next we work with

$$x^3 + 2x + 2 = 0$$

Let us begin by trying some values. Clearly $0$ does not work. If we plug in $1$ we get

$$1 + 2 + 2 = 5 \neq 0$$

Next $2$ gives us

$$8 + 4 + 2 = 14 \equiv 0$$

so $2$ is a solution.

$$27 + 6 + 2 = 35 \equiv 0$$
$$64 + 8 + 2 = 74 \equiv 4 \neq 0$$

and so on to find the rest of the zeros. □

**Problem 14.10** (Problem 27 Fraleigh's p.208). *Let $F$ be a characteristic zero field and let $D$ be the formal polynomial differentiation map, so that*

$$D\left(\sum_{k=0}^{n} a_k x^k\right) = \sum_{k=1}^{n} k a_k x^{k-1}$$

*a) Show that $D : F[x] \to F[x]$ is a group homomorphism of $\langle F[x], + \rangle$ onto itself. Is $D$ a ring homomorphism?*
*b) Find the kernel of $D$.*
*c) Find the image of $F[x]$ under $D$.*

*Proof of (a).* Let $f, g \in F[x]$, i.e.

$$f(x) := \sum_{k=0}^{n} a_k x^k, \quad g(x) := \sum_{k=0}^{n} b_k x^k$$

Then

$$\begin{aligned} D(f+g) &= D\left(\left[\sum_{k=0}^{n} a_k x^k\right] + \left[\sum_{k=0}^{n} b_k x^k\right]\right) \\ &= D\left(\sum_{k=0}^{n} (a_k + b_k) x^k\right) \\ &= \sum_{k=1}^{n} k(a_k + b_k) x^{k-1} \\ &= \sum_{k=1}^{n} k(a_k) x^{k-1} + \sum_{k=1}^{n} k(b_k) x^{k-1} \\ &= D(f) + D(g) \end{aligned}$$

It is not a ring homomorphism. Consider

$$x(x^2+1)$$

Then

$$D(x(x^2+1)) = D(x^3+x) = 3x^2+1$$

but

$$D(x)D(x^2+1) = 2x$$

□

*Solution of (b).* The preimage of the zero polynomial under differentiation is simply all the constant polynomials i.e.

$$c \in F$$

□

*Solution of (c).* Differentiation is surjective so the image of $F[x]$ is simply $F[x]$. Take any $f \in F[x]$, i.e. $f = \sum_{k=0}^{n} a_k x^k$. Then

$$D^{-1}(f) = c + \sum_{k=0}^{n} \frac{a_k}{k+1} x^{k+1}$$

for any $c \in F$. So $f \in F[x]$ has precisely $|F|$ elements in the preimage.

□

# 15 Polynomial Rings

**Definition 15.1.** *Let $R$ be a ring.*
*A polynomial over $R$ is*

$$\sum_{k=0}^{n} a_k x^k$$

*where $a_k \in R$ and $n \geq 0$.*
*The set of all such polynomials is denoted $R[x]$, and is called the polynomial ring over $R$.*

Two polynomials $\sum_{k=0}^{n} a_k x^k$ and $\sum_{k=0}^{n} b_k x^k$ are equal iff $a_k = b_k, \forall k \in [n]$.

With addition, they are very nicely behaved:

$$\sum_{k=0}^{n} a_k x^k + \sum_{k=0}^{n} b_k x^k = \sum_{k=0}^{n} (a_k + b_k) x^k$$

while multiplication looks like:

$$\sum_{k=0}^{n} a_k x^k \cdot \sum_{k=0}^{n} b_k x^k = \sum_{k} \left( \sum_{i+j=k} a_i b_j \right) x^k$$

Notably, if $R$ is a ring then $R[x]$ is a ring. Furthermore commutativity, identity and being and integral domain carry over. Note that not all properties transfer, e.g. being a field does not always carry over.

## 15.1 Degree of a Polynomial

**Definition 15.2.** *Let*

$$f(x) = a_n x^n + \cdots + a_0$$

*with $a_n \neq 0$. The **degree** of $f$ is*

$$\deg(f) = n.$$

*The degree of the zero polynomial is defined to be $-\infty$.*

If $R$ is an integral domain then

$$\deg(fg) = \deg(f) + \deg(g).$$

### 15.2 Units in Polynomial Ring

If $R$ is an integral domain, the only units in $R[x]$ are the constant polynomials whose coefficients are units in $R$.

For example, the only units in $\mathbb{Z}[x]$ are $\pm 1$.

### 15.3 Evaluation Homomorphisms

For $c \in R$ define

$$\varphi_c : R[x] \to R$$

by

$$\varphi_c(f(x)) = f(c).$$

This map is a ring homomorphism and is called the **evaluation homomorphism**.

### 15.4 Roots

**Definition 15.3.** *Let $f(x) \in R[x]$ and $c \in R$. We say $c$ is a **root** of $f$ if*

$$f(c) = 0.$$

**Theorem 15.1** (Factor Theorem). *If $c$ is a root of $f(x)$ in $R$, then*

$$(x - c) \mid f(x).$$

If $R$ is an integral domain and $\deg(f) = n$, then $f$ has at most $n$ roots in $R$.

## 16 Polynomial Factorisation

Polynomial arithmetic behaves especially well over fields.

**Theorem 16.1** (Division Algorithm). *Let $F$ be a field and $f, g \in F[x]$ with $g \neq 0$. Then there exist unique polynomials $q, r \in F[x]$ such that*

$$f(x) = q(x)g(x) + r(x)$$

*where either $r = 0$ or*

$$\deg(r) < \deg(g).$$

This allows us to compute greatest common divisors of polynomials using the Euclidean algorithm.

**Definition 16.1.** *A polynomial* $f(x) \in F[x]$ *is* ***irreducible*** *if*

- $\deg(f) \geq 1$
- *whenever* $f = gh$ *with* $g, h \in F[x]$*, one of* $g, h$ *is a unit.*

**Theorem 16.2.** *If* $F$ *is a field then* $F[x]$ *is a unique factorization domain. In particular, every polynomial can be written as a product of irreducible polynomials, uniquely up to order and multiplication by units.*

## 16.1 Useful Irreducibility Criterion

If $f(x) \in F[x]$ has degree $2$ or $3$, then

$$f \text{ is irreducible} \iff f \text{ has no root in } F.$$

Thus to test irreducibility of such polynomials it suffices to check whether they have roots.

## 17 Problems from week of March 22

**Problem 17.1** (Problem 23.9 Fraleigh's p.218). *The polynomial $x^4+4$ can be factored into linear factors in $\mathbb{Z}_5[x]$. Find this factorization.*

*Solution.* We work with

$$f(x) = x^4 + 4 = (x^2 + 2x + 2)(x^2 - 2x + 2)$$

We see that

$$x^2 + 2x + 2 \equiv x^2 + 2x - 3 = (x + 3)(x - 1) \equiv (x + 3)(x + 4)$$

and

$$x^2 - 2x + 2 \equiv x^2 + 3x + 2 = (x + 2)(x + 1)$$

so

$$f(x) \equiv (x + 4)(x + 3)(x + 2)(x + 1)$$

□

**Problem 17.2** (Problem 23.14 Fraleigh's p.218). *Show that $f(x) = x^2 + 8x - 2$ is irreducible over $\mathbb{Q}$. Is $f(x)$ irreducible over $\mathbb{R}$?*

*Proof.* We consider $f(x) \in \mathbb{Q}[x]$.

$$f(x) = x^2 + 8x - 2$$

Since $f(x)$ is degree $2$ it is irreducible in $\mathbb{Q}$ if and only if it has no root in $\mathbb{Q}$.

$$\begin{aligned} f(x) &= 0 \\ x^2 + 8x - 2 &= 0 \\ (x + 4 - 3\sqrt{2})(x + 4 + 4\sqrt{2}) &= 0 \end{aligned}$$

$\sqrt{2} \notin \mathbb{Q}[x]$ so $f(x)$ is irreducible in $\mathbb{Q}$.

As $\sqrt{2} \in \mathbb{R}$ and $f(x)$ is degree $2$ we conclude that $f(x)$ is reducible in $\mathbb{R}[x]$. □

**Problem 17.3** (Problems 18 & 19 Fraleigh's p.219). *Determine whether the given polynomials satisfy the Eisenstein criterion for irreducibility over $\mathbb{Q}$.*

$$f(x) = x^2 - 12$$
$$g(x) = 8x^3 + 6x^2 - 9x + 24$$

*Solution.* Recall that for some polynomial

$$\sum_{k=0}^{n} a_k x^k$$

If there exists a prime $p$ which divdes $a_0, \dots, a_{n-1}$ and doesn't divide $a_n$ and $p^2$ doesn't divide $a_0$, then it is irreducible over the rationals.

For

$$f(x) = x^2 - 12$$

we have $a_0 = -12$, $a_1 = a_n = 1$. No prime number divides $1$ so that condition is fulfilled. Then we see that $3|12$, but $3^2$ does not divide $-12$. We conclude that $f(x)$ is irreducible in $\mathbb{Q}[x]$.

Next look at

$$g(x)$$

Observe that for $p = 3$ we have

$$\begin{aligned} p &\mid 24 \\ p &\mid -9 \\ p &\mid 6 \end{aligned}$$

but $p$ does not divide $8$ and $p^2 = 9$ does not divide $24$ so $g(x)$ is irreducible over the rationals. □

**Problem 17.4** (Problem 23.34 Fraleigh's p.219). *Show that for a prime $p$, the polynomial $x^p + a$ in $\mathbb{Z}_p[x]$ is not irreducible for any $a \in \mathbb{Z}_p$*

*Proof.* Let $p$ be prime and

$$x^p + a \in \mathbb{Z}_p[x]$$

By Fermat's Little Theorem $a = a^p$. Furthermore observe that

$$x^p + a = x^p + a^p = (x + a)^p$$

So $x^p + a$ is reducible to

$$\prod_{[p]} (x + a) \in \mathbb{Z}_p[x]$$

□

**Problem 17.5** (Problem 23.35 Fraleigh's p.219). *If $F$ is a field and $a \neq 0$ is a zero of $f(x) = a_0 + a_1 x + \cdots x_n x^n$ in $F[x]$, show that*

*$1/a$ is a zero of $a_n + a_{n-1}x + \cdots + a_0x^n$.*

*"Thinking out loud" proof.* Let $F$ be a field and let

$$f(x) := \sum_{i=0}^{n} a_i x^i$$

be a polynomial in $F[x]$. Assume $a \neq 0$ is a root of $f$, i.e.

$$f(a) = \sum_{i=0}^{n} a_i a^i = 0.$$

Consider the expression

$$\frac{f(x)}{x^n} = \sum_{i=0}^{n} a_i x^{i-n} = a_0 x^{-n} + a_1 x^{1-n} + \cdots + a_{n-1}x^{-1} + a_n.$$

This is not a polynomial in $F[x]$, but it suggests reversing the powers of $x$.

To obtain a genuine polynomial, define

$$g(x) := x^n f\left(\frac{1}{x}\right).$$

Then

$$\begin{aligned} g(x) &= x^n \sum_{i=0}^{n} a_i \left(\frac{1}{x}\right)^i \\ &= \sum_{i=0}^{n} a_i x^{n-i} \\ &= a_0 x^n + a_1 x^{n-1} + \cdots + a_{n-1}x + a_n. \end{aligned}$$

Thus $g(x) \in F[x]$ and it is the polynomial obtained by reversing the coefficients of $f$.

Now evaluate $g$ at $1/a$:

$$\begin{aligned} g\left(\frac{1}{a}\right) &= \left(\frac{1}{a}\right)^n f(a) \\ &= \frac{f(a)}{a^n} \\ &= 0. \end{aligned}$$

Hence $1/a$ is a root of $g$ whenever $a \neq 0$ is a root of $f$. $\square$

# 18 Homomorphisms of rings, Prime/maximal ideals

**Definition 18.1.** *A* ***ring homomorphism***

$$\phi : R \to S$$

*between rings satisfies*

$$\phi(a+b) = \phi(a) + \phi(b)$$
$$\phi(ab) = \phi(a)\phi(b)$$

*and typically*

$$\phi(1_R) = 1_S$$

**Theorem 18.1.** *Let $R, S$ be rings and $\phi : R \to S$ a homomorphism. Then $\ker \phi$ is an ideal of $R$.*

*Proof.* Let $a, b \in \ker \phi$.

$$\phi(a-b) = 0$$

so closed under subtraction. For any $r \in R$ we have

$$\phi(ra) = \phi(r)0 = 0$$

so $ra \in \ker \phi$.

Therefore kernels behave like normal subgroups. □

**Theorem 18.2** (First Isomorphism Theorem for Rings). *If*

$$\phi : R \to S$$

*then*

$$R/\ker \phi \cong \operatorname{Im} \phi$$

*meaning every homomorphic image of a ring is a quotient by an ideal.*

**Example 18.1.** *Take*

$$\phi : \mathbb{Z} \to \mathbb{Z}_n$$

*as*

$$\phi(k) = k \mod n$$

*The kernel is then*

$$\ker \phi = n\mathbb{Z}$$

*so*

$$\mathbb{Z}/n\mathbb{Z} \cong \mathbb{Z}_n$$

**Definition 18.2.** *If $I$ is an ideal of a ring $R$ we define*

$$R/I$$

*just like quotient groups, with operations*

$$(a+I)+(b+I)=(a+b)+I$$
$$(a+I)(b+I)=ab+I$$

## 18.1 Prime Ideals

**Definition 18.3.** *An ideal $P \subset R$ is **prime** if*

$$ab \in P \Rightarrow a \in P \text{ or } b \in P$$

**Example 18.2.** *In $\mathbb{Z}$ all ideals are $(n)$ and the prime ideals are $(p)$ where $p$ is a prime number. If $a, b \in (p)$ then $p|ab$ so $p|a$ or $p|b$.*

**Theorem 18.3.** *Suppose $P \subset R$ is an ideal. Then*

$$P \text{ prime} \Leftrightarrow R/P \text{ is an integral domain}$$

## 18.2 Maximal Ideals

**Definition 18.4.** *An ideal $M \subset R$ is **maximal** if $M \neq R$ and there is no ideal strictly between $M$ and $R$, meaning*

$$M \subset I \subset R$$

*implies $M = I$ or $R = I$.*

**Theorem 18.4.** *Let $M \subset R$ be an ideal.*
*Then*

$$M \text{ maximal} \Leftrightarrow R/M \text{ is a field}$$

An important note is that maximal $\Rightarrow$ prime, but not necessarily the other way around, so being maximal is a strictly stronger condition. This is easy to see since a field is also an integral domain, but not vice versa.

## 19 Problems problems and more problems

**Problem 19.1** (Problem 26.12 Fraleigh's p.244). *Give an example to show that a factor ring of an integral domain may be a field.*

*Solution.* Recalling that an integral domain is a nonzero, commutative, unital ring, we give the following example.

Consider the integral domain $\mathbb{Z}$. Take some prime $p \in \mathbb{Z}$ and examine the factor ring $\mathbb{Z}/\langle p \rangle$. It is known that

$$\mathbb{Z}/\langle p \rangle \cong \mathbb{Z}_p$$

The integers modulo a prime is a field. □

**Problem 19.2** (Problem 26.13 Fraleigh's p.244). *Give an example to show that a factor ring of an integral domain may have $0$ divisors.*

*Solution.* Consider again the integers $\mathbb{Z}$. Take the ideal generated by $6$

$$\langle 6 \rangle = \{\dots, -6, 0, 6, \dots\}$$

Notice that

$$\mathbb{Z}/\langle 6 \rangle \cong \mathbb{Z}_6$$

and $2, 3 \in \mathbb{Z}_6$, but

$$2 \cdot 3 \equiv 0$$

□

**Problem 19.3** (Problem 26.25 Fraleigh's p.244). *Show that if $R$ is a ring with unity and $N$ is an ideal of $R$ such that $N \neq R$, then $R/N$ is a ring with unity.*

*Proof.* Let $R$ be a unital ring with ideal $N \neq R$.

We study the quotient ring

$$R/N \neq \{N\}$$

Let $(a + N) \in R/N$. Consider $(1 + N) \in R/N$ Then

$$(a + N)(1 + N) = (a \cdot 1) + N = (1 \cdot a) + N = (1 + N)(a + N)$$

□

**Problem 19.4** (Problem 26.26 Fraleigh's p.245). *Let $R$ be a commutative ring and let $a \in R$. Show that $I_a = \{x \in R \mid ax = 0\}$ is an*

*ideal of $R$.*

*Proof.* We consider, for some arbitrary element of a commutative ring $R$, the set of right (and correspondingly left) zero divisors of $a$

$$I_a = \{x \in R \mid ax = 0\}$$

We know $I_a \neq \emptyset$ as $a \cdot 0 = 0$. Let $b, c \in I_a$. Then $ab = ac = 0$, so

$$a(b-c) = ab - ac = 0 - 0 = 0$$

so $b - c \in I_a$. Therefore $(I_a, +) \leq (R, +)$. Let $r \in R$. As stated $ab = 0$ and furthermore

$$a(rb) = a(br) = (ab)r = 0r = 0$$

so $rb \in I_a$. Thus $I_a$ is an ideal of $R$. □

**Problem 19.5** (Problem 26.30 Fraleigh's p.245). *An element $a$ of a ring $R$ is nilpotent if $a^n = 0$ for some $n \in \mathbb{Z}^+$. Show that the collection of all nilpotent elements in a commutative ring $R$ is an ideal, the nilradical of $R$.*

*Proof.* Let $R$ be a commutative ring and let $A \subseteq R$ be the set

$$A = \{a \in R \mid \exists n \in \mathbb{Z}^+ : a^n = 0\}$$

$A \neq \emptyset$ as $0^1 = 0$. Let $a, b \in A$. Then there is some $m, n \in \mathbb{Z}^+$ such that

$$a^m = 0 \text{ and } b^n = 0$$

Let $k = m + n \in \mathbb{Z}^+$.

$$\begin{aligned}
(a-b)^k &= \sum_{i=0}^{k} \binom{k}{i} a^i (-b)^{k-i} \\
&= \sum_{i=0}^{m+n} \binom{m+n}{i} a^i (-b)^{k-i} \\
&= \sum_{i=m+1}^{m+n} \binom{m+n}{i} a^i (-b)^{k-i} \\
&+ \sum_{i=0}^{m} \binom{m+n}{i} a^i (-b)^{k-i}
\end{aligned}$$

Notice that $\sum_{i=m+1}^{m+n} \binom{m+n}{i} a^i (-b)^{k-i}$ has $i \geq m+1$ for every term so $a^i = 0$ for all terms. Hence every term is $0$. For $\sum_{i=0}^{m} \binom{m+n}{i} a^i (-b)^{k-i}$ we have $k - i \geq n$ so $(-b)^{k-i} = 0$ for every term. Therefore

$$\sum_{i=m+1}^{m+n} \binom{m+n}{i} a^i (-b)^{k-i} + \sum_{i=0}^{m} \binom{m+n}{i} a^i (-b)^{k-i} = 0 + 0 = 0$$

Therefore $(a-b) \in A$. Now let $r \in R$ and observe that

$$(ra)^m = r^m a^m = r^m \cdot 0 = 0$$

so $ra \in A$. Thus $A$ constitutes an ideal of $R$. $\square$

**Problem 19.6** (Problem 26.34 Fraleigh's p.245). *Let $R$ be a commutative ring and $N$ an ideal of $R$. Show that the set $\sqrt{N}$ of all $a \in R$, such that $a^n \in N$ for some $n \geq \mathbb{Z}^+$, is an ideal of $R$, the* ***radical ideal*** *of $N$.*

*Proof.* Let $N$ be an ideal of some commutative ring $R$.

Let

$$\sqrt{N} = \{a \in R \mid \exists n \in \mathbb{Z}^+ \text{ s.t. } a^n \in N\}$$

First, observe that $\sqrt{N} \neq \emptyset$ as $0 = 0^1 \in N$. Next, let $a, b \in \sqrt{N}$. We show that $(a-b)^k \in N$ for some $k$, i.e. $a - b \in \sqrt{N}$. By assumption there exists $m, n \in \mathbb{Z}^+$ so that $a^m, b^n \in N$. Let $k = m + n$. Then

$$\begin{aligned}
(a-b)^k &= \sum_{i=0}^{k} \binom{k}{i} a^i (-b)^{k-i} \\
&= \sum_{i=0}^{m+n} \binom{m+n}{i} a^i (-b)^{m+n-i} \\
&= \sum_{i=m+1}^{m+n} \binom{m+n}{i} a^i (-b)^{m+n-i} \\
&\quad + \sum_{i=0}^{m} \binom{m+n}{i} a^i (-b)^{m+n-i}
\end{aligned}$$

The sum $\sum_{i=m+1}^{m+n} \binom{m+n}{i} a^i (-b)^{m+n-i}$ has $i \geq m+1$ so each term is of the form $a^m (a \cdots a)(-b)^{m+n-i}$. By assumption $a^m \in N$ and as $R$ is commutative we can rearange terms and see that since $N$ is an ideal of $R$ and $a \cdots a(-b)^{m+n-i} \in R$ we have that each term is in $N$ by the multiplication property of the ideal. As $N$ is a group under the addition, the sum of these terms is in $N$. Apply this same argument to the sum from $i = 0$ to $m$, observing that $m + n - i \geq n$, to find that $\sum_{i=0}^{m} \binom{m+n}{i} a^i (-b)^{m+n-i} \in N$. Both sums are in $N$ so

$$(a-b)^k \in N$$

i.e. $a - b \in \sqrt{N}$.

Lastly let $r \in R$ and see that

$$(ra)^m = r^m a^m \in N$$

as $N$ is an ideal. Thus $ra \in \sqrt{N}$.

Therefore $\sqrt{N}$ is an ideal of $R$. $\square$

**Problem 19.7** (Problem 27.5 Fraleigh's p.252). *Find all $c \in \mathbb{Z}_3$ such that*

$$\mathbb{Z}_3[x]/\langle x^2 + c\rangle$$

*is a field.*

*Proof.* Since $\mathbb{Z}_3$ is a field, the quotient

$$\mathbb{Z}_3[x]/\langle x^2 + c\rangle$$

is a field if and only if $x^2 + c$ is irreducible in $\mathbb{Z}_3[x]$.

The elements of $\mathbb{Z}_3$ are $0, 1, 2$. We check each case.

If $c = 0$, then $x^2$ is reducible since $x^2 = x \cdot x$.

If $c = 1$, we check whether $x^2 + 1$ has a root in $\mathbb{Z}_3$. The squares in $\mathbb{Z}_3$ are

$$0^2 = 0, \quad 1^2 = 1, \quad 2^2 = 1.$$

Thus $x^2$ never equals $2$, so $x^2 + 1$ has no root and is irreducible.

If $c = 2$, then

$$x^2 + 2 = x^2 - 1 = (x - 1)(x + 1),$$

so it is reducible.

Therefore the quotient ring is a field only when $c = 1$. □

**Problem 19.8** (Problem 27.15/16 Fraleigh's p.253). *Find a maximal ideal of $\mathbb{Z} \times \mathbb{Z}$.*
*Find a prime ideal of $\mathbb{Z}^2$ which is not maximal.*

*Solution.* We want to find the the proper ideal $M \subseteq \mathbb{Z} \times \mathbb{Z}$ such that for any ideal $I$, if $M \subseteq I \subseteq \mathbb{Z} \times \mathbb{Z}$ then $I = \mathbb{Z} \times \mathbb{Z}$ or $I = M$.

The maximal ideals of $\mathbb{Z}^2$ is $\langle p\rangle \times \mathbb{Z}$ and $\mathbb{Z} \times \langle p\rangle$ where $p$ is some prime. $\langle p\rangle^2$ does not work as the quotient would be $\mathbb{Z}_p^2$. Both factors are nontrivial so this is not a field. □

**Problem 19.9** (Problem 18 Fraleigh's p.253). *Is $\mathbb{Q}[x]/\langle x^2 - 5x + 6\rangle$ a field? Why?*

*Proof.* Recall that the ideal generated by $x^2 - 5x + 6$ is

$$\langle x^2 - 5x + 6\rangle = \{f(x)(x^2 - 5x + 6) : f(x) \in \mathbb{Q}[x]\}$$

We apply a theorem

$$\mathbb{Q}/\langle p\rangle \text{ is a field} \Leftrightarrow p \text{ is irreducible over } \mathbb{Q}$$

Notice that $x^2 - 5x + 6 = (x-2)(x-3)$ and $x-2, x-3 \in \mathbb{Q}[x]$, so

$$\mathbb{Q}[x]/\langle x^2 - 5x + 6 \rangle$$

is **not** a field. $\square$

**Problem 19.10** (Problem 19 Fraleigh's p.253). *Is* $\mathbb{Q}[x]/\langle x^2 - 6x + 6 \rangle$ *a field? Why?*

*Proof.* Notice that

$$x^2 - 6x + 6 = (x - (3 - \sqrt{3}))(x - (3 + \sqrt{3}))$$

Notice that $\sqrt{3} \notin \mathbb{Q}$, i.e. irreducible since the polynomial is a quadratic, so this quotient ring is a field. ($\cong \mathbb{Q}(\sqrt{3})$) $\square$

## 20 Tackling some extension field problems early

**Problem 20.1** (Problem 29.1/3/4 Fraleigh's p.272). *For each following $\alpha$, show that the given number $\alpha \in \mathbb{C}$ is algebraic over $\mathbb{Q}$ by finding $f(x) \in \mathbb{Q}$ such that $f(\alpha) = 0$.*

$$\alpha = 1 + \sqrt{2}$$
$$\alpha = 1 + i$$
$$\alpha = \sqrt{1 + \sqrt[3]{2}}$$

*Solutions.* We begin by considering $\alpha = 1 + \sqrt{2}$. Let $x = 1 + \sqrt{2}$. Then

$$\begin{aligned} x &= 1 + \sqrt{2} \\ \sqrt{2} &= x - 1 \\ 2 &= (x-1)^2 \\ x^2 - 2x + 1 &= 2 \\ x^2 - 2x - 1 &= 0 \end{aligned}$$

Thus $x^2 - 2x - 1 \in \mathbb{Q}$ is zero for $x = \alpha = 1 + \sqrt{2}$, hence $1 + \sqrt{2}$ is algebraic over $\mathbb{Q}$.

Next we look at $\alpha = 1 + i$. Set $x = 1 + i$.

$$\begin{aligned} i &= x - 1 \\ -1 &= (x-1)^2 \\ -1 &= x^2 - 2x + 1 \\ x^2 - 2x + 2 &= 0 \end{aligned}$$

$f(x) = x^2 - 2x + 2 \in \mathbb{Q}[x]$ and $f(\alpha) = 0$ so $\alpha = 1 + i$ is algebraic over $\mathbb{Q}$.

Lastly let $\alpha = \sqrt{1 + \sqrt[3]{2}}$.

$$\begin{aligned} x &= \sqrt{1 + \sqrt[3]{2}} \\ x^2 &= 1 + 2^{1/3} \\ x^2 - 1 &= 2^{1/3} \\ (x^2 - 1)^3 &= 2 \\ (x^2 - 1)^3 - 2 &= 0 \end{aligned}$$

As above ... $\alpha$ is algebraic over $\mathbb{Q}$. $\square$

**Problem 20.2** (Problem 29.18 Fraleigh's p.272). *a. Show that $x^2+1$ is irreducible in $\mathbb{Z}_3[x]$.*
*b. Let $\alpha$ be a zero of $x^2+1$ in an extension field of $\mathbb{Z}_3$. As in Example 29.19, give the multiplication and additions tables for the nine elements of $\mathbb{Z}_3[\alpha]$, written in the order $0, 1, 2, \alpha, 2\alpha, 3\alpha, 1+\alpha, 1+2\alpha, 2+\alpha$ and $2+2\alpha$.*

*Proof of (a).* Let $f(x) = x^2+1$. $f(x)$ is a classical introductory example to motivate the complex numbers, having $\alpha = \pm i \in \mathbb{C} \setminus \mathbb{R}$ as zero.

As $f$ is a degree $2$ polynomial with no zeros in $\mathbb{Z}_3$ we conclude that it is irreducible in $\mathbb{Z}_3[x]$. $\square$

## 21 Trying an earlier exam

**Problem 21.1** (Problem 1a, Exam 2023). *Prove that the set*

$$S = \{2^n \mid n \in \mathbb{Z}\}$$

*equipped with the usual operation of multiplication is a group.*

*Proof.* It suffices to show that $(S, \cdot)$ is isomorphic to a known group. Let $\phi : \mathbb{Z} \to S$ be defined by

$$\phi(n) = 2^n$$

It is clear that $\phi$ is a bijection as the preimage of $2^n$ is $n$ and $2^n = 2^m$ iff $n = m$.

Lastly we show $\phi$ to be a homomorphism.

$$\begin{aligned} \phi(n+m) &= 2^{n+m} \\ &= 2^n \cdot 2^m \\ &= \phi(n)\phi(m) \end{aligned}$$

Thus $(S, \cdot)$ is isomorphic to $(\mathbb{Z}, +)$ which is a known group. □

**Problem 21.2** (Problem 1b, Exam 2023). *List all finite abelian groups of order $20$, up to isomorphism.*

*Solution.* The finite abelian groups of order $20$ are trivially finitely generated so the fundamental theorem of finitely generated abelian groups applies.

I.e.

$$G \cong \left( \prod_{i=1}^{n} \mathbb{Z}_{p_i^{e_i}} \right) \times \mathbb{Z}^k$$

We are asked to find those of order $20$ (finite) so

$$G \cong \prod_{i=1}^{n} \mathbb{Z}_{p_i^{e_i}}$$

where $\prod_{i=1}^{n} p_i^{e_i} = 20$.

$20 = 4 \cdot 5 = 2 \cdot 2 \cdot 5$. Thus there exists exactly two abelian groups of order $20$, namely

$$\mathbb{Z}_4 \times \mathbb{Z}_5$$

and

$$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_5$$

□

**Problem 21.3** (Problem 1c, Exam 2023). *Let $A_4$ be the alternating group on $4$ letters, that is the subgroup of $S_4$ consisting of even permutations. Find elements $\sigma_1, \sigma_2, \sigma_3 \in A_4$ such that $\sigma_i$ has order $i$ (for $i = 1, 2, 3$).*

*Solution.* $\sigma_1$ is the trivial permutation, i.e. the one which sends

$$1 \mapsto 1, 2 \mapsto 2, \text{ etc.}$$

Next a candidate for $\sigma_2$ is any permutation which is the composition of two transpositions, e.g.

$$(1\ 2)(3\ 4)$$

Notice that

$$(1\ 2)(3\ 4) \circ (1\ 2)(3\ 4)$$

sends

$$1 \to 2 \to 1, 2 \to 1 \to 2, 3 \to 4 \to 3, 4 \to 3 \to 4$$

So $(1\ 2)(3\ 4)$ has order $2$.

An exampele of $\sigma_3$ is something like

$$(1\ 2\ 3) = (1\ 3)(1\ 2)$$

□

**Problem 21.4** (Problem 1d, Exam 2023). *Prove that $A_4$ is not abelian.*

*Proof.* Observe that $(1\ 2)(3\ 4)$ and $(1\ 3)(1\ 2)$ do not commute.

$$(1\ 2)(3\ 4) \circ (1\ 3)(1\ 2)$$

sends

$$\begin{aligned} 1 &\mapsto 2 \mapsto 1 \\ 2 &\mapsto 3 \mapsto 4 \\ 3 &\mapsto 1 \mapsto 2 \\ 4 &\mapsto 4 \mapsto 3 \end{aligned}$$

So

$$(1\ 2)(3\ 4) \circ (1\ 3)(1\ 2) = (2\ 4\ 3)$$

while

$$(1\ 3)(1\ 2) \circ (1\ 2)(3\ 4)$$

sends

$$3 \mapsto 4 \mapsto 4$$

So we can already see that

$$(1\ 3)(1\ 2) \circ (1\ 2)(3\ 4) \neq (1\ 2)(3\ 4) \circ (1\ 3)(1\ 2)$$

Hence $A_4$ is not abelian. □

**Problem 21.5** (Problem 2a, Exam 2023). *Determine the $0$-divisors in the ring $\mathbb{Z}_{10}$.*

*Solution.* $\mathbb{Z}_{10}$ is commutative so left- and right divisors coincide.

$$\begin{aligned}(2)(5) &= 10 \equiv 0\\ (4)(5) &= 20 \equiv 0\\ (6)(5) &= 30 \equiv 0\\ (8)(5) &= 40 \equiv 0\end{aligned}$$

so the $0$-divisrors in $\mathbb{Z}_{10}$ are $5$ with any power of $2$, which just so happens to be the nonzero elements of $\mathbb{Z}_{10}$ which are not coprime to $\mathbb{Z}_{10}$. □

**Problem 21.6** (Problem 2b, Exam 2023). *Let $R$ be the ring $\mathbb{Z}\times\mathbb{Z}$, where addition and multiplication is given by, for $(a,b),(c,d) \in \mathbb{Z}\times\mathbb{Z}$, setting*

$$\begin{aligned}(a,b)+(c,d) &= (a+c, b+d)\\ (a,b)(c,d) &= (ac, bd)\end{aligned}$$

*Let*

$$I = \{(a,0) | a \in \mathbb{Z}\} \subset R$$

*Prove that $I$ is an ideal of $R$.*

*Proof.* We show that $I$ is a subgroup of $(R,+)$ with the proper behavior w.r.t left-multiplication.

First we begin by observing that $(0,0) \in I \neq \emptyset$. Let $(a,0),(b,0) \in I$. Then we see that

$$\begin{aligned}(a,0)-(b,0) &= (a,0)+(-b,0)\\ &= (a-b, 0+0)\\ &= (a-b,0) \in I\end{aligned}$$

So $I$ is a subgroup of the additive group of $R$.

Lastly let $(c,d) \in R$. Then

$$\begin{aligned}(c,d)(a,0) &= (ca, d0)\\ &= (ca,0) \in I\end{aligned}$$

Thus $I$ is an ideal of $R$. □

**Problem 21.7** (Problem 2c, Exam 2023). *Prove that $I$ is a prime ideal and that $I$ is not a maximal ideal.*

*Proof.* It is clear that $I \neq R$ as $(1,1) \in R \setminus I$.

Now suppose $(ab, 0) \in I$. Then

$$(ab, 0) = (a, c)(b, d)$$

with $a, b, c, d \in \mathbb{Z}$ and $cd = 0$. In $\mathbb{Z}$ there are no zero-divisors, thus

$$c = 0 \text{ or } d = 0$$

and hence $(a, c) \in I$ or $(b, d) \in I$.

We show that $I$ is not maximal by giving some ideal $J$ so that

$$I \subseteq J \subseteq R$$

with $I \neq J$ and $J \neq R$.

Let $J$ be the set of $(a, b) \in R$ so that $b$ is even. Formally

$$J = \{(a, b) \in R \mid b = 2n \text{ for some } n \in \mathbb{Z}\}$$

It is trival to check that this is an ideal and that

$$I \subsetneq J \subsetneq R$$

$\square$

**Problem 21.8** (Problem 3a, Exam 2023). *Let $p$ be a prime number, and let $f(x) = x^p - x \in \mathbb{Z}_p[x]$. Determine (with justification) the zeros of $f(x)$ in $\mathbb{Z}_p[x]$.*

*Solution.* The zeros are all numbers in $\mathbb{Z}_p$. We can apply Fermat's little theorem

$$a^p \equiv a \mod p$$

to get that

$$f(x) = x^p - x^p = x - x = 0$$

□

**Problem 21.9** (Problem 3b, Exam 2023). *Consider $x^2+1 \in \mathbb{Z}_3[x]$. Prove that the ring $\mathbb{Z}_3[x]/\langle x^2+1\rangle$ is a field, and determine its number of elements.*

*Proof.* We use the following theorem:

For some field $F$ and $f \in F[x]$

$$F[x]/\langle f(x)\rangle$$

is a field if and only if $f(x)$ is irreducible in $F[x]$. Since $f(x) = x^2+1$ is a degree $2$ polynomial we see that $f$ is irreducible in $\mathbb{Z}_3[x]$ if and only if $f$ has no zeros in $\mathbb{Z}_3[x]$.

It is known that $f(x)$ is irreducible since it is the motivating introductory example for the complex numbers with solution $x = \pm i$.

We quickly check that $f(0) \neq 0$, $f(1) \neq 0$ and $f(2) \neq 0$ to confirm this.

As $3$ is prime we see that $\mathbb{Z}_3$ is a field.

We conclude therefore that

$$\mathbb{Z}_3[x]/\langle x^2+1\rangle$$

is a field. □

## 22 Extension fields & Vector spaces

**Definition 22.1.** *A **field extension** of a field $F$ is a field $K$ with $F \subseteq K$.*
*It is common to write*

$$K/F$$

Some basic examples of the above include $\mathbb{Q} \subseteq \mathbb{R}$, $\mathbb{Q} \subseteq \mathbb{C}$.

As we have seen earlier,

$$\mathbb{Z}_3 \subseteq \mathbb{Z}_3[x]/\langle x^2+1\rangle \cong \mathbb{Z}_3(i)$$

**Definition 22.2.** *Given a field $F$ and an element $\alpha \notin F$, define*

$$F(\alpha)$$

*to be the smallest field containing both $F$ and $\alpha$.*

**Theorem 22.1.** *If $\alpha$ satisfies a polynomial over $F$, then*

$$F(\alpha) \cong F[x]/\langle f(x)\rangle$$

*where $f$ is the minimal polynomial.*
*In other words, if $\alpha$ is algebraic over $F$ then this holds.*

**Definition 22.3.** *The minimal polynomial of $\alpha$ over $F$ is the monic polynomial of smallest degree in $F[x]$ such that $\alpha$ is a zero of said polynomial.*

**Definition 22.4.** *The degree of an extension $[K:F]$ is the dimension of $K$ as a vector space over $F$.*

**Example 22.1.** $[\mathbb{Q}(\sqrt{2}):\mathbb{Q}] = 2$ *with basis* $\{1, \sqrt{2}\}$.

**Theorem 22.2.** *If $K/F$ is a field extension then $K$ is a vector space over $F$.*

**Theorem 22.3.** *If minimal polynomial has degree $n$ then*

$$[F(\alpha):F] = n$$

**Theorem 22.4** (Tower Law). *If*

$$F \subseteq K \subseteq L$$

*then*

$$[L : F] = [L : K][K : F]$$

**Problem 22.1** (Problem 4a, Exam 2023). *Let $f(x) = x^5 - 8x^4 + 2x + 2 \in \mathbb{Q}[x]$. You may use w/o proof that $f$ is irreducible.*
*Let $\alpha \in \overline{\mathbb{Q}}$ be a zero of $f$. Compute $[\mathbb{Q}(\alpha) : \mathbb{Q}]$, and give a basis for $\mathbb{Q}(\alpha)$ as a vector space over $\mathbb{Q}$.*

*Proof.* Since $f(x) \in \mathbb{Q}[x]$ is irreducible and $\alpha$ is a root, $f$ is the minimal polynomial of $\alpha$ over $\mathbb{Q}$.

Therefore,

$$[\mathbb{Q}(\alpha) : \mathbb{Q}] = \deg f = 5.$$

A basis for $\mathbb{Q}(\alpha)$ over $\mathbb{Q}$ is

$$\{1, \alpha, \alpha^2, \alpha^3, \alpha^4\}.$$

□

**Problem 22.2** (Problem 4b, Exam 2023). *Let $\beta \in \mathbb{Q}(\alpha) \setminus \mathbb{Q}$. Prove that $\mathbb{Q}(\beta) = \mathbb{Q}(\alpha)$.*

*Proof.* $\beta \in \mathbb{Q}(\alpha)$ and $\beta \notin \mathbb{Q}$, by assumption. So

$$\mathbb{Q} \subseteq \mathbb{Q}(\beta) \subseteq \mathbb{Q}(\alpha)$$

We know that

$$[\mathbb{Q}(\alpha) : \mathbb{Q}] = [\mathbb{Q}(\alpha) : \mathbb{Q}(\beta)] \cdot [\mathbb{Q}(\beta) : \mathbb{Q}]$$

$5$ is prime so one of the factors is $5$ and the other is $1$. It can't be the case that

$$[\mathbb{Q}(\alpha) : \mathbb{Q}(\beta)] = 5$$

since that would imply $\mathbb{Q}(\beta) = \mathbb{Q}$ which is not the case, by assumption. Hence $\mathbb{Q}(\alpha) = \mathbb{Q}(\beta)$. □

## 23 Some problems relevant to Fraleigh's sections 29-33

**Problem 23.1.** *Let $\alpha = \sqrt{2} + \sqrt{5}$. Find the minimal polynomial for $\alpha$ over $\mathbb{Q}$. Then compute degree and identify a basis.*

*Solution.*

$$\begin{aligned} x &= \sqrt{2} + \sqrt{5} \\ x^2 &= 7 + 2\sqrt{10} \\ x^2 - 7 &= 2\sqrt{10} \\ \frac{x^2 - 7}{2} &= \sqrt{10} \\ \frac{(x^2-7)(x^2-7)}{4} &= 10 \\ \frac{x^4 - 14x^2 + 49 - 40}{4} &= 0 \\ \frac{x^4 - 14x^2 + 9}{4} &= 0 \\ x^4 - 14x^2 + 9 &= 0 \end{aligned}$$

is the minimal polynomial of $\alpha$. It is degree $r$ and thus a basis of $Q(\alpha)$ over $\mathbb{Q}$ is $\{1, \alpha, \alpha^2, \alpha^3\}$. □

**Problem 23.2.** *Give an example of a field extension which is finitely generated and not finite.*

*Solution.* Consider $\mathbb{C}/\mathbb{R}$. $\mathbb{C}$ is a 2-dimensional vector space over $\mathbb{R}$, but as a field $|\mathbb{C}| = \aleph_1$ and

$$|\mathbb{C} \setminus \mathbb{R}| = \aleph_1$$

□

**Problem 23.3.** *Let $\alpha, \beta$ be algebraic over $F$. Prove*

$$[F(\alpha, \beta) : F] \leq [F(\alpha) : F][F(\beta) : F]$$

*and explain when equality is achieved.*

*Proof.* Suppose $\alpha, \beta$ algebraic over some field $F$. Then there exists minimal polynomials

$$f, g \in F[x]$$

so that $f(\alpha) = 0$ and $g(\beta) = 0$. With

$$F(\alpha) = \left\{ \sum_{i=0}^{\deg f - 1} c_i \alpha_i \,\middle|\, c_i \in F \right\}$$

and

$$F(\beta) = \left\{ \sum_{i=0}^{\deg g - 1} c_i \beta_i \,\middle|\, c_i \in F \right\}$$

Then

$$[F(\alpha) : F][F(\beta) : F] = (\deg f - 1)(\deg g - 1)$$

It is clear that the two expressions are equal when $g$ is irreducible in $F(\alpha)$, recalling

$$F \subseteq F(\alpha) \subseteq F(\alpha, \beta)$$

A nice example is $\mathbb{Q}(\sqrt{2}, \sqrt{5})$.

$F(\alpha, \beta)$ is an extension field of $F(\alpha)$ so if $\beta$ and $\alpha$ are independent in their minimal polynomials then we get equality. □

**Problem 23.4** (Problem 31.3/5/7 Fraleigh's p.291). *Find the degree and basis for the given field extensions. Be prepared to justify your answers.*

3. $\mathbb{Q}(\sqrt{2}, \sqrt{3}, \sqrt{18})$ *over* $\mathbb{Q}$
5. $\mathbb{Q}(\sqrt{2}, \sqrt[3]{2})$ *over* $\mathbb{Q}$
7. $\mathbb{Q}(\sqrt{2}, \sqrt{3})$ *over* $\mathbb{Q}$

*Solution.* Notice that

$$\mathbb{Q}(\sqrt{2}, \sqrt{3}, \sqrt{18}) = \mathbb{Q}(\sqrt{2}, \sqrt{3})$$

since $\sqrt{18} = \sqrt{9}\sqrt{2} = 3\sqrt{2} \in \mathbb{Q}(\sqrt{2})$. Now notice that

$$[\mathbb{Q}(\sqrt{2}) : \mathbb{Q}] = 2$$

since $x^2 - 2$ is irreducible over $\mathbb{Q}$. Lastly we conclude that $\sqrt{3} \notin \mathbb{Q}(\sqrt{2})$ since

$$\begin{aligned} \sqrt{3} &= a + b\sqrt{2} \\ 3 &= a^2 + 2ab\sqrt{2} + 2b^2 \end{aligned}$$

Thus $2ab = 0$ so $a = 0$ or $b = 0$. Both cases work out to show that $a, b \notin \mathbb{Q}$. Therefore

$$[\mathbb{Q}(\sqrt{2}, \sqrt{3}) : \mathbb{Q}(\sqrt{2})] = 2$$

By the tower law:

$$[\mathbb{Q}(\sqrt{2}, \sqrt{3}) : \mathbb{Q}] = 4$$

with a standard basis being $\{1, \sqrt{2}, \sqrt{3}, \sqrt{6}\}$.

Next up we examine $\mathbb{Q}(\sqrt{2}, \sqrt[3]{2})$.

$$\begin{aligned}\sqrt[3]{2} &= a + b\sqrt{2} \\ 2 &= (a + b\sqrt{2})(a^2 + 2b^2 + 2ab\sqrt{2}) \\ 2 &= a^3 + 2ab^2 + 2a^2b\sqrt{2} + a^2b\sqrt{2} + 2b^3\sqrt{2} + 2ab^2\end{aligned}$$

The left side is rational so we require $b = 0$, but then $a = \sqrt[3]{2} \notin \mathbb{Q}(\sqrt{2})$. Hence we conclude that

$$[\mathbb{Q}(\sqrt{2}, \sqrt[3]{2}) : \mathbb{Q}] = 6$$

For problem 7 notice that we have already shown it for problem 3. □

**Problem 23.5** (Problem 31.23 Fraleigh's). *Show that if $E$ is a finite extension of a field $F$ and $[E : F]$ is prime, then $E$ is a simple extension of $F$ and, indeed $E = F(\alpha)$ for every $\alpha \in E \setminus F$.*

*Proof.* We assume $E$ is a finite extension of $F$ and in particular

$$[E : F] = p$$

for some prime $p$. Let $\alpha \in E \setminus F$ then $\alpha$ is algebraic over $F$ so $\alpha$ has a minimal polynomial over $F$ and

$$[F(\alpha) : F] = \deg m_\alpha(x)$$

Now note that

$$F \subseteq F(\alpha) \subseteq E$$

since $\alpha \in E$. By the tower law:

$$[E : F] = [E : F(\alpha)][F(\alpha) : F] = p$$

so one side of the product must be $1$. Since $\alpha \notin F$ it must be the case that

$$[F(\alpha) : F] > 1$$

so $[F(\alpha) : F] = p$, i.e.

$$E = F(\alpha)$$

□

**Problem 23.6** (Problem 31.24 Fraleigh's p.292). *Prove that $x^2 - 3$ is irreducible over $\mathbb{Q}(\sqrt[3]{2})$.*

*Proof.* We can make the simple observation that $\sqrt{3} \notin \mathbb{Q}(\sqrt[3]{2})$. Let $\alpha = \sqrt[3]{2}$. Then

$$\mathbb{Q}(\alpha) = \{a + b\alpha + c\alpha^2 : a, b, c \in \mathbb{Q}\}$$

So if $\sqrt{3} \in \mathbb{Q}(\alpha)$ then

$$\sqrt{3} = a + b\alpha + c\alpha^2$$

Suppose this holds. Since $\mathbb{Q}(\sqrt{3}) \subseteq \mathbb{Q}(\alpha)$ then by the tower law

$$[\mathbb{Q}(\alpha)] = [\mathbb{Q}(\alpha) : \mathbb{Q}(\sqrt{3})] \cdot [\mathbb{Q}(\sqrt{3}) : \mathbb{Q}]$$

But then

$$3 = [\mathbb{Q}(\alpha) : \mathbb{Q}(\sqrt{3})] \cdot 2$$

but $2$ does not divide $3$ so this is impossible. □

**Problem 23.7** (Problem 31.29 Fraleigh's p.292). *Let $E$ be a finite extension of $F$, and let $p(x) \in F[x]$ be irreducible over $F$ and have degree that is not a divisor of $[E : F]$. Show that $p(x)$ has no zeros in $E$.*

*Proof.* Let $\alpha \in E$ be a hypothetical zero of $p(x)$. Since $p$ is irreducible in $F$ and $p(\alpha) = 0$, $p$ must be the minimal polynomial of $\alpha$.

Thus

$$[F(\alpha) : F] = \deg p(x)$$

Using the tower law

$$[E : F] = [E : F(\alpha)][F(\alpha) : F]$$

But then $\deg p$ divides $[E : F]$ which, by assumption, is not the case. Hence we have a contradiction. □

**Problem 23.8** (Problem 31.30 Fraleigh's p.292). *Let $E$ be an extension of $F$. Let $\alpha \in E$ be algebraic of odd degree over $F$. Show that $\alpha^2$ is algebraic of odd degree over $F$, and $F(\alpha) = F(\alpha^2)$.*

*Proof.* We fix some $\alpha \in E$ algebraic of odd degree over $F$. Then

$$[F(\alpha) : F] = 2n + 1, \quad n \in \mathbb{N}$$

Crucially $\alpha \notin F$, by assumption. Now suppose $\alpha^2 \in F$. Then $\alpha^2 \notin F(\alpha) \backslash F$ so $[F(\alpha) : F] = 2$ (since $x^2 - \alpha^2 \in F[x]$ would be a minmal polynomial of $\alpha$), which contradicts our assumption. Thus $\alpha^2 \notin F$. Now since $\alpha^2 \in F(\alpha)$ we see that $\alpha$ satisfies

$$x^2 - \alpha^2 \in F(\alpha^2)[x]$$

Hence $[F(\alpha) : F(\alpha^2)] \leq 2$. By the tower law

$$[F(\alpha) : F] = [F(\alpha) : F(\alpha^2)][F(\alpha^2) : F]$$

We see that then $[F(\alpha) : F(\alpha^2)]$ must be $1$, hence

$$F(\alpha) = F(\alpha^2)$$

□

**Problem 23.9.** *Let $\alpha = \sqrt{2} + \sqrt{3}$.*
*Show that $\alpha$ is algebraic over $\mathbb{Q}$.*
*Show that $\mathbb{Q}(\alpha) = \mathbb{Q}(\sqrt{2}, \sqrt{3})$.*
*Deduce that $\mathbb{Q}(\sqrt{2}, \sqrt{3})$ is a simple extension.*

*Proof.* Clearly $\sqrt{2} + \sqrt{3} \notin \mathbb{Q}$. Now it remains to find a polynomial in $\mathbb{Q}[x]$ for which $\alpha$ is a zero.

$$\begin{aligned} x &= \sqrt{2} + \sqrt{3} \\ x^2 &= 2 + 2\sqrt{2}\sqrt{3} + 3 \\ x^2 - 5 &= 2\sqrt{2}\sqrt{3} \\ \frac{x^2 - 5}{2} &= \sqrt{6} \\ \frac{(x^2-5)(x^2-5)}{4} &= 6 \\ \frac{x^4 - 10x^2 + 25}{4} - 6 &= 0 \\ x^4 - 10x^2 + 1 &= 0 \end{aligned}$$

and $x^4 - 10x^2 + 1 \in \mathbb{Q}[x]$ so $\alpha$ is algebraic over $\mathbb{Q}$.

Thus

$$[\mathbb{Q}(\alpha) : \mathbb{Q}] = [\mathbb{Q}(\sqrt{2}, \sqrt{3}) : \mathbb{Q}] = 4$$

The standard basis for $\mathbb{Q}(\sqrt{2}, \sqrt{3})$ is $\{1, \sqrt{2}, \sqrt{3}, \sqrt{2}\sqrt{3}\}$ which just so happens to be the basis for $\mathbb{Q}(\alpha)$. We can also take some arbitrary element from either extension field and show how to express it in terms of the other to show that they are indeed equal.

Since $\mathbb{Q}(\alpha)$ is simple, it follows from the above observation that

$$\mathbb{Q}(\sqrt{2}, \sqrt{3})$$

is indeed simple. □

## 24 Some problems from Section 33

**Problem 24.1** (Problem 33.4 Fraleigh's). *Find the number of primitive $8$th roots of unity in $GF(9)$ (also known as $\mathbb{F}_9$).*

*Proof.* The multiplicative group $\mathbb{F}_9^x$ has order $8$ since $0$ is excluded. The multiplicative group of a finite field is cyclic so

$$\mathbb{F}_9^x \cong \mathcal{C}_8$$

The $8$th roots of unity are the elements $x \in \mathcal{C}_8$ for which

$$x^8 = 1$$

Since the whole group has order $8$, every nonzero element satisfies this. A primitive $8$th root of unity has order exactly $8$ and the Euler totient function gives

$$\phi(8) = 4$$

□

**Problem 24.2** (Problem 33.9 Fraleigh's p.305). *Let $\overline{\mathbb{Z}}_2$ be the algebraic closure of $\mathbb{Z}_2$, and let $\alpha, \beta \in \overline{\mathbb{Z}}_2$ be zeros of $x^3 + x^2 + 1$ and of $x^3 + x + 1$, respectively. Using the results of this section, show that $\mathbb{Z}_2(\alpha) = \mathbb{Z}_2(\beta)$.*

*Proof.* We begin by showing that the polynomials $f(x) = x^3 + x^2 + 1$ and $g(x) = x^3 + x + 1$ are irreducible over $\mathbb{Z}_2$. These are degree $3$ polynomials and $\mathbb{Z}_2$ is a field so they are irreducible if and only if they have no zeros in $\mathbb{Z}_2$.

$$f(0) = 1 \text{ and } f(1) = 1,$$

and

$$g(0) = 1 \text{ and } g(1) = 1.$$

Thus $f, g$ are irreducible over $\mathbb{Z}_2$. Hence

$$[\mathbb{Z}_2(\alpha) : \mathbb{Z}_2] = [\mathbb{Z}_2(\beta) : \mathbb{Z}_2] = 3$$

So both $\mathbb{Z}_2(\alpha)$ and $\mathbb{Z}_2(\beta)$ are field with degree $2^3$. Since $2$ is prime we can use the finite-field theorem and conclude that these two fields are isomorphic. Furthermore in an algebraic closure of some field $\mathbb{F}_p$, there is exactly one subfield with $p^n$ elements, namely the set of roots of

$$x^{p^n} - x.$$

Thus

$$\mathbb{Z}_2(\alpha) = \mathbb{Z}_2(\beta).$$

□

**Problem 24.3** (Problem 33.10 Fraleigh's p.305). *Show that every irreducible polynomial in $\mathbb{Z}_p[x]$ is a dvisor of $x^{p^n} - x$ for some $n$.*

*Proof.* Let $\alpha \in \overline{\mathbb{Z}}_p$ be the root of some $f(x) \in \mathbb{Z}_p$, irreducible. Then

$$[\mathbb{Z}_p(\alpha) : \mathbb{Z}_p] = \deg f$$

So $\mathbb{Z}_p(\alpha)$ is a finite field with $p^m$ elements. If a field has $p^m$ elements then $a^{p^m} = a$ for all $a$ in said field. Equivalently every elements is a root of

$$x^{p^m} - x.$$

Then

$$\alpha^{p^m} - \alpha = 0.$$

If a polynomial $g$ is the minimal polynomial of $\alpha$ over $\mathbb{Z}_p$, and $\alpha$ is also a root of another polynomial $f$ then

$$g \mid f$$

□

**Problem 24.4** (Problem 3a Exam 2016). *What is an integral domain? Show that $\mathbb{Z}[x]/(x^2+1)$ is an integral domain. Is $\mathbb{Z}[x]/(x^2-1)$ an integral domain? Give reasons for your answer.*

*Solution and Proof.* An integral domain is a non-zero ring with multiplicative identity ($1 \neq 0$) that has no zero divisors.

For $f(x) \in F[x]$, $F[x]/\langle f(x)\rangle$ is an integral domain iff $f(x)$ is irreducible in $F[x]$. Since $x^2+1$ is a degree $2$ polynomial it is irreducible if and only if it has no zeros in $\mathbb{Z}$. $f(x) = x^2+1$ is famously a motivating example for complex numbers with zeros $\pm i \in \mathbb{C}$. A quick sanity check gives $f(0) = 1$, $f(\pm 1) = 2$ and generally $f(n) = f(-n) < f(n+1) = f(-(n+1))$ so it ceirtainly does not have any zeros in $\mathbb{Z}$ (can be proven with induction or standard techniques if necessary). Thus $f$ is irreducible and so $\mathbb{Z}[x]/\langle f(x)\rangle$ is an integral domain.

Using this same procedure we see that $x^2-1$ has zeros $\pm 1 \in \mathbb{Z}$ and is therefore not irreducible so

$$\mathbb{Z}[x]/\langle x^2-1\rangle$$

is not an integral domain. □

# 25 Automorphisms of Field Extensions & Isomorphism Extension

**Definition 25.1** (Field Automorphism). *A **field automorphism** of $E$ is a bijection*

$$\sigma : E \to E$$

*such that $\forall x, y \in E$:*

1. $\sigma(x+y) = \sigma(x) + \sigma(y)$
2. $\sigma(xy) = \sigma(x)\sigma(y)$
3. $\sigma(1) = 1$

If $F \subseteq E$ is a field extension we write

$$\mathrm{Aut}(E/F) := \{\sigma : E \to E \mid \sigma \text{ is a field automorphism and } \sigma|_F = \mathrm{id}_F\}.$$

**Proposition 25.1.** *A field automorphism $\sigma$ preserves algebraic structure:*

1. $\sigma(0) = 0$
2. $\sigma(-x) = -\sigma(x)$
3. $\sigma(x^{-1}) = \sigma(x)^{-1}$ *for* $x \neq 0$

*Proof Sketch.* Straightforward from homomorphism properties. For instance:

$$1 = \sigma(1) = \sigma(xx^{-1}) = \sigma(x)\sigma(x^{-1})$$

and so on. □

**Proposition 25.2.** $\mathrm{Aut}(E/F)$ *is a group under composition.*

**Theorem 25.1.** *Let $E/F$ be an extension, and let $\alpha \in E$ be algebraic over $F$. Let $m_\alpha(x) \in F[x]$ be a minimal polynomial.*
*Then for any $\sigma \in \mathrm{Aut}(E/F)$,*

$$m_\alpha(\sigma(\alpha)) = 0.$$

*Proof.* Write $m_\alpha(x) = \sum_{i=0}^{n} a_i x^i,\ a_i \in F\ (a_n = 1)$.

Since $\sigma$ fixes $F$, we have $\sigma(a_i) = a_i$. Then:

$$0 = \sigma(m_\alpha(\alpha)) = \sigma\left(\sum a_i \alpha^i\right) = \sum a_i \sigma(\alpha)^i = m_\alpha(\sigma(\alpha))$$

□

**Proposition 25.3.** *Let $E = F(\alpha)$, and let $\sigma \in \mathrm{Aut}(E/F)$. Then $\sigma$ is completely determined by $\sigma(\alpha)$.*

**Theorem 25.2.** *Let $E = F(\alpha)$, where $\alpha$ is algebraic over $F$ with minimal polynomial $m_\alpha(x)$.*
*Then*

$$|\mathrm{Aut}(E/F)| \leq \deg m_\alpha.$$

**Definition 25.2.** *A field embedding*

$$\sigma : F \to E$$

*is an injective field homomorphism.*

**Theorem 25.3.** *Let $F \subseteq E = F(\alpha)$, $\sigma : F \to K$ be a field homomorphism, $m_\alpha(x)$ the minimal polynomial of $\alpha$. Then there exists an extension*

$$\tilde{\sigma} : E \to K$$

*with $\tilde{\sigma}(\alpha) = \beta$ if and only if $\beta$ is a root of $\sigma(m_\alpha)(x)$.*

**Theorem 25.4** (Isomorphism Extension Theorem)**.** *Let $\sigma : F \to K$ be a field isomorphism and $\alpha$ algebraic over $F$ with minimal polynomial $m_\alpha(x)$.*
*Let $\beta \in \overline{K}$ be a root of $\sigma(m_\alpha)(x)$.*
*Then $\sigma$ is extends to an isomorphism*

$$\tilde{\sigma} : F(\alpha) \to K(\beta)$$

*with*

$$\tilde{\sigma}(\alpha) = \beta.$$

**Theorem 25.5** (Automorphisms of Finite Fields)**.** *Let $E = \mathbb{F}_{p^n}$. Then*

$$\mathrm{Aut}(E/\mathbb{F}_p) = \langle \phi \rangle$$

*where*

$$\phi(x) = x^p$$

*and*

$$|\mathrm{Aut}(E/\mathbb{F}_p)| = n.$$

## 26 Exam Practice

Henceforth these notes will be mostly revision, even though I haven't covered all topics yet. There will also probably be a lot of problems repeated.

**Problem 26.1** (1 Exam 2019). *a) Is $\mathbb{Z}_{\geq 0}$ equipped with addition a group? Justify your answer.*

*b) What are the abelian groups of order $18$ up to isomorphism?*

*c) What are the cosets of the subgroup $\langle (0,3) \rangle$ in $\mathbb{Z}_2 \times \mathbb{Z}_9$? Describe the quotient*

$$\frac{\mathbb{Z}_2 \times \mathbb{Z}_9}{\langle (0,3) \rangle}$$

*as a product of finite cyclic groups.*

*Proof of (a).* No it is not a group because there are additive inverses for all elements except $0$.

A single counter-example is $1 \in \mathbb{Z}_{\geq 0}$ which has no inverse.

$$1 + x = 0 \Rightarrow x = -1 \notin \mathbb{Z}_{\geq 0}.$$

We can prove it rigorously as follows. $0$ is obviously not the additive inverse of $1$. Suppose $n \in \mathbb{Z}_{\geq 0}$ is not an inverse of $1$. Assume, for a contradiction, that $n+1$ is an inverse of $1$. Then

$$\begin{aligned} 1 + (n+1) &= 0 \\ n &= -2 \end{aligned}$$

From this we conclude that the inverse of $1$ is the inverse of $2$, i.e

$$-1 = -2$$

so $1 = 2$. Hence $n+1$ is not an inverse of $1$. By induction no element of $\mathbb{Z}_{\geq 0}$ is an inverse of $1$. □

*Solution of (b).* We use the fundamental theorem of finitely generated abelian groups. $18$ is finite so we can simply find all ways of expressing $18$ as products of prime powers and we'll be done. We see that

$$18 = 2 \cdot 9 = 2 \cdot 3^2 = 2 \cdot 3 \cdot 3$$

so there are two (up to isomprhism) abelian groups of order $18$, namely

$$\mathbb{Z}_2 \times \mathbb{Z}_9 \text{ and } \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_3.$$

□

*Solution of (c).* $\langle(0,3)\rangle = \{(0,0),(0,3),(0,6)\}$. The left and right cosets coincide since this group is abelian and they are precisely

$$\{x+y : y \in \langle(0,3)\rangle\},\ \ x \in \mathbb{Z}_2 \times \mathbb{Z}_9.$$

Since for $(a,b)$, $b$ is taken mod $9$ only $b$ mod 3 matters. So the left cosets are

$$(a,b) + \langle(0,3)\rangle$$

where $a \in \mathbb{Z}_2$ and $b \in \mathbb{Z}_3$. Denote $H := \langle(0,3)\rangle$. That gives us cosets

$$\begin{gathered}(0,0)+H,\ \ (0,1)+H,\ \ (0,2)+H\\ (1,0)+H,\ \ (1,1)+H,\ \ (1,2)+H\end{gathered}$$

i.e. $6$ cosets, which is what we expect because the order of $\mathbb{Z}_2 \times \mathbb{Z}_9$ is $18$ and $18/|H| = 6$. The quotient groups is abelian and $6 = 2 \cdot 3$ so necessarily

$$\frac{\mathbb{Z}_2 \times \mathbb{Z}_9}{\langle(0,3)\rangle} \cong \mathbb{Z}_2 \times \mathbb{Z}_3.$$

We can also argue by observing the surjective homomorphism $\phi : \mathbb{Z}_2 \times \mathbb{Z}_9 \to \mathbb{Z}_2 \times \mathbb{Z}_3$, $\phi(a,b) = (a, b \mod 3)$ has kernel $\ker\phi = H$ so the result follows by the first isomorphism theorem. $\square$

**Problem 26.2** (2 Exam 2019). *a) Let $\sigma = (2\ 6\ 1\ 4\ 3)$ and $\tau = (5\ 2\ 1\ 3)$ be permutations in $S_6$. Express the composition $\sigma\tau$ in disjoint cycle notation. What is the order of $\sigma\tau$?*

*b) Show that if $\sigma \in S_n$ is a cycle of odd length $2k+1$, then $\sigma^2$ is also a cycle of length $2k+1$.*

*c) Let $G$ be a subgroup of the symmetric group $S_n$ such that there exist permutations $\sigma_2, \dots, \sigma_n \in G$ satisfying $\sigma_i(1) = i$ for $2 \le i \le n$. Show that $G$ acts* ***transitively*** *on $\{1, \dots, n\}$. In other words, show that for each pair $1 \le i, j \le n$ there exists a permutation $\sigma \in G$ such that $\sigma(i) = j$.*

*Solution of (a).* $\sigma$ is $2 \to 6 \to 1 \to 4 \to 3 \to 2$ while $\tau$ is $5 \to 2 \to 1 \to 3$. We work out $\sigma\tau$ like this:

$$\begin{gathered}5 \to 2 \to 6 \Rightarrow 5 \to 6\\ 2 \to 1 \to 4 \Rightarrow 2 \to 4\\ 1 \to 3 \to 2 \Rightarrow 1 \to 2\\ 3 \to 5 \to 5 \Rightarrow 3 \to 5\\ 4 \to 4 \to 3 \Rightarrow 4 \to 3\\ 6 \to 6 \to 1 \Rightarrow 6 \to 1\end{gathered}$$

so $\sigma\tau = (1\ 2\ 4\ 3\ 5\ 6)$. The order of an $n$-cycle is $n$ so $|\sigma\tau| = 6$. $\square$

*Proof of (b).* Suppose $\sigma$ is a cycle of length $2k+1$, i.e.

$$\sigma = (a_1 \ a_2 \ \cdots \ a_{2k+1}).$$

Composing $\sigma^2$ sends $a_1 \to a_3$, $a_2 \to a_4$ and so on. In particular $a_i \to a_{(i+2)}$ where $i+2$ is taken mod $2k+1$. Then all odd elements are taken to odd elements, but the $2k+1$-th element is sent to $a_2$ which is sent to $a_4$ and so on. Even elements are sent to even elements until $a_{2k} \to a_1$ so

$$\sigma^2 = (a_1 \ a_3 \ \cdots \ a_{2k+1} \ a_2 \ a_4 \ \cdots \ a_{2k})$$

which has length $2k+1$. □

*Proof of (c).* Since $G$ is a group, each permutation $\sigma_k$ has an inverse $\sigma_k^{-1}$ which sends $k$ to $1$. Also $e \in G$ so for $1 \leq i, j \leq n$ we can send $i$ to $j$ by sending $i$ to $1$ and then $1$ to $j$ so

$$(\sigma_j \circ \sigma_i^{-1})(i) = \sigma_j(1) = j.$$

This works well for the case where neither $i, j$ are $1$. If $i$ is one then let $\sigma_i := e$ and likewise for $j$. □

**Problem 26.3** (3 Exam 2019). *a) Show that if $n$ is not prime, then $n\mathbb{Z}$ is not a prime ideal of $\mathbb{Z}$.*

*b) Show that the quotient*

$$\frac{\mathbb{Z}_2[x]}{\langle x^3 + x + 1 \rangle}$$

*is a field. How many elements does it contain?*

*Proof.* Suppose $n$ is not prime. Then $n = ab$ for some integers $a, b$ with $1 < a < n$ and $1 < b < n$. Hence

$$ab = n \in n\mathbb{Z}.$$

However, $a \notin n\mathbb{Z}$ and $b \notin n\mathbb{Z}$, since neither $a$ nor $b$ is divisible by $n$. Thus there exist $a, b \in \mathbb{Z}$ such that $ab \in n\mathbb{Z}$ but $a, b \notin n\mathbb{Z}$. Therefore $n\mathbb{Z}$ is not a prime ideal of $\mathbb{Z}$. □

*Proof of (b).* We will use the fact that the quotient (in this case) is a field if and only if $f(x) = x^3 + x + 1$ is irreducible over $\mathbb{Z}_2$. Notice that

$$f(0) = f(1) = 1 \neq 0$$

and $f(x)$ is degree $3$ hence irreducible over $\mathbb{Z}_2$. Thus the quotient is a field. There are $2^3 = 8$ elements. □