

# Analysis of silicone contamination of batches

Thom Hurks	<i>0828691</i>
Puck Mulders	<i>0737709</i>
Marijn van Knippenberg	<i>0676548</i>
Rik Coenders	<i>0777004</i>

May 24, 2016

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Methods</b>	<b>3</b>
2.1	Critical value . . . . .	3
2.2	Normal distribution . . . . .	3
2.3	Non-parametric tests . . . . .	4
2.3.1	Wilcoxon . . . . .	4
2.3.2	Serial correlation test on the means . . . . .	4
<b>3</b>	<b>Results</b>	<b>4</b>
3.1	QQ Plot and Shapiro-Wilk . . . . .	4
3.2	Normal distribution . . . . .	6
3.2.1	Shapiro-Wilk on log-transformed data . . . . .	6
3.3	Skewness and Kurtosis . . . . .	8
3.3.1	T-Test . . . . .	9
3.3.2	Homogeneity tests . . . . .	10
3.4	Non-parametric tests . . . . .	11
3.4.1	Serial correlation tests with means . . . . .	11
3.4.2	Wald-Wolfowitz Runs Test . . . . .	11
3.4.3	Rank Serial Correlation Test . . . . .	11
<b>4</b>	<b>Conclusion</b>	<b>11</b>

## 1 Introduction

The health inspection has detected a silicone leakage in a machine used in a chemical production process. The leakage was detected in close vicinity of packed product, so the question is whether the product has been affected by

the leakage. The machines are regularly inspected, and data on a batch produced before the previous inspection is available and is denoted as batch B0. Batches denoted B1 to B7 have been produced between the previous inspection and the current inspection and may have been affected by the leakage. The data consists of measurements of samples of units from the batches. The silicone that leaked is already part of the product itself, so it will always be present in the measured units to some extent. The goal of this report is to possibly identify the batches that were affected by the silicone leakage.

In the first section we describe which methods we use to find the affected batches. In the next section we will describe what the results of these methods are. Lastly, we will conclude which of the batches are affected. In this section we will also describe what the weaknesses and risks are of our investigation.

## 2 Methods

To decide which of the batches are affected, if any, we perform both parametric and non-parametric tests. The parametric tests assume that the data is normally distributed, which means we must test for normality and if necessary transform the data. Non-parametric tests are generally less powerful than parametric tests, but can always be applied regardless of normality. With parametric tests one must be careful to draw conclusions about the original non-transformed data if the data needs to be transformed in order to meet the normality assumption. As a start, we will plot the data and perform a Shapiro-Wilk test to see whether the non-transformed data is normal. After that, we will perform a Wald-Wolfowitz Runs Test which has as a null hypothesis that each element in the sequence is independently drawn from the same random distribution; if this is not the case, that could be an indicator that some of the batches are affected by silicones. We reject the null hypothesis if  $p < 0.01$ .

### 2.1 Critical value

Most of our hypothesis include multiple comparisons. Because of this reason, we have to adjust the critical value to make sure that it will not reject batches too soon. Applying Bonferroni will give us an  $\alpha = \frac{0.05}{7} = 0.00714$ , but since this value is quite low, it can be the case that too many batches will pass the test. Because we are dealing with dangerous materials, we will raise the alpha value slightly to prevent batches from passing the test even though they could be infected. To be a bit more strict, we will set the critical value to 0.01.

### 2.2 Normal distribution

To decide whether the data is normally distributed, we will plot the data to get an insight in how the data is distributed. We will then perform the Shapiro-Wilk tests on each of the batches to check whether the data is now normally distributed. We formulate the hypothesis as following:

$$\begin{cases} H_0 : B_i \sim N & \text{for } 1 \leq i \leq 7 \\ H_1 : B_i \not\sim N \end{cases} \quad (1)$$

We will reject  $H_0$  if  $p < 0.01$ . Besides a Shapiro-Wilk test, we will perform skewness and kurtosis tests to test on where normality is violated. The values for skewness should be between -0.03 and 0.3, while the values of kurtosis should be between -0.5 and 1.5. We will perform a Jarque-Bera test to see if it is appropriate to reject a normal distribution based on skewness and kurtosis. The null hypothesis is the same as with the Shapiro-Wilk test, which can be found in equation 1. If, however, the data is not normally distributed, we will apply a transformation to remove skewness and bring the data closer to a normal distribution. If a transformation is necessary, we must be careful to draw conclusions about the original untransformed data, however. Because batch 0

is produced before the previous inspection where no leakage was discovered, we know that the mean and variance of the amount of silicones found in each batch should be similar to the amount of silicone in batch 0. Therefore our hypothesis is the following:

$$\begin{cases} H_0 : \mu_0 = \mu_i & \text{for } 1 \leq i \leq 7 \\ H_1 : \mu_0 \neq \mu_i & \text{for } 1 \leq i \leq 7 \end{cases} \quad (2)$$

If the result of the t-test is less than 0.01, we will reject the null hypothesis.

Besides performing a t-test, we will also perform some homogeneity tests to see whether the data also share the same characteristics. To check if this is the case, we will perform a Barlett's test, which checks whether all the variances of the batches are equal to each other. The null hypothesis of that test is that all variances are equal, and thus that the sets are homogeneous. We will also perform an F-test to see which batches have similar properties as batch 0.

## 2.3 Non-parametric tests

### 2.3.1 Wilcoxon

### 2.3.2 Serial correlation test on the means

We will perform a serial correlation test on the means of the batches. The serial correlation test is Von Neumann with lag-1 autocorrelation. The null hypothesis is that the means of the batches are random. If this indeed is the case, this is an indication that none of the batches is affected by the silicones. We will reject the null hypothesis if  $p < 0.05$ .

## 3 Results

### 3.1 QQ Plot and Shapiro-Wilk

The first step is to generate a normal QQ plot for batch 0 as an easy visual confirmation of normality.




Figure 1: QQ Plot of batch 0

The normal QQ plot already seems to suggest that batch 0 is not normally distributed. When we also perform the Shapiro-Wilk normality test we can see the  $p\text{-value} = 0.0013049 < 0.01$  so we can reject the null-hypothesis that the outcome is normally distributed.

Next we can generate a normal QQ plot for the other, possible affected, batches 1 to 7. Of course generating this plot for multiple batches at once assumes that the units are somewhat consistent across different batches.




Figure 2: QQ Plot of batches 1 to 7

The normal QQ plot here too seems to suggest that the data is not normally distributed. When we again perform the Shapiro-Wilk normality test we can see the  $p\text{-value} = 8.1716549 \times 10^{-12} < 0.01$  so we can reject the null-hypothesis that the outcome of batches 1 to 7 is normally distributed.

## 3.2 Normal distribution

### 3.2.1 Shapiro-Wilk on log-transformed data

In the previous section we observed that the data does not seem to follow a normal distribution. In order to perform parametric tests that assume normality, we must then transform the data. We will transform the data using the

well-known log transformation and then check for each batch what the p-value of the Shapiro-Wilk test is in order to confirm that the transformation had the desired effect.

	batch	p.value
1	B0	0.04
2	B1	0.14
3	B2	0.62
4	B3	0.66
5	B4	0.90
6	B5	0.06
7	B6	0.21
8	B7	0.96

No of the batches reject normality, so using the Shapiro-Wilk test we cannot reject normality for all of the batches. We can generate a normal QQ-plot to get more insight into the log transformed batch 0.




Figure 3: QQ Plot of the log of batch 0

### 3.3 Skewness and Kurtosis



	kurtosis	skewness
B0	2.596	1.780
B1	0.369	1.254
B2	-0.025	1.043
B3	-0.128	0.904
B4	2.681	1.479
B5	3.552	1.965
B6	6.339	2.429
B7	1.835	1.377

	Jarque.Bera
B0	0.06
B1	0.30
B2	0.41
B3	0.50
B4	0.13
B5	0.02
B6	0.00
B7	0.26

After performing the Shapiro-Wilk test, we have performed skewness and kurtosis tests. We can compute the skewness and kurtosis metrics and observe that for batch 0 they are respectively. This means that although the kurtosis is in the appropriate range to be normally distributed, the skewness of the data is a bit too high. This kind of test is also formalized in the Jarque-Bera test, which tests if the skewness is close to zero and the kurtosis close to three. The p-value of the Jarque-Bera test is , so it cannot reject the normality hypothesis based on skewness and kurtosis. The data is skewed right, which is probably the reason that the Shapiro-Wilk test rejects the assumption of a normal distribution. The p-value of the Shapiro-Wilk test is not very low and the skewness, kurtosis and Jarque-Bera tests seem to suggest a close-to-normal shape. In combination with the fact that for all the other batches we cannot reject the normality null hypothesis, we will assume that batch 0 is normally distributed for the following tests. It must also be noted that the t-test is robust against data that deviates somewhat from normality, so our assumption at least for that test is reasonable.

### 3.3.1 T-Test

We can use a t-test to compare the batches, since we could not reject the null hypothesis that the log-transformed batches are normally distributed. To see which t-test variant we need to perform, we need to analyse the variance of all the log-transformed batches.

The variance differs between batches, as can be seen in the preceding table. Because of that, we will perform a t-test that assumes unequal variances. It must be noted that batch seven has one missing value.

	batch	variance
1	B0	0.940
2	B1	0.315
3	B2	1.447
4	B3	0.683
5	B4	0.156
6	B5	0.212
7	B6	0.383
8	B7	0.437

The results of the t-tests can be found in the following table:

	batch	p.value
1	B1	0.40
2	B2	0.03
3	B3	0.00
4	B4	0.12
5	B5	0.12
6	B6	0.47
7	B7	0.03

The results indicate that batch three has a mean that is significantly different from the mean of the unaffected batch, since its p-value is less than 0.01. This could indicate that this batch is affected.

### 3.3.2 Homogeneity tests

Performing the Barlett's tests gives us a p-value of  $4.5232441 \times 10^{-15} < 0.01$ . This rejects the null hypothesis that all the batches have the same variance. After performing an F-test that compares all the batches with batch 0, we have found the following results:

	F.test
B1	0.083
B2	0.486
B3	0.605
B4	0.006
B5	0.020
B6	0.152
B7	0.239

These results indicate that batch four has different properties from the other batches.

### 3.4 Non-parametric tests

#### 3.4.1 Serial correlation tests with means

We have performed a Rank von Neumann Test for lag-1. The p-value is  $0.4895337 < 0.01$  and we therefore cannot reject the null hypothesis. This indicates that no significant pattern can be found. This could indicate that the silicones have affected only one or two batches and not all batches from the point the leakage has started.

#### 3.4.2 Wald-Wolfowitz Runs Test

We have performed a Wald-Wolfowitz Runs test. We see that the p-value  $= 7.1225335 \times 10^{-5} < 0.01$  so we can reject that null hypothesis. This is an interesting result, because we know the data is obtained from the same source. This indeed indicates that some of the outcomes

#### 3.4.3 Rank Serial Correlation Test

Here we perform the rank serial autocorrelation test at lag 1 using the rank von Neumann ratio. This tests the null hypothesis that the lag- $k$  autocorrelation is 0 for all values of  $k$  greater than 0 (i.e., the time series is purely random). The procedure that we use emits some warnings because the dataset contains ties, and with this specific test that can make the p-value less accurate. With that warning in mind, we present the table containing the rank serial autocorrelation test p-values of each batch:

	batch	p.value
1	B0	0.99
2	B1	0.13
3	B2	0.26
4	B3	0.36
5	B4	0.68
6	B5	0.21
7	B6	0.49
8	B7	0.37

As one can observe, the p-values are all large, which means that we cannot reject the null hypothesis that the batches are random using this test.

## 4 Conclusion