
Australian Government
IP Australia

Plant Varieties Journal - Optimised for Screen Viewing



Plant Varieties Journal

Official Journal of Plant Breeder's Rights Office, IPAustralia

Quarter Four 2007

Volume 20 Number 4

ISSN: 1030-9748

Date of Publication : 6 Mar 2008

- Home
- Part 1 General Information
- Part 2 Public Notices
- Part 3 Appendices
- Subscribe

Australian Government
IP Australia

Part 1 General Information

Part 1 of ***Plant Varieties Journal*** provides the link with the General Information about the Plant Breeder's Rights scheme, the procedures for objections and revocations, UPOV developments, Important Changes etc. The General Information pages of ***Plant Varieties Journal*** **(Vol. 20 Issue 4)** are listed below:

- Home
- Interactive Variety Description System (IVDS)
- Objections and revocations
- Report on Breeding Issues
- Use of Overseas Data
- PBR Infringement
- On-line Database for PBR Varieties
- Cumulative Index to Plant Varieties Journal
- Applying for Plant Breeder's Rights
- Requirement to Supply Comparative Varieties
- UPOV Developments
- European Developments
- Obligation under the International Convention for the Protection of New Varieties of Plants 1991 (UPOV91)
- Instructions to Qualified Persons
- Current PBR Forms

Interactive Variety Description System (IVDS)

For preparing the detailed description, the Plant Breeder's Rights Office (PBRO) has released the Interactive Variety Description System (IVDS) in the Internet (https://pbr-ivds.ipaustralia.plantbreeders.gov.au/pbr_ivds/) for the Qualified Persons (QPs).

In the beginning of April 2005, all QPs have officially been notified of this new system giving them access to IVDS with their individual user name and password. The main purpose of the system is to harmonise variety descriptions at both national and international level and make the PBR application process as smooth and efficient as possible.

The IVDS allows QPs to fill in descriptions on-line by accessing relevant test guidelines and selecting specific characteristics with their various states of expressions from the options provided. The IVDS incorporated all of the approved UPOV test guidelines (and some national equivalents where a UPOV test guideline is not available) into interactive forms with easy to use drop-down menus. QPs can "build" their own additional/special characteristics if they are not available in the guideline. The IVDS also accepts statistical information.

The IVDS emphasises the use of "grouping characteristics" in selecting comparator varieties. Finally, it allows QPs to lodge the completed variety descriptions on-line. There is a minimum typing involved in the process.

The PBRO anticipates that the QPs had the opportunity to familiarise themselves with IVDS during the testing and demonstration phase (August – Dec 2004) and could operate the system comfortably. There are step by step on-screen instructions with examples in each step of IVDS, which will assist the QPs to complete the process smoothly. In addition, PBRO is ready to help QPs, if they encounter any problem. Please send an e-mail to pbr@ipaustralia.gov.au if there is a problem in completing the description using IVDS.

Objections and revocations

Objections to Applications and Requests for Revocation of a Grant or of a Declaration that a Plant Variety is Essentially Derived from Another Plant Variety

The Plant Breeder's Rights scheme is administered consistent with the model law of the *International Convention for the Protection of New Plant Varieties 1991* (UPOV 91), that is, applicants are entitled to protection, in the absence of proof to the contrary.

The Plant Breeder's Rights Office (PBRO) is not required to advocate for the views, assertions, and opinions of persons challenging an application for plant breeder's rights. Those objecting to applications, requesting revocation of a grant, or seeking a declaration that a plant variety is essentially derived from another plant variety should provide sufficient probative evidence to enable the Secretary to be satisfied of their validity of their claims. It cannot be stressed too strongly that all available evidence ought to accompany the application for objection/revocation/declaration at the outset.

Occasionally the PBRO receives comments on applications. The PBRO seeks to give effect to the processes set out in the PBR Act. The Act provides for a formal objection process, and comments are not formal objections. Where members of the public genuinely believe their commercial interests would be affected and that PBR for a proposed variety ought not to be granted, they are encouraged to use the Act's processes, eg. lodging an objection. Comments are simply informal information from the public to a governmental decision maker. The PBRO will generally not engage in further communication with the commentator regarding their comment, although the comment may be valuable in alerting the PBRO to an important matter of which it was previously unaware.

Objections to Applications

A person may make objections to applications for PBR if (i) their commercial interests would be affected adversely, and (ii) the application will not fulfil all the conditions required by the Plant Breeder's Rights Act.

Objections to applications must be lodged with the Registrar no later than six months after the date the description of the variety is published in this journal. The objector must provide evidence of adverse affect on their commercial interests and that the application should not be granted.

The Registrar of the Plant Breeder's Rights Office (PBRO) is required to give a copy of the objection to the applicant. The objection is also available to the general public on request. The applicant has the opportunity to respond to the evidence presented. The Registrar then decides whether or not the objection will be upheld and, subsequently, whether the application will be granted. The PBRO is under no obligation to enter into further dialogue regarding an objection or to communicate reasons why an objection is not upheld. If an objection is upheld it will be notified in this journal.

A payment of $100 is required on lodgement of the objection. Additional costs of $75 per hour for work undertaken in relation to the objection will be billed to the objector.

Requests for Revocation, (where an individual's interests are affected) of:

· a Grant

· a Declaration that a Plant Variety is Essentially Derived

A person may, when their interests are affected adversely, apply for the revocation of:

· a grant of PBR; or

· a declaration that a plant variety is essentially derived from another plant variety.

The person requesting revocation is required to lodge a revocation payment fee of $500. The person seeking revocation of a grant or declaration that a plant variety is essentially derived from another plant, must provide conclusive evidence of adverse affect on their interests and that the grant should be revoked.

The PBRO also accepts information regarding revocation of grants and declarations of essentially derived plant varieties. Such information must demonstrate conclusively that a grant or declaration should not have been made. All written information will be acknowledged. The PBRO is under no obligation to enter into further communication regarding information provided.

Report on Breeding Issues

A report providing greater clarification of certain ‘difficult’ and sometimes controversial plant breeding issues has been finalised by a panel of experts. The report defines ‘discovery’, ‘selective propagation’ and ‘eligible breeding’ methodologies as well as canvassing questions and answers to a range of situations. The principal areas covered are the source population and associated issues relating to ownership, location, homogeneity, parentage, boundaries, and selection from variable material. The issue of essentially derived varieties and the relationship between the first and the second breeder(s) is also explored. The final report of the expert panel is available now.

Use of Overseas Data

Overseas Testing/Data

The PBR Act allows DUS data produced in other countries (overseas data) be used in lieu of conducting a comparative trial in Australia provided certain conditions are met; relating to the filing of applications, sufficiency of the data and the likelihood that the candidate variety will express the distinctive characteristic(s) in the same way when grown locally. Briefly the overseas data could be considered where:

- The first PBR application relating to the candidate variety has been lodged overseas, and
- the variety has previously been test grown in a UPOV member country using official UPOV test guidelines and test procedures, (i.e. equivalent to a comparative trial in Australia) and
- either, all the most similar varieties of common knowledge (including those in Australia) have been included in the overseas DUS trial, or
- the new overseas variety is so clearly distinct from all the Australian varieties of common knowledge that further DUS test growing is not warranted, and
- sufficient data and descriptive information is available to publish a description of the variety in an accepted format in Plant Varieties Journal; and to satisfy the requirements of the PBR Act.

Taxa that must be trailled in Australia

It is the policy of PBR office to not accept overseas data for the following taxa due to the wide genotype by environment interactions that have been previously experienced. Varietal descriptions from overseas trials have consistently been different from those obtained from trials grown under Australian conditions. Consequently, for the following taxon a full PBR trial must be conducted in Australia:

Solanum tuberosum Potato

The Qualified Person, in consultation with the agent/applicant, and perhaps other specialists and taxonomists, will need to evaluate the overseas data, test report and photographs to see if the application does fulfil all PBR Office requirements, and then advise the agent/applicant:

- either, to submit Part 2 incorporating a description for publication, any additional data and photographs and to pay the examination fee;
- or, to conduct a DUS trial in Australia, recommending to the applicant/agent which additional varieties of common knowledge to include;

- or, submit Part 2 including additional data (information about similar varieties in Australia to show that they are clearly distinct from the candidate variety that a further DUS test growing including the similar varieties is not warranted and that the variety displays the distinctive characteristics when grown in Australia)

Please note that the PBR office does not obtain overseas DUS test reports on behalf of applicants. It is the sole responsibility of the applicants to obtain these reports directly from the relevant overseas testing authorities. Where applicants already have the report they are advised to submit a certified true copy of the report with the Part 1 application. Applicants, or those duly authorised, may certify the copy.

If you do not have the test report available at the time of Part-1 application then you are advised to submit the Part-1 application without the test report. However, you should make arrangements to procure the DUS test report directly from the relevant testing authority. When the report becomes available, a certified copy should be supplied to the QP and the PBR office.

When the trial is based on an UPOV technical guideline and test report in an official UPOV language (English, German or French), it can be lodged in support of the application. In other cases the test reports must be in English.

The applicant/agent and Qualified Person should use the overseas test report to complete Part 2 of the application, making a decision on how to proceed in view of the completeness of the information, the comparators (if any) used in the overseas DUS trial and their knowledge of similar Australian varieties that may not have been included in the overseas test report.

If a description is based on an overseas test report, Australian PBR will not be granted until after the decision to grant PBR in the country producing the DUS test is made. The final decision on the acceptability of overseas data rests with the PBR office.

PBR Infringement

Grantees should be aware of recent revisions to infringement provisions of the *Plant Breeder's Rights Act 1994* (see section 54) and related provisions of the Federal Court Rules (see order 58 rule 27) both of which can be found at the ComLaw site

On-line Database for PBR Varieties

The PBR Office has a comprehensive service for Internet users ~ a searchable database for all Australian PBR varieties, both past and present. The database features a detailed description and image for every variety granted full rights and basic information for other PBR varieties. Searches by genus, species, common name, variety name and titleholder are some of its many advantages. Varieties for which an application has been lodged but not yet accepted in the PBR scheme are not included in this database. Please browse the Plant Breeder's Rights on-line database and provide your feedback.

Cumulative Index to Plant Varieties Journal

The cumulative index to the *Plant Varieties Journal* has been updated to include variety information from all hardcopy versions up to volume 16 issue 3. After that issue the Plant Varieties Journal is only published in the electronic format and there is no need for a cumulative index, as the variety information can be easily searched in the PBR online database and also by downloading the *Plant Varieties Journal* electronically.

The final updated version of the cumulative index is available in PBR website. This document has information up to Plant Varieties Journal volume 16 issue 3. The PBR office recommends use its PBR online database to get most updated information on variety registration. The online database is updated on a weekly basis.

Applying for Plant Breeder's Rights

Applications are accepted from the original breeder of a new variety (from their employer if the breeder is an employee) or from a person who has acquired ownership from the original breeder. Overseas breeders need to appoint an agent to represent their interests in Australia. Interested parties should contact the PBR office and an accredited Qualified Person experienced in the plant species in question.

Steps in Applying for Plant Breeder's Rights

- Obtain from the breeder a signed Authorisation to act as their agent in Australia for the variety in question if your role is as the Australian agent of an overseas breeder;
- Complete Part 1 of the application form, supplying a photograph of the new variety, paying the application fee, nominating an accredited 'Qualified Person' and, if the variety is an Australian species, despatch as soon as possible a herbarium specimen;
- Engage the services of the nominated accredited 'Qualified Person' to plan and supervise the comparative growing trial;
- Conduct a comparative growing trial to demonstrate Distinctness, Uniformity and Stability (DUS), complete Part 2 of the application form and paying the examination fee;
- Deposit propagating material in a Genetic Resources Centre.
- Examination of the application by the PBR Office, which may include a field examination of the comparative growing trial; and including
- Publication of a description and photograph comparing the new variety with similar varieties in Plant Varieties Journal, followed by a six-month period for objection or comment.
- Upon successful completion of all the requirements, resolution of objections (if any) and payment of certificate fee, the applicant(s) receive a Certificate of Plant Breeder's Rights.

Requirement to Supply Comparative Varieties

Once an application has been accepted by the PBR office, it is covered by provisional protection. Also it immediately becomes a ‘variety of common knowledge’ and thus may be required by others as a comparator for their applications with a higher application number.

Applicants are reminded that they are required to release propagative material for comparative testing provided that the material is used for no other purpose and all material relating to the variety is returned when the trial is complete. The expenses incurred in the provision of material for comparative trials are borne by those conducting the trials.

As the variety is already under provisional protection, any use outside the conditions outlined above would qualify as an infringement and would be dealt with under section 53 of the *Plant Breeder’s Rights Act 1994.*

Applicants having difficulties procuring varieties for use in comparative trials are urged to contact the PBR office immediately

UPOV Developments

The UPOV Convention provides the international legal framework for the granting of plant breeders' rights which are a key element in encouraging breeders to pursue and enhance their search for improved varieties with benefits such as higher yield and quality and better resistance to pests and diseases. Plant breeders' rights thereby help to enhance sustainable agriculture, productivity, income, international trade and economic development in general.

The members of UPOV are (as of November 18, 2007):

Albania, Argentina, Australia, Austria, Azerbaijan, Belarus, Belgium, Bolivia, Brazil, Bulgaria, Canada, Chile, China, Colombia, Croatia, Czech Republic, Denmark, Dominican Republic, Ecuador, European Community, Estonia, Finland, France, Germany, Hungary, Iceland, Ireland, Israel, Italy, Japan, Jordan, Kenya, Kyrgyzstan, Latvia, Lithuania, Mexico, Morocco, Netherlands, New Zealand, Nicaragua, Norway, Panama, Paraguay, Poland, Portugal, Republic of Korea, Republic of Moldova, Romania, Russian Federation, Singapore, Slovakia, Slovenia, South Africa, Spain, Sweden, Switzerland, Trinidad and Tobago, Turkey, Tunisia, Ukraine, United Kingdom, United States of America, Uruguay, Uzbekistan and Vietnam. (Total 65).

On October 18, 2007 Turkey deposited with the Office of the Union its instrument of accession to the 1991 Act of the UPOV Convention. The 1991 Act entered into force for Turkey on November 18, 2007. On that day, Turkey became the 65th member state of UPOV.

Further Information on UPOV and its activities is available on the website located at http://www.upov.int

The adopted UPOV Technical Guidelines (TG) for testing different plant species are now available for this website at
http://www.upov.int/en/publications/tg-rom/index.html

European Developments

Community plant variety rights within the European Union are administered by the Community Plant Variety Office (CPVO) in Angers, France. With more than 2,600 applications per year, the CPVO receives the highest number of requests for variety protection among the 63 members of UPOV. The CPVO provides for one application, one examination and one title of protection that is valid and enforceable in all 25 members of the European Union.

The potential applicants for Plant Variety Rights within European Union are requested to consult Notes for Applicants published by the Community Plant Variety Office (CPVO). This note aims to answer legal, administrative and financial questions that one may have when requesting Community plant variety rights. Further information is available from CPVO website.

Obligation under the International Convention for the Protection of New Varieties of Plants 1991 (UPOV91)

Consistent with Australia's membership of UPOV 1991, the criteria for the granting of protection under the *Plant Breeder's Rights Act 1994* (PBRA) is that the variety: has a breeder; is new, distinct, uniform and stable; has an acceptable name; and that application formalities are completed and relevant fees payed.

Applicants for protection need to be aware of the existence of any other Australian legislation, which could impact on their intended use of the registered variety. Administrators of other Australian legislation may have an interest in applications for registration notified in this journal.

It is feasible for a new variety to be registered under the PBRA, but, as the PBRA co-exists with other laws of the land, the exercise of the breeder's right may be restricted by such legislation. For example, current legislation may prohibit the use of that variety in food, or, the growing of that variety as a noxious weed.

The Plant Breeder's Rights Office (PBRO) advises that it is the responsibility of the applicant and of administrators of legislation to take these matters up directly between the responsible parties and not with the PBRO.

Instructions to Qualified Persons

Instruction to Qualified Persons: Interactive Variety Description System (IVDS) for Preparing Detailed Description for Plant Varieties Journal

For preparing the detailed description, the Plant Breeder's Rights Office (PBRO) has released the Interactive Variety Description System (IVDS) in the Internet (https://pbr-ivds.ipaustralia.plantbreeders.gov.au/pbr_ivds/) for the Qualified Persons (QPs).

In the beginning of April 2005, all QPs have officially been notified of this new system giving them access to IVDS with their individual user name and password. The main purpose of the system is to harmonise variety descriptions at both national and international level and make the PBR application process as smooth and efficient as possible.

The IVDS allows QPs to fill in descriptions on-line by accessing relevant test guidelines and selecting specific characteristics with their various states of expressions from the options provided. The IVDS incorporated all of the approved UPOV test guidelines (and some national equivalents where a UPOV test guideline is not available) into interactive forms with easy to use drop-down menus. QPs can "build" their own additional/special characteristics if they are not available in the guideline. The IVDS also accepts statistical information.

The IVDS emphasises the use of "grouping characteristics" in selecting comparator varieties. Finally, it allows QPs to lodge the completed variety descriptions on-line. There is a minimum typing involved in the process.

The PBRO anticipates that the QPs had the opportunity to familiarise themselves with IVDS during the testing and demonstration phase (August – Dec 2004) and could operate the system comfortably. There are step by step on-screen instructions with examples in each step of IVDS, which will assist the QPs to complete the process smoothly. In addition, PBRO is ready to help QPs, if they encounter any problem. Please send an e-mail to pbr@ipaustralia.gov.au if there is a problem in completing the description using IVDS.

The detailed descriptions are accepted only in the IVDS format.

Also, please note that the after finalising the description through IVDS, the QPs will still need to submit the signed hardcopies of the Part 2 documentations in order to complete the application process. Please contact the PBRO (pbr@ipaustralia.gov.au) for further information.

Current PBR Forms

As part of a comprehensive review of PBR forms, several are now available in fillable WORD format and can be completed electronically and saved. Currently, only the Part 1 Application, Supplementary Pages to Part 1 Application, Authorisation of Agent and Nomination of Qualified Person forms are available in fillable WORD.

We are endeavouring to have all forms in both fillable WORD and fillable PDF in the near future and will continue to update this list. Please check regularly for updates.

The remainder of the forms and publications are static PDFs and may be viewed using Acrobat Reader. The electronic forms are available from the IP Australia Website at http://www.ipaustralia.gov.au/pbr/forms.shtml

Please Do Not Use Old Forms

To avoid processing delays, it is recommended that the most recent version of a form be submitted. Refer to the PBR website for the latest version of the forms. Please note applications submitted on old forms will be returned so they can be submitted on current forms for assessment.

Australian Government
IP Australia

Part 2 Public Notices (Acceptances, Descriptions, Grants, Variations etc)

This part of the *Plant Varieties Journal* provides public notices on Acceptances, Variety Descriptions, Grants, Variations etc. The Part 2 Public Notices pages of ***Plant Varieties Journal*** (**Vol. 20 Issue 4**) are listed below:

- **Home**
- **Acceptances**
- **Variety Descriptions**
- **Grants**
- **Denomination Changed**
- **Synonym Changed**
- **Applicant's Name Amended**
- **Change of Agent**
- **Assignment of Rights**
- **Transfer of Rights**
- **Grants Surrendered**
- **Applications Withdrawn**
- **Corrigenda**

ACCEPTANCES

The following varieties are under provisional protection from the date of acceptance:

Actinotus helianthi

FLANNEL FLOWER

‘Shooting-Star’

Application No: 2007/301 Accepted: 12 December, 2007
Applicant: **Louise (AKA Lana) Helena Mitchell**, Gundaroo, NSW.

Anthurium andraeanum

FLAMINGO FLOWER

‘ANTHEPCI’ syn Amis

Application No: 2007/313 Accepted: 21 December, 2007
Applicant: **Anthura b.v.**.
Agent: **Sprint Horticulture Pty Ltd**, Wamberal, NSW.

Avena sativa

OATS

‘Dawson’

Application No: 2007/241 Accepted: 7 November, 2007
Applicant: **NDSU Research Foundation**.
Agent: **Pacific Seeds Pty Ltd**, Toowoomba, QLD.

Calothamnus quadrifidus

ONE SIDED BOTTLEBRUSH

‘Calgreen1GL’

Application No: 2007/250 Accepted: 24 October, 2007
Applicant: **George A Lullfitz**, Wanneroo, WA.

‘CalgreyGL’

Application No: 2007/248 Accepted: 25 October, 2007
Applicant: **George A Lullfitz**, Wanneroo, WA.

‘CalredGL’

Application No: 2007/247 Accepted: 24 October, 2007

Applicant: **George A Lullfitz**, Wanneroo, WA.

Citrullus lanatus

WATERMELON

‘SP-4’

Application No: 2007/233 Accepted: 26 November, 2007
Applicant: **Syngenta Crop Protection AG**.
Agent: **Syngenta Seeds Pty Ltd**, Dandenong South, VIC.

Citrus aurantifolia

LIME

‘Sublime’

Application No: 2007/152 Accepted: 7 October, 2007
Applicant: **Darwin Plant Wholesalers**.
Agent: **Greenhills Propagation Nursery Pty Ltd**, Tynong, VIC.

Citrus clementina x *sinensis*

MANDARIN

‘Alkantara’

Application No: 2007/243 Accepted: 28 November, 2007
Applicant: **Giuseppe Reforgiato Recupero, Giuseppe Russo & Santo Recupero**.
Agent: **Australian Nurserymen's Fruit Improvement Company Ltd (ANFIC)**, Bathurst, NSW.

Citrus reticulata x *deliciosa*

MANDARIN

‘Mandalate’

Application No: 2007/244 Accepted: 28 November, 2007
Applicant: **Giuseppe Reforgiato Recupero, Giuseppe Russo & Santo Recupero**.
Agent: **Australian Nurserymen's Fruit Improvement Company Ltd (ANFIC)**, Bathurst, NSW.

Correa reflexa

NATIVE FUCHSIA

‘Multi Bella’

Application No: 2007/255 Accepted: 26 October, 2007
Applicant: **Friends of Warrandyte State Park (FOWSP)**.

Agent: **Australflora Pty Ltd**, Yarra Glen, VIC.

Crowea saligna

WAX FLOWER, WILLOW-LEAVED CROWEA

‘PPCS1’

Application No: 2007/259 Accepted: 22 November, 2007
Applicant: **Prestige Plants Pty Ltd**.
Agent: **Greenhills Propagation Nursery Pty Ltd**, Tynong, VIC.

Cuphea ignea x *lanceolata*

CUPHEA

‘Everbloom Purple’

Application No: 2007/302 Accepted: 12 December, 2007
Applicant: **Unique Plants**.
Agent: **Plants Management Australia Pty. Ltd.**, Dodges Ferry, TAS.

Dianella caerulea

BLUE FLAX-LILY, UMBRELLA DRACAENA

‘Newpladia1’ syn Stampede

Application No: 2007/236 Accepted: 19 November, 2007
Applicant: **Ian Angus Stewart**, Ourimbah, NSW.

Dianella ensifolia

FLAX LILY

‘DarwinGold’

Application No: 2007/229 Accepted: 1 November, 2007
Applicant: **Darwin Plant Wholesalers**, Winnellie, NT.

Dietes iridioides

AFRICAN IRIS, FORTNIGHT LILY, MOREA IRIS

‘White Tiger’

Application No: 2007/232 Accepted: 12 December, 2007
Applicant: **Nursery Australia Pty. Ltd.**.
Agent: **Plants Management Australia Pty. Ltd.**, Dodges Ferry, TAS.

Euphorbia characias

SPURGE

‘Tasmanian Tiger’

Application No: 2007/276 Accepted: 16 November, 2007
Applicant: **Sally Hohannsohn & Barbara Jennings**.
Agent: **Plants Management Australia Pty. Ltd.**, Dodges Ferry, TAS.

Ficus obliqua

SMALL LEAVED FIG

‘Fig-A-Row’

Application No: 2007/282 Accepted: 10 December, 2007
Applicant: **Agbiz Holdings Pty Ltd and Southern Advanced Plants Pty Ltd**.
Agent: **Southern Advanced Plants Pty Ltd**, Dromana, VIC.

Fragaria xananassa

STRAWBERRY

‘JULIETTE’

Application No: 2007/268 Accepted: 1 November, 2007
Applicant: **Agriculture Victoria Services Pty Ltd**, Attwood, VIC.

Glycine max

SOYBEAN

‘Fraser’

Application No: 2007/305 Accepted: 27 November, 2007
Applicant: **Commonwealth Scientific and Industrial Research Organisation and Grains Research and Development Corporation**, Canberra, ACT.

Gossypium barbadense

PIMA COTTON, SEA ISLAND COTTON

‘Sipima 280’

Application No: 2007/287 Accepted: 19 November, 2007
Applicant: **Commonwealth Scientific and Industrial Research Organisation**, Canberra, ACT.

Gossypium hirsutum

COTTON

‘Sicot 71BRF’

Application No: 2007/285 Accepted: 16 November, 2007
Applicant: **Commonwealth Scientific and Industrial Research Organisation**, Canberra, ACT.

‘Sicot 75’

Application No: 2007/286 Accepted: 16 November, 2007
Applicant: **Commonwealth Scientific and Industrial Research Organisation**, Canberra, ACT.

Grevillea hybrid

GREVILLEA

‘Carpet Layer’

Application No: 2007/261 Accepted: 9 November, 2007
Applicant: **Vaughans Australian Plants**.
Agent: **Humphris Nursery**, Mooroollbark, VIC.

Hardenbergia comptoniana

FALSE SARSPARILLA

‘LittleGL’

Application No: 2007/251 Accepted: 25 October, 2007
Applicant: **George A Lullfitz**, Wanneroo, WA.

Hardenbergia violacea

FALSE SARSPARILLA

‘Mystic Marvel’

Application No: 2007/317 Accepted: 19 December, 2007
Applicant: **Courtney Peter Whitton**, Junee, NSW.

Imperata cylindrica

BLADY GRASS, COGONGRASS

‘ICL200’

Application No: 2007/231 Accepted: 25 October, 2007
Applicant: **Ozbreed Pty Ltd**, Richmond, NSW.

Kalanchoe blossfeldiana

KALANCHOE

‘JACKIE’

Application No: 2007/207 Accepted: 7 October, 2007
Applicant: **Knud Jepson A/S**.
Agent: **Ball Australia Pty. Ltd.**, Keysborough, VIC.

‘JENNA’

Application No: 2007/205 Accepted: 7 October, 2007
Applicant: **Knud Jepson A/S**.
Agent: **Ball Australia Pty. Ltd.**, Keysborough, VIC.

‘JODIE’

Application No: 2007/206 Accepted: 7 October, 2007
Applicant: **Knud Jepson A/S**.
Agent: **Ball Australia Pty. Ltd.**, Keysborough, VIC.

‘MONA’

Application No: 2007/210 Accepted: 7 October, 2007
Applicant: **Knud Jepson A/S**.
Agent: **Ball Australia Pty. Ltd.**, Keysborough, VIC.

‘ROSEFLOWER-LEA’

Application No: 2007/209 Accepted: 7 October, 2007
Applicant: **Knud Jepson A/S**.
Agent: **Ball Australia Pty. Ltd.**, Keysborough, VIC.

‘SARAH’

Application No: 2007/208 Accepted: 7 October, 2007
Applicant: **Knud Jepson A/S**.
Agent: **Ball Australia Pty. Ltd.**, Keysborough, VIC.

Lupinus albus

WHITE LUPIN

‘WALAB2008’

Application No: 2007/200 Accepted: 7 October, 2007
Applicant: **Western Australian Agriculture Authority, Grains Research and Development Corporation, Council of grain Grower Organisations Ltd.**, South Perth, WA.

Malus domestica

APPLE

‘Burkitt Gala’ syn Cherry Gala

Application No: 2007/258 Accepted: 26 November, 2007
Applicant: **BMA TRUST c/-Dr Mark Burkitt**.
Agent: **Australian Nurserymen's Fruit Improvement Company Ltd (ANFIC)**, Bathurst, NSW.

‘Fugachee Fuji’

Application No: 2007/257 Accepted: 26 November, 2007
Applicant: **Brandt's Fruit Trees Inc.**.
Agent: **Australian Nurserymen's Fruit Improvement Company Ltd (ANFIC)**, Bathurst, NSW.

Melaleuca huegelii

CHENILLE HONEYMYRTLE

‘HuegflatGL’

Application No: 2007/249 Accepted: 24 October, 2007
Applicant: **George A Lullfitz**, Wanneroo, WA.

Melaleuca lanceolata

ROTTNEST TEATREE

‘Short1GL’

Application No: 2007/253 Accepted: 25 October, 2007
Applicant: **George A Lullfitz**, Wanneroo, WA.

Pennisetum clandestinum

KIKUYU GRASS

‘KIK01’

Application No: 2007/199 Accepted: 30 October, 2007
Applicant: **Ozbreed Pty Ltd**, Richmond, NSW.

Phormium cookianum

NEW ZEALAND MOUNTAIN FLAX

‘Storm Edition’

Application No: 2007/260 Accepted: 22 November, 2007
Applicant: **Greenhills Propagation Nursery Pty Ltd**, Tynong, VIC.

Pimelea ferruginea

PIMELEA

‘WhiteferruGL’

Application No: 2007/254 Accepted: 25 October, 2007
Applicant: **George A Lullfitz**, Wanneroo, WA.

Pimelea physodes

QUALUP BELL

‘QualredGL’

Application No: 2007/246 Accepted: 24 October, 2007
Applicant: **George A Lullfitz**, Wanneroo, WA.

Pisum sativum

FIELD PEA

‘XP 08530727’

Application No: 2007/224 Accepted: 26 October, 2007
Applicant: **Seminis Vegetable Seeds, Inc.**.
Agent: **Seminis Vegetable Seeds New Zealand Ltd.**, Ivanhoe, VIC.

Ricinocarpos tuberculatus

WEDDING BUSH

‘RicpenGL’

Application No: 2007/252 Accepted: 25 October, 2007
Applicant: **George A Lullfitz**, Wanneroo, WA.

Rosa hybrid

ROSE

‘Grandemufrap’

Application No: 2007/309 Accepted: 12 December, 2007
Applicant: **Mr H Schreuders**.
Agent: **Grandiflora Nurseries Pty Ltd**, Skye, VIC.

‘Grandhonemo’

Application No: 2007/311 Accepted: 12 December, 2007
Applicant: **Mr H Schreuders**.
Agent: **Grandiflora Nurseries Pty Ltd**, Skye, VIC.

‘Grandshanla’

Application No: 2007/310 Accepted: 12 December, 2007
Applicant: **Mr H Schreuders**.
Agent: **Grandiflora Nurseries Pty Ltd**, Skye, VIC.

‘Grandtinifa’

Application No: 2007/312 Accepted: 12 December, 2007
Applicant: **Mr H Schreuders**.
Agent: **Grandiflora Nurseries Pty Ltd**, Skye, VIC.

‘Poulcs010’

Application No: 2007/280 Accepted: 16 November, 2007
Applicant: **Poulsen Roser A/S**.
Agent: **Griffith Hack**, Perth, WA.

‘Poulcs012’

Application No: 2007/279 Accepted: 16 November, 2007
Applicant: **Poulsen Roser A/S**.
Agent: **Griffith Hack**, Perth, WA.

‘Poultc004’

Application No: 2007/277 Accepted: 16 November, 2007
Applicant: **Poulsen Roser A/S**.
Agent: **Griffith Hack**, Perth, WA.

‘Poultw003’

Application No: 2007/278 Accepted: 16 November, 2007
Applicant: **Poulsen Roser A/S**.
Agent: **Griffith Hack**, Perth, WA.

Senecio hybrid

SENECIO, CINERARIA

‘Sunsenebabu’ syn Baby Blue

Application No: 2007/184 Accepted: 8 November, 2007
Applicant: **Suntory Flowers Limited**.
Agent: **Oasis Horticulture Pty Limited**, Winmalee, NSW.

‘Sunsenebapiba’ syn Baby Magenta Bicolour

Application No: 2007/183 Accepted: 8 November, 2007
Applicant: **Suntory Flowers Limited**.
Agent: **Oasis Horticulture Pty Limited**, Winmalee, NSW.

Solanum tuberosum

POTATO

‘Romeo’

Application No: 2007/281 Accepted: 10 December, 2007
Applicant: **Irish Potato Marketing Ltd**.
Agent: **Bright Harvest**, Virginia, SA.

Stenotaphrum secundatum

BUFFALO GRASS, ST AUGUSTINE GRASS

‘TF01’

Application No: 2007/245 Accepted: 12 November, 2007
Applicant: **Transvaal Park Pty Ltd**, Beaudessert, QLD.

Syzygium australe

LILLY PILLY

‘Little Miss-Elegance’

Application No: 2007/202 Accepted: 16 November, 2007
Applicant: **Brent Edwin Wilson**, Logan Reserve, QLD.

‘PIP SQUEAK’

Application No: 2007/203 Accepted: 16 November, 2007
Applicant: **Brent Edwin Wilson**, Logan Reserve, QLD.

‘SUNSET’

Application No: 2007/204 Accepted: 12 December, 2007
Applicant: **Brent Edwin Wilson**, Logan Reserve, QLD.

Triticum aestivum

WHEAT

‘EGA Bounty’

Application No: 2007/303 Accepted: 21 December, 2007
Applicant: **State of Queensland through its Department of Primary Industries & Fisheries, Department of Primary Industries for and on behalf of the State of New South Wales and Grains Research and Development Corporation**, Brisbane, QLD.

‘EGA Stampede’

Application No: 2007/304 Accepted: 21 December, 2007
Applicant: **State of Queensland through its Department of Primary Industries & Fisheries, Department of Primary Industries for and on behalf of the State of New South Wales, The University of Queensland, Grains Research and Development Corporation**, Brisbane, QLD.

‘LongReach Bullet’ syn LPB0423

Application No: 2007/238 Accepted: 7 October, 2007
Applicant: **LongReach Plant Breeders Management Pty Ltd**, Bundoora, VIC.

‘WAWHT2726’

Application No: 2007/291 Accepted: 29 November, 2007
Applicant: **Western Australian Agriculture Authority, Grains Research and Development Corporation**, South Perth, WA.

Triticum turgidum ssp *turgidum*

DURUM WHEAT

‘HYPERNO’

Application No: 2007/300 Accepted: 12 December, 2007
Applicant: **Australian Grain Technologies Pty Ltd**, Urrbrae, SA.

Vaccinium hybrid

SOUTHERN HIGHBUSH BLUEBERRY

‘Abundance’

Application No: 2007/264 Accepted: 10 December, 2007
Applicant: **Florida Foundation Seed Producers, Inc**.

Agent: **BerryExchange (a division of CostaExchange Ltd)**, Corindi Beach, NSW.

‘C00-09’

Application No: 2007/269 Accepted: 16 November, 2007
Applicant: **BerryExchange (a division of CostaExchange Ltd)**, Range Rd, NSW.

‘C01-43’

Application No: 2007/272 Accepted: 16 November, 2007
Applicant: **BerryExchange (a division of CostaExchange Ltd)**, Range Rd, NSW.

‘C95-115’

Application No: 2007/270 Accepted: 16 November, 2007
Applicant: **BerryExchange (a division of CostaExchange Ltd)**, Range Rd, NSW.

‘C95-12’

Application No: 2007/271 Accepted: 16 November, 2007
Applicant: **BerryExchange (a division of CostaExchange Ltd)**, Range Rd, NSW.

‘C97-41’

Application No: 2007/273 Accepted: 16 November, 2007
Applicant: **BerryExchange (a division of CostaExchange Ltd)**, Range Rd, NSW.

‘FL92-84’

Application No: 2007/266 Accepted: 10 December, 2007
Applicant: **Florida Foundation Seed Producers, Inc**.
Agent: **BerryExchange (a division of CostaExchange Ltd)**, Corindi Beach, NSW.

‘Snowchaser’

Application No: 2007/265 Accepted: 10 December, 2007
Applicant: **Florida Foundation Seed Producers, Inc**.
Agent: **BerryExchange (a division of CostaExchange Ltd)**, Corindi Beach, NSW.

‘Springhigh’

Application No: 2007/263 Accepted: 10 December, 2007
Applicant: **Florida Foundation Seed Producers, Inc**.
Agent: **BerryExchange (a division of CostaExchange Ltd)**, Corindi Beach, NSW.

‘Sweetcrisp’

Application No: 2007/262 Accepted: 10 December, 2007
Applicant: **Florida Foundation Seed Producers, Inc**.
Agent: **BerryExchange (a division of CostaExchange Ltd)**, Corindi Beach, NSW.

Vigna unguiculata

COWPEA

‘BlackStallion’

Application No: 2007/284 Accepted: 22 November, 2007
Applicant: **B.W. Algate & Co Pty Ltd trading as J.W. Koek & Company, Blue Ribbon Seed & Pulse Exporters Pty Ltd & Champion Seeds Pty Ltd**, Burbank, QLD.

x*Triticosecale*

TRITICALE

‘Hawkeye’

Application No: 2007/234 Accepted: 10 October, 2007
Applicant: **Australian Grain Technologies Pty Ltd**, Urrbrae, SA.

‘Jaywick’

Application No: 2007/235 Accepted: 10 October, 2007
Applicant: **Australian Grain Technologies Pty Ltd**, Urrbrae, SA.

Zantedeschia spp.

CALLA LILY

‘Rosa BLZ’

Application No: 2007/141 Accepted: 10 December, 2007
Applicant: **BLOOMZ Ltd**.
Agent: **Rural Funds Management Flower Fund**, Nurioopta, SA.

Zoysia macrantha

PRICKLY COUCH, COAST COUCH, AUSTRALIAN ZOYSIA

‘MAC03’ syn Nara

Application No: 2007/275 Accepted: 30 November, 2007
Applicant: **Ozbreed Pty Ltd**, Richmond, NSW.

Australian Government
IP Australia
Plant Varieties Journal

Variety Descriptions

Common (Genus Species)	Variety	Title Holder
Bower Wattle *(Acacia cognata)*	Goldcog	Peter Goldup
Lilly Pilly *(Acmena smithii)*	DOW30	Downes Wholesale Nursery Pty Ltd
Willow Myrtle *(Agonis flexuosa)*	Jedda's Dream	James F Koppman and Jaqueline A Koppman
Bugle Bells *(Ajuga reptans)*	Black Scallop	Mike Tristram
Peruvian Lily *(Alstroemeria hybrid)*	Konimpa	Konst Breeding B.V.
Pineapple *(Ananas comosus)*	Aus-Jubilee	State of Queensland through its Department of Primary Industries and Fisheries
Pineapple *(Ananas comosus)*	Aus-Carnival	State of Queensland through its Department of Primary Industries and Fisheries
Kangaroo Paw *(Anigozanthos hybrid)*	Regal Velvet	George A Lullfitz
Peanut *(Arachis hypogaea)*	Georgia Hi/OL	The University of Georgia Research Foundation, Inc.
Oats *(Avena sativa)*	Dawson	NDSU Research Foundation

Oats *(Avena sativa)*	Monty	New Zealand Institute for Crop & Food Research Limited
Everlasting Daisy *(Bracteantha bracteata)*	Ohdrejumwhi	Bonza Botanicals Pty Limited
Canola *(Brassica napus)*	Argyle	Canola Breeders Western Australia Pty Ltd
Spider Plant *(Chlorophytum comosum)*	Ocean	Koning Smit IPR S.A.
Cordyline *(Cordyline australis)*	Jel01	Geoff Jewelll
Cordyline *(Cordyline australis)*	Kau01	Kauri Park Nursereis Ltd
Cordyline *(Cordyline fruticosa)*	BRA01	Peter Brauns
Cordyline *(Cordyline hybrid)*	Uto01	Utopia Palms and Cycads
Cordyline *(Cordyline hybrid)*	Tana	Evan David Lloyd
Flax lily *(Dianella tasmanica)*	TAS300	Wyeena Nurseries Pty Ltd
Flax lily *(Dianella tasmanica)*	TAS100	Ozbreed Pty Ltd
Coneflower *(Echinacea purpurea)*	Fragrant Angel	Terra Nova Nurseries, Inc
Strawberry *(Fragaria xananassa)*	Cal Giant 5	California Giant, Inc.
Grevillea *(Grevillea hybrid)*	Blood Orange	Christopher John Hughes

Barley *(Hordeum vulgare)*	Urambie	Department of Primary Industries for and on behalf of the State of New South Wales and Grains Research and Development Corporation
Barley *(Hordeum vulgare)*	Pacific Ranger	Her Majesty the Queen in Right of Canada as represented by the Minister of Agriculture and Agri-Food Canada
Wart-stemmed Pincushion *(Leucospermum cuneiforme)*	LS005A01	Proteaflora Enterprises Pty Ltd
Spiny Headed Mat Rush *(Lomandra hystrix)*	LHCOM	Ozbreed Pty Ltd
Spiny Headed Mat Rush *(Lomandra hystrix)*	LHBYF	Ozbreed Pty Ltd
Mango *(Mangifera indica)*	Minijac	Herminia and Jacinto Lay
Barrel Medic *(Medicago truncatula x littoralis)*	Cheetah	Pristine Forage Technologies Pty Ltd
Barrel Medic *(Medicago truncatula x littoralis)*	Lynx	Pristine Forage Technologies Pty Ltd
Spanish Cherry *(Mimusops elengi)*	Street Snow	Darwin Plant Wholesalers
Long Leaved Waxflower *(Philotheca myoporoides)*	Bournda Gold	Lystare Pty Ltd trading as Bournda Plants

New Zealand Mountain Flax *(Phormium cookianum)*	Storm Edition	Greenhills Propagation Nursery Pty Ltd
New Zealand Flax *(Phormium tenax)*	PHORD1	Ozbreed Pty Ltd
Field Pea *(Pisum sativum)*	Bundi	Agriculture Victoria Services Pty Ltd and Grains Research and Development Corporation
Pittosporum *(Pittosporum tenuifolium)*	EMERALDSTAR	Grant Farmer McKechnie
Pittosporum *(Pittosporum tenuifolium)*	Golf Ball	M & R Fyfe
Polygala *(Polygala xDalmaisiana)*	Whitepol	Chris Cristou
Sweet Cherry *(Prunus avium)*	Arodel	Societe Anonyme des Pepinieres et Roseraies GEORGES DELBARD
Sweet Cherry *(Prunus avium)*	Dame Nancy	Minister for Agriculture, Food and Fisheries
Interspecific Plum *(Prunus hybrid)*	Black Kat	Zaiger's Inc. Genetics
Rose *(Rosa hybrid)*	Grandtang	Mr H Schreuders
Rose *(Rosa hybrid)*	Kribigpea	Lux Riviera S.r.l.
Raspberry *(Rubus idaeus)*	Cardinal	Driscoll Strawberry Associates, Inc
Raspberry *(Rubus idaeus)*	Maravilla	Driscoll Strawberry Associates, Inc
Sage *(Salvia hybrid)*	Heatwave Blaze	Plant Growers Australia Pty. Ltd.

Sage *(Salvia hybrid)*	Heatwave Sizzle	Plant Growers Australia Pty. Ltd.
Buffalo Grass *(Stenotaphrum secundatum)*	TF01	Transvaal Park Pty Ltd
(Strobilanthes anisophyllus)	Goldust	Valdis and Solveiga Schutz
Lilly Pilly *(Syzygium australe)*	AATS	John Crump
Small Leaf Lilly Pilly *(Syzygium smithii)*	Cherry Surprise	Wirreanda Nursery
Small Leaf Lilly Pilly *(Syzygium smithii)*	Sunrise	Wirreanda Nursery
Kanooka *(Tristaniopsis laurina)*	DOW10	Downes Wholesale Nursery Pty Ltd
Wheat *(Triticum aestivum)*	Axe	Australian Grain Technologies Pty Ltd
Wheat *(Triticum aestivum)*	Gladius	Australian Grain Technologies Pty Ltd
Wheat *(Triticum aestivum)*	Espada	Australian Grain Technologies Pty Ltd
Durum Wheat *(Triticum turgidum ssp turgidum)*	HYPERNO	Australian Grain Technologies Pty Ltd
Field Bean *(Vicia faba)*	Doza	Department of Primary Industries for and on behalf of the State of New South Wales and Grains Research and Development Corporation

Weeping Lilly Pilly *(Waterhousea floribunda)*	DOW20	Downes Wholesale Nursery Pty Ltd
Triticale *(xTriticosecale)*	Hawkeye	Australian Grain Technologies Pty Ltd
Triticale *(xTriticosecale)*	Jaywick	Australian Grain Technologies Pty Ltd

Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

(Strobilanthes anisophyllus)

Variety:	'Goldust'
Synonym:	N/A
Application no:	2007/111
Current status:	ACCEPTED
Certificate no:	N/A
Received:	10-Apr-2007
Accepted:	01-May-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Valdis and Solveiga Schutz
Agent:	N/A
Telephone:	0296511458
Fax:	0296513856

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Barley *(Hordeum vulgare)*

Variety:	'Urambie'
Synonym:	N/A
Application no:	2005/349
Current status:	ACCEPTED
Certificate no:	N/A
Received:	14-Dec-2005
Accepted:	09-Feb-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Department of Primary Industries for and on behalf of the State of New South Wales and Grains Research and Development Corporation
Agent:	N/A
Telephone:	0263913550
Fax:	0263913563

View the detailed description of this variety.



Australian Government
IP Australia


Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Barley *(Hordeum vulgare)*

Variety:	'Pacific Ranger'
Synonym:	AC Ranger
Application no:	2006/299
Current status:	ACCEPTED
Certificate no:	N/A
Received:	16-Nov-2006
Accepted:	05-Feb-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Her Majesty the Queen in Right of Canada as represented by the Minister of Agriculture and Agri-Food Canada
Agent:	Pacific Seeds Pty Ltd
Telephone:	0746902663
Fax:	0746301063

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Barrel Medic *(Medicago truncatula x littoralis)*

Variety:	'Cheetah'
Synonym:	N/A
Application no:	2007/195
Current status:	ACCEPTED
Certificate no:	N/A
Received:	01-Aug-2007
Accepted:	05-Sep-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Pristine Forage Technologies Pty Ltd
Agent:	N/A
Telephone:	0881770558
Fax:	0881770558

View the detailed description of this variety.



Australian Government
IP Australia


Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Barrel Medic *(Medicago truncatula x littoralis)*

Variety:	'Lynx'
Synonym:	N/A
Application no:	2007/194
Current status:	ACCEPTED
Certificate no:	N/A
Received:	01-Aug-2007
Accepted:	05-Sep-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Pristine Forage Technologies Pty Ltd
Agent:	N/A
Telephone:	0881770558
Fax:	0881770558

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Bower Wattle *(Acacia cognata)*

Variety:	'Goldcog'
Synonym:	N/A
Application no:	2005/354
Current status:	ACCEPTED
Certificate no:	N/A
Received:	03-Jan-2006
Accepted:	09-Feb-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Peter Goldup
Agent:	Bushland Flora
Telephone:	0397364364
Fax:	0397364716

View the detailed description of this variety.



Australian Government
IP Australia


Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Buffalo Grass *(Stenotaphrum secundatum)*

Variety:	'TF01'
Synonym:	N/A
Application no:	2007/245
Current status:	ACCEPTED
Certificate no:	N/A
Received:	20-Sep-2007
Accepted:	12-Nov-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Transvaal Park Pty Ltd
Agent:	N/A
Telephone:	0755436090
Fax:	0755436097

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Bugle Bells *(Ajuga reptans)*

Variety:	'Black Scallop'
Synonym:	N/A
Application no:	2006/030
Current status:	ACCEPTED
Certificate no:	N/A
Received:	21-Feb-2006
Accepted:	24-Mar-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Mike Tristram
Agent:	Plants Management Australia
Telephone:	0397221444
Fax:	0397221018

View the detailed description of this variety.


Black Scallop
Evening Glow

Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Canola *(Brassica napus)*

Variety:	'Argyle'
Synonym:	N/A
Application no:	2007/058
Current status:	ACCEPTED
Certificate no:	N/A
Received:	22-Feb-2007
Accepted:	08-Mar-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Canola Breeders Western Australia Pty Ltd
Agent:	N/A
Telephone:	(08) 9285 8087
Fax:	0893874388

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Coneflower *(Echinacea purpurea)*

Variety:	'Fragrant Angel'
Synonym:	N/A
Application no:	2007/030
Current status:	ACCEPTED
Certificate no:	N/A
Received:	19-Jan-2007
Accepted:	13-Feb-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Terra Nova Nurseries, Inc
Agent:	Lifetech Laboratories Ltd
Telephone:	(02) 4381 0051
Fax:	(02) 4381 0071

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Cordyline *(Cordyline australis)*

Variety:	'Jel01'
Synonym:	N/A
Application no:	2005/063
Current status:	ACCEPTED
Certificate no:	N/A
Received:	07-Mar-2005
Accepted:	21-Apr-2005
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Geoff Jewelll
Agent:	Anthony Tesselaar Plants Pty Ltd
Telephone:	0397377921
Fax:	0397379899

View the detailed description of this variety.



Australian Government
IP Australia
Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Cordyline *(Cordyline australis)*

Variety:	'Kau01'
Synonym:	N/A
Application no:	2006/126
Current status:	ACCEPTED
Certificate no:	N/A
Received:	08-Jun-2006
Accepted:	05-Aug-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Kauri Park Nurseries Ltd
Agent:	Greenhills Propagation Nursery Pty Ltd
Telephone:	0356292443
Fax:	0656292822

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Cordyline *(Cordyline fruticosa)*

Variety:	'BRA01'
Synonym:	N/A
Application no:	2004/133
Current status:	ACCEPTED
Certificate no:	N/A
Received:	15-Apr-2004
Accepted:	22-Apr-2005
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Peter Brauns
Agent:	Anthony Tesselaar Plants Pty Ltd
Telephone:	N/A
Fax:	N/A

View the detailed description of this variety.


BRA01
Nigra

Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Cordyline *(Cordyline hybrid)*

Variety:	'Uto01'
Synonym:	N/A
Application no:	2005/121
Current status:	ACCEPTED
Certificate no:	N/A
Received:	05-May-2005
Accepted:	26-Oct-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Utopia Palms and Cycads
Agent:	N/A
Telephone:	0754466205
Fax:	0754466205

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Cordyline *(Cordyline hybrid)*

Variety:	'Tana'
Synonym:	Renegade
Application no:	2007/010
Current status:	ACCEPTED
Certificate no:	N/A
Received:	02-Jan-2007
Accepted:	25-Jan-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Evan David Lloyd
Agent:	Greenhills Propagation Nursery Pty Ltd
Telephone:	0356292443
Fax:	0356292822

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Durum Wheat *(Triticum turgidum ssp turgidum)*

Variety:	'HYPERNO'
Synonym:	N/A
Application no:	2007/300
Current status:	ACCEPTED
Certificate no:	N/A
Received:	05-Nov-2007
Accepted:	12-Dec-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4

Title Holder:	Australian Grain Technologies Pty Ltd
Agent:	N/A
Telephone:	0883036861
Fax:	0883036865

View the detailed description of this variety.


'ECA BELLAROU
'TAMARA'

Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Everlasting Daisy *(Bracteantha bracteata)*

Variety:	'Ohdrejumwhi'
Synonym:	Jumbo White
Application no:	2007/214
Current status:	ACCEPTED
Certificate no:	N/A
Received:	17-Aug-2007
Accepted:	26-Sep-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Bonza Botanicals Pty Limited
Agent:	Oasis Horticulture Pty Limited
Telephone:	0247541422
Fax:	0147544260

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Field Bean *(Vicia faba)*

Variety:	'Doza'
Synonym:	N/A
Application no:	2007/161
Current status:	ACCEPTED
Certificate no:	N/A
Received:	18-Jun-2007
Accepted:	09-Jul-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Department of Primary Industries for and on behalf of the State of New South Wales and Grains Research and Development Corporation
Agent:	N/A
Telephone:	0263913550
Fax:	0263913563

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Field Pea *(Pisum sativum)*

Variety:	'Bundi'
Synonym:	N/A
Application no:	2006/026
Current status:	ACCEPTED
Certificate no:	N/A
Received:	16-Feb-2006
Accepted:	24-Mar-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Agriculture Victoria Services Pty Ltd and Grains Research and Development Corporation
Agent:	N/A
Telephone:	0392174200
Fax:	0392174161

View the detailed description of this variety.



Australian Government
IP Australia


Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Flax lily *(Dianella tasmanica)*

Variety:	'TAS300'
Synonym:	N/A
Application no:	2007/097
Current status:	ACCEPTED
Certificate no:	N/A
Received:	16-Mar-2007
Accepted:	26-Apr-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Wyeena Nurseries Pty Ltd
Agent:	Ozbreed Pty Ltd
Telephone:	0245780866
Fax:	0245780855

View the detailed description of this variety.



Australian Government
IP Australia


Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Flax lily *(Dianella tasmanica)*

Variety:	'TAS100'
Synonym:	N/A
Application no:	2007/021
Current status:	ACCEPTED
Certificate no:	N/A
Received:	17-Jan-2007
Accepted:	05-Feb-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Ozbreed Pty Ltd
Agent:	N/A
Telephone:	0245780866
Fax:	0245780855

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Grevillea *(Grevillea hybrid)*

Variety:	'Blood Orange'
Synonym:	N/A
Application no:	2006/218
Current status:	ACCEPTED
Certificate no:	N/A
Received:	07-Aug-2006
Accepted:	05-Oct-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Christopher John Hughes
Agent:	N/A
Telephone:	0266884189
Fax:	0266884383

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Interspecific Plum *(Prunus hybrid)*

Variety:	'Black Kat'
Synonym:	N/A
Application no:	2003/375
Current status:	ACCEPTED
Certificate no:	N/A
Received:	25-Dec-2003
Accepted:	05-May-2004
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4

Title Holder:	Zaiger's Inc. Genetics
Agent:	Fleming's Nurseries & Associates Pty Ltd
Telephone:	0397566105
Fax:	0397520005

View the detailed description of this variety.


Black Kat

Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Kangaroo Paw *(Anigozanthos hybrid)*

Variety:	'Regal Velvet'
Synonym:	N/A
Application no:	2006/012
Current status:	ACCEPTED
Certificate no:	N/A
Received:	01-Feb-2006
Accepted:	22-Feb-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	George A Lullfitz
Agent:	N/A
Telephone:	0894051607
Fax:	0893062933

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Kanooka *(Tristaniopsis laurina)*

Variety:	'DOW10'
Synonym:	N/A
Application no:	2005/288
Current status:	ACCEPTED
Certificate no:	N/A
Received:	18-Aug-2005
Accepted:	24-Oct-2005
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Downes Wholesale Nursery Pty Ltd
Agent:	Ozbreed Pty Ltd
Telephone:	0245780866
Fax:	0245780855

View the detailed description of this variety.



Australian Government
IP Australia
Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Lilly Pilly *(Syzygium australe)*

Variety:	'AATS'
Synonym:	N/A
Application no:	2006/127
Current status:	ACCEPTED
Certificate no:	N/A
Received:	09-Jun-2006
Accepted:	31-Aug-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	John Crump
Agent:	Ozbreed Pty Ltd
Telephone:	0245780866
Fax:	0245780855

View the detailed description of this variety.



AATS Bronzed Aussie

Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Lilly Pilly *(Acmena smithii)*

Variety:	'DOW30'
Synonym:	N/A
Application no:	2005/317
Current status:	ACCEPTED
Certificate no:	N/A
Received:	17-Oct-2005
Accepted:	29-Apr-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Downes Wholesale Nursery Pty Ltd
Agent:	Ozbreed Pty Ltd
Telephone:	0245780866
Fax:	0245780855

View the detailed description of this variety.



DOW30 *A. smithii* parent form

Australian Government
IP Australia
Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Long Leaved Waxflower *(Philotheca myoporoides)*

Variety:	'Bournda Gold'
Synonym:	N/A
Application no:	2005/072
Current status:	ACCEPTED
Certificate no:	N/A
Received:	11-Mar-2005
Accepted:	14-Jun-2005
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Lystare Pty Ltd trading as Bournda Plants
Agent:	Greenhills Propagation Nursery Pty Ltd
Telephone:	0356292443
Fax:	0356292822

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Mango *(Mangifera indica)*

Variety:	'Minijac'
Synonym:	N/A
Application no:	2000/301
Current status:	ACCEPTED
Certificate no:	N/A
Received:	04-Oct-2000
Accepted:	30-Nov-2000
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Herminia and Jacinto Lay
Agent:	N/A
Telephone:	0889816112
Fax:	0889812892

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

New Zealand Flax *(Phormium tenax)*

Variety:	'PHORD1'
Synonym:	N/A
Application no:	2004/250
Current status:	ACCEPTED
Certificate no:	N/A
Received:	26-Aug-2004
Accepted:	21-Sep-2004
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Ozbreed Pty Ltd
Agent:	N/A
Telephone:	0245780866
Fax:	0245780855

View the detailed description of this variety.


PHORD 1
Merlot

Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

New Zealand Mountain Flax *(Phormium cookianum)*

Variety:	'Storm Edition'
Synonym:	N/A
Application no:	2007/260
Current status:	ACCEPTED
Certificate no:	N/A
Received:	02-Oct-2007
Accepted:	22-Nov-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Greenhills Propagation Nursery Pty Ltd
Agent:	N/A
Telephone:	0356292443
Fax:	0356292822

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Oats *(Avena sativa)*

Variety:	'Dawson'
Synonym:	N/A
Application no:	2007/241
Current status:	ACCEPTED
Certificate no:	N/A
Received:	17-Sep-2007
Accepted:	07-Nov-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	NDSU Research Foundation
Agent:	Pacific Seeds Pty Ltd
Telephone:	0746902663
Fax:	0746301063

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Oats *(Avena sativa)*

Variety:	'Monty'
Synonym:	N/A
Application no:	2007/150
Current status:	ACCEPTED
Certificate no:	N/A
Received:	24-May-2007
Accepted:	26-Jun-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	New Zealand Institute for Crop & Food Research Limited
Agent:	Heritage Seeds Pty Ltd
Telephone:	0260265288
Fax:	0260265268

View the detailed description of this variety.


Graza 50
Monty

Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Peanut *(Arachis hypogaea)*

Variety:	'Georgia Hi/OL'
Synonym:	Reid
Application no:	2006/002
Current status:	ACCEPTED
Certificate no:	N/A
Received:	16-Jan-2006
Accepted:	08-May-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	University of Georgia Research Foundation, Inc.
Agent:	Peanut Company of Australia Limited
Telephone:	0741626311
Fax:	0741624402

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Peruvian Lily *(Alstroemeria hybrid)*

Variety:	'Konimpa'
Synonym:	N/A
Application no:	2006/084
Current status:	ACCEPTED
Certificate no:	N/A
Received:	26-Apr-2006
Accepted:	08-May-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Konst Breeding B.V.
Agent:	David Nichols - postal address for service of notice on the applicant Konst Breeding BV
Telephone:	0359774755
Fax:	0359774921

View the detailed description of this variety.



Australian Government
IP Australia


Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Pineapple *(Ananas comosus)*

Variety:	'Aus-Jubilee'
Synonym:	Jubilee
Application no:	2005/353
Current status:	ACCEPTED
Certificate no:	N/A
Received:	23-Dec-2005
Accepted:	09-Feb-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	State of Queensland through its Department of Primary Industries and Fisheries
Agent:	N/A
Telephone:	0732390802
Fax:	0732393948

View the detailed description of this variety.


73-50
Australian


Australian Government
IP Australia
Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Pineapple *(Ananas comosus)*

Variety:	'Aus-Carnival'
Synonym:	N/A
Application no:	2007/036
Current status:	ACCEPTED
Certificate no:	N/A
Received:	23-Jan-2007
Accepted:	26-Feb-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	State of Queensland through its Department of Primary Industries and Fisheries
Agent:	N/A
Telephone:	0732390802
Fax:	0732393948

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Pittosporum *(Pittosporum tenuifolium)*

Variety:	'EMERALDSTAR'
Synonym:	N/A
Application no:	2003/080
Current status:	ACCEPTED
Certificate no:	N/A
Received:	14-Apr-2003
Accepted:	15-May-2003
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Grant Farmer McKechnie
Agent:	Greenhills Propagation Nursery Pty Ltd
Telephone:	0356292443
Fax:	0356292822

View the detailed description of this variety.


Green Pillar

Australian Government
IP Australia
Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Pittosporum *(Pittosporum tenuifolium)*

Variety:	'Golf Ball'
Synonym:	N/A
Application no:	2006/213
Current status:	ACCEPTED
Certificate no:	N/A
Received:	31-Jul-2006
Accepted:	26-Oct-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	M & R Fyfe
Agent:	Greenhills Propagation Nursery Pty Ltd
Telephone:	0356292443
Fax:	0356292822

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Polygala *(Polygala xDalmaisiana)*

Variety:	'Whitepol'
Synonym:	N/A
Application no:	2006/087
Current status:	ACCEPTED
Certificate no:	N/A
Received:	26-Apr-2006
Accepted:	01-Aug-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Chris Cristou
Agent:	N/A
Telephone:	0397421828
Fax:	0397421183

View the detailed description of this variety.



Australian Government
IP Australia


Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Raspberry *(Rubus idaeus)*

Variety:	'Cardinal'
Synonym:	N/A
Application no:	2003/339
Current status:	ACCEPTED
Certificate no:	N/A
Received:	27-Nov-2003
Accepted:	05-Mar-2004
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Driscoll Strawberry Associates, Inc
Agent:	Phillips Ormonde & Fitzpatrick
Telephone:	(03) 9614 1944
Fax:	(03) 9614 1867

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal


Plant Varieties Journal - Search Result Details

Raspberry *(Rubus idaeus)*

Variety:	'Maravilla'
Synonym:	N/A
Application no:	2003/338
Current status:	ACCEPTED
Certificate no:	N/A
Received:	27-Nov-2003
Accepted:	05-Mar-2004
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4

Title Holder: Driscoll Strawberry Associates, Inc
Agent: Phillips Ormonde & Fitzpatrick
Telephone: (03) 9614 1944
Fax: (03) 9614 1867

View the detailed description of this variety.


Maravilla

Australian Government
IP Australia


Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Rose *(Rosa hybrid)*

Variety:	'Grandtang'
Synonym:	N/A
Application no:	2006/115
Current status:	ACCEPTED
Certificate no:	N/A
Received:	18-May-2006
Accepted:	30-May-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Mr H Schreuders
Agent:	Grandiflora Nurseries Pty Ltd
Telephone:	0397822777
Fax:	0397822576

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Rose *(Rosa hybrid)*

Variety:	'Kribigpea'
Synonym:	N/A
Application no:	2004/012
Current status:	ACCEPTED
Certificate no:	N/A
Received:	19-Jan-2004
Accepted:	03-Mar-2004
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Lux Riviera S.r.l.
Agent:	Grandiflora Nurseries Pty Ltd
Telephone:	0397822777
Fax:	0397822576

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Sage *(Salvia hybrid)*

Variety:	'Heatwave Blaze'
Synonym:	N/A
Application no:	2007/059
Current status:	ACCEPTED
Certificate no:	N/A
Received:	26-Feb-2007
Accepted:	09-Mar-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Plant Growers Australia Pty. Ltd.
Agent:	Plants Management Australia Pty. Ltd.
Telephone:	0362659920
Fax:	0362659919

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Sage *(Salvia hybrid)*

Variety:	'Heatwave Sizzle'
Synonym:	N/A
Application no:	2007/060
Current status:	ACCEPTED
Certificate no:	N/A
Received:	26-Feb-2007
Accepted:	21-Mar-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4

Title Holder:	Plant Growers Australia Pty. Ltd.
Agent:	Plants Management Australia Pty. Ltd.
Telephone:	0362659920
Fax:	0362659919

View the detailed description of this variety.



Australian Government
IP Australia


Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Small Leaf Lilly Pilly *(Syzygium smithii)*

Variety:	'Cherry Surprise'
Synonym:	N/A
Application no:	2006/297
Current status:	ACCEPTED
Certificate no:	N/A
Received:	16-Nov-2006
Accepted:	16-Mar-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Wirreanda Nursery
Agent:	N/A
Telephone:	0294501400
Fax:	0294502664

View the detailed description of this variety.



Australian Government
IP Australia


Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Small Leaf Lilly Pilly *(Syzygium smithii)*

Variety:	'Sunrise'
Synonym:	N/A
Application no:	2006/298
Current status:	ACCEPTED
Certificate no:	N/A
Received:	16-Nov-2006
Accepted:	16-Mar-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Wirreanda Nursery
Agent:	N/A
Telephone:	0294501400
Fax:	0294502664

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Spanish Cherry *(Mimusops elengi)*

Variety:	'Street Snow'
Synonym:	N/A
Application no:	2001/229
Current status:	ACCEPTED
Certificate no:	N/A
Received:	30-Aug-2001
Accepted:	04-Sep-2001
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Darwin Plant Wholesalers
Agent:	N/A
Telephone:	0889881888
Fax:	0889882110

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Spider Plant *(Chlorophytum comosum)*

Variety:	'Ocean'
Synonym:	N/A
Application no:	2007/146
Current status:	ACCEPTED
Certificate no:	N/A
Received:	24-May-2007
Accepted:	11-Jul-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Koning Smit IPR S.A.
Agent:	Ramm Botanicals Pty Ltd
Telephone:	0243512099
Fax:	0243531875

View the detailed description of this variety.



Australian Government
IP Australia


Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Spiny Headed Mat Rush *(Lomandra hystrix)*

Variety:	'LHCOM'
Synonym:	N/A
Application no:	2006/088
Current status:	ACCEPTED
Certificate no:	N/A
Received:	28-Apr-2006
Accepted:	30-May-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Ozbreed Pty Ltd
Agent:	N/A
Telephone:	0245780866
Fax:	0245780855

View the detailed description of this variety.



Australian Government
IP Australia


Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Spiny Headed Mat Rush *(Lomandra hystrix)*

Variety:	'LHBYF'
Synonym:	N/A
Application no:	2006/270
Current status:	ACCEPTED
Certificate no:	N/A
Received:	03-Oct-2006
Accepted:	26-Oct-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Ozbreed Pty Ltd
Agent:	N/A
Telephone:	0245780866
Fax:	0245780855

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Strawberry *(Fragaria xananassa)*

Variety:	'Cal Giant 5'
Synonym:	Galexia
Application no:	2005/340
Current status:	ACCEPTED
Certificate no:	N/A
Received:	28-Nov-2005
Accepted:	22-Dec-2005
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4

Title Holder:	California Giant, Inc.
Agent:	State of Queensland through its Departrment of Primary Industries and Fisheries
Telephone:	0738969401
Fax:	0732393948

View the detailed description of this variety.


Cal Giant 5
Cal Giant 3

Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Sweet Cherry *(Prunus avium)*

Variety:	'Arodel'
Synonym:	N/A
Application no:	2002/008
Current status:	ACCEPTED
Certificate no:	N/A
Received:	07-Jan-2002
Accepted:	27-Jun-2003
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Societe Anonyme des Pepinieres et Roseraies GEORGES DELBARD
Agent:	Australian Nurserymen's Fruit Improvement Company Limited
Telephone:	0263326960
Fax:	0263326962

View the detailed description of this variety.



Australian Government
IP Australia


Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Sweet Cherry *(Prunus avium)*

Variety:	'Dame Nancy'
Synonym:	N/A
Application no:	2003/148
Current status:	ACCEPTED
Certificate no:	N/A
Received:	18-Jun-2003
Accepted:	07-Jul-2003
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Minister for Agriculture, Food and Fisheries
Agent:	Australian Nurseryman's Fruit Improvement Company Limited
Telephone:	0263326960
Fax:	0263326962

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Triticale *(xTriticosecale)*

Variety:	'Hawkeye'
Synonym:	N/A
Application no:	2007/234
Current status:	ACCEPTED
Certificate no:	N/A
Received:	12-Sep-2007
Accepted:	10-Oct-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4

Title Holder:	Australian Grain Technologies Pty Ltd
Agent:	N/A
Telephone:	0883036861
Fax:	0883036865

View the detailed description of this variety.


KOSCIUSZKO
TIOGA
HAWKEYE

Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Triticale *(xTriticosecale)*

Variety:	'Jaywick'
Synonym:	N/A
Application no:	2007/235
Current status:	ACCEPTED
Certificate no:	N/A
Received:	12-Sep-2007
Accepted:	10-Oct-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Australian Grain Technologies Pty Ltd
Agent:	N/A
Telephone:	0883036861
Fax:	0883036865

View the detailed description of this variety.


'JAYWICK'
'TAHARA'
'JAYWICK'
'TAHARA'

Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Wart-stemmed Pincushion *(Leucospermum cuneiforme)*

Variety:	'LS005A01'
Synonym:	N/A
Application no:	2007/001
Current status:	ACCEPTED
Certificate no:	N/A
Received:	02-Jan-2007
Accepted:	25-Jan-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Proteaflora Enterprises Pty Ltd
Agent:	N/A
Telephone:	0397567233
Fax:	0397566948

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Weeping Lilly Pilly *(Waterhousea floribunda)*

Variety:	'DOW20'
Synonym:	N/A
Application no:	2005/289
Current status:	ACCEPTED
Certificate no:	N/A
Received:	18-Aug-2005
Accepted:	29-Apr-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Downes Wholesale Nursery Pty Ltd
Agent:	Ozbreed Pty Ltd
Telephone:	0245780866
Fax:	0245780855

View the detailed description of this variety.



Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Wheat *(Triticum aestivum)*

Variety:	'Axe'
Synonym:	N/A
Application no:	2007/117
Current status:	ACCEPTED
Certificate no:	N/A
Received:	26-Apr-2007
Accepted:	18-May-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Australian Grain Technologies Pty Ltd
Agent:	N/A
Telephone:	0883036861
Fax:	0883036865

View the detailed description of this variety.


'RAC1192' - generation 1
'SILVERSTAR'
'RAC1192' - generation 2

Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Wheat *(Triticum aestivum)*

Variety:	'Gladius'
Synonym:	N/A
Application no:	2006/302
Current status:	ACCEPTED
Certificate no:	N/A
Received:	27-Nov-2006
Accepted:	17-Jan-2007
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Australian Grain Technologies Pty Ltd
Agent:	N/A
Telephone:	0883037835
Fax:	0883037964

View the detailed description of this variety.



Australian Government
IP Australia


Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Wheat *(Triticum aestivum)*

Variety:	'Espada'
Synonym:	N/A
Application no:	2007/322
Current status:	ACCEPTED
Certificate no:	N/A
Received:	20-Dec-2007
Accepted:	17-Jan-2008
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	Australian Grain Technologies Pty Ltd
Agent:	N/A
Telephone:	0883036861
Fax:	0883036865

View the detailed description of this variety.


'GLADIUS' 'ESPADA' / 'CORRELL'

Australian Government
IP Australia

Plant Varieties Journal

Plant Varieties Journal - Search Result Details

Willow Myrtle *(Agonis flexuosa)*

Variety:	'Jedda's Dream'
Synonym:	N/A
Application no:	2006/222
Current status:	ACCEPTED
Certificate no:	N/A
Received:	09-Aug-2006
Accepted:	15-Aug-2006
Granted:	N/A
Description published in Plant Varieties Journal:	Volume 20, Issue 4
Title Holder:	James F Koppman and Jaqueline A Koppman
Agent:	N/A
Telephone:	0244478432
Fax:	0244478032

View the detailed description of this variety.



Details of Application

Application Number	2007/111
Variety Name	'Goldust'
Genus Species	*Strobilanthes anisophyllus*
Common Name	Strobilanthes
Synonym	Nil
Accepted Date	1 May 2007
Applicant	Valdis and Solveiga Schutz, Arcadia, NSW
Agent	N/A
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Dural, NSW.
Descriptor	General Descriptor (for plant varieties with no specific descriptor available) PBR GEN-DES.
Period	Summer-autumn 2007.
Conditions	Trial conducted in a opens beds, plants propagated from cuttings, rooted cuttings planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random.
RHS Chart - edition	1995.

Origin and Breeding

Spontaneous mutation: parent *S. anisophyllus*. The parent is characterised by an absence of leaf variegation and leaf colour predominated by greyed purple and brown tones over a dark yellow green base. Selection took place in Arcadia, NSW in 1999. Selection criteria: leaf variegation and colour. Propagation: vegetative cuttings were found to be uniform and stable. Breeder: Val Schutz, Arcadia, NSW.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	type	shrub
Plant	growth habit	bushy
Plant	size	medium
Plant	time of beginning of flowering	medium
Leaf	type	simple
Leaf	size	medium
Leaf	shape	lanceolate
Leaf	incision of margin	present

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
S. anisophyllus	parent variety (variegation absent)

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Goldust'	*S. anisophyllus*
☐ Plant: type	shrub	shrub
☐ Plant: growth habit	bushy	bushy
☐ Plant: size	medium	medium
☐ Plant: height	medium	medium
☐ Plant: width	medium	medium
☐ Plant: time of beginning of flowering	medium	medium
☐ Stem: presence of anthocyanin in new growth	present	present
☐ Young shoot: anthocyanin colouration	strong	very strong
☐ Leaf: leaf type	simple	simple
☐ Leaf: size	medium	medium
☐ Leaf: attitude	horizontal	horizontal
☐ Leaf: arrangement	opposite and decussate	opposite and decussate
☐ Leaf: length of blade	medium	medium
☐ Leaf: width of blade	medium	medium
☐ Leaf: length of petiole	short	short
☐ Leaf: shape	lanceolate	lanceolate
☐ Leaf: shape of apex	acute	acute
☐ Leaf: shape of base	cuneate	cuneate
☐ Leaf: incision of margin	present	present
☐ Leaf: depth of incision	very shallow	very shallow
☐ Leaf: type of incision	toothed	toothed
☐ Leaf: undulation of the margin	very weak	very weak
☐ Leaf: shape of cross-section	flat	flat
☐ Leaf: curvature of longitudinal axis	straight	straight
☐ Leaf: glossiness of upper side	medium to strong	medium to strong
☐ Leaf: green colour	medium to dark	medium to dark
☑ Leaf: presence of variegation	present	absent
☐ Leaf: degree of variegation	medium	
☐ Leaf: primary colour (RHS colour chart)	147A	147A
☐ Leaf: secondary colour (RHS colour chart)	10C to 10D	
☐ Leaf: border between colours	not clearly defined	

☑ Leaf colour: number of colours	two	one

Statistical Table

Organ/Plant Part: Context	**'Goldust'**	***S. anisophyllus***
☐ Leaf: length (mm)		
Mean	69.90	65.40
Std. Deviation	5.70	6.00
LSD/sig	6.7	ns
☐ Leaf: width (mm)		
Mean	18.00	17.40
Std. Deviation	2.10	2.00
LSD/sig	2.38	ns

Prior Applications and Sales

Nil.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW

Details of Application

Application Number	2005/349
Variety Name	'Urambie'
Genus Species	*Hordeum vulgare*
Common Name	Barley
Synonym	Nil
Accepted Date	9 Feb 2006
Applicant	Department of Primary Industries for and on behalf of the State of New South Wales, Orange, NSW and Grains Research and Development Corporation, Barton, ACT
Agent	N/A
Qualified Person	Ross Downes

Details of Comparative Trial

Location	Temora Research Station.
Descriptor	Barley (*Hordeum vulgare*) TG 19/10.
Period	Sown in May for winter-spring crop 2007.
Conditions	Because of drought conditions plots were irrigated from the time of ear emergence until maturity.
Trial Design	Randomised block, 6 reps sown, plots 20 x 1.2 m, entries 2 generations of 'Urambie' and comparators 'Tantangara', 'Tilga' and 'Yagan'.
Measurements	Taken 6 Sep 07, 9 Oct 07, and 13 Nov 07.
RHS Chart - edition	Nil

Origin and Breeding

Controlled pollination: The first cross 'Yagan'/'Ulandra' was made in spring 1987 and the F_1 was grown in the glasshouse the next year, when it was back-crossed in spring to 'Ulandra' (cross XB1106). 'Yagan' is a very early maturing semi-dwarf feed-grain type with large grain but very susceptible to leaf scald. 'Ulandra' is a late maturing winter variety, resistant to leaf scald and having a vernalisation requirement (period of cold growing conditions) for head initiation. 'Ulandra' has high yield potential but is too late for local growing areas. Both parental varieties have strong straw in the absence of diseases. Twelve BC_1F_1 plants were grown in the glasshouse over summer and then the 12 BC_1F_2 families were grown in separate field plots in 1989 in a mass selection trial. These were advanced in 1990 as F_3 mass selection plots and single head selections were taken from 6 of these plots. Four of the 1990 plots were selected and advanced as F_4 bulks in the 1991 mass selection trial, when single head selections were again taken. Single head selections were grown as hill plots at Wagga Wagga in 1991 and 1992 and selected hill plots harvested to provide seed for single observation and seed increase plots in an early sown stage 0 experiment in 1994. The early sown stage 0 experiment, W94, was sown in pedigree order with a grid of check varieties 'Ulandra', 'Franklin' and 'Skiff'. W94%175 was one of 7 selections derived from the same BC1F1 plant. It was identified as being an early maturing semi-dwarf with good straw strength and promoted to stage 1 early-sown trials. In 1995 stage 1 early-sown trials, sown on 3 sites, W94%175 was one of 84 selections derived from the back-cross XB1106. It was selected on the basis of its yield performance in an across-sites statistical analysis and promoted to stage 2 testing, 2 sites in 1996 and 3 in 1997. On the basis of its yield it was promoted to early-sown stage 3 testing in 1998, including some grazed experiments. W94%175 out-yielded 'Tantangara' and 'Gairdner' in an across-sites analysis which included 9 stage 3 experiments from 1998 and the stage 2 experiments from 1997. In 1999 W94%175 was given the synonym WB234, promoted to stage 4 early sown trials and included in the elite barley disease screening nursery (EBDSN). In 2000 to 2003 it continued to be tested in stage 4 early sown trials. Concurrent testing for grain quality was carried out by the Wagga Wagga Agricultural Institute malting laboratory.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Lowest leaves	hairiness of leaf sheaths	absent
Sterile spikelet	attitude	divergent
Grain	rachilla hair type	short
Grain	husk	present
Ear	number or rows	two
Time of	maturity	early
Season	type	spring type

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Yagan'	parent
'Tilga'	
'Tantangara'	

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comment
'Ulandra'	Time of	maturity	early	late	parental variety

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Urambie'	'Tantangara'	'Tilga'	'Yagan'
☐ *Plant: growth habit	semi-prostrate to prostrate	prostrate	erect	erect
☐ *Lowest leaves: hairiness of leaf sheaths	absent	absent	absent	absent
☑ *Flag leaf: anthocyanin colouration of auricles	absent	absent	present	present
☐ Plant: frequency of plants with recurved flag leaves	medium	medium	medium	medium
☐ Flag leaf: glaucosity of sheath	medium	medium	medium to strong	medium
☐ *Time of: ear emergence	early to medium	early	early to medium	early
☑ *Awns: anthocyanin colouration of tips	absent	present	present	present
☐ *Ear: glaucosity	absent or very weak	weak to medium	medium	weak
☐ Ear: attitude	semi-erect to horizontal	semi-erect	horizontal	semi-erect
☑ *Plant: length	short	short	long	medium
☐ *Ear: number of rows	two	two	two	two
☑ Ear: shape	tapering	parallel	parallel	tapering

*Ear: density	medium	medium to dense	dense	medium
Ear: length	medium	short	long	long
*Awn: length	medium	medium	medium	short
Rachis: length of first segment	short	medium	medium	medium
Rachis: curvature of first segment	absent or very weak	weak	very weak to weak	medium
*Sterile spikelet: attitude	divergent	divergent	divergent	divergent
Median spikelet: length of glume and its awn relative to grain	longer	shorter	shorter	equal
*Grain: rachilla hair type	short	short	short	short
*Grain: husk	present	present	present	present
Grain: anthocyanin colouration of nerves of lemma	absent or very weak	absent or very weak	absent or very weak	absent or very weak
Grain: spiculation of inner lateral nerves of dorsal side of lemma	absent or very weak	absent or very weak	absent or very weak	absent or very weak
*Grain: hairiness of ventral furrow	absent	absent	absent	absent
Grain: disposition of lodicules	frontal	frontal	frontal	frontal
Kernel: colour of aleurone layer	weakly coloured	weakly coloured	whitish	weakly coloured
*Season: type	spring type	spring type	spring type	spring type

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Urambie'	'Tantangara'	'Tilga'	'Yagan'
Flag leaf: length	very short	short	long	medium
Awn: length relative to ear	long	long	short to medium	short
Time of: maturity	early	early	early	early
Head: length	medium to long	medium	long	medium

Statistical Table

Organ/Plant Part: Context	'Urambie'	'Tantangara'	'Tilga'	'Yagan'
Flag leaf: length (mm)				
Mean	70.80	70.60	116.50	93.50
Std. Deviation	23.4	20.30	29.10	26.00
LSD/sig	21.0	ns	P≤0.01	P≤0.01
Plant: length (cm)				
Mean	67.40	64.50	82.30	74.70
Std. Deviation	2.4	4.40	3.60	6.80
LSD/sig	3.6	ns	P≤0.01	P≤0.01
Ear: length (mm)				
Mean	78.30	68.70	94.90	94.10
Std. Deviation	8.6	5.50	7.90	11.80
LSD/sig	6.9	P≤0.01	P≤0.01	P≤0.01

☑ Ear plus awn: length (mm)				
Mean	197.30	189.50	221.80	195.80
Std. Deviation	18.5	11.40	11.60	27.90
LSD/sig	14.9	ns	P≤0.01	ns
☑ Awn: length (mm)				
Mean	117.90	120.70	126.90	101.70
Std. Deviation	12.1	8.40	10.00	26.30
LSD/sig	12.4	ns	ns	P≤0.01
☑ Awn/ear: ratio				
Mean	1.49	1.76	1.35	1.11
Std. Deviation	0.14	0.15	0.17	0.34
LSD/sig	0.17	P≤0.01	ns	P≤0.01

Prior Applications and Sales

Nil.

Description: **Ross Downes,** Moruya, NSW.

Details of Application

Application Number	2006/299
Variety Name	'Pacific Ranger'
Genus Species	*Hordeum vulgare*
Common Name	Barley
Synonym	AC Ranger
Accepted Date	5 Feb 2007
Applicant	Her Majesty the Queen in Right of Canada as represented by the Minister of Agriculture and Agri-Food Canada
Agent	Pacific Seeds Pty Ltd, Toowoomba, QLD
Qualified Person	Peter Stuart

Details of Comparative Trial

Location	Gatton, QLD.
Descriptor	Barley (*Hordeum vulgare*) TG /19/10.
Period	Winter – Spring 2007. Sown 26 Apr 2007.
Conditions	The trial was sown into a well prepared seedbed at the Pacific Seeds Research Station, located at Gatton in the Lockyer Valley in South East Queensland. The trial was conducted under irrigated conditions using a row spacing of 76cm.
Trial Design	The trial design was a randomised complete block with four replications, four rows per plot, plots five metres long.
Measurements	Measurements were taken from 20 plants selected randomly from over 2,500 plants.
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: AC Ranger, is a six-row forage barley tested as EX 467-5 from 1993-1999, was developed from a single cross which was made in 1992 between PC 11 and AC Rosser. PC 11 is a selection from CIMMYT, Mexico City, Mexico, with resistance to QCCJ stem rust, *Puccinia graminis* Zhuk. Using standard pedigree selection, individual F_1 seed were grown to F_2 plants in the greenhouse. These, in turn, were individually harvested and grown as double 5 m F_3 in the field at Brandon. Individual heads were selected from each F_3 row and planted in the field as head rows, in Brandon. AC Ranger originated from a single F_4 head row selection, (Section 35) from the cross EX 467, with the designation EX 467-5. This row was selected on the basis of vigorous growth, straw strength, showing few disease symptoms, with dense foliage and numerous fertile spikes. EX 467-5, along with other selected lines, was then grown in replicated yield trials in the F_5 and F_6 at Brandon and Hamiota, Manitoba. Selection was on the basis of forage and feed quality (including percent crude protein, acid detergent fibre (ADF) and neutral detergent fibre (NDF)), height, lodging, maturity, test and thousand kernel weights, kernel plumpness and general disease resistance. Breeder: Dr. Mario Therrien, Agriculture and Agri-Food Canada, Brandon Research Centre, Brandon, MB, Canada.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Lowest leaves	hairiness of leaf sheaths	absent
Awns	anthocyanin colouration of tips	present
Awns	intensity of anthocyanin colouration of tips	medium
Grain	hairiness of ventral furrow	absent
Grain	husk	present
Season	type	spring

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Kaputar'	
'Corvette'	
'Mackay'	
'Dictator'	6 row barley, commercially used for hay and silage.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Pacific Ranger'	'Corvette'	'Dictator'	'Kaputar'	'Mackay'
☐ *Plant: growth habit	erect	erect	erect	erect	semi-erect
☐ *Lowest leaves: hairiness of leaf sheaths	absent	absent	absent	absent	absent
☑ *Flag leaf: anthocyanin colouration of auricles	absent	present	absent	absent	present
☐ Plant: frequency of plants with recurved flag leaves	high to very high	very high	very high	medium	high
☐ Flag leaf: glaucosity of sheath	medium	strong	strong	strong	very strong
☑ *Time of: ear emergence	very early to early	early	medium	medium	medium
☑ *Awns: anthocyanin colouration of tips	present	present		present	present
☑ *Awns: intensity of anthocyanin colouration of tips	medium	medium		medium	medium
☐ *Ear: glaucosity	weak	weak	absent or very weak	weak	weak
☐ Ear: attitude	semi-erect	semi-erect to horizontal	semi-erect	semi-erect to horizontal	semi-erect to horizontal
☐ *Plant: length	medium to long	short to medium	long to very long	very short to short	short to medium

☑ *Ear: number of rows	more than two	two	more than two	two	two
☑ Ear: shape	tapering	tapering	parallel	tapering	parallel
☐ *Ear: density	lax to medium	medium	medium	medium	medium
☑ Ear: length	short	long	short to medium	long	long to very long
☐ *Awn: length	short to medium	short to medium	n/a	short	short
☐ Rachis: length of first segment	short	short	long	short	short
☐ Rachis: curvature of first segment	weak to medium	weak to medium	medium to strong	weak to medium	weak to medium
☑ Median spikelet: length of glume and its awn relative to grain	longer	equal	equal	equal	shorter
☐ *Grain: rachilla hair type	long	long	short	long	long
☐ *Grain: husk	present	present	present	present	present
☐ *Grain: hairiness of ventral furrow	absent	absent	absent	absent	absent
☑ *Season: type	spring type	spring type	spring type	spring type	spring type

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Pacific Ranger'	'Corvette'	'Dictator'	'Kaputar'	'Mackay'
☑ Flag leaf: length	long to very long	medium	long	medium	short
☑ Flag leaf: intensity of anthocyanin coloration of auricles	absent to very weak	weak	absent	absent to very weak	medium
☑ Sterile spikelet: attitude (mid third of ear)	absent	divergent	absent	parallel to weakly divergent	divergent
☑ Flag leaf: width	wide to very wide	medium	wide to very wide	medium	narrow

Statistical Table

Organ/Plant Part: Context	'Pacific Ranger'	'Corvette'	'Dictator'	'Kaputar'	'Mackay'
☑ Plant (stem, ear, awns): length (cm)					
Mean	101.40	87.60	120.10	80.60	88.40
Std. Deviation	4.21	2.95	4.22	3.79	4.29
LSD/sig	1.79	P≤0.01	P≤0.01	P≤0.01	P≤0.01
☑ Ear: length (mm)					
Mean	94.70	120.10	98.40	120.20	138.30
Std. Deviation	7.49	10.14	9.88	7.31	12.13
LSD/sig	2.79	P≤0.01	P≤0.01	P≤0.01	P≤0.01

☑ Awn: length (mm)					
Mean	73.60	88.10	n/a	81.65	90.40
Std. Deviation	7.18	13.64	n/a	6.13	6.83
LSD/sig	6.5	P≤0.01	n/a	P≤0.01	P≤0.01
☑ Flag leaf: length (mm)					
Mean	270.90	199.60	239.00	196.70	180.20
Std. Deviation	22.76	44.36	27.76	42.42	22.13
LSD/sig	20.6	P≤0.01	P≤0.01	P≤0.01	P≤0.01
☑ Flag leaf: width (mm)					
Mean	23.10	17.00	24.00	16.30	13.40
Std. Deviation	1.61	3.10	3.82	2.94	1.57
LSD/sig	1.5	P≤0.01	ns	P≤0.01	P≤0.01

Prior Applications and Sales

Prior applications nil.

First sold in Canada in Jan 2003 under the name ‘AC Ranger’

Description: **Peter Stuart,** Pacific Seeds Pty Ltd, Toowoomba, QLD.

Details of Application

Application Number	2007/195
Variety Name	'Cheetah'
Genus Species	*Medicago truncatula* x *Medicago littoralis*
Common Name	Barrel Medic
Synonym	Nil
Accepted Date	5 Sep 2007
Applicant	Pristine Forage Technologies Pty Ltd, Daw Park, SA
Agent	N/A
Qualified Person	Andrew Lake

Details of Comparative Trial

Location	Currency Creek, SA.
Descriptor	Medics (new) (*Medicago* (excluding *M. sativa*)) TG/228/1
Period	Jun – Dec 2007.
Conditions	Seed was sown into jiffies in late Jun and transplanted into the field at Currency Creek, SA, in mid Jul. The soil was a moderately fertile, free draining sandy loam of approximately pH 6. Lime was added to the soil prior to planting to slightly raise pH. A mixed fertiliser (mainly P and trace elements) was used at plant out. Dacthal herbicide was applied two weeks post plant out. The trial was sprayed for red legged earthmite twice (early and late Aug). Plots were also sprayed with Fusilade for grass control (Sep) and hand weeded as required. The season had a wet start but a dry early finish with some hand watering carried out, followed by late rain. NB. The comparator 'Lynx' flowered earlier in this trial than has been observed in most other trials. Usually 'Lynx' flowers noticeably later than 'Cheetah'.
Trial Design	Randomised complete block with four replicates in single rows of 12 plants per replicate. 40 cm between rows; 20 cm between plants in rows.
Measurements	On individual plants or whole rows.
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: a pod holding selection from the species *M. littoralis* (MM 127) was crossed with a plant selected from a 'Caliph' based breeding population (~75% 'Caliph' parentage). The resultant F_1 plants were then crossed with 'Caliph'. Multiple selections of this cross were then further backcrossed to 'Caliph'. Progeny of these crosses were then allowed to self and were grown on to the F_2 and F_3 for selection. MZ-7 was a single plant selection (selection criteria; early flowering, plant type and vigour, pod holding) from the F_3 of one of the crosses produced in this way; code numbered MX-93. All crosses were carried out by hand with full emasculation to prevent selfing. At each stage of the process, progeny testing and pedigree selection were used to select for and track the (recessive) pod-holding gene, as well as to monitor and select for other characteristics and traits. Breeder: Andrew Lake and Ricki Drewry, Pristine Forage Technologies Pty Ltd, Daw Park, SA.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaflet	presence of marks	present on both sides
Leaflet	type of marks on upper side	flecked
Leaflet	position of marks on upper side	over whole surface
Leaflet	pubescence on upper side	present
Leaflet	pubescence on lower side	present
Pod	texture of whorl edges	spined
Mature pod	shedding	present
Mature leaf	shedding	present
Plant	aphid resistance	Blue Green Aphid (BGA) resistant

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Caliph'	
'Lynx'	new pod holding variety also in trial.
'Mogul'	included in trial as comparator for 'Lynx'

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Jaguar'	Leaf markings	no blotch	prominent blotch	This is one of a number of significant differences between the candidate and 'Jaguar'.
'Cyprus'	Plant aphid resistance	BGA resistant	BGA susceptible	'Cyprus' also not a pod or leaf holder.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Cheetah'	'Caliph'	'Lynx'	'Mogul'
☐ *Leaflet: presence of marks	present on both sides	present on both sides	present on both sides	present on both sides
☐ *Leaflet: type of marks on upper side	flecked	flecked	flecked	flecked
☐ *Leaflet: position of marks on upper side	over whole surface	over whole surface	over whole surface	over whole surface
☑ Leaflet: number of marks on upper side (varieties with spot or fleck type of marks on upper side only)	very few	very few	few	few
☑ Leaflet: number of marks on lower side (varieties with marks on lower side only)	very few to few	very few to few	many to very many	many to very many
☑ *Time of: flowering	very early	very early	early to medium	early to medium
☐ Plant: length of longest stem	long	long	medium to long	medium to long

☑ Plant: length of internode	long	long	medium	medium
☑ Runner: pubescence	medium to dense	medium to dense	very sparse to sparse	very sparse to sparse
☑ Leaflet: length	medium to long	medium to long	short to medium	short to medium
☑ Leaflet: width	narrow to medium	narrow to medium	medium to broad	medium to broad
☑ Leaflet: ratio length/width	medium to large	medium to large	small to medium	small to medium
☑ Leaflet: shape of base	narrow acute	narrow acute	broad acute	broad acute
☐ Leaflet: shape of apex	rounded	rounded	rounded	rounded
☑ Leaflet: serration of margin	coarse	coarse	medium	fine to medium
☐ *Leaflet: pubescence on upper side	present	present	present	present
☐ Leaflet: density of pubescence on upper side	dense	dense	dense	dense
☐ *Leaflet: pubescence on lower side	present	present	present	present
☐ Leaflet: density of pubescence on lower side	dense	dense	dense	dense
☐ Petiole: length	short to medium	short to medium	short	short
☐ Petiole: thickness	medium	medium	medium to thick	medium to thick
☐ Stipule: size	small	small	small to medium	small to medium
☐ Stipule: length of teeth	short	short	short to medium	short to medium
☐ Inflorescence: predominant number of florets	three	three	three	three
☐ Flower: intensity of yellow colour of petal	medium	medium	medium	medium
☐ Flower: marks on calyx	absent	absent	absent	absent
☑ Time of: physiological ripening of pods	late	early	late	medium to late
☑ Pod: length	medium	medium	short	short
☑ *Pod: shape	cylindrical	cylindrical	ovoid	ovoid
☐ Pod: compactness of whorls (excluding varieties with sickle-shaped pods)	compact	medium to compact	compact	medium to compact
☑ Pod: direction of whorls	clockwise	clockwise	clockwise	anti-clockwise
☐ Pod: number of whorls (excluding varieties with sickle-shaped pods)	three to five	three to five	three to five	three to five
☐ *Pod: texture of whorl edges (excluding varieties with sickle-shaped pods)	spined	spined	spined	spined

☐ Pod: length of spines (varieties with spined texture of whorl edges only)	short	short to medium	short	short to medium
☐ Pod: attitude of spines (varieties with spined texture of whorl edges only)	adpressed	adpressed	adpressed	oblique
☐ Pod: presence of apical hook on spines (varieties with spined texture of whorl edges only)	absent	absent	absent	absent
☑ Seed: 1000 seed weight	medium	low to medium	low	low

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Cheetah'	'Caliph'	'Lynx'	'Mogul'
☑ Mature pod: shedding	very low	high	low	very high
☑ Mature leaf: shedding	low	high	low	high

Statistical Table

Organ/Plant Part: Context	'Cheetah'	'Caliph'	'Lynx'	'Mogul'
☑ Flower: days to first flower				
Mean	69.49	71.13	70.97	82.38
Std. Deviation	1.41	1.72	0.82	1.09
LSD/sig	2.29	ns	ns	P≤0.01
☑ Pod: weight of 100 pods (g)				
Mean	12.00	12.20	9.16	10.50
Std. Deviation	0.83	1.04	0.60	0.62
LSD/sig	1.374	ns	P≤0.01	P≤0.01

Prior Applications and Sales

Nil.

Description: **Andrew Lake,** Pristine Forage Technologies Pty Ltd, Daw Park, SA.

Details of Application

Application Number	2007/194
Variety Name	'Lynx'
Genus Species	*Medicago truncatula* x *Medicago littoralis*
Common Name	Barrel Medic
Synonym	Nil
Accepted Date	5 Sep 2007
Applicant	Pristine Forage Technologies Pty Ltd, Daw Park, SA
Agent	N/A
Qualified Person	Andrew Lake

Details of Comparative Trial

Location	Currency Creek, SA.
Descriptor	Medics (new) (*Medicago* (excluding *M. sativa*)) TG/228/1
Period	Jun – Dec 2007
Conditions	Seed was sown into jiffies in late Jun and transplanted into the field at Currency Creek, SA, in mid Jul. The soil was a moderately fertile, free draining sandy loam of approximately pH 6. Lime was added to the soil prior to planting to slightly raise pH. A mixed fertiliser (mainly P and trace elements) was used at plant out. Dacthal herbicide was applied two weeks post plant out. The trial was sprayed for red legged earthmite twice (early and late Aug). Plots were also sprayed with Fusilade for grass control (Sep) and hand weeded as required. The season had a wet start but a dry early finish with some hand watering carried out, followed by late rain. NB. Lynx flowered comparatively earlier in this trial than has been observed in most other trials. Usually Lynx flowers noticably later than Cheetah.
Trial Design	Randomised complete block with four replicates in single rows of 12 plants per replicate. 40cm between rows; 20cm between plants in rows.
Measurements	On individual plants or whole rows.
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: a pod holding selection from the species *M. littoralis* (MM 126, subsequently registered for PBR protection as 'Jaguar') was crossed with a plant selected from a 'Mogul' barrel medic based breeding population (~87% 'Mogul' parentage). The resultant F_1 plants were then crossed with Mogul. Multiple selections of this cross were then further backcrossed to Mogul. Progeny of these crosses were then allowed to self and were grown on to the F_2 and F_3 for selection. MZ-8 was a single plant selection (selection criteria; flowering time, plant type and vigour, pod holding) from the F_3 of one of the crosses produced in this way; code numbered MX-101. All crosses were carried out by hand with full emasculation to prevent selfing. At each stage of the process, progeny testing and pedigree selection were used to select for and track the (recessive) pod-holding gene, as well as to monitor and select for other characteristics and traits. Breeder: Andrew Lake and Ricki Drewry, Pristine Forage Technologies Pty Ltd, Daw Park, SA.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaflet	presence of marks	present on both sides
Leaflet	type of marks on upper side	flecked
Leaflet	position of marks on upper side	over whole surface
Leaflet	pubescence on upper side	present
Leaflet	pubescence on lower side	present
Pod	texture of whorl edges	spined
Mature pod	shedding	present
Mature leaf	shedding	present
Plant	aphid resistance	Blue Green Aphid (BGA) resistant

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Mogul'	
'Caliph'	Included in trial as comparator for 'Cheetah'.
'Cheetah'	New pod holding variety also in trial.

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Jaguar'	Leaf markings	no spot, dense flecking	spot, sparse flecking	This is one of a number of significant differences between the candidate and 'Jaguar'.
'Borung'	Plant aphid resistance	BGA resistant	BGA susceptible	'Borung' also not a pod or leaf holder.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Lynx'	'Caliph'	'Cheetah'	'Mogul'
☐ *Leaflet: presence of marks	present on both sides	present on both sides	present on both sides	present on both sides
☐ *Leaflet: type of marks on upper side	flecked	flecked	flecked	flecked
☐ *Leaflet: position of marks on upper side	over whole surface	over whole surface	over whole surface	over whole surface
☑ Leaflet: number of marks on upper side (varieties with spot or fleck type of marks on upper side only)	few	very few	very few	few
☑ Leaflet: number of marks on lower side (varieties with marks on lower side only)	many to very many	very few to few	very few to few	many to very many
☑ *Time of: flowering	early to medium	very early	very early	early to medium
☐ Plant: length of longest stem	medium to long	long	long	medium to long

☑ Plant: length of internode	medium	long	long	medium
☑ Runner: pubescence	very sparse to sparse	medium to dense	medium to dense	very sparse to sparse
☑ Leaflet: length	short to medium	medium to long	medium to long	short to medium
☑ Leaflet: width	medium to broad	narrow to medium	narrow to medium	medium to broad
☑ Leaflet: ratio length/width	small to medium	medium to large	medium to large	small to medium
☑ Leaflet: shape of base	broad acute	narrow acute	narrow acute	broad acute
☐ Leaflet: shape of apex	rounded	rounded	rounded	rounded
☑ Leaflet: serration of margin	medium	coarse	coarse	fine to medium
☐ *Leaflet: pubescence on upper side	present	present	present	present
☐ Leaflet: density of pubescence on upper side	dense	dense	dense	dense
☐ *Leaflet: pubescence on lower side	present	present	present	present
☐ Leaflet: density of pubescence on lower side	dense	dense	dense	dense
☐ Petiole: length	short	short to medium	short to medium	short
☐ Petiole: thickness	medium to thick	medium	medium	medium to thick
☐ Stipule: size	small to medium	small	small	small to medium
☐ Stipule: length of teeth	short to medium	short	short	short to medium
☐ Inflorescence: predominant number of florets	three	three	three	three
☐ Flower: intensity of yellow colour of petal	medium	medium	medium	medium
☐ Flower: marks on calyx	absent	absent	absent	absent
☑ Time of: physiological ripening of pods	late	early	late	medium to late
☑ Pod: length	short	medium	medium	short
☑ *Pod: shape	ovoid	cylindrical	cylindrical	ovoid
☐ Pod: compactness of whorls (excluding varieties with sickle-shaped pods)	compact	medium to compact	compact	medium to compact
☑ Pod: direction of whorls	clockwise	clockwise	clockwise	anti-clockwise
☐ Pod: number of whorls (excluding varieties with sickle-shaped pods)	three to five	three to five	three to five	three to five
☐ *Pod: texture of whorl edges (excluding varieties with sickle-shaped pods)	spined	spined	spined	spined

☐ Pod: length of spines (varieties with spined texture of whorl edges only)	short	short to medium	short	short to medium
☐ Pod: attitude of spines (varieties with spined texture of whorl edges only)	adpressed	adpressed	adpressed	oblique
☐ Pod: presence of apical hook on spines (varieties with spined texture of whorl edges only)	absent	absent	absent	absent
☑ Seed: 1000 seed weight	low	low to medium	medium	low

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	**'Lynx'**	**'Caliph'**	**'Cheetah'**	**'Mogul'**
☑ Mature pod: shedding	low	high	very low	very high
☑ Mature leaf: shedding	low	high	low	high

Statistical Table

Organ/Plant Part: Context	**'Lynx'**	**'Caliph'**	**'Cheetah'**	**'Mogul'**
☑ Pod: weight of 100 pods (g)				
Mean	9.16	12.20	12.00	10.50
Std. Deviation	0.60	1.04	0.83	0.62
LSD/sig	1.374	P≤0.01	P≤0.01	ns
☑ Flower: days to first flower				
Mean	70.97	71.13	69.49	82.38
Std. Deviation	0.82	1.72	1.41	1.09
LSD/sig	2.29	ns	ns	P≤0.01

Prior Applications and Sales

Nil.

Description: **Andrew Lake,** Pristine Forage Technologies Pty Ltd, Daw Park, SA.

Details of Application

Application Number	2005/354
Variety Name	'Goldcog'
Genus Species	*Acacia cognata*
Common Name	Bower Wattle
Synonym	Nil
Accepted Date	9 Feb 2006
Applicant	Peter Goldup, Mt Evelyn, VIC
Agent	Bushland Flora, Mt Evelyn, VIC
Qualified Person	Mark Lunghusen

Details of Comparative Trial

Location	Mt Evelyn, VIC.
Descriptor	Acacia (*Acacia*) PBR ACAC.
Period	Autumn to spring 2007.
Conditions	Plants were grown in 14cm pots in full sun in commercial pine bark based potting mix with controlled release fertiliser. Plants were grown on benches with overhead watering.
Trial Design	10 plants in block design.
Measurements	Leaf measurements taken from largest leaves.
RHS Chart - edition	2005.

Origin and Breeding

Seedling selection: a compact seedling was selected from a batch of seedlings of *Acacia cognata* in 2000. The seed parent is characterised by tall plant height. Cuttings were taken from this seedling, established, and then another generation of cuttings were taken from the young plants. This was repeated to determine distinctness, uniformity and stability. To date, the plant has been grown through three generations with no off-types being recorded. Selection criteria: leaf colour. Propagation: vegetative. Breeder: Peter Goldup, Mt Evelyn, VIC.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Phyllode	width	medium
Plant	type	shrub
Plant	growth habit	bushy

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Bower of Beauty'	Closest variety based on all grouping characteristics.

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Limelight'	Phyllodes width	medium	narrow	'Limelight' has much finer leaves, is a smaller plant and the stems are more arching.
'Green Mist'	Phyllodes width	medium	narrow	'Green Mist' has narrower phyllodes and the stems are more arching.
'River Cascade'	Phyllodes width	medium	narrow	'River Cascade' has narrower phyllodes and the stems are more arching.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Goldcog'	'Bower of Beauty'
☐ Plant: type	shrub	shrub
☐ Plant: growth habit	bushy	bushy
☑ Plant: attitude of branches	semi-upright to upright	upright to spreading
☐ Plant: density of branches	strong	medium to strong
☐ Phyllode: shape	falcate	falcate
☑ Phyllode: colour of new growth (RHS colour chart)	green 144A	green N144A
☑ Phyllode: colour of mature leaf (RHS colour chart)	green 137A	green 139A
☐ Phyllode: variegation	absent	absent

Statistical Table

Organ/Plant Part: Context	'Goldcog'	'Bower of Beauty'
☑ Leaf: length (mm)		
Mean	49.74	47.04
Std. Deviation	4.24	4.16
LSD/sig	1.50	P≤0.01
☑ Leaf: width (mm)		
Mean	2.14	2.29
Std. Deviation	0.16	0.33
LSD/sig	0.11	P≤0.01
☐ Internode: length (mm)		
Mean	8.66	6.66
Std. Deviation	2.21	1.00
LSD/sig	2.03	ns
☐ Plant: height (mm)		
Mean	202.00	173.00
Std. Deviation	14.76	13.37
LSD/sig	31.79	ns

Plant: width (mm)		
Mean	320.00	324.00
Std. Deviation	32.66	32.04
LSD/sig	4.50	ns

Prior Applications and Sales

Nil.

Description: **Mark Lunghusen**, World Select Plants, Cranbourne, VIC.

Details of Application

Application Number	2007/245
Variety Name	'TF01'
Genus Species	*Stenotaphrum secundatum*
Common Name	Buffalo Grass
Synonym	Nil
Accepted Date	12 Nov 2007
Applicant	Transvaal Park Pty Ltd, Beadessert, QLD
Agent	N/A
Qualified Person	Matthew Roche

Details of Comparative Trial

Location	Queensland Department of Primary Industries & Fisheries, Redlands Research Station, Cleveland, QLD (Latitude 27°32′S, 153°15′E, elevation 25 masl).
Descriptor	Stenotaphrum (*Stenotaphrum secundatum*) PBR STEN
Period	13 Feb. 2006 – 15 Dec. 2006.
Conditions	Individual propagules (four per tube) were grown in 40x40mm tubes until covered and planted on a red volcanic (krasnozem) soil 13 Dec 2006; plants not defoliated; armyworm control by cyfluthrin 19 Oct 2006, weed control by pre-emergence oxadiazon and nutrition maintained by slow release fertiliser (18-10-9) at time of planting.
Trial Design	Thirty (30) spaced plants of each cultivar ('Sir James', University of Western Australia, 'ST-26', 'Sir Walter', 'Shademaster', 'Matilda', 'Sapphire', 'Kings Pride', 'Velvet', 'Palmetto', 'ST-135', 'EB-2', 'Marine', 'ST-91', 'ST-85', 'Ned Kelly') arranged in six (6) randomised blocks with five (5) plants per plot; 1.5m between plots, 1.5m between plants within plots.
Measurements	Four (4) diameter of spread measurements were taken per plant (11-12 Apr, 26 Apr and 11 May 2006); two (2) stolons per plant were collected 13-27 Jul 2006 and stolon and leaf characteristics were measured; two (2) shoot and inflorescence measurements per plant were taken 28 Nov to 14 Dec 2006; average sward height per plant 6 Nov 2006; inflorescence density (0.1125m2) per plant 15 Dec 2006; exposed stolon and leaf colour 18 Aug 2006.
RHS Chart - edition	2001.

Origin and Breeding

Chance seedling: 'TF01' was selected by the breeder, John Powell, as an isolated and distinctive plant of buffalo grass (*Stenotaphrum secundatum*) growing among kikuyu grass on the banks of the Bellinger River along its tidal reaches where it was occasionally inundated by brackish water during king tides. It showed shorter internodes than existing buffalo grass varieties of comparable texture within the breeder's knowledge, and showed good colour retention during periods of drought. Initially designated 'TF01', the buffalo grass cultivar was trialled for turf adaptation by Turf Force on their Beaudesert turf farm and characterised in a national buffalo grass project coordinated by the Queensland Department of Primary Industries and Fisheries Turf Research group initiated in 2005.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Turf	texture	coarse
Stigma	colour	purple

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'B12'	
'Kings Pride'	
'Marine'	
'Matilda'	
'Ned Kelly'	
'Shademaster'	
'Sir Walter'	
'ST-26'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'TF01'	'B12'	'Kings Pride'	'Marine'	'Matilda'	'Ned Kelly'	'Shademaster'	'Sir Walter'	'ST-26'
☐ Plant: habit	creeping								
☐ Plant: type	mat-forming								
☐ Plant: height	short								
☐ Plant: longevity	perennial								
☐ Plant: spreading	laterally by stolons								
☐ Stolon: nodes	compound nodes with 2 leaves								
☐ Stolon: internode length	medium to long								
☐ Stolon: internode thickness	medium to thick								
☑ Stolon: colour when exposed to sunlight	purple to brown (RHS N186C)	RHS N199A	RHS N186C	RHS 199B	RHS N186C	RHS N186C	RHS N186C	RHS N186C	RHS N199A
☐ Unmown culms: habit	decumbent								
☐ Unmown culms: branching	present								
☐ Unmown culms: length	medium								
☐ Unmown culms: leaves	distichous								

Leaf blade: texture of surface	glabrous									
Leaf blade: shape	linear									
Leaf blade: appearance	conduplicate									
Leaf blade: apex	obtuse									
Leaf blade: length	medium to short									
Leaf blade: width	medium to narrow									
Leaf blade: colour	green (RHS 137A)	RHS 137A	RHS 137A	RHS 137A	RHS 137A	RHS 137A	RHS 137B (outer margins RHS N186C)	RHS 137A	RHS 137A	
Leaf sheath: appearance	tightly compressed and keeled									
Leaf sheath: texture of surface	glabrous									
Ligule: hairs	fringe of hairs (ca 0.4-0.6 mm long)									
Inflorescence: position	terminal or axillary									
Inflorescence: type	laterally compressed solid panicle									
Inflorescence: central axis	flattened									

Inflorescence: texture	corky
Inflorescence: toughness	tough
Inflorescence: length of racemes	very short
Inflorescence: number of sessile spikelets per raceme	(1-) 3
Inflorescence: appearance of racemes	unilateral false spikes, sunken into central inflorescence axis
Spikelets: type	deciduous with accessory inflorescence branch structure
Spikelets: colour of stigmas	purple
Peduncle: length	medium
Peduncle: thickness	medium to coarse

Statistical Table

Organ/Plant Part: Context	'TF01'	'B12'	'Kings Pride'	'Marine'	'Matilda'	'Ned Kelly'	'Shademaster'	'Sir Walter'	'ST-26'
☑ Plant: mean diameter after 118 days									
Mean	174.50	121.20	167.40	107.40	160.00	159.60	129.70	142.00	100.00
Std. Deviation	30.19	12.97	30.52	15.96	18.20	37.06	35.86	50.66	5.93
LSD/sig	34.0	P≤0.01	ns	P≤0.01	ns	ns	P≤0.01	P≤0.01	P≤0.01
☑ Stolon node: first stolon node with a second lateral branch (spaced plants)									
Mean	1.18	1.10	1.40	1.40	1.83	1.50	1.73	1.12	0.95
Std. Deviation	0.81	0.75	0.69	0.83	0.81	0.75	0.73	0.67	0.73
LSD/sig	0.45	ns	ns	ns	P≤0.01	ns	P≤0.01	ns	ns
☑ Stolon node: first stolon node with a third lateral branch (spaced plants)									
Mean	2.05	2.07	2.08	2.22	2.40	2.10	2.25	2.02	2.00
Std. Deviation	0.43	0.31	0.28	0.56	0.62	0.44	0.51	0.29	0.37
LSD/sig	0.25	ns	ns	ns	P≤0.01	ns	ns	ns	ns
☑ Stolon node: first stolon node with a fourth lateral branch (spaced plants)									
Mean	2.25	2.32	2.10	2.65	2.57	2.27	2.43	2.08	2.17
Std. Deviation	0.47	0.47	0.30	0.73	0.56	0.63	0.53	0.38	0.42
LSD/sig	0.27	ns	ns	P≤0.01	P≤0.01	ns	ns	ns	ns
☑ Stolon node: first stolon node with a fifth lateral branch (spaced plants)									
Mean	2.43	2.38	2.03	2.88	2.77	2.25	2.78	2.13	2.18
Std. Deviation	0.50	0.64	0.37	0.96	0.65	0.65	0.72	0.39	0.57
LSD/sig	0.27	ns	P≤0.01	P≤0.01	P<=0.01	ns	ns	P≤0.01	ns
☑ Stolon node: first stolon node with a sixth lateral branch (spaced plants)									
Mean	2.52	2.58	2.28	3.65	2.68	2.57	3.23	2.07	2.48
Std. Deviation	0.60	0.85	0.58	1.56	0.83	1.05	1.41	0.45	0.75
LSD/sig	0.55	ns	ns	P≤0.01	ns	ns	P<=0.01	ns	ns
☑ Internode: length of fourth internode from stolon tip (mm)									
Mean	65.03	52.93	60.70	50.05	55.23	60.68	43.65	64.73	47.09
Std. Deviation	14.37	11.33	12.64	9.12	10.09	12.99	8.51	11.22	5.76
LSD/sig	5.54	P≤0.01	ns	P≤0.01	P≤0.01	ns	P≤0.01	ns	P≤0.01

☑ Internode: diameter of fourth internode from stolon tip (mm)									
Mean	2.72	2.87	2.90	2.97	2.77	2.96	3.04	2.82	3.10
Std. Deviation	0.40	0.38	0.52	0.30	0.49	0.49	0.39	0.50	0.47
LSD/sig	0.26	ns	ns	ns	ns	ns	P≤0.01	ns	P≤0.01
☑ Leaf sheath: length of leaf sheath on fourth visible node from stolon tip (mm)									
Mean	18.24	17.40	19.58	17.67	16.97	18.68	16.23	21.29	16.97
Std. Deviation	2.49	2.25	3.10	2.92	2.65	3.23	2.91	2.37	2.07
LSD/sig	1.36	ns	ns	ns	ns	ns	P≤0.01	P≤0.01	ns
☐ Flag leaf: length of blade on flag leaf on flowering tillers (mm)									
Mean	28.17	28.67	24.36	33.75	33.37	27.93	25.56	30.17	32.17
Std. Deviation	12.97	12.73	8.96	14.05	12.09	12.78	12.29	14.80	13.31
LSD/sig	10.79	ns	ns	ns	ns	ns	ns	ns	ns
☐ Flag leaf: width of blade on flag leaf on flowering tillers (mm)									
Mean	6.25	6.14	5.81	6.26	6.67	6.42	5.31	6.68	6.16
Std. Deviation	0.99	1.46	1.54	1.17	1.09	1.53	1.40	1.24	1.42
LSD/sig	0.98	ns	ns	ns	ns	ns	ns	ns	ns
☐ Flag leaf: length: width ratio of flag leaf blade on flowering tillers									
Mean	4.43	4.60	4.22	5.46	4.99	4.21	4.85	4.41	5.24
Std. Deviation	1.70	1.62	1.21	2.51	1.61	1.35	2.09	1.68	1.64
LSD/sig	1.47	ns	ns	ns	ns	ns	ns	ns	ns
☑ Leaf: length of sheath on fourth leaf on flowering tillers (mm)									
Mean	36.19	38.45	34.59	31.03	39.93	36.59	26.24	32.98	38.52
Std. Deviation	10.90	9.95	8.04	7.62	10.55	7.90	8.57	8.78	11.12
LSD/sig	8.33	ns	ns	ns	ns	ns	P≤0.01	ns	ns
☑ Leaf blade: length of blade on fourth leaf on flowering tillers (mm)									
Mean	80.08	69.47	66.49	65.06	85.57	75.94	46.58	75.01	74.20
Std. Deviation	28.81	25.44	17.76	20.00	25.52	26.25	21.64	29.26	26.20
LSD/sig	20.33	ns	ns	ns	ns	ns	P≤0.01	ns	ns
☐ Leaf blade: width of blade on fourth leaf on flowering tillers (mm)									
Mean	6.75	7.39	7.10	6.11	6.61	7.37	5.80	7.19	7.23
Std. Deviation	0.76	1.29	1.34	1.00	1.11	1.26	1.44	1.17	1.84

LSD/sig	1.18	ns	ns	ns	ns	ns	ns	ns	ns
☑ Leaf blade: length: width ratio of fourth leaf blade on flowering tillers									
Mean	12.08	9.66	9.62	10.94	13.41	10.56	8.26	10.67	10.54
Std. Deviation	4.79	4.22	2.97	3.71	4.00	4.11	3.76	4.41	3.65
LSD/sig	2.94	ns	ns	ns	ns	ns	P≤0.01	ns	ns
☑ Peduncle: length of peduncle (mm) on flowering tillers (mm)									
Mean	55.18	93.60	60.20	77.45	77.82	67.28	74.87	69.78	83.16
Std. Deviation	16.94	23.36	20.29	23.28	25.65	24.48	22.05	27.49	22.60
LSD/sig	17.74	P≤0.01	ns	P≤0.01	P≤0.01	ns	P≤0.01	ns	P≤0.01
☑ Peduncle: diameter of peduncle on flowering tillers (mm)									
Mean	1.37	1.44	1.63	1.32	1.44	1.36	1.62	1.64	1.52
Std. Deviation	0.18	0.36	0.46	0.24	0.22	0.20	0.41	0.38	0.30
LSD/sig	0.25	ns	P≤0.01	ns	ns	ns	P≤0.01	P≤0.01	ns
☑ Spike: mean primary spike length (mm)									
Mean	78.18	88.56	81.54	65.82	86.28	90.31	74.54	85.95	84.82
Std. Deviation	10.68	10.32	11.61	13.46	12.65	16.15	12.59	14.98	9.59
LSD/sig	9.05	P≤0.01	ns	P≤0.01	ns	P≤0.01	ns	ns	ns
☑ Spike: mean primary spike width (smaller) (mm)									
Mean	2.17	2.16	2.47	2.19	2.34	2.28	2.41	2.49	2.36
Std. Deviation	0.31	0.28	0.39	0.28	0.36	0.34	0.39	0.32	0.28
LSD/sig	0.23	ns	P≤0.01	ns	ns	ns	P≤0.01	P≤0.01	ns
☑ Spike: mean primary spike breadth (wider) (mm)									
Mean	4.15	4.61	4.55	4.30	4.32	4.47	4.43	4.76	4.64
Std. Deviation	0.48	0.59	0.70	0.53	0.68	0.60	0.66	0.55	0.75
LSD/sig	0.47	ns	ns	ns	ns	ns	ns	P≤0.01	P≤0.01
☐ Spike: number of spikes on tiller									
Mean	2.57	2.75	2.72	2.53	2.38	2.77	2.43	2.73	2.83
Std. Deviation	0.72	0.68	0.72	0.62	0.58	0.67	0.59	0.66	0.76
LSD/sig	0.43	ns	ns	ns	ns	ns	ns	ns	ns
☐ Inflorescence: count (0.1125m2 quadrat) 15 December 2006									
Mean	46.10	117.80	48.40	42.70	62.70	43.80	22.50	53.90	59.60

Std. Deviation	29.60	44.41	55.53	28.32	27.13	30.59	20.50	30.36	31.95
LSD/sig	33.85	P≤0.01	ns	ns	ns	ns	ns	ns	ns
☑ Sward: height (cm)									
Mean	35.90	30.40	34.81	26.76	37.65	30.98	21.57	35.81	24.47
Std. Deviation	4.72	9.92	9.35	6.53	7.59	14.98	9.42	6.78	4.15
LSD/sig	12.09	ns	ns	ns	ns	ns	P≤0.01	ns	ns
☑ Leaf blade: length of leaf blade on fourth visible node from stolon tip (mm)									
Mean	16.72	14.20	21.41	15.78	19.58	19.98	17.03	23.95	14.49
Std. Deviation	2.90	4.19	4.12	4.07	3.43	5.47	3.29	5.28	3.06
LSD/sig	2.39	P≤0.01	P≤0.01	ns	P≤0.01	P≤0.01	ns	P≤0.01	ns
☑ Leaf blade: width of leaf blade on fourth visible node from stolon tip (mm)									
Mean	6.47	5.81	7.01	6.42	6.65	6.61	6.30	7.36	5.76
Std. Deviation	0.62	1.11	0.90	0.89	0.78	1.12	0.94	0.86	0.94
LSD/sig	0.19	P≤0.01	P≤0.01	ns	P≤0.01	ns	ns	P≤0.01	P≤0.01
☑ Leaf blade: length:width ratio of leaf blade on fourth visible node from stolon tip									
Mean	2.58	2.44	3.05	2.48	2.97	2.99	2.72	3.24	2.52
Std. Deviation	0.35	0.50	0.45	0.84	0.54	0.50	0.47	0.53	0.32
LSD/sig	0.32	ns	P≤0.01	ns	P≤0.01	P≤0.01	ns	P≤0.01	ns
☑ Flag leaf: length of sheath on flag leaf on flowering tillers (mm)									
Mean	43.21	48.51	41.14	46.90	50.25	43.18	38.27	42.45	54.16
Std. Deviation	8.80	6.77	7.26	7.50	9.35	9.82	7.01	8.57	9.76
LSD/sig	6.99	ns	ns	ns	P≤0.01	ns	ns	ns	P≤0.01

Prior Applications and Sales

Nil.

Description: **M.B. Roche** and **D.S. Loch**, DPI&F Redlands Research Station, Cleveland, QLD.

Details of Application

Application Number	2006/030
Variety Name	'Black Scallop'
Genus Species	*Ajuga reptans*
Common Name	Bugle Bells
Synonym	Nil
Accepted Date	24 Mar 2006
Applicant	Mike Tristram, West Sussex, UK
Agent	Plants Management Australia, Wonga Park, VIC
Qualified Person	Steve Eggleton

Details of Comparative Trial

Location	Wonga Park VIC.
Descriptor	Ajuga (*Ajuga*) PBR AJUG.
Period	Feb 2007 to Oct 2007.
Conditions	Trial conducted in the open, plants propagated from cuttings during Feb 2007, transferred from plugs to 140mm pots in Apr 2007. Pots filled with soilless, pinebark based mix with controlled release fertilizers. Appropriate pest and disease treatments were applied as required .
Trial Design	Twelve pots of each variety in a completely randomised design.
Measurements	From ten plants randomly selected.
RHS Chart - edition	1995.

Origin and Breeding

Spontaneous mutation: was first observed as a whole plant at Binsted Nursery, Binsted, Arundel, West Sussex, England during 1998. It occurred in a batch of *Ajuga reptans* 'Braunherz' which had been grown on from tissue culture, produced in the breeder's own laboratory. This single plant was selected, isolated and grown on until 2000 when the first cuttings were taken. Selection criteria: leaf shape rounded, leaf colour very dark purple, plant density dense. Propagation: Since this initial propagation it has been regularly reproduced via cuttings. More than ten subsequent generations have all been found to be uniform and stable. Breeder: Mike Tristram, West Sussex, UK.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	presence of variegation	absent
Leaf	predominant colour of upper side	brown
Plant	growth habit	spreading
Petal	predominant colour of upper side	violet blue

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Braunhertz'	parental variety
'Evening Glow'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Black Scallop'	'Braunhertz'	'Evening Glow'
☐ *Plant: growth habit	spreading	spreading	spreading
☑ Leaf: shape	obovate	spathulate	elliptic
☑ Leaf: shape of apex	obtuse	obtuse	acute
☑ Leaf: shape of base	obtuse	attenuate	cuneate
☐ Leaf: incision of margin	present	present	present
☑ Leaf: depth of incision	medium	very shallow to shallow	very shallow to shallow
☐ Leaf: type of incision	crenate	crenate	crenate
☐ Leaf: undulation of the margin	weak	weak	very weak
☑ Leaf: glossiness of upper side	very strong	medium	weak
☐ Leaf: presence of variegation	absent	absent	absent
☐ Leaf: predominant colour of upper side (RHS colour chart)	brown 200A	brown 200A	brown 200A + 147A yellow - green
☐ Bract: shape	ovate	ovate	ovate
☑ Inflorescence: length of internode	short to very short	short to very short	medium
☑ Petal: predominant colour of upper side (RHS colour chart)	violet-blue 90A	violet-blue 90A	violet-blue 93B

Prior Applications and Sales

Country	Year	Current Status	Name Applied
US	2004	Granted	'Black Scallop'

First sold in USA in Sep 2003. First Australian sale Mar 2005.

Description: **Steve Eggleton,** Wonga Park, VIC.

Details of Application

Application Number	2007/058
Variety Name	'Argyle'
Genus Species	*Brassica napus*
Common Name	Canola
Synonym	Nil
Accepted Date	8 Mar 2007
Applicant	Canola Breeders Western Australia Pty Ltd, Shenton Park, WA
Agent	N/A
Qualified Person	Milton Sanders

Details of Comparative Trial

Location	Shenton Park, Perth, WA.
Descriptor	Canola/Rape Seed (*Brassica napus*) TG/36/6+corr.
Period	25 May 2007 – 7 Nov 2007.
Conditions	Seeds were sown into the ground and then grown under normal winter-spring conditions, following normal agronomic practices for canola in Perth, Western Australia.
Trial Design	Randomised complete block design with 3 replicates with at least 70 plants per replicate sown in 8 m rows.
Measurements	Measurements were made on 20 random plants per replication, over three replications.
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: The cross 02N199 was made in 2002 in Perth, Western Australia. During 2003, doubled haploid progeny were developed by microspore tissue culture from the F1 of this cross. Doubled haploid progeny were selected for blackleg resistance in a disease nursery and pure seed was increased in pollination bags over winter 2004. Progeny were further selected for oil and protein in seed, and selections were bulked in pollination tents over summer 2004/05. One of the doubled haploid progeny, N03D-0339, was tested for yield and quality in replicated field trials at 10 locations across Southern Australia in 2005 and 2006, and for blackleg resistance in parallel blackleg disease nurseries. N03D-0339 was among the highest yielding and highest seed oil lines in these trials, with moderate blackleg resistance and tolerance to triazine herbicides. Pure seed production of N03D-0339 continued in a large pollination tent over summer 2006/07, and in a 1-ha Pre-Basic seed production block in 2007, where <0.1% tall late types were observed. The variety is early-mid season flowering with medium height. Breeder: Wallace A Cowling, Canola Breeders Western Australia Pty Ltd, Shenton Park, WA.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	height	medium
Plant	herbicide tolerance	triazine tolerant
Seed	size	small
Time of	flowering	medium/medium to late

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Surpass 501TT'	
'Lantern'	
'Thunder TT'	
'ATR-Beacon'	
'Tribune'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Argyle'	'ATR-Beacon'	'Lantern'	'Surpass 501TT'	'Thunder TT'	'Tribune'
☐ *Seed: erucic acid	absent	absent	absent	absent	absent	absent
☑ *Leaf: green colour	dark	medium	medium	medium	medium	dark
☐ *Leaf: lobes	present	present	present	present	present	present
☐ *Leaf: number of lobes	very few to few	medium	very few to few	few	very few to few	very few to few
☑ *Leaf: dentation of margin	weak	weak	medium	weak to medium	medium	medium
☐ *Time of: flowering	medium	early to medium	medium	medium to late	medium	medium
☐ *Flower: colour of petals	yellow	yellow	yellow	yellow	yellow	yellow
☐ Flower: length of petals	long	long	long	long	medium to long	long
☐ Flower: width of petals	medium to broad	broad	medium to broad	medium to broad	broad	medium to broad
☑ Plant: height at full flowering	medium	medium	tall	tall to very tall	tall	medium
☐ *Plant: total length including side branches	medium to long	medium	medium	medium to long	medium	medium to long
☑ Siliqua: length	long	long	long	long	long to very long	long to very long
☑ Siliqua: length of beak	short	short	medium	short	medium	long

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Argyle'	'ATR-Beacon'	'Lantern'	'Surpass 501TT'	'Thunder TT'	'Tribune'
☐ Plant: herbicide tolerance	triazine tolerant	triazine tolerant	triazine sensitive	triazine tolerant	triazine tolerant	triazine tolerant
☐ Seed: oil quality	canola quality	canola quality	canola quality	canola quality	canola quality	canola quality
☑ Plant: blackleg resistance	moderate to high	moderate	moderate	low to moderate	moderate to high	moderate
☑ Seed: colour	brown	black	black	black	black	black

Statistical Table

Organ/Plant Part: Context	'Argyle'	'ATR-Beacon'	'Lantern'	'Surpass 501TT'	'Thunder TT'	'Tribune'
☑ Flower: petal length (mm)						
Mean	16.47	16.18	16.10	16.37	15.47	16.12
Std. Deviation	1.20	1.07	1.08	1.13	1.20	1.06
LSD/sig	0.72	ns	ns	ns	P≤0.01	ns
☑ Flower: petal width (mm)						
Mean	8.05	9.18	7.38	8.50	9.42	7.53
Std. Deviation	0.87	1.10	0.96	0.79	0.70	0.81
LSD/sig	0.80	P≤0.01	ns	ns	P≤0.01	ns
☑ Plant: height (cm)						
Mean	97.50	117.00	135.10	150.20	130.70	92.25
Std. Deviation	9.40	19.00	15.50	15.50	12.60	15.56
LSD/sig	21.63	ns	P≤0.01	P≤0.01	P≤0.01	ns
☑ Plant: length (cm)						
Mean	80.80	57.67	56.18	65.12	58.82	72.87
Std. Deviation	14.92	17.43	14.65	18.37	14.29	16.66
LSD/sig	17.27	P≤0.01	P≤0.01	ns	P≤0.01	ns
☑ Siliqua: length (mm)						
Mean	68.07	63.38	72.15	69.52	77.77	77.77
Std. Deviation	5.12	5.39	7.47	5.25	8.39	6.32
LSD/sig	5.3	ns	ns	ns	P≤0.01	P≤0.01
☑ Siliqua: length of beak (mm)						
Mean	10.78	11.32	14.30	10.97	15.72	17.40
Std. Deviation	1.84	2.20	2.31	1.94	2.34	2.12
LSD/sig	1.59	ns	P≤0.01	ns	P≤0.01	P≤0.01

Prior Applications and Sales

Nil.

Description: **Wallace Cowling and Rozlyn Ezzy,** Canola Breeders Western Australia Pty Ltd, Shenton Park, WA

Details of Application

Application Number	2007/030
Variety Name	'Fragrant Angel'
Genus Species	*Echinacea purpurea*
Common Name	Coneflower
Synonym	Nil
Accepted Date	13 Feb 2007
Applicant	Terra Nova Nurseries, Inc, Tigard, Oregon, USA
Agent	Lifetech Laboratories Ltd, Kincumber, NSW
Qualified Person	Ian Paananen

Details of Comparative Trial

Overseas Testing Authority	US Patent and Trademark Office
Overseas Data Reference Number	PP16,054
Location	Macmasters Beach, NSW
Descriptor	Echinacea (*Echinacea purpurea*) PBR CONE
Period	Summer 2006-2007.
Conditions	Trial conducted in a open beds, plants propagated from micropropagation, planted into 140mm pots filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by capillary method, pest and disease treatments as required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random.
RHS Chart - edition	2001.

Origin and Breeding

Induced mutation: maternal parent 'Ruby Giant'. The parent is characterised by a light pink ray floret colour. Selection took place in Canby, Oregon, USA in 2002. Selection criteria: Flower: colour white. Propagation: vegetative divisions and micropropagation were found to be uniform and stable. Breeder: Harini Korlipara, Canby, Oregon, USA.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Ray floret	colour	white
Ray floret	attitude	horizontal
Disc floret	Colour	yellow orange

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Alba'	

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Prima Donna White'	Ray florets	number of rows	two	one
'Kim's Mop Head'	ray floret	attitude	horizontal	drooping
'White Swan'	Ray florets	number of rows	two	one
'Cygnet White'	Ray florets	number of rows	two	one

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Fragrant Angel'	'Alba'
☑ Plant: height	medium	tall
☐ Plant: 2. number of flower heads per stem	more than two	
☐ Basal leaf: length	medium to long	medium to long
☐ Basal leaf: width	medium	
☐ Basal leaf: shape	ovate	
☐ Basal leaf: margin	dentate	
☐ Basal leaf: pubescence (lower side)	absent or very weakly expressed	
☐ Basal leaf: colour (upper side)	medium green	
☐ Flower head: height	medium	medium
☐ Flower head: diameter	medium to large	medium to large
☐ Flower head: length of peduncle	medium	medium to long
☐ Ray floret: attitude	horizontal	horizontal
☐ Ray floret: length	short to medium	
☐ Ray floret: main colour (RHS Colour Chart)	155D	155D
☐ Ray floret: greenish colour of apex	present	
☐ Disc floret: colour	yellow orange	yellow orange
☐ Disc floret: time of beginning of flowering	summer to autumn	
☐ Flower: fragrance	present	
☐ Anther: colour	yellow	
☐ Peduncle: colour (RHS colour chart)	146C-D	
☑ Ray florets: number	many	few
☑ Ray florets: number of rows	two	one
☐ Disc: shape	convex to conic	

Prior Applications and Sales

Country	Year	Current Status	Name Applied
New Zealand	2005	Applied	'Fragrant Angel'
EU	2004	Granted	'Fragrant Angel'
USA	2004	Granted	'Fragrant Angel'

First sold in USA in Jul 2004. First Australian sale Feb 2006.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW.

Details of Application

Application Number	2005/063
Variety Name	'Jel01'
Genus Species	*Cordyline australis*
Common Name	Cordyline
Synonym	Nil
Accepted Date	21 Apr 2005
Applicant	Geoff Jewelll, Otaki, New Zealand
Agent	Anthony Tesselaar Plants Pty Ltd, Silvan, VIC
Qualified Person	Christopher Prescott

Details of Comparative Trial

Location	145 Moores road, Clyde, VIC (Latitude 38º09' South, elevation 16m).
Descriptor	Cordyline (*Cordyline* spp.) PBR CORD.
Period	2006/2007.
Conditions	The trial was carried out on two to three year old plants in the soil. Maintenance was unnecessary irrigation as plants required. The examination data was collected on 28 Nov 2007.
Trial Design	Plants were set out in blocks, 9 plants of 'Jel01', 9 plants of 'Kau01' and 6 plants of 'Dominator'.
Measurements	Measurements were taken at random with the assistance of examiners from the PBR office on 28 Nov 2007 after first flowering (with the exception of 'Kau01' which has not flowered in the duration of the trial).
RHS Chart - edition	2001.

Origin and Breeding

Spontaneous mutation: *Cordyline australis* 'Jel01' was selected in Otaki, New Zealand as a sport of a wild population of *Cordyline australis* 'Purpurea', by Geoff Jewell. The new variety was selected from amongst thousands of seedlings that had been cultivated from seeds collected in the wild. Selection criteria: upright growth habit, foliage colour. Propagation: all future generations have been propagated by tissue culture, and have remained true to type with no recordings of variation from the initial selection. Breeder: All work has been conducted by Geoff Jewell, settlement Road, Otaki, new Zealand.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	colour	burgundy

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Kau01'	
'Dominator'	

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Purpurea'	leaf colour	burgundy	brown

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Jel01'	'Dominator'	'Kau01'
☐ Stem: branching	absent	absent	absent
☐ Leaf: number of colours on upper side	one	one	one
☑ Leaf: main colour of upper side (RHS Colour Chart)	N200A	200A + redder	N200A
☐ Leaf: attitude of bottom half of leaf	erect to semi-erect	semi-erect	erect to semi-erect
☑ Leaf: attitude of top half of leaf	weeping	semi-weeping	semi-weeping
☐ Plant: suckering	absent	absent	absent
☑ Leaf: glossiness of upper side	weak	medium	weak

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Jel01'	'Dominator'	'Kau01'
☑ Leaf: mid rib colour on under side (RHS)	187A	187B	200A
☐ Young leaf: anthocyanin colouration	reddish brown	green	green
☑ Leaf: ridging on upper side	weak	medium	weak
☑ Plant: size	large	medium	small

Statistical Table

Organ/Plant Part: Context	'Jel01'	'Dominator'	'Kau01'
☑ Plant: height (cm)			
Mean	259.88	226.17	192.44
Std. Deviation	24.20	11.00	35.05
LSD/sig	49.27	ns	P≤0.01
☑ Leaf : length (cm)			
Mean	95.95	106.08	98.61
Std. Deviation	3.43	11.22	5.12
LSD/sig	3.95	P≤0.01	ns

Prior Applications and Sales

Country	Year	Current Status	Name Applied
New Zealand	2006	Applied	'Jel01'
EU	2006	Applied	'Jel01'
USA	2006	Applied	'Jel01'

Prior sale nil.

Description: **Christopher Prescott,** Prescott Roses Pty Ltd, Clyde, VIC.

Details of Application

Application Number	2006/126
Variety Name	'Kau01'
Genus Species	*Cordyline australis*
Common Name	Cordyline
Synonym	Nil
Accepted Date	5 Aug 2006
Applicant	Kauri Park Nurseries Ltd, Maungaturoto, New Zealand
Agent	Greenhills Propagation Nursery Pty Ltd, Tynong, VIC
Qualified Person	Christopher Prescott

Details of Comparative Trial

Location	145 Moores Road, Clyde, VIC (Latitude 38º09' South, elevation 16m).
Descriptor	Cordyline (*Cordyline* spp.) PBR CORD.
Period	2006/2007.
Conditions	The trial was carried out on two to three year old plants in the soil. Maintenance was unnecessary irrigation as plants required. The examination data was collected on 28 Nov 2007.
Trial Design	Plants were set out in blocks, 9 plants of 'Jel01', 9 plants of 'Kau01' and 6 plants of 'Dominator'.
Measurements	Measurements were taken at random with the assistance of examiners from the PBR office on 28 Nov 2007 after first flowering (with the exception of 'Kau01' which has not flowered in the duration of the trial) .
RHS Chart - edition	2001.

Origin and Breeding

Spontaneous Mutation: *Cordyline australis* 'Kau01' was selected in Maungaturoto, New Zealand as a seedling mutation in a population of *Cordyline australis* 'Purpurea' by Vern Wearmouth in 2001. The new variety was selected from amongst thousands of seedlings that had been cultivated from seeds collected from plants of the parent, in the nursery at Kauri Park Nurseries Ltd. Selection criteria: strong growth, foliage colour. Propagation: Future generations have been propagated by tissue culture to build stock and then by cuttings, and has remained true to type with no recordings of variation from the initial selection. Breeder: All work has been conducted by Vern Wearmouth, Maungaturoto, New Zealand.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	colour	burgundy

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Jel01'	
'Dominator'	

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Purpurea'	Leaf colour	burgundy	brown

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Kau01'	'Dominator'	'Jel01'
☐ Stem: branching	absent	absent	absent
☐ Leaf: number of colours on upper side	one	one	one
☑ Leaf: main colour of upper side (RHS Colour Chart)	N200A	200A + redder	N200A
☐ Leaf: attitude of bottom half of leaf	erect to semi-erect	semi-erect	erect to semi-erect
☑ Leaf: attitude of top half of leaf	semi-weeping	semi-weeping	weeping
☐ Plant: suckering	absent	absent	absent
☑ Leaf: glossiness of upper side	weak	medium	weak

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Kau01'	'Dominator'	'Jel01'
☑ Leaf: ridging on upper side	weak	medium	weak
☑ Plant: size	small	medium	large
☑ Leaf: mid rib colour on under side (RHS)	200A	187B	187A
☐ Young leaf: anthocyanin colouration	green	green	reddish brown

Statistical Table

Organ/Plant Part: Context	'Kau01'	'Dominator'	'Jel01'
☑ Leaf: length (cm)			
Mean	98.61	106.08	95.95
Std. Deviation	5.12	11.22	3.43
LSD/sig	3.95	P≤0.01	ns
☑ Plant: height (cm)			
Mean	192.44	226.17	259.88
Std. Deviation	35.05	11.00	24.20
LSD/sig	49.27	ns	P≤0.01
☐ Leaf: width at widest part (cm)			
Mean	6.52	6.28	6.08
Std. Deviation	0.39	0.48	0.91
LSD/sig	1.50	ns	ns

Prior Applications and Sales

Prior applications nil. First sold in Australia in June 2005.

Description: **Christopher Prescott,** Prescott Roses Pty Ltd, Clyde, VIC.

Details of Application

Application Number	2004/133
Variety Name	'BRA01'
Genus Species	*Cordyline fruticosa*
Common Name	Cordyline
Synonym	Nil
Accepted Date	22 Apr 2005
Applicant	Peter Brauns, Edmonton, QLD
Agent	Anthony Tesselaar Plants Pty Ltd, Silvan, VIC
Qualified Person	Christopher Prescott

Details of Comparative Trial

Location	145 Moores Road, Clyde, VIC (Latitude 38º09' South, elevation 16m).
Descriptor	Cordyline (*Cordyline* spp.) PBR CORD.
Period	2006/2007.
Conditions	The trial was conducted in a controlled environment polyhouse with shade, temperature ranged between 15 and 36 degrees Celsius within the 6 weeks prior to examination with plants on their own roots planted into 210mm pots (1 to 2 plants per pot) filled with a co-co coir mix, nutrition was maintained as part of a commercial hydroponic system, pest and disease treatments applied as required.
Trial Design	Pots were on hydroponic benches in a dual row, 6 plants of Cordyline 'Bra01' and 6 plants of Cordyline 'Nigra'.
Measurements	Measurements were taken from 2 year old plants at random on 28/11/2007.
RHS Chart - edition	2001.

Origin and Breeding

Spontaneous mutation: 'Bra01' was selected as a purple/black mutation of a wild green *Cordyline fruiticosa* by Peter Brauns of Plant Source Australia. Propagation of the new variety is by cutting, and has remained true to type over several generations. Selection criteria: compact growth habit, foliage colour. Propagation: vegetative. Breeder: Peter Brauns, Edmonton, QLD.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	colour	blackish

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Nigra'	

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Rubra'	leaf	colour	black look	red to burgundy	'Rubra' was the closest *C. fruticosa* in colour to *C.* 'Bra01' but was easily distinguished as being a different variety.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'BRA01'	'Nigra'
☐ Stem: branching	absent	absent
☐ Leaf: number of colours on upper side	one	one
☐ Leaf: main colour of upper side (RHS Colour Chart)	202A with reddish hue	202A with reddish hue
☐ Leaf: attitude of bottom half of leaf	erect	erect to semi-erect
☑ Leaf: attitude of top half of leaf	semi-erect	horizontal
☑ Plant: suckering	present	absent
☐ Leaf: glossiness of upper side	medium	medium

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'BRA01'	'Nigra'
☑ Leaf: cross section	convex	slightly concave
☑ Leaf: undulation of margin	strong	weak

Statistical Table

Organ/Plant Part: Context	'BRA01'	'Nigra'
☐ Plant: height (cm)		
Mean	105.72	131.47
Std. Deviation	12.37	11.18
LSD/sig	26.47	ns
☑ Leaf: length (cm)		
Mean	43.70	88.75
Std. Deviation	3.19	3.25
LSD/sig	5.89	P≤0.01
☑ Leaf: width at broadest part (cm)		
Mean	8.30	12.43
Std. Deviation	0.65	0.72
LSD/sig	1.26	P≤0.01

Prior Applications and Sales

Country	Year	Current Status	Name Applied
New Zealand	2004	Granted	'BRA01'
EU	2005	Applied	'BRA01'

First sold in Australia in Sep 2003 under the name 'Cobra'.

Description: **Christopher Prescott,** Prescott Roses Pty Ltd, Clyde, VIC**.**

Details of Application

Application Number	2005/121
Variety Name	'Uto01'
Genus Species	*Cordyline* hybrid
Common Name	Cordyline
Synonym	Nil
Accepted Date	26 Oct 2006
Applicant	Utopia Palms and Cycads, Valdora, QLD
Agent	N/A
Qualified Person	Christopher Prescott

Details of Comparative Trial

Location	145 Moores road, Clyde, VIC (Latitude 38º09' South, elevation 16m).
Descriptor	Cordyline (*Cordyline* spp.) PBR CORD.
Period	2006/2007.
Conditions	The trial was conducted in a controlled environment polyhouse with shade, temperature ranged between 15 and 36 degrees Celsius within the 6 weeks prior to examination with plants on their own roots planted into 210mm pots (1 to 2 plants per pot) filled with a co-co coir mix, nutrition was maintained as part of a commercial hydroponic system, pest and disease treatments applied as required.
Trial Design	Pots were on hydroponic benches in a dual row, 6 plants of Cordyline 'Uto01', 6 plants of Cordyline 'Bra01' and 6 plants of Cordyline 'Nigra'.
Measurements	Measurements were taken from 2 year old plants at random on 28 Nov 2007.
RHS Chart - edition	2001.

Origin and Breeding

Controlled pollination: Cordyline 'Uto01' was the result of a cross between *Cordyline fruticose* 'Bra01' (seed parent) and a *Cordyline terminalis* seedling (pollen parent) at the end of Apr 2000. Subsequent generations have been shown to be stable, with no off types noted. Selection criteria: upright growth habit, foliage colour. Breeder: All work has been conducted by Clayton Hank York, proprietor of Utopia Palms & Cycads, Valdora, QLD.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	colour	blackish

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Bra01'	seed parent
'Nigra'	pollen parent

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Stricta'	leaf colour	black look	green	'Stricta' was considered due to similar leaf width, but was rejected due to distinct colour difference.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Uto01'	'Bra01'	'Nigra'
☐ Stem: branching	absent	absent	absent
☐ Leaf: number of colours on upper side	one	one	one
☐ Leaf: main colour of upper side (RHS Colour Chart)	202A with reddish hue	202A with reddish hue	202A with reddish hue
☐ Leaf: attitude of bottom half of leaf	erect to semi-erect	erect	erect to semi-erect
☑ Leaf: attitude of top half of leaf	horizontal	semi-erect	horizontal
☑ Plant: suckering	present	present	absent
☐ Leaf: glossiness of upper side	medium	medium	medium

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Uto01'	'Bra01'	'Nigra'
☑ Leaf: cross section	slightly concave	convex	slightly concave
☑ Leaf: undulation of margin	weak	strong	weak

Statistical Table

Organ/Plant Part: Context	'Uto01'	'Bra01'	'Nigra'
☑ Plant: height (cm)			
Mean	146.62	105.72	131.47
Std. Deviation	18.66	12.37	11.18
LSD/sig	50.44	P≤0.01	ns
☑ Leaf: length (cm)			
Mean	70.65	43.70	88.75
Std. Deviation	3.46	3.19	3.25
LSD/sig	5.61	P≤0.01	P≤0.01
☑ Leaf: width at broadest part (cm)			
Mean	2.47	8.30	12.43
Std. Deviation	0.45	0.65	0.72
LSD/sig	1.47	P≤0.01	P≤0.01

Prior Applications and Sales

Nil.

Description: **Christopher Prescott,** Prescott Roses Pty Ltd, Clyde, VIC.

Details of Application

Application Number	2007/010
Variety Name	'Tana'
Genus Species	*Cordyline* hybrid
Common Name	Cordyline
Synonym	Renegade
Accepted Date	25 Jan 2007
Applicant	Evan David Lloyd, Ashhurst, New Zealand.
Agent	Greenhills Propagation Nursery Pty Ltd, Tynong, VIC.
Qualified Person	Mark Lunghusen

Details of Comparative Trial

Location	Greenhills Propagation Nursery, Tynong, VIC.
Descriptor	Cordyline (*Cordyline* spp.) PBR CORD.
Period	Spring/summer 2007.
Conditions	Plants were grown in 14cm pots in full sun in commercial pine bark based potting mix with controlled release fertiliser. Plants were grown on benches with overhead watering.
Trial Design	10 plants in block design.
Measurements	Leaf measurements taken from widest part of leaf
RHS Chart - edition	2005.

Origin and Breeding

Seedling selection: a seedling was selected from a batch of seedlings of *Cordyline australis* in 2000. The seed parent is characterised by green to bronze foliage colour. Divisions were taken from this seedling, established, and then another generation of divisions were taken from the young plants. This was repeated to determine distinctness, uniformity and stability. To date, the plant has been grown through six generations with no off-types being recorded. Selection criteria: leaf colour. Propagation: vegetative. Breeder: Evan Lloyd, Ashhurst, New Zealand.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	colour	brown

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Purple Sensation'	
'Red Star'	

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Red Chocolate'	Plant height	medium	large	Also known as 'New Red', or 'Cardinal' and other names.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Tana'	'Purple Sensation'	'Red Star'
☐ Stem: branching	absent	absent	absent
☑ Leaf: number of colours on upper side	one	two	two
☑ Leaf: main colour of upper side (RHS Colour Chart)	brown 200A	brown 200B	177A
☑ Leaf: secondary colour of upper side (RHS Colour Chart)	brown 200A	greyed red 178C	178D
☐ Leaf: attitude of bottom half of leaf	semi-erect	semi-erect	semi-erect
☐ Leaf: attitude of top half of leaf	horizontal	semi-erect	semi-erect
☐ Plant: suckering	absent	absent	absent
☑ Leaf: glossiness of upper side	medium	weak	weak

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Tana'	'Purple Sensation'	'Red Star'
☑ Leaf: stiffness	weak	strong	medium
☑ Plant: size	small	medium	small

Statistical Table

Organ/Plant Part: Context	'Tana'	'Purple Sensation'	'Red Star'
☑ Leaves: number per plant (mm)			
Mean	14.30	18.70	32.40
Std. Deviation	1.57	1.83	4.06
LSD/sig	6.64	ns	P≤0.01
☑ Plant: height (mm)			
Mean	423.00	689.00	405.00
Std. Deviation	36.50	38.43	18.41
LSD/sig	40.15	P≤0.01	ns
☑ Leaf: width (mm)			
Mean	20.29	17.51	15.99
Std. Deviation	1.63	1.38	1.35
LSD/sig	1.80	P≤0.01	P≤0.01

Prior Applications and Sales

Country	Year	Current Status	Name Applied
New Zealand	2006	Applied	'Tana'

Prior sale nil.

Description: **Mark Lunghusen,** World Select Plants, Cranbourne, VIC.

Details of Application

Application Number	2007/300
Variety Name	'HYPERNO'
Genus Species	*Triticum turgidum* ssp *turgidum*
Common Name	Durum Wheat
Synonym	Nil
Accepted Date	12 Dec 2007
Applicant	Australian Grain Technologies Pty Ltd, Glen Osmond, SA
Agent	N/A
Qualified Person	Gil Hollamby

Details of Comparative Trial

Location	Mintaro, South Australia.
Descriptor	Durum wheat (*Triticum durum*) TG/120/3.
Period	2007.
Conditions	The trial was grown in a black self mulching soil which had been pasture in 2006 and wheat in 2005. The area was sprayed with Roundup Power Max (1.2L/ha)+Goal CT(75ml/ha) on 24 May 2007 and direct drilled at 2-4cm in slightly moist conditions on 25 May at 200 plants/m2 and with 90kg/ha DAP and 80kg/ha Urea. During the winter months moisture was adequate and the trial grew well. In crop weeds were controlled with 2,4-D amine 625(1.5l/ha) on 6 Sep. Spring was dry and some moisture stress occurred. Harvest took place on 11 Dec about two weeks earlier than normal. There were no diseases of note. A similar trial was planted at Roseworthy.
Trial Design	Randomised Block Design of 3 blocks and 20 entries consisting of comparators and potential candidates. Sown in 12 ranges of 5 plots wide, block 1 being in ranges 1 to 4 and so on. Plots were 1.25m wide (5 rows) and 3.2m long. There were approx. 1000 plants per plot.
Measurements	Heading times were recorded on the same trial planted at Roseworthy 2007, but this trial later was abandoned due to a heavy infestation of Crown Rot. All other measurements and observations were recorded on plant samples taken from the Mintaro trial. At anthesis 5 primary tillers were sampled from each plot in each replicate and flag leaf measurements made. Glaucosity and leaf angle was observed at this time. After maturity plant heights to the top of the awns were recorded at 10 random locations in replicate 2 and 3 only. Twenty heads were also sampled at random from each plot in replicates 2 and 3 for head descriptions and measurements. Measurements were performed on 10 intact heads. Statistical analyses were completed using GENSTAT software. Quality data (semolina colour) are from independent tests performed on grain from field trials in SA and NSW over 3 years (6 tests in all). A paired t test was used to determine significance.

RHS Chart - edition

Origin and Breeding

The maternal parent was a breeder's line derived from the complex cross: 'Lingzhi' 'Baimong' 'Baidamai'/2*'Yallaroi'//RH88009///'Wollaroi' (Derived from the same cross as 'Kalka') and the paternal parent 'Tamaroi'. The cross was completed in 1994 with the F_1 grown as a row over summer in 1994/95 and the F_2 grown as a plot over winter of 1995. Single heads were selected from F_2 plants with individual head hills grown over the summer of 1995/96 at the University of Adelaide, Waite Campus. F_4 plots were grown over the winter of 1996 where F_4 derived F_5 heads were selected and grown over summer of 1996/97. The F_5 bulks were trialled for yield, disease resistance and quality in field plots at a number of sites from 1997 to 2002. A promising F_5 bulk designated (WLYY9Tm) 2/3/1. This line entered advanced trials in 2002 where it was designated the name WID22209. From 2003 to 2007 it was tested for yield, disease resistance and quality across the national trial network conducted by Australian Grain Technologies which enabled the evaluation of its performance in the major durum growing areas of Australia. In 2006, WID22209 entered the National Variety Trials. WID22209 has also been evaluated for a range of semolina and pasta quality traits. Breeder: Tony Rathjen, The University of Adelaide and Jason Reinheimer, Australian Grain Technologies.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Ear	distribution of awns	fully awned
Ear	glume colour	white
Plant	time of ear emergence	257 to 263 Julian days
Plant	height	>80 cm

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Kalka'	Related variety.
'Tamaroi'	Grown in the expected area of adoption.
'EGA Bellaroi'	Grown in the expected area of adoption.

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Yallaroi'	Plant height	95.0 cm	83.5 cm	LSD=3.1 (P=1%), significantly shorter.
'Wallaroi'	Plant time of ear emergence	259.2 Julian Days	255.7	LSD=2.7 (P=1%), significantly earlier.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'HYPERNO'	'EGA Bellaroi'	'Kalka'	'Tamaroi'
☐ *Flag leaf: glaucosity of sheath	weak to medium	medium to strong	medium to strong	medium to strong
☐ *Flag leaf: glaucosity of blade	weak to medium	absent or very weak to weak	medium to strong	weak to medium
☐ Awn: anthocyanin colouration	absent or very weak	absent or very weak	absent or very weak	
☐ *Culm: glaucosity of neck	medium	medium	medium to strong	medium to strong
☐ *Ear: glaucosity	medium	medium to strong	medium	medium
☐ Ear: distribution of awns	whole length	whole length	whole length	whole length
☑ *Awns at tip of ear: length in relation to ear	shorter	longer	shorter	equal
☐ Lower glume: shape	elongated	elongated	elongated	elongated
☑ Lower glume: shape of shoulder	straight	elevated with 2nd beak present	straight	elevated with 2nd beak present
☐ Lower glume: shoulder width	narrow	narrow	very narrow	medium
☑ *Lower glume: length of beak	short	medium	short	medium to long
☐ Lower glume: shape of beak	slightly curved	slightly curved	slightly curved	slightly curved
☐ *Lower glume: hairiness on external surface	absent	absent	absent	absent
☐ *Straw: pith in cross section	thin	thin to medium	thin to medium	medium
☑ *Awn: colour	brown	whitish	whitish	black
☑ Ear: hairiness of margin of first rachis segment	strong		strong	absent or very weak
☐ *Ear: colour at maturity	white	white	white	white
☐ Ear: shape in profile view	parallel sided	parallel sided	tapering	parallel sided
☐ *Ear: density	medium to dense	medium	medium	medium
☐ Grain: shape	ovoid to semi-elongated	elongated	ovoid to semi-elongated	elongated
☐ Grain: length of brush hair in dorsal view	very short	short	short	very short
☐ *Grain: colouration with phenol	nil or very light	nil or very light	nil or very light	nil or very light
☐ *Season: type	spring type	spring type	spring type	spring type

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'HYPERNO'	'EGA Bellaroi'	'Kalka'	'Tamaroi'
☑ Roots: boron tolerance	intolerant		tolerant	
☐ Grain: black point tolerance	moderately tolerant			

Statistical Table

Organ/Plant Part: Context	'HYPERNO'	'EGA Bellaroi'	'Kalka'	'Tamaroi'
☑ Flag leaf: length (mm)				
Mean	185.40	179.40	208.00	231.40
Std. Deviation	28.30	21.70	32.20	31.60
LSD/sig	31.5	P≤0.01	ns	P≤0.01
☐ Flag leaf: width (mm)				
Mean	16.20	14.80	17.20	18.70
Std. Deviation	1.40	1.80	1.90	1.90
LSD/sig	1.7	ns	ns	ns
☑ Flag leaf: sheath length				
Mean	175.10	164.70	192.30	188.30
Std. Deviation	12.50	15.10	15.20	13.10
LSD/sig	14.3	ns	P≤0.01	ns
☑ Plant: height (cm)				
Mean	95.00	81.00	93.40	91.40
Std. Deviation	3.70	37.90	3.90	28.80
LSD/sig	3.1	P≤0.01	ns	P≤0.01
☑ Ear: length (mm)				
Mean	86.20	71.90	91.10	86.70
Std. Deviation	5.10	6.80	6.30	3.00
LSD/sig	10.3	P≤0.01	ns	ns
☐ Ear: density (rachis internode)				
Mean	3.41	3.21	3.53	3.37
Std. Deviation	0.17	0.28	0.23	0.23
LSD/sig	0.31	ns	ns	ns
☐ Plant: time of ear emergence (Julian days)				
Mean	259.20		258.00	258.00
Std. Deviation	1.70		0.00	0.00
LSD/sig	2.7		ns	ns
☐ Grain: semolina colour (compared with 'EGA Bellaroi') (Minolta b*)				
Mean	27.50	26.30		
Std. Deviation	4.46	4.35		
LSD/sig	2.19	ns		
Method Used	paired t test			
☑ Grain: semolina colour (compared with 'Kalka') (Minolta b*)				
Mean	28.06		24.50	
Std. Deviation	5.00		5.00	
LSD/sig	3.25		P<=0.01	
Method Used	paired t test			

Prior Applications and Sales

Nil.

Description: **Gil Hollamby,** Williamstown, SA.

Details of Application

Application Number	2007/214
Variety Name	'Ohdrejumwhi'
Genus Species	*Bracteantha bracteata*
Common Name	Everlasting Daisy
Synonym	Jumbo White
Accepted Date	26 Sep 2007
Applicant	Bonza Botanicals Pty Limited, Winmalee, NSW
Agent	Oasis Horticulture Pty Limited, Winmalee, NSW
Qualified Person	Tim Angus

Details of Comparative Trial

Overseas Testing Authority	Canada
Overseas Data Reference Number	05-4574
Location	Overseas data was verified under local conditions in Winmalee, NSW, Australia.
Descriptor	Strawflower (*Bracteantha*) TG/205/1
Period	Dec 2006 to Apr 2007
Conditions	Trial conducted in outside commercial production area, rooted cuttings (propagated from stock plants grown at Winmalee) potted into 150mm standard pots in commercial potting mix, nutrients supplied by slow release and liquid feed fertilizer applications, plant protection treatments applied as necessary. No pinching or other plant shaping treatments were applied.
Trial Design	10 plants of the candidate variety were grown to confirm overseas test report data.
Measurements	Taken at random from 10 plants.
RHS Chart - edition	2001.

Origin and Breeding

Controlled pollination: seed parent proprietary breeding line 02-7 x pollen parent variety 'OHB003970' in a planned breeding program. Seed parent is characterised by Foliage: height medium, Flower head: diameter large. Pollen parent is characterised by Involucre bract: colour yellow. Selection criteria: Plant: habit, Foliage: colour, Flower: habit, Flower: colour. Selection was done at Winmalee, NSW, Australia in 2003. Propagation: by vegetative tip cuttings, no off types occurred in at least three successive vegetative generations during the selection process and in numerous vegetative generations since selection. 'Ohdrejumwhi' will be commercially propagated by vegetative tip cuttings. 'Ohdrejumwhi' was selected from the progeny of this cross in Feb 2003 in a controlled environment in Winmalee, NSW, Australia. Asexual reproduction by terminal stem cuttings taken since Feb 2003 at Winmalee and other locations has shown the characteristics of 'Ohdrejumwhi' are stable and true to type over many generations. Breeder: Dr Andrew Bernuetz, Winmalee, NSW, Australia.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	height of foliage	very short
Plant	density	dense to very dense
Leaf	length	short
Leaf	width	narrow to medium
Leaf	shape of apex	acute
Leaf	main colour of upper side	medium green
Leaf	hairiness of upper side	strong
Flower head	diameter	medium

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'OHB00-37.90'	pollen parent - morphologically very similar, except for the flower colour

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Rredbrawhi'	Leaf	main colour upper side	medium green	yellow green
'Redbrawhi'	Leaf	hairiness of upper side	strong	weak to absent
'Redbrawhi'	Flower bud	colour	white	yellow
'Dargan Hill Monarch'	Plant	height of foliage	very short	medium
'Dargan Hill Monarch'	Flower head	predominant position above foliage	above to far above	slightly above to slightly below

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Ohdrejumwhi'	'OHB00-37.90'
☐ *Plant: type	basal clusters	basal clusters
☐ Plant: growth habit (bushy plant types only)	semi-upright	spreading
☐ Plant: height including flowers	very short	very short to short
☐ Plant: height of foliage	very short	very short
☐ Plant: density	dense to very dense	very dense
☐ Stem: hairiness	medium	medium
☐ Leaf: length	short	short
☐ Leaf: width	narrow to medium	narrow to medium
☐ Leaf: position of broadest part	middle third	upper third
☐ Leaf: shape of apex	acute	acute
☐ *Leaf: variegation	absent	absent
☐ Leaf: main colour of upper side	medium green	medium green

☐ Leaf: hairiness of upper side	strong	strong
☐ Leaf: hairiness of lower side	medium	strong
☐ Leaf: undulation of margin	absent or weak	absent or weak
☐ Flowering shoot: length	very short	very short to short
☐ Flowering shoot: branching	absent or weak	absent or weak
☐ Flower bud: profile of apex	rounded	rounded
☑ Flower bud: main colour (RHS colour chart)	white 155A	yellow 6A
☐ Flower head: diameter	medium	medium
☐ Flower head: side view of lower part	flat	flat
☐ Flower head: side view of upper part	flat	convex
☐ *Involucre: number of colours	only one	only one
☑ *Involucre: main colour	white	yellow
☐ Bract: length	short to medium	short to medium
☐ Bract: width	narrow	narrow
☐ Bract: ratio length/width	three times as long as broad	three times as long as broad
☑ Bract: main colour of lower third of bract from inner third of involucre (RHS colour chart)	white 155A	yellow 6A
☑ Bract: main colour of middle third of bract from inner third of involucre (RHS colour chart)	white 155A	yellow 6A
☑ Bract: main colour of upper third of bract from inner third of involucre (RHS colour chart)	white 155A	yellow 6A
☑ Bract: main colour of lower third of bract from middle third of involucre (RHS colour chart)	white 155A	yellow 6A
☑ Bract: main colour of middle third of bract from middle third of involucre (RHS colour chart)	white 155A	yellow 6A
☑ Bract: main colour of upper third of bract from middle third of involucre (RHS colour chart)	white 155A	yellow 6A
☑ Bract: main colour of lower third of bract from outer third of involucre (RHS colour chart)	white 155A	yellow 6A
☑ Bract: main colour of middle third of bract from outer third of involucre (RHS colour chart)	white 155A	yellow 6A
☑ Bract: main colour of upper third of bract from outer third of involucre (RHS colour chart)	white 155A	yellow 6A

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	**'Ohdrejumwhi'**
☐ Pappus: colour	yellow orange

Statistical Table

Organ/Plant Part: Context	**'Ohdrejumwhi'**
☐ Leaf: length (mm)	

Mean	135.10
Std. Deviation	12.95
☐ Leaf: width (mm)	
Mean	21.20
Std. Deviation	1.69
☐ Flower head: diameter (mm)	
Mean	44.90
Std. Deviation	1.76

Prior Applications and Sales

Country	Year	Current Status	Name Applied
Canada	2005	Applied	‘Ohdrejumwhi’
EU	2005	Applied	‘Ohdrejumwhi’

First sold in USA in Dec 2004. First Australian sale Aug 2006.

Description: **Tim Angus**, Wellington, NZ

Details of Application

Application Number	2007/161
Variety Name	'Doza'
Genus Species	*Vicia faba*
Common Name	Field Bean
Synonym	Nil
Accepted Date	9 Jul 2007
Applicant	Department of Primary Industries for and on behalf of the State of New South Wales, Orange, NSW and Grains Research and Development Corporation, Barton, ACT
Agent	N/A
Qualified Person	Ross Downes

Details of Comparative Trial

Location	DPI Temora Research Station.
Descriptor	Broad Bean (*Vicia faba*) TG/206/1.
Period	Winter-spring 2007.
Conditions	Seed was sown in soil in plots 10m x 1m. Plants grew poorly before flowering because of drought conditions. Plots were watered after flowering. Two generations of 'Doza' were grown with comparators 'Cairo', 'Farah' and 'Fiesta'.
Trial Design	Randomised block, 4 replications, data processed from 3 replications only.
Measurements	Data were collected on 6 Sep, 9 Oct and 13 Nov 2007.
RHS Chart - edition	

Origin and Breeding

Single Plant Selection: 'Doza', synonym SP1040, originated as a single plant progeny selected in 2001, at the Australian Cotton Research Institute Narrabri, from an outcrossed population of 'SP98066'. This originated as a single plant progeny selected in 1998 from an outcrossed population of 'SP9558'. 'SP9558' was in turn selected from an outcrossed population based on the cross of 'Accession 383'/'Sudan Triple White' in 1995. The seedlot used for selection in 1995 was harvested from regional variety trial plots in 1994. These plots had been subject to outcrossing by bees. Selection during self pollination was for seed appearance and seed size. Populations were rogued to remove plants with white flowers or susceptible rust reactions. Open pollinated seed production in isolation commenced in 2006 in parallel with further self pollinated seed production. Breeder: Dr Ian Rose, NSW Department of Primary Industries, Narrabri, NSW.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth type	determinate
Wing	melanin spot	present
Dry seed	colour of testa	beige

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Cairo'	
'Farah'	
'Fiesta'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Doza'	'Cairo'	'Farah'	'Fiesta'
☐ *Plant: growth type	determinate	determinate	determinate	determinate
☑ *Plant: height	short	medium to tall	medium to tall	short
☐ *Plant: number of stems	few	very few to few	very few to few	very few
☐ Stem: number of nodes up to and including first flowering node	medium	medium	medium	medium
☐ Stem: anthocyanin colouration	absent	present	absent	present
☐ Foliage: greyish hue of green colour	absent	absent	absent	absent
☐ Foliage: intensity of green colour	medium	medium	medium	medium
☐ *Leaflet: length	medium	medium to long	medium to long	medium
☐ *Leaflet: width	medium	medium to broad	medium to broad	medium
☐ *Leaflet: position of maximum width	at middle	at middle	at middle	at middle
☐ Leaflet: folding	medium	medium	medium	medium
☐ *Raceme: number of flowers	medium	medium	medium	medium
☑ *Time of: flowering	very early to early	early to medium	early to medium	early
☐ Flower: length	medium	medium to long	medium to long	medium to long
☐ *Wing: melanin spot	present	present	present	present
☐ *Wing: colour of melanin spot	brown	brown	brown	brown
☐ Standard: melanin spot	absent	absent	absent	absent
☐ *Standard: anthocyanin colouration	present	present	present	present
☐ Standard: extent of anthocyanin colouration	small	small	small	small
☐ Truss: number of pods	few	few	very few	medium
☐ *Pod: attitude	semi-erect	semi-erect	semi-erect	semi-erect
☐ *Pod: length	medium	medium to long	medium to long	medium
☐ *Pod: width	narrow to medium	medium	medium	narrow to medium
☐ Pod: degree of curvature at green shell stage	weak	weak	weak	weak

☐ Pod: intensity of green colour	medium	medium	medium	medium
☐ Pod: number of ovules	few	few	few	few
☐ Pod: thickness of pod wall	medium	medium	medium	medium
☐ Dry seed: shape of median longitudinal section	elliptic	elliptic	elliptic	elliptic
☐ Dry seed: shape of cross section	elliptic	elliptic	elliptic	elliptic
☑ *Dry seed: weight	low	medium	high	high
☐ *Dry seed: colour of testa	beige	beige	beige	beige
☐ Dry seed: black pigmentation of hilum	present	present	present	present
☑ Time of: full development of pod	early	medium	medium	medium

Statistical Table

Organ/Plant Part: Context	**'Doza'**	**'Cairo'**	**'Farah'**	**'Fiesta'**
☑ Dry seed: weight (g)/100 seeds				
Mean	52.0	61.6	71.2	71.4
Std Deviation	0.8	3.6	1.7	6.0
LSD/sig 7.9 (.01)	7.9	P≤0.01	P≤0.01	P≤0.01

Prior Applications and Sales

Nil.

Description: **Ross Downes,** Moruya, NSW.

.

Details of Application

Application Number	2006/026
Variety Name	'Bundi'
Genus Species	*Pisum sativum*
Common Name	Field Pea
Synonym	Nil
Accepted Date	24 Mar 2006
Applicant	Agriculture Victoria Services Pty Ltd, Atwood, VIC and Grains Research and Development Corporation, Barton, ACT
Agent	N/A
Qualified Person	Antonio Leonforte

Details of Comparative Trial

Location	Horsham, VIC
Descriptor	Pea (*Pisum sativum*) TG/7/9
Period	2007
Conditions	Winter sown; lower than average rainfall year.
Trial Design	Randomised complete block design.
Measurements	Plant height at flowering (mm) and time to flowering (days).
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: 'Bundi' is derived from a cross made in 1989 (89-036) between breeding lines PS772 and PS770. 'Bundi' was developed following a pedigree selection program. Single plant selections were taken from the F_2 population (89-036-9) and again from an F_2 derived F_6 population (89-036*9-8). 'Bundi' was primarily selected on the basis of high yield potential, particularly for the medium to lower rainfall regions. 'Bundi' was selected as an early flowering and maturing line with excellent pod shatter resistance at maturity and a semi-erect plant habit. 'Bundi' was also selected with high resistance to downy mildew. 'Bundi' produces medium to large spherical white seed with a yellow cotyledon which is suitable for both human consumption and stockfeed markets. Breeder: Tony Lenoforte, Department of Primary Industry, VIC.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	leaflets	absent
Seed	anthocyanin	absent
Seed	cotyledon	yellow
Pod	parchment layer	partially absent

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Moonlight'	Semi-dwarf, semi-leafless, white seeded, reduced pod parchment, mid season flowering time.

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Excell'	Seed	cotyledon colour	yellow	green
'Snowpeak'	Pod	parchment layer	present	partially absent
'Sturt'	Leaflets	absence	absent	present
'Kaspa'	Plant	anthocyanin	absent	present

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Bundi'	'Moonlight'
☐ Seed: shape	spherical	spherical
☐ *Seed: shape of starch grain	simple	simple
☐ *Seed: colour of cotyledon	yellow	yellow
☐ *Seed: marbling of testa (varieties with anthocyanin only)	absent	absent
☐ *Seed: violet or pink spots on testa (varieties with anthocyanin only)	absent	absent
☐ *Seed: black colour of hilum	absent	absent
☐ Seed: dimpled cotyledons (varieties with unwrinkled seed and simple starch grains only)	absent	absent
☐ *Plant: anthocyanin colouration	absent	absent
☑ Plant: height	short to medium	tall
☐ Stem: fasciation	absent	absent
☑ *Stem: length	short to medium	long
☑ Stem: number of nodes up to and including first fertile node	few to medium	many
☐ Stem: anthocyanin colouration of axil (varieties with anthocyanin only)	absent	absent
☐ *Foliage: colour	green	green
☐ Foliage: intensity of colour (excluding yellow-green and blue-green varieties)	medium	medium
☐ *Leaf: leaflets	absent	absent
☐ Leaf: waxiness of surface of upper leaflet	present	present
☐ *Stipule: type of development	well developed	well developed
☐ Stipule: 'rabbit-eared stipules'	absent	absent
☐ Stipule: waxiness of surface of upper stipule	present	present
☐ Stipule: length	medium	medium
☐ Stipule: width	medium	medium
☐ *Stipule: flecking	present	present
☐ Stipule: maximum density of flecking	sparse	sparse
☐ Petiole: length (varieties without leaflets only)	medium	medium
☐ *Time of: flowering	early	medium

*Plant: maximum number of flowers per node (non-fasciated varieties only)	two	two
Flower: colour of standard (varieties without anthocyanin only)	white	white
Flower: maximum width of standard	medium	medium
Flower: shape of base of standard	level	level
Flower: intensity of undulation of standard	medium	medium
Flower: width of sepal	medium	medium
Flower: shape of apex of upper sepal	accuminate	accuminate
Flower: length of peduncle from stem to first flower	medium	medium
*Pod: length	medium to long	medium
*Pod: maximum width	medium	medium
Pod: parchment	partially absent	partially absent
Pod: thickened wall (varieties with no or partial parchment only)	absent	absent
*Pod: degree of curvature	weak	weak
*Pod: type of curvature	concave	concave
*Pod: shape of distal part (varieties without thickened pod wall only)	blunt	blunt
*Pod: colour	green	green
Pod: intensity of green colour	medium	medium
Pod: strings of suture (varieties with no or partial parchment only)	absent or rudimentary	absent or rudimentary
Pod: anthocyanin colouration of suture (varieties with anthocyanin only)	absent	absent
Pod: spots of anthocyanin colouratin on outer wall (varieties with anthocyanin only)	absent	absent
*Pod: number of ovules	medium to many	few to medium
Pod: intensity of green colour of immature seed	medium	medium
Seed: time of maturity	early	medium
Seed: wrinkling of cotyledon	absent	absent
*Seed: weight	medium to large	large to very large
Resistance to: *Fusarium oxysporum* f. sp. pisi race 1	present	
Resistance to: *Erysiphe pisi* Syd.	absent	absent
Resistance to: seed-borne mosaic virus (SbmV), strain P1	absent	absent

Statistical Table

Organ/Plant Part: Context	**'Bundi'**	**'Moonlight'**
Plant height: height at flowering (mm)		
Mean	467.00	548.00
Std. Deviation	36.00	66.00
LSD/sig	22.0	P≤0.01

☑ Flower: flowering time (days)		
Mean	120.00	127.00
Std. Deviation	1.50	1.50
LSD/sig	2.0	P≤0.01

Prior Applications and Sales

Nil.

Description: **Tony Lenoforte**, Victorian Institute for Dryland Agriculture, Horsham, VIC.

Details of Application

Application Number	2007/097
Variety Name	'TAS300'
Genus Species	*Dianella tasmanica*
Common Name	Flax lily
Synonym	Nil
Accepted Date	26 Apr 2007
Applicant	Wyeena Nurseries Pty Ltd, Smiths Gully, VIC
Agent	Ozbreed Pty Ltd, Clarendon, NSW
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Clarendon, NSW.
Descriptor	Dianella (*Dianella*) PBR DIAN
Period	Autumn 2007 - spring 2007.
Conditions	Trial conducted in open beds, plants propagated from cuttings, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random.
RHS Chart - edition	2001.

Origin and Breeding

Seedling selection: seed parent *D. tasmanica*. The seed parent is characterised by an absence of leaf blade variegation . Selection took place in Wyeena Nurseries Pty Ltd, Smiths Gully, VIC in 1998. Selection criteria: presence of variegation, stable reproduction. Propagation: vegetative, micropropagation is found to be uniform and stable. Breeder: Kahn Franke, Wyeena Nurseries Pty Ltd, Smiths Gully, VIC.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf blade	presence of variegation	present
Leaf blade	glaucosity of upper side	medium to strong

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'TAS100'	
D. tasmanica variegated common form	

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Splice'	Leaf blade	glaucosity of upper side	medium-strong	absent or very weak
'Rainbow'	Leaf blade	glaucosity of upper side	medium-strong	weak-medium
'Rainbow'	Leaf blade	width	medium-broad	narrow-medium

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'TAS300'	*D. tasmanica* variegated common form	'TAS100'
☐ Plant: growth habit	semi-erect	semi-erect	semi-erect
☑ Plant: height	medium to tall	medium	short to medium
☑ Plant: density of shoots	sparse to medium	sparse	medium
☐ Stem: length of internodes	very short	very short	very short
☐ Leaf: attitude	semi-erect	semi-erect	semi-erect
☑ Leaf: arching	weak	medium	medium
☐ Leaf: width	medium	medium	medium
☐ Leaf: glaucosity of upper side	medium to strong	medium to strong	medium to strong
☐ Leaf: colour of upper side (waxiness removed) (RHS colour chart)	147A	147A	147A
☑ Leaf: colour of lower side (waxiness removed) (RHS colour chart)	191A	191A	147B
☐ Leaf: variegation	present	present	present
☑ Leaf: secondary colour of upper side (variegated leaves only) (RHS colour chart)	12D	12D	11D
☐ Leaf: shape of blade	linear	linear	linear
☐ Leaf: shape of apex	acute	acute	acute
☐ Leaf: cross-section	concave	concave	concave
☐ Leaf: spines on margin	present	present	present
☐ Leaf: prominence of spines on margin	weak to medium	weak to medium	weak to medium
☐ Leaf: spines on lower side of midrib	present	present	present
☐ Leaf: prominence of spines on lower side of midrib	weak to medium	weak to medium	weak to medium
☐ Basal leaf sheath: anthocyanin colouration (in summer)	red-purple	red-purple	red-purple
☐ Basal leaf sheath: intensity of anthocyanin colouration	strong	strong	strong

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'TAS300'	*D. tasmanica* variegated common form	'TAS100'
☑ Leaf blade: degree of variegation	small	small	medium

Statistical Table

Organ/Plant Part: Context	'TAS300'	*D. tasmanica* variegated common form	'TAS100'
☑ Plant: height (mm)			
Mean	357.50	267.00	246.50
Std. Deviation	52.90	51.90	54.60
LSD/sig	60.62	P≤0.01	P≤0.01
☑ Leaf: width (mm)			
Mean	24.76	26.12	28.52
Std. Deviation	2.00	3.30	2.50
LSd/sig	3.04	ns	P≤0.01

Prior Applications and Sales

Nil.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW.

Details of Application

Application Number	2007/021
Variety Name	'TAS100'
Genus Species	*Dianella tasmanica*
Common Name	Flax lily
Synonym	Nil
Accepted Date	5 Feb 2007
Applicant	Ozbreed Pty Ltd, Clarendon, NSW
Agent	N/A
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Clarendon, NSW.
Descriptor	Dianella (*Dianella*) PBR DIAN
Period	Autumn 2007 - spring 2007.
Conditions	Trial conducted in open beds, plants propagated from cuttings, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random.
RHS Chart - edition	2001.

Origin and Breeding

Seedling selection: seed parent *D. tasmanica* variegated form. The seed parent is characterised by a variegated leaf blade with medium prominence and an unstable expression of this trait during propagation. Selection took place in Clarendon, NSW in 2005. Selection criteria: stable reproduction, enhanced variegation prominence and compact plant growth habit. Propagation: vegetative, micropropagation is found to be uniform and stable. Breeder: Todd Layt, Clarendon, NSW.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	semi-erect
Leaf blade	presence of variegation	present
Leaf blade	glaucosity of upper side	medium to strong

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'TAS300'	
D. tasmanica variegated common form	parent of candidate

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Splice'	Leaf blade	glaucosity of upper side	medium-strong	absent or very weak
'Splice'	Leaf	attitude	semi-erect	erect to semi-erect
'Splice'	Leaf blade	secondary colour of upper side (RHS)	11D	N144A
'Rainbow'	Leaf blade	glaucosity of upper side	medium-strong	weak-medium
'Rainbow'	Leaf	attitude	semi-erect	erect to semi-erect
'Rainbow'	Leaf blade	width	medium-broad	narrow-medium
D. tasmanica common from from South Australia	Leaf	attitude	semi-erect	erect to semi-erect
D. tasmanica common from from South Australia	Leaf blade	width	medium-broad	narrow-medium
D. tasmanica common from from South Australia	Leaf blade	degree of variegation	weak	strong
D. tasmanica common from from South Australia	Basal leaf sheath	intensity of anthocyanin coloration	strong	medium

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'TAS100'	*D. tasmanica* variegated common form	'TAS300'
☐ Plant: growth habit	semi-erect	semi-erect	semi-erect
☑ Plant: height	short to medium	medium	medium to tall
☑ Plant: density of shoots	medium	sparse	sparse to medium
☐ Stem: length of internodes	very short	very short	very short
☐ Leaf: attitude	semi-erect	semi-erect	semi-erect
☑ Leaf: arching	medium	medium	weak
☐ Leaf: width	medium	medium	medium
☐ Leaf: glaucosity of upper side	medium to strong	medium to strong	medium to strong
☐ Leaf: colour of upper side (waxiness removed) (RHS colour chart)	147A	147A	147A
☑ Leaf: colour of lower side (waxiness removed) (RHS colour chart)	147B	191A	191A
☐ Leaf: variegation	present	present	present
☑ Leaf: secondary colour of	11D	12D	12D

upper side (variegated leaves only) (RHS colour chart)			
☐ Leaf: shape of blade	linear	linear	linear
☐ Leaf: shape of apex	acute	acute	acute
☐ Leaf: cross-section	concave	concave	concave
☐ Leaf: spines on margin	present	present	present
☐ Leaf: prominence of spines on margin	weak to medium	weak to medium	weak to medium
☐ Leaf: spines on lower side of midrib	present	present	present
☐ Leaf: prominence of spines on lower side of midrib	weak to medium	weak to medium	weak to medium
☐ Basal leaf sheath: anthocyanin colouration (in summer)	red-purple	red-purple	red-purple
☐ Basal leaf sheath: intensity of anthocyanin colouration	strong	strong	strong

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	**'TAS100'**	***D. tasmanica* variegated common form**	**'TAS300'**
☑ Leaf blade: degree of variegation	medium	small	small

Statistical Table

Organ/Plant Part: Context	**'TAS100'**	***D. tasmanica* variegated common form**	**'TAS300'**
☑ Plant: height (mm)			
Mean	246.50	267.00	357.50
Std. Deviation	54.60	51.90	52.90
LSD/sig	60.62	ns	P≤0.01
☑ Leaf blade: width (mm)			
Mean	28.52	26.12	24.76
Std. Deviation	2.50	3.30	2.00
LSD/sig	3.04	ns	P≤0.01

Prior Applications and Sales

Prior applications nil. First sold in Australia in Apr 2006.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW.

Details of Application

Application Number	2006/218
Variety Name	'Blood Orange'
Genus Species	*Grevillea* hybrid
Common Name	Grevillea
Synonym	Nil
Accepted Date	5 Oct 2006
Applicant	Christopher John Hughes, Federal, NSW
Agent	N/A
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Federal, NSW.
Descriptor	Grevillea (*Grevillea*) PBR GREV
Period	Summer 2007 - winter 2007.
Conditions	Trial conducted with mature plants in ground, plants originally propagated by cuttings, potted to 200mm containers filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Ten plants of each variety arranged in a completely randomised design.
Measurements	From ten plants.
RHS Chart - edition	1995.

Origin and Breeding

Open pollination: 'Honey Gem'. The parent is characterised by a predominantly orange inflorescence colour. Selection took place in Federal, NSW. Selection criteria: colour of inflorescence. Propagation: vegetative cuttings were found to be uniform and stable. Breeders: Christopher Hughes, Federal, NSW.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	division of blade	all leaves on plant divided
Leaf	degree of division	first order
Leaf	depth of division of blade	greater than two thirds of way to midrib

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Honey Gem'	parent variety with same foliage type
'Bird Song'	similar foliage type

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Blood Orange'	'Bird Song'	'Honey Gem'
☑ Plant: growth habit	upright	bushy	upright
☐ Plant: attitude of branches	erect to semi-erect	erect to semi-erect	erect to semi-erect
☑ Plant: height	medium (1-3m)	medium (1-3m)	tall (> 3m)
☐ Plant: density (assessment of foliage at flowering)	medium	medium to dense	medium
☑ Young stem: colour	greyed orange	greyed orange	brown
☑ Stem: colour	brown	greyed orange	brown
☐ Stem: hairiness	medium to strong	medium to strong	strong
☐ Petiole: length	short to medium	short to medium	medium
☐ Leaf: length	very long (> 20cm)	long (15-20cm)	very long (> 20cm)
☐ Leaf: width at widest point	very broad (> 20cm)	very broad (> 20cm)	very broad (> 20cm)
☐ Leaf: attitude to stem	semi-erect	semi-erect to horizontal	semi-erect
☐ Leaf: curvature of margin	flat or slightly recurved, under surface on either side of the mid-vein wholly exposed	flat or slightly recurved, under surface on either side of the mid-vein wholly exposed	flat or slightly recurved, under surface on either side of the mid-vein wholly exposed
☐ Leaf: colour of upper side (including hairs)	dark green	dark green	dark green
☐ Leaf: degree of hairiness on upper side	very weak	very weak	very weak
☐ Leaf: degree of hairiness on lower side	medium	medium	weak to medium
☐ Leaf: colour of hairiness on lower side	white	white	white
☐ Leaf: undulation of margin	weak	weak	weak
☐ Leaf: division of blade	some or all leaves on plant divided	some or all leaves on plant divided	some or all leaves on plant divided
☐ Leaf: degree of division of blade (varieties with division of blade present only)	first order	first order	first order
☐ Leaf: depth of division of blade (varieties with division of blade present only)	sinus greater than two thirds of way to midrib	sinus greater than two thirds of way to midrib	sinus greater than two thirds of way to midrib
☐ Leaf: number of lobes (varieties with division of blade present only)	many (> 20)	many (> 20)	many (> 20)
☐ Leaf: regularity of lobing (varieties with division of blade present only)	regular	regular	regular
☐ Leaf: attitude of longitudinal axis of	semi-erect	semi-erect	semi-erect

lobes to longitudinal axis of midrib (varieties with division of blade present only)			
☐ Leaf: attitude of longitudinal axis of lobes to one another on same side of leaf (varieties with division of blade present only)	parallel	parallel	parallel
☐ Leaf: shape of apex of sinus (varieties with division of blade present only)	pointed	pointed	pointed
☐ Lobe: width (varieties with division of blade present only)	narrow	narrow	narrow
☐ Lobe: shape of apex of ultimate lobe (varieties with division of blade present only)	pointed	pointed	pointed
☐ Flowering branch: position of inflorescence	both terminal and axillary	both terminal and axillary	both terminal and axillary
☐ Inflorescence: length	medium to long	medium	medium to long
☐ Inflorescence: width	medium	medium	medium
☑ Inflorescence: predominant colour	pink	orange	orange
☐ Inflorescence: density of florets	dense	dense	dense
☑ Inflorescence: number of flowers	many to very many	many	many to very many
☑ Inflorescence: attitude	semi-erect to horizontal	semi-erect	semi-erect
☐ Inflorescence: form	cylindrical	cylindrical	cylindrical
☐ Inflorescence: branching	absent or very weak	absent or very weak	absent or very weak
☐ Inflorescence: sequence of opening of the flowers	centripetal	centripetal	centripetal
☐ Rachis: length	medium to long	medium	medium to long
☐ Bud: colour of perianth	green	green	green
☑ Bud: colour of limb	green	orange	yellow
☐ Bud: attitude of limb in relation to longitudinal axis of bud (late bud prior to anthesis)	drooping	drooping	drooping
☐ Flower: attitude of pedicel in relation to rachis	leaning away from inflorescence peduncle	leaning away from inflorescence peduncle	leaning away from inflorescence peduncle
☑ Flower: length of pedicel	short	very short to short	short to medium
☑ Perianth: colour	pink	orange	orange
☐ Perianth: degree of hairiness (outside of perianth including limb)	medium	weak to medium	medium
☐ Perianth: colour of hairs	white	white	white
☐ Perianth: length	short to medium	medium	medium

Perianth: width	medium	medium	medium
Perianth: ratio length/width	medium	medium	medium
Perianth: coherence of tepals on dorsal side	less than one third	less than one third	less than one third
Perianth: coherence of tepals on ventral side	less than one third	less than one third	less than one third
Tepal: flanging at margin	weak	weak	weak
Nectary: colour	yellow	white	yellow
Ovary: colour	white	white	white
Ovary: hairiness	strong	strong	strong
Style: colour	red	orange	orange
Style: curvature (after anthesis before dehiscence of perianth)	gently curved	gently curved	gently curved
Style: position of curve	top half	top half	top half
Style: hairiness	absent or very weak	absent or very weak	absent or very weak
Pistil: length	long	long	long
Pistil: length in relation to length of perianth	much longer	much longer	much longer
Stigma: colour	yellow	yellow	yellow
Pollen presenter: attitude to style	lateral	lateral	lateral
Pollen presenter: colour	yellow	yellow	yellow
Pollen presenter: shape	cone	cone	cone
Pollen: colour	yellow	yellow	yellow

Prior Applications and Sales

No prior applications. First sold in Australia in Jun 2006 under the name Grevillea ‘Blood Orange’.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW

Details of Application

Application Number	2003/375
Variety Name	'Black Kat'
Genus Species	*Prunus* hybrid
Common Name	Interspecific Plum
Synonym	Nil
Accepted Date	5 May 2004
Applicant	Zaiger's Inc. Genetics, Modesto, CA, USA
Agent	Fleming's Nurseries & Associates Pty Ltd, Monbulk, VIC
Qualified Person	Lisa Corcoran

Details of Comparative Trial

Overseas Testing Authority	US Patent and Trademark Office
Overseas Data Reference Number	U.S PP 13,134
Descriptor	Japanese Plum (*Prunus salicina*) TG/84/3
Conditions	Where possible the US plant patent data was verified under local conditions in Yellingbo, Victoria. The US Plant Patent data was converted into the standard UPOV descriptors.

Origin and Breeding

Cross pollination: the present new and distinct interspecific plum was developed by Zaiger Inc Genetics at their experimental orchard at Modesto, California, as a first generation cross between two seedlings with field identification numbers 73ED135 and 72GC211. A large number of these first generation crosses were planted and maintained on their own roots. In 1994 the present variety was observed to have desirable fruiting characteristics and was selected for asexual propagation and commercialisation. Breeder: Zaiger Inc Genetics, Modesto, California USA.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Fruit	flesh colour	yellow
Fruit	skin colour	blue to black

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Flavorich'	Matures approximately 10-12 days later than 'Black Kat'. The fruit of 'Black Kat' is more round than that of 'Flavorich'.

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Angelino'	Fruit	skin colour	blue to black	red

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Black Kat'	'Flavorich'
☐ Tree: vigour	strong	strong

Tree: density of the head	medium	medium
One year old shoot: intensity of colour	light	
*Leaf blade: shape	broad obovate	broad obovate
*Leaf blade: angle of the tip	pointed	pointed
Leaf blade: green colour of upper side	medium to dark	medium to dark
Leaf blade: incisions of margin	serrate	serrate
*Petiole: length	medium	medium
Leaf: position of glands	on both leaf base and petiole	on both leaf base and petiole
Flowers: size	small	medium
Petal: size	medium	medium
*Petal: shape	obovate	
Petal: undulation of margin	weak	
Stigma: position as compared with anthers	same level to above	
*Fruit: size	large	large
*Fruit: general shape	rounded-flattened	rounded-flattened
*Fruit: ground colour of skin	dark blue	violet blue
*Fruit: colour of flesh	yellow	yellow
Fruit: firmness of flesh	firm	firm
*Fruit: degree of adherence of stone to flesh	semi-adherent	fully adherent
*Stone: size	small	medium
*Stone: general shape in profile	round-elliptical	round-elliptical
*Time of: flowering	medium	medium
*Time of: ripening	late to very late	very late[1]

[1]matures approximately 10-12 days later than 'Black Kat'

Prior Applications and Sales

Country	Year	Current Status	Name Applied
Chile	2005	Granted	'Black Kat'
USA	2001	Granted	'Black Kat'

First sold in the USA in Oct 2002.

Description: **Lisa Corcoran, Fleming's Nurseries & Associates Pty Ltd,** Monbulk, VIC.

Details of Application

Application Number	2006/012
Variety Name	'Regal Velvet'
Genus Species	*Anigozanthos* hybrid
Common Name	Kangaroo Paw
Synonym	Nil
Accepted Date	22 Feb 2006
Applicant	George A Lullfitz, Wanneroo, WA
Agent	N/A
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Macmasters Beach, NSW.
Descriptor	Kangaroo Paw (*Anigozanthos*) TG/175/3.
Period	Autumn 2007 to spring 2007.
Conditions	Trial conducted in open beds, plants propagated from cuttings, planted into 140mm pots filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random.
RHS Chart - edition	1995.

Origin and Breeding

Controlled pollination: seed parent *Anigozanthos manglesii* x pollen parent *A. flavidus*. The seed parent is characterised by a short-medium plant height, large flower size and flower colour consisting of green perianth tube with red ovary. The pollen parent is characterised by a tall plant height, medium flower size and flower colour predominantly greenish to red. Selection took place in Waneroo, WA in 2003. Selection criteria: strong plant growth habit, attractive flower colour. Propagation: vegetative by micropropagation is found to be uniform and stable. Breeder: Keith Oliver, Hammersley, WA.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Perianth tube	predominant colour	green
Ovary	colour of hairs	red
Flower	predominant colour	green and red

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Bush Games'	

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Big Red'	Plant	height	medium	tall
'Big Red'	Perianth tube	predominant colour	green	greyed-purple
'Big Red'	Flower	predominant colour	green and red	red

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Regal Velvet'	'Bush Games'
☑ *Plant: height	medium	short
☑ Plant: number of inflorescences	medium	few
☐ Leaf: length	medium	short to medium
☐ Leaf: width	medium	medium
☐ *Leaf: attitude	semi-erect	semi-erect
☑ Leaf: degree of curvature	straight	slightly curved
☐ Leaf: colour	green	green
☐ Leaf: glaucosity	very weak	very weak
☐ Leaf: degree of hairiness of margin	weakly expressed	absent or very weakly expressed
☑ *Inflorescence: ramification	present	absent
☑ Inflorescence: degree of ramification	tertiary	absent
☑ Inflorescence: length of lowest lateral	medium	
☑ Inflorescence: number of flowers	medium to many	few
☐ Pedicel: colour of hairs (RHS colour chart)	53A	53A
☑ Perianth tube: length	short	long
☑ Perianth tube: width	narrow	medium to broad
☑ Perianth tube: profile	flared distally	broadening evenly
☐ *Perianth tube: predominant colour	green	green
☐ Perianth tube: number of colours of hair	one	one
☐ Perianth tube: colour of tip of hairs (RHS colour chart)	187A	187A
☐ Perianth tube: colour of middle third of hairs (RHS colour chart)	187A	187A
☐ Perianth lobe: length of longest	medium	medium
☐ *Perianth lobes: reflexing	strong	strong
☑ Flower: number of anthers at top of perianth	six	four
☐ Ovary: colour of hairs (RHS colour chart)	53A	53A
☐ Flower: position of stigma in relation to anthers	above	above
☑ Time of: beginning of flowering	medium	early

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Regal Velvet'	'Bush Games'
☐ Flower: number of colour	two	two
☐ Flower: predominant colour	green and red	green and red

Prior Applications and Sales

Prior applications nil. First sold in Australia in Jun 2005 as 'Regal Velvet'

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW.

Details of Application

Application Number	2005/288
Variety Name	'DOW10'
Genus Species	*Tristaniopsis laurina*
Common Name	Kanooka
Synonym	Nil
Accepted Date	24 Oct 2005
Applicant	Downes Wholesale Nursery Pty Ltd, Rossmore, NSW
Agent	Ozbreed Pty Ltd, Clarendon, NSW
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Theresa Park, NSW.
Descriptor	Lilly Pilly (*Acmena smithii*/*Syzygium* sp) PBR LILL.
Period	Summer – Autumn 2007.
Conditions	Trial conducted in open beds, plants propagated from cuttings, planted into 45L bags filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random.
RHS Chart - edition	1995.

Origin and Breeding

Seedling selection: seed parent *Tristaniopsis laurina*. The seed parent is characterised by a narrow leaf width, oblanceolate to lanceolate leaf shape and a dark brown bark colour on mature stems prior to shedding. In 2002 approximately 4000 seedlings arising from open-pollinated seed of *T. laurina* were grown in an open bed. In 2003, 4 seedlings were selected due to their larger leaf size, dark, shiny leaf form and attractive leaf shape. In 2004, a final single selection was made from these due to its extreme differences to the parent form and also due to its faster rate of growth. Selection took place in Tuckombil, NSW in 2003. Selection criteria: broad leaf width, light brown bark colour on mature stems prior to shedding and vigorous plant growth rate. Propagation: vegetative cuttings were found to be uniform and stable. Breeders: Greg Hellyar and Stuart Nolan , Tuckombil, NSW.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	upright
Plant	height	tall
Leaf blade	presence of variegation	absent

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
T. laurina	parent of candidate

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Hot Tips'	Leaf blade	presence of variegation	absent	present
'Golden Summers'	Leaf blade	presence of variegation	absent	present

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'DOW10'	*T. laurina*
☐ Plant: growth habit	upright	upright
☐ Plant: branch density	medium	medium
☐ Stem: branch angle	acute	acute
☐ Stem: internode length	medium	medium
☐ Stem: basal diameter	medium	medium
☐ Stem: colour of mature stem (RHS colour chart)	199D	199D
☐ Stem: colour of new growth (RHS colour chart)	146C	146D
☐ Leaf: blade length	medium	medium
☐ Leaf: blade width	very broad	medium
☐ Leaf: petiole length	long	medium
☑ Leaf: shape of blade	elliptic	oblanceolate
☐ Leaf: shape of apex	acute	acute
☑ Leaf: shape of base	attenuate	cuneate
☐ Leaf: glossiness	medium	medium
☐ Leaf: stiffness	medium	medium
☐ Mature leaf: primary colour of upper side (RHS colour chart)	147A	147A
☑ Mature leaf: primary colour of lower side (RHS colour chart)	146B	147C
☐ Partly mature leaf: primary colour of upper side (RHS colour chart)	146A	146A
☐ Partly mature leaf: primary colour of lower side (RHS colour chart)	146D	146D
☑ Newly emerged: upper side (RHS colour chart)	ca 176A	ca 177A
☐ Leaf: variegation	absent	absent

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'DOW10'	*T. laurina*
☑ Semi-mature stem: colour (RHS)	166A	200B

Statistical Table

Organ/Plant Part: Context	**'DOW10'**	***T. laurina***
Leaf: length (mm)		
Mean	123.90	125.20
Std. Deviation	9.40	7.90
LSD/sig	9.92	ns
Leaf: width (mm)		
Mean	52.50	27.20
Std. Deviation	7.00	2.90
LSD/sig	6.08	P≤0.01

Prior Applications and Sales

Nil.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW

Details of Application

Application Number	2006/127
Variety Name	'AATS'
Genus Species	*Syzygium australe*
Common Name	Lilly Pilly
Synonym	Nil
Accepted Date	31 Aug 2006
Applicant	John Crump, Newrybar, NSW
Agent	Ozbreed Pty Ltd, Clarendon, NSW
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Clarendon, NSW.
Descriptor	Lilly Pilly (*Acmena smithii*/*Syzygium* sp) PBR LILL.
Period	Spring 2006 – autumn 2007.
Conditions	Trial conducted in open beds, plants propagated from cuttings, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random.
RHS Chart - edition	1995.

Origin and Breeding

Seedling selection: seed parent *Syzygium australe*. The seed parent is characterised by a bushy and spreading plant growth habit. Approximately 200 seeds were collected and sown in 2003. From the resulting progeny a single selection was made which had a distinctly columnar growth habit compared to the parent tree. Selection took place in Newrybar, NSW in 2004. Selection criteria: columnar plant growth habit. Propagation: vegetative cuttings were found to be uniform and stable. Breeder: John Crump, Newrybar, NSW.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	branch density	medium
Stem	branch angle	acute
Leaf	length	short

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Bronzed Aussie'	similar foliage

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'AATS'	'Bronzed Aussie'
☑ Plant: growth habit	strongly upright	bushy to upright
☐ Plant: height	medium to tall	medium to tall
☐ Plant: branch density	medium	medium
☐ Stem: branch angle	acute	acute
☐ Stem: internode length	medium	medium
☐ Stem: colour of mature stem (RHS colour chart)	199B	199B
☑ Stem: colour of new growth (RHS colour chart)	183A-B	182B and 153A
☐ Leaf: blade length	short	short
☐ Leaf: blade width	narrow	narrow
☑ Leaf: petiole length	medium	long
☐ Leaf: shape of blade	elliptic	elliptic
☑ Leaf: shape of apex	acute	acuminate
☐ Leaf: shape of base	cuneate	cuneate
☐ Leaf: glossiness	medium	medium
☐ Leaf: shape of cross section	flat to concave	concave
☐ Leaf: shape of longitudinal section	convex	convex to flat
☑ Leaf: stiffness	medium	strong
☐ Leaf: prominence of midrib on lower surface	prominent	prominent
☐ Mature leaf: primary colour of upper side (RHS colour chart)	147A	147A
☑ Mature leaf: primary colour of lower side (RHS colour chart)	146B	147B
☐ Partly mature leaf: primary colour of upper side (RHS colour chart)	146A	146A
☐ Partly mature leaf: primary colour of lower side (RHS colour chart)	146C	146C
☑ Newly emerged: upper side (RHS colour chart)	175A	153A
☐ Leaf: variegation	absent	absent
☑ Leaf: petiole colour (RHS colour chart)	146B	153A

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'AATS'	'Bronzed Aussie'
☐ Stem: basal branching	medium	medium

Statistical Table

Organ/Plant Part: Context	'AATS'	'Bronzed Aussie'
☐ Leaf: length (mm)		

Mean	41.20	37.80
Std. Deviation	3.60	4.00
LSD/sig	4.32	ns
☑ Leaf: width (mm)		
Mean	18.30	15.60
Std. Deviation	1.10	1.90
LSD/sig	1.75	P≤0.01

Prior Applications and Sales

Nil.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW

Details of Application

Application Number	2005/317
Variety Name	'DOW30'
Genus Species	*Acmena smithii*
Common Name	'Lilly Pilly'
Synonym	Nil
Accepted Date	29 Apr 2006
Applicant	Downes Wholesale Nursery Pty Ltd, Rossmore, NSW
Agent	Ozbreed Pty Ltd, Clarendon, NSW
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Theresa Park, NSW.
Descriptor	Lilly Pilly (*Acmena smithii*/*Syzygium* sp) PBR LILL.
Period	Summer - autumn 2007.
Conditions	Trial conducted in open beds, plants propagated from cuttings, planted into 100L bags filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random.
RHS Chart - edition	2001.

Origin and Breeding

Seedling selection: seed parent *Acmena smithii*. The seed parent is characterised by a medium plant height and width and reddish immature leaf colour. In 2003, approximately 1000 seedlings arising from open-pollinated *Acmena smithii* were grown in an open-bed. A single seedling was selected due to its distinctive lime green colour. Selection took place in Tuckombil, NSW in 2003. Selection criteria: green immature leaf colour. Propagation: vegetative cuttings were found to be uniform and stable. Breeder: Greg Hellyar, Tuckombil, NSW.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	bushy to upright
Plant	branch density	dense to very dense

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
A. smithii	parent of candidate

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'DOW30'	*A. smithii*
☐ Plant: growth habit	bushy to upright	bushy to upright
☐ Plant: height	medium	medium

☐ Plant: branch density	dense to very dense	dense to very dense
☐ Stem: branch angle	acute	acute
☐ Stem: internode length	medium to long	medium
☐ Stem: colour of mature stem (RHS colour chart)	198A and 166C	198A and 166C
☑ Stem: colour of new growth (RHS colour chart)	144B	199A to 200D
☑ Leaf: blade length	long	medium
☑ Leaf: blade width	medium to broad	medium
☑ Leaf: petiole length	long	medium
☐ Leaf: shape of blade	elliptic	elliptic
☐ Leaf: shape of apex	acuminate	acuminate
☐ Leaf: shape of base	acuminate	acuminate
☐ Leaf: glossiness	medium	medium
☑ Leaf: shape of cross section	strongly concave	flat to concave
☐ Leaf: shape of longitudinal section	convex	convex to flat
☐ Leaf: stiffness	medium	medium
☐ Leaf: prominence of midrib on lower surface	not prominent	not prominent
☐ Mature leaf: primary colour of upper side (RHS colour chart)	147A	147A
☑ Mature leaf: primary colour of lower side (RHS colour chart)	146C	146B
☐ Partly mature leaf: primary colour of upper side (RHS colour chart)	ca 146A	146A
☑ Partly mature leaf: primary colour of lower side (RHS colour chart)	146D	152A-B
☑ Newly emerged: upper side (RHS colour chart)	N144A	ca 165A
☐ Leaf: variegation	absent	absent
☐ Leaf: petiole colour (RHS colour chart)	ca 153A	ca 153A

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	**'DOW30'**	***A. smithii***
☑ Stem: basal branching	absent or very weak	strong

Statistical Table

Organ/Plant Part: Context	**'DOW30'**	***A. smithii***
☑ Leaf: length (mm)		
Mean	65.70	46.30
Std. Deviation	8.70	6.60
LSD/sig	8.82	P≤0.01
☑ Leaf: width (mm)		
Mean	31.60	21.40

Std. Deviation	6.00	3.00
LSD/sig	5.37	P≤0.01

Prior Applications and Sales

Prior applications nil. First sold in Australia in Aug 2005 under the name DOW30

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW

Details of Application

Application Number	2005/072
Variety Name	'Bournda Gold'
Genus Species	*Philotheca myoporoides*
Common Name	Long Leaved Waxflower
Synonym	Nil
Accepted Date	14 Jun 2005
Applicant	Lystare Pty Ltd trading as Bournda Plants
Agent	Greenhills Propagation Nursery Pty Ltd, Tynong, VIC
Qualified Person	Mark Lunghusen

Details of Comparative Trial

Location	Greenhills Propagation Nursery, Tynong, VIC.
Descriptor	Philotheca (*Philotheca*) PBR PHIL.
Period	Spring/summer 2007.
Conditions	Plants were grown in 14cm pots in full sun in commercial pine bark based potting mix with controlled release fertiliser. Plants were grown on benches with overhead watering.
Trial Design	10 plants in block design.
Measurements	Leaf measurements taken from middle third of stem.
RHS Chart - edition	2005.

Origin and Breeding

Spontaneous mutation: a variegated sport was selected from *Philotheca myoporoides* 'Bournda Beauty' in 2000. Cuttings were taken from this sport, established, and then a number of generations of cuttings were taken from the young plants. This was repeated a number of times to determine distinctness, uniformity and stability. To date, the plant has been grown through four generations with no off-types being recorded. Selection criteria: leaf colour. Propagation: vegetative. Breeder: Dave Theobald, Merimbula, NSW.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	variegation	present

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Moon Shadow'	Closest variegated variety.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Bournda Gold'	'Moon Shadow'
☐ Plant: growth habit	bushy	bushy
☑ Plant: height	medium to tall	short
☑ Plant: width	medium	narrow
☐ Plant: density	medium	medium
☐ Stem: length of internode	medium	short
☐ Young leaf: variegation	present	present
☐ Leaf: variegation	present	present
☐ Leaf: main colour of upper side (RHS Colour Chart)	yellow green 147B	yellow green 147C
☐ Leaf: secondary colour of upper side (RHS Colour Chart)	yellow 10B	yellow 11B
☑ Leaf: shape	oblanceolate	elliptical
☐ Leaf: shape of apex	acute	acute
☑ Leaf: shape of base	attenuate	cuneate
☐ Leaf: shape in cross section	concave	concave
☐ Leaf: undulation of margin	absent or weak	absent or weak
☐ Flower bud: colour (RHS Colour Chart)	75B	75D
☐ Flowers: arrangement	clusters	clusters
☐ Flower: main colour	whitish	whitish
☐ Petal: main colour (RHS Colour Chart)	white 155C	white 155C
☐ Petal: length	short	short
☐ Petal: shape	elliptic	elliptic
☐ Peduncle: length	short	short
☐ Pedicel: length	short	short
☐ Pedicel: colour (RHS Colour Chart)	green	green

Statistical Table

Organ/Plant Part: Context	'Bournda Gold'	'Moon Shadow'
☑ Leaf: length (mm)		
Mean	29.73	53.40
Std. Deviation	3.40	3.53
LSD/sig	18.53	P≤0.01
☑ Leaf: width (mm)		
Mean	7.57	13.10
Std. Deviation	0.44	0.99
LSD/sig	4.59	P≤0.01
☑ Leaf: length to width ratio (mm)		
Mean	3.93	4.09
Std. Deviation	0.44	0.38
LSD/sig	0.12	P≤0.01

Prior Applications and Sales

Prior applications nil. First sold in Australia in Sep 2004 under the name 'Bournda Gold'

Description: **Mark Lunghusen**, World Select Plants, Cranbourne, VIC.

Details of Application

Application Number	2000/301
Variety Name	'Minijac'
Genus Species	*Mangifera indica*
Common Name	Mango
Synonym	Nil
Accepted Date	30 Nov 2000
Applicant	Herminia and Jacinto Lay, Colton Park Trading Pty Ltd, Darwin, NT
Agent	N/A
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Noonamah, NT
Descriptor	Mango (new) (*Mangifera indica*) TG/112/4
Period	Spring 2007
Conditions	Trial conducted with mature trees under a typical orchard system and with typical management with uniform growing conditions.
Trial Design	Five plants of each variety; no formal design used as plants were from within a standard block planting.
Measurements	Randomly selected ten fruits.
RHS Chart - edition	1995

Origin and Breeding

Open pollination: The new variety originated as an open-pollinated seedling of 'Nam Dok Mai'. The parent is characterised by a green and yellow mature fruit skin colour and an absence of pink blush on skin of immature fruit. The seedling fruited in 1992 and the unique and attractive features of the fruits were noted in 1994. Selection took place in Noonamah, NT. Selection criteria: colour of skin of fruit. Propagation: vegetative grafts were found to be uniform and stable. Breeders: Herminia and Jacinto Lay, Noonamah, NT.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Mature fruit	presence of sinus	present
Mature fruit	shape of dorsal shoulder	sloping downward
Mature fruit	point at stylar scar	medium
Ripe fruit	turpentine flavour	absent
Seed	embryony	polyembryonic
Time of	beginning of flowering	medium
Time of	fruit maturity	medium

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Nam Dok Mai'	

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
'TPP1'	immature fruit	colour of skin	green with pink blush	green only
'TPP1'	mature fruit	colour of skin	green with pink blush	predominantly green

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Minijac'	'Nam Dok Mai'
☐ *Tree: attitude of main branches	erect	erect
☑ *Young leaf: intensity of anthocyanin colouration	medium	weak
☑ Leaf blade: length	medium	long
☑ Leaf blade: width	narrow to medium	medium to broad
☐ *Leaf blade: ratio length/width	large	large to very large
☐ Leaf blade: shape	elliptic	elliptic
☐ Leaf blade: colour	medium green	medium green
☐ Leaf blade: twisting	present	present
☐ Leaf blade: spacing of secondary veins	medium	medium
☑ Leaf blade: undulation of margin	absent or weak	medium
☐ Leaf blade: shape of base	acute	acute
☑ Leaf blade: shape of apex	acute	acuminate
☐ Petiole: attitude in relation to shoot	semi erect	semi erect
☐ Petiole: length	medium	short to medium
☐ *Inflorescence: length	medium	medium
☐ *Mature fruit: length	medium to long	medium to long
☑ *Mature fruit: width	narrow	medium
☑ *Mature fruit: ratio length/width	large to very large	medium to large
☐ *Mature fruit: shape in cross section	broad elliptic	broad elliptic
☑ *Mature fruit: colour of skin	green and pink	green and yellow
☐ Mature fruit: density of lenticels	sparse to medium	sparse to medium
☑ Mature fruit: colour contrast between lenticels and skin	weak to medium	very weak
☐ Mature fruit: size of lenticels	small to medium	small to medium
☐ Mature fruit: roughness of surface	absent	absent
☐ Mature fruit: stalk cavity	absent or shallow	absent or shallow
☐ Mature fruit: presence of neck	absent	absent
☑ *Mature fruit: shape of ventral shoulder	sloping downward	rounded downward
☐ *Mature fruit: shape of dorsal shoulder	sloping downward	sloping downward
☐ Mature fruit: length of groove in ventral shoulder	absent or short	absent or short
☐ Mature fruit: bulging on ventral shoulder	absent	absent

*Mature fruit: presence of sinus	present	present
*Mature fruit: depth of sinus	medium	shallow
*Mature fruit: bulging proximal of stylar scar	absent or weak	medium
Mature fruit: point at stylar scar	medium	medium
Mature fruit: diameter of stalk attachment	medium	medium
*Ripe fruit: predominant colour of skin	yellow and orange	yellow
Ripe fruit: speckling of skin	weak	weak
Ripe fruit: thickness of skin	thin to medium	medium
Ripe fruit: adherence of skin to flesh	weak	medium to strong
Ripe fruit: main colour of flesh	medium orange	light yellow
Ripe fruit: firmness of flesh	soft	medium
Ripe fruit: juiciness	medium	low to medium
Ripe fruit: texture of flesh	fine	fine to medium
*Ripe fruit: amount of fiber attached to stone	low	very low to low
Ripe fruit: amount of fiber attached to skin	medium	medium
*Ripe fruit: turpentine flavor	absent	absent
Stone: relief of surface	grooved	ridged
Seed: shape in lateral view	reniform	reniform
*Seed: embryony	polyembryonic	polyembryonic
Time of: beginning of flowering	medium	medium
*Time of: fruit maturity	medium	medium

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	**'Minijac'**	**'Nam Dok Mai'**
Immature fruit: presence of pink blush	present	absent

Statistical Table

Organ/Plant Part: Context	**'Minijac'**	**'Nam Dok Mai'**
Leaf blade: length (mm)		
Mean	188.00	230.50
Std. Deviation	14.40	21.20
LSD/sig	20.67	P≤0.01
Leaf blade: width (mm)		
Mean	45.20	53.40
Std. Deviation	4.40	5.90
LSD/sig	5.96	P≤0.01
Leaf blade: ratio length/width		
Mean	4.18	4.33
Std. Deviation	0.40	0.30
LSD/sig	0.35	ns
Petiole: length (mm)		
Mean	36.70	30.60

Std. Deviation	5.90	4.20
LSD/sig	5.88	P≤0.01
☐ Mature fruit: length		
Mean	145.50	151.60
Std. Deviation	10.60	13.70
LSD/sig	13.96	ns
☑ Mature fruit: width (mm)		
Mean	64.20	76.50
Std. Deviation	3.30	6.20
LSD/sig	5.68	P≤0.01
☑ Mature fruit: ratio length/width		
Mean	2.27	1.98
Std. Deviation	0.10	0.10
LSD/sig	0.15	P≤0.01

Prior Applications and Sales

Nil.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW.

Details of Application

Application Number	2004/250
Variety Name	'PHORD1'
Genus Species	*Phormium tenax*
Common Name	New Zealand Flax
Synonym	Nil
Accepted Date	21 Sep 2004
Applicant	Ozbreed Pty Ltd, Clarendon, NSW
Agent	N/A
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Clarendon, NSW.
Descriptor	Phormium (*Phormium tenax*) PBR PHOR
Period	Autumn 2007 - spring 2007.
Conditions	Trial conducted in open beds, plants propagated from cuttings, planted into 140mm pots filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random.
RHS Chart - edition	2001.

Origin and Breeding

Seedling selection: In 1999 a batch of commercial seed (most likely *P. tenax purpurea*) were sown and approximately 140000 plants were grown. The seed parent is characterised by a purple leaf colour and weak resistance to phytophthora. In late 2000, about 200 red leaf forms were selected and grown on. During 2001 they were exposed to phytophthora and 130 survived. Finally, in early 2002 a single plant was selected due to its red leaf colour and short height. This selection was later named 'PHORD1'. Selection took place in Clarendon, NSW in 2002. Selection criteria: red leaf colour and strong resistance to phytophthora. Propagation: vegetative, micropropagation is found to be uniform and stable. Breeder: Todd Layt, Clarendon, NSW.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf blade	predominant colour	dark red

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Merlot'	

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	
'Dark Delight'	Leaf blade	predominant colour	dark red	lighter red
'Anna Red'	Leaf blade	presence of red margin	absent	present

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'PHORD1'	'Merlot'
☑ Plant: height	short	medium
☑ Plant: width	narrow	medium
☐ Plant: number of leaves	medium to many	medium
☐ Plant: main colour	red	red
☐ Leaf: length	short	short to medium
☐ Leaf: width at broadest part	narrow	narrow
☐ Young leaf: main colour of middle zone on upper side (RHS colour chart)	darker than N77A	ca N77A
☐ Young leaf: main colour of margin zone on upper side (RHS colour chart)	darker than N77A	ca N77A
☐ Young leaf: main colour of middle zone on lower side (RHS colour chart)	darker than N77A	ca N77A
☐ Young leaf: secondary colour of margin zone on lower side (RHS colour chart)	darker than N77A	ca N77A
☐ Leaf: main colour of middle zone on upper side (RHS colour chart)	ca 187A	ca 187A
☐ Leaf: width of middle zone on upper side	from two thirds to full width of leaf	from two thirds to full width of leaf
☑ Leaf: colour of edge on upper side (RHS colour chart)	187C	ca 202A
☑ Leaf: main colour of middle zone on lower side (RHS colour chart)	ca N77A	ca N200A
☑ Leaf: colour of edge on lower side (RHS colour chart)	187C	ca 202A

Statistical Table

Organ/Plant Part: Context	'PHORD1'	'Merlot'
☑ Plant: height (mm)		
Mean	266.00	376.50
Std. Deviation	31.80	74.70
LSD/sig	65.52	P≤0.01
☑ Plant: width (mm)		
Mean	245.00	542.50
Std. Deviation	36.10	109.20
LSD/sig	92.78	P≤0.01
☑ Leaf: length (mm)		
Mean	242.70	364.60
Std. Deviation	28.80	81.50

LSD/sig	69.77	P≤0.01
Leaf: width (mm)		
Mean	18.50	14.40
Std. Deviation	2.90	3.60
LSD/sig	3.74	ns

Prior Applications and Sales

Nil.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW.

Details of Application

Application Number	2007/260
Variety Name	'Storm Edition'
Genus Species	*Phormium cookianum*
Common Name	New Zealand Mountain Flax
Synonym	Nil
Accepted Date	22 Nov 2007
Applicant	Greenhills Propagation Nursery Pty Ltd, Tynong, VIC
Agent	N/A
Qualified Person	Mark Lunghusen

Details of Comparative Trial

Location	Tynong, VIC.
Descriptor	Phormium (*Phormium tenax*) PBR PHOR.
Period	Autumn to spring 2007.
Conditions	Plants were grown in 14cm pots in full sun in commercial pine bark based potting mix with controlled release fertiliser. Plants were grown on benches with overhead watering.
Trial Design	10 plants in block design.
Measurements	Leaf measurements taken from largest leaves.
RHS Chart - edition	2005.

Origin and Breeding

Spontaneous mutation: a dark purple sport was selected from green form of *Phormium cookianum* in 2004. Divisions were taken from this sport, established, and then a number of generations of divisions were taken from the young plants. This was repeated two further times to determine distinctness, uniformity and stability. To date, the plant has been grown through four generations with no off-types being recorded. Selection criteria: leaf colour. Propagation: vegetative. Breeder: Robert Harrison, Tynong VIC.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	colour	purple

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Purple Haze'	
'Chocolate Cookie'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Storm Edition'	'Chocolate Cookie'	'Purple Haze'
☑ Plant: height	short to medium	medium	medium
☑ Plant: width	medium to broad	medium	medium
☐ Plant: number of suckers	very few	very few	very few
☐ Plant: main colour	brown	brown	brown

☐ Leaf: width at broadest part		medium	medium
☐ Leaf: main colour of margin zone on upper side (RHS colour chart)	ca200A	200A	200A
☑ Leaf: main colour of middle zone on lower side (RHS colour chart)	200A	N200A	N200A

Statistical Table

Organ/Plant Part: Context	**'Storm Edition'**	**'Chocolate Cookie'**	**'Purple Haze'**
☑ Plant: height (mm)			
Mean	460.00	577.70	512.00
Std. Deviation	46.67	48.25	31.55
LSD/sig	53.06	P≤0.01	P≤0.01
☑ Plant: width (mm)			
Mean	71.00	47.20	59.50
Std. Deviation	4.59	2.49	5.50
LSD/sig	5.42	P≤0.01	P≤0.01
☑ Plant: number of shoots			
Mean	13.00	2.88	7.10
Std. Deviation	2.62	1.10	1.66
LSD/sig	2.35	P≤0.01	P≤0.01
☑ Leaf: width (mm)			
Mean	21.55	24.23	26.07
Std. Deviation	2.30	2.76	2.45
LSD/sig	3.11	ns	P≤0.01

Prior Applications and Sales

Prior applications nil.

First sold in Australia in Nov 2006 under the name 'Storm Edition'.

Description: **Mark Lunghusen,** World Select Plants, Cranbourne, VIC.

Details of Application

Application Number	2007/241
Variety Name	'Dawson'
Genus Species	*Avena sativa*
Common Name	Oats
Synonym	Nil
Accepted Date	7 Nov 2007
Applicant	NDSU Research Foundation, Fargo, ND, USA
Agent	Pacific Seeds Pty Ltd, Toowoomba, QLD
Qualified Person	Peter Johnson

Details of Comparative Trial

Location	Gatton, QLD.
Descriptor	Oats (*Avena sativa*) TG/20/10.
Period	Winter – spring 2007. Sown 26 Apr 2007.
Conditions	The trial was sown into a well prepared seedbed at the Pacific Seeds Research Station, located at Gatton in the Lockyer Valley of South East Queensland. The trial was conducted under irrigated conditions using a row spacing of 76 cm.
Trial Design	The trial design was a randomized complete block with four replications, four rows per plot, five metres long.
Measurements	Measurements were taken from 20 plants selected at random from over 2,500 plants.
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: ND 020290 ('Dawson') resulted from a cross made in the fall greenhouse season of 1998 with the pedigree MN 97112 x ND 971454. The F_1 was grown in the 1999 spring greenhouse and F_2 populations were grown in the field at Fargo in 1999. Single plant selections were made from the F_2. F_3 selections were screened for seedling resistance to crown and stem rust. F_4 planted in hill plots in 2000. Single panicle selections were made from the hillplots to produce the $F_{4:5}$ breeding line, which became the source of ND 020290. This seed was planted in four-row plots for evaluation in the field at Fargo in 2002. ND 020290 was evaluated in a preliminary yield trial at two locations in 2003, and advanced yield trials at four locations in 2004. During all stages of development, ND 020290 was subjected to stringent selection for resistance to races of crown rust and stem rust prevalent in North Dakota. Breeder: NDSU Research Foundation, Fargo, ND, USA.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Panicle	orientation of branches	equilateral
Panicle	attitude of spikelets	pendulous
Stem	hairiness of top node	absent
Grain	husk	present

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Volta'	Commercial forage variety.
'Drover'	Forage oat variety released by Pacific Seeds in 2006.
'Taipan'	Forage oat variety from Pacific Seeds.
'Genie'	Commercial slow rusting forage oat variety released in 2006.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Dawson'	'Drover'	'Genie'	'Taipan'	'Volta'
☐ Plant: growth habit	erect	intermediate	erect	erect	erect to semi-erect
☐ Lowest leaves: hairiness of sheaths	absent or very weak	absent or very weak	very weak to weak	absent or very weak	very weak to weak
☐ *Leaf blade: hairiness of margins of leaf below flag leaf	absent or very weak	absent or very weak	absent or very weak	absent or very weak	absent or very weak
☐ Plant: frequency of plants with recurved flag leaves	low	low to medium	medium	low to medium	low to medium
☐ *Time of: panicle emergence	medium to late	medium to late	late	late	medium
☐ *Stem: hairiness of uppermost node	absent	absent	absent	absent	absent
☐ Panicle: orientation of branches	equilateral	equilateral	equilateral	equilateral	equilateral
☐ Panicle: attitude of branches	semi-erect	semi-erect	semi-erect	semi-erect	horizontal
☐ Panicle: attitude of spikelets	pendulous	pendulous	pendulous	pendulous	pendulous
☐ Glumes: glaucosity	weak	medium	weak to medium	weak	weak to medium
☐ Glumes: length	medium	medium	medium	medium to long	long
☐ *Primary grain: glaucosity of lemma	absent	absent	absent	absent	absent
☐ *Plant: length	long	medium	long	long	long
☑ Panicle: length	medium	short to medium	long to very long	long to very long	short to medium
☐ *Grain: husk	present	present	present	present	present
☑ Primary grain: tendency to be awned	weak	absent or very weak	weak	strong	weak
☐ Primary grain: length of lemma	medium	medium	medium	medium	medium

*Grain: colour of lemma	yellow	yellow	yellow	yellow	yellow
Primary grain: hairiness of back of lemma	absent	absent	absent	absent	present
Primary grain: hairiness of base	absent or very weak	absent or very weak	medium to strong	weak	very strong
Primary grain: length of basal hairs	very short	very short	medium	medium	long
Primary grain: length of rachilla	medium	medium	medium	medium	medium

Statistical Table

Organ/Plant Part: Context	**'Dawson'**	**'Drover'**	**'Genie'**	**'Taipan'**	**'Volta'**
Plant (stem & panicle): height (cm)					
Mean	176.40	160.00	183.00	176.90	179.20
Std. Deviation	8.96	5.53	8.14	4.90	9.28
LSD/sig	2.49	P≤0.01	P≤0.01	ns	P≤0.01
Flag leaf: width (mm)					
Mean	32.10	29.90	31.00	31.60	23.30
Std. Deviation	2.61	3.17	2.26	3.19	2.96
LSD/sig	1.53	P≤0.01	ns	ns	P≤0.01
Flag leaf: length (mm)					
Mean	186.70	247.30	312.90	276.50	223.10
Std. Deviation	36.62	45.47	50.99	54.60	42.36
LSD/sig	6.16	P≤0.01	P≤0.01	P≤0.01	P≤0.01
Panicle: length (mm)					
Mean	364.20	330.80	455.60	441.60	331.10
Std. Deviation	36.25	56.76	50.47	30.57	36.19
LSD/sig	5.95	P≤0.01	P≤0.01	P≤0.01	P≤0.01

Prior Applications and Sales

Nil.

Description: **Peter Johnson**, Pacific Seeds Pty Ltd, Toowoomba, QLD.

Details of Application

Application Number	2007/150
Variety Name	'Monty'
Genus Species	*Avena sativa*
Common Name	Oats
Synonym	Nil
Accepted Date	26 Jun 2007
Applicant	New Zealand Institute for Crop & Food Research Limited, Christchurch, NZ
Agent	Heritage Seeds Pty Ltd, Howlong, NSW
Qualified Person	Allen Newman

Details of Comparative Trial

Location	Heritage Seeds Research, "Shrublands", Riverina Highway, Howlong, NSW 2643 (Latitude 35o60' South, elevation 150m), autumn-summer 200
Descriptor	Oats (*Avena sativa*) TG/20/10
Period	June - December 2007
Conditions	Trial was sown into a moist, cultivated seed bed in good condition. The trial was sown using a cone seeder. Normal agronomic practices were applied to the trial.
Trial Design	Three replicates laid out in a randomised block design.
Measurements	Measurements were taken from at least five plants per replicate (15 plants per entry), selected at random.
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: parentage Pg16//OT240. In 1992/93 F_2 population was selected from 'Aorangi' research site close to Palmerston North. During 1994 to 1998, F_3 to F_7 selections were carried out on CFR site located near Gore, New Zealand. Two further selections (F_8 and F_9) were carried out in New Zealand for forage production and disease resistance. Panicles were harvested from superior lines. F_{10} panicle rows sown and increased under NZ Australian quarantine protocols for shipment to Australia. Re-selected seed lines harvested as individual bulks and shipped to Heritage seed in NSW for field evaluation. Between 2001 -06 evaluated as "CDA28,01/G4" by Heritage Seeds for forage potential using a parallel system of small forage plot trials, seed multiplication for on-farm evaluation and pure seed production. Selection criteria: grain type, forage production, Barley yellow dwarf virus resistance, lodging resistance. Propagation: seed. Breeder: New Zealand Institute for Crop & Food Research Limited, Christchurch, NZ.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Lowest leaves	hairiness of sheaths	absent or very weak
Plant	frequency of plants with recurved flag leaves	absent or very low
Plant	time of panicle emergence	late/very late
Panicle	orientation of branches	equilateral
Panicle	attitude of spikelets	pendulous
Primary grain	glaucosity of lemma	absent
Primary grain	hairiness of back of lemma	absent
Grain	husk	present

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Graza 50'	
'Graza 51'	
'Graza 68'	
'Graza 80'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Monty'	'Graza 50'	'Graza 51'	'Graza 68'	'Graza 80'
☐ Plant: growth habit	erect to semi-erect	semi-erect	semi-erect	semi-erect	semi-erect to intermediate
☐ Lowest leaves: hairiness of sheaths	absent or very weak	absent or very weak	absent or very weak	absent or very weak	absent or very weak
☑ *Leaf blade: hairiness of margins of leaf below flag leaf	absent or very weak	absent or very weak	medium	weak	weak
☐ Plant: frequency of plants with recurved flag leaves	absent or very low	absent or very low	absent or very low	absent or very low	absent or very low
☑ *Time of: panicle emergence	late	late	very late	late	late to very late
☑ *Stem: hairiness of uppermost node	absent	absent	present	present	present
☐ Panicle: orientation of branches	equilateral	equilateral	equilateral	equilateral	equilateral
☐ Panicle: attitude of branches	erect to semi-erect	semi-erect to horizontal	semi-erect	semi-erect	semi-erect to horizontal
☐ Panicle: attitude of spikelets	pendulous	pendulous	pendulous	pendulous	pendulous
☑ Glumes: glaucosity	very strong	medium to strong	weak to medium	medium	medium
☑ Glumes: length	medium	medium to long	medium to long	long	medium to long

☐ *Primary grain: glaucosity of lemma	absent	absent	absent	absent	absent
☐ *Plant: length	long to very long	long	long to very long	long	long
☐ Panicle: length	medium	medium to long	medium to long	medium	medium
☐ *Grain: husk	present	present	present	present	present
☑ Primary grain: tendency to be awned	absent or very weak	absent or very weak	absent or very weak	medium	absent or very weak
☐ Primary grain: length of lemma	medium	medium	medium	medium to long	medium
☑ *Grain: colour of lemma	white	white	white	yellow	white
☐ Primary grain: hairiness of back of lemma	absent	absent	absent	absent	absent
☑ Primary grain: hairiness of base	absent or very weak	strong	strong	absent or very weak	strong
☐ Primary grain: length of rachilla	medium	medium	medium to long	medium	medium

Statistical Table

Organ/Plant Part: Context	**'Monty'**	**'Graza 50'**	**'Graza 51'**	**'Graza 68'**	**'Graza 80'**
☐ Plant: height (cm)					
Mean	803.67	716.00	805.67	725.33	738.00
Std. Deviation	44.96	52.31	42.78	45.44	36.95
LSD/sig	97.80	ns	ns	ns	ns
☑ Flag leaf: length (mm)					
Mean	91.50	84.80	92.30	106.10	106.90
Std. Deviation	11.69	9.47	10.83	8.93	14.83
LSD/sig	14.4	ns	ns	P≤0.01	P≤0.01
☑ Flag leaf: width (mm)					
Mean	10.93	9.27	12.20	10.40	10.47
Std. Deviation	2.43	0.90	1.34	1.66	0.97
LSD/sig	1.61	P≤0.01	P≤0.01	ns	ns

Prior Applications and Sales

Nil.

Description: **Allen Newman**, Heritage Seeds Pty Ltd, Howlong, NSW.

Details of Application

Application Number	2006/002
Variety Name	'Georgia Hi/OL'
Genus Species	*Arachis hypogaea*
Common Name	Peanut
Synonym	Reid
Accepted Date	8 May 2006
Applicant	University of Georgia Research Foundation, Inc., Athens, GA, USA
Agent	Peanut Company of Australia Limited, Kingaroy, QLD
Qualified Person	Grant Baker

Details of Comparative Trial

Location	Chinchilla and Kingaroy QLD.
Descriptor	Peanut (*Arachis*) TG/93/3.
Period	Summer 2005 until late autumn 2006.
Conditions	Chinchilla trial was an irrigated trial, with incidences of Pod Rot later in the season. Plot size was 2x5 to 6 m rows with 3 replicates. this Chinchilla trial included 24 entries which included both the candidate and the comparator. The Kingaroy trial was grown under well irrigated conditions. Plot size was 2x5 to 6m rows with 3 replicates. This trial included 14 entries including once again both the candidate and the comparator.
Trial Design	Experimental designs employed were row – column designs, row lattices and randomised complete block design.
Measurements	Establishment, pod yield, kernel yield, total kernel percentage, graded outturn and estimated crop value and kernel counts.
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: 'Georgia Hi-O/L' is the result of an initial cross between 'GA-C330A' and 'GA-T2636'. Pedigree selection was undertaken through the F_2-F_5 and yield tests were performed from the F_6-F_8. Selection criteria: high oleic to linoleic acid ratio. Breeder: Dr. William D. Branch, University of Georgia, Department of Crop and Soil Sciences, Coastal Plain Experimental Station, Tifton, GA, USA.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	**Context**	**State of Expression in Group of Varieties**
Plant	commercial grouping	runner
Plant	time of maturity	late
Kernel	oleic to linoleic acid ratio	high
Kernel	colour of mature uncured testa	pink
Kernel	weight per 1000 kernels	medium to high

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Holt'	most similar variety based on the above grouping criteria

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Menzies'	Kernel weight per 1000 kernels	medium to high	low to medium	Trials in 2003/2004 as stated in application No. 2003/317 and 2005/ 2006 yield performance trials.
'Scullin'	Kernel colour of mature uncured testa	pink	flesh	
'Bruce'	Kernel colour of mature uncured testa	pink	flesh	
'GA-C330A'	Kernel oleic to linoleic acid ratio	high	low	As per part 1.
'GA-T2636'	Kernel size	medium	small	As per part 1.
'SO95R'	Kernel weight per 1000 kernels	medium to high	low to medium	Trials in 2003/2004 as stated in application No. 2003/317 and 2005/2006 yield performance trials.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Georgia Hi/OL'	'Holt'
☑ *Plant: growth habit	prostrate	semi-erect
☐ Main stem: growth habit (prostrate varieties only)	erect	erect
☐ *Time of: maturity	late	late
☑ Leaflet: colour	dark green	medium green
☐ *Flowering: general pattern	sequential	sequential
☑ *Pod: constrictions	deep	medium
☐ *Pod: prominence of beak	absent or very inconspicuous to inconspicuous	absent or very inconspicuous
☐ *Pod: shape of beak	curved	curved
☐ *Kernel: colour of uncured mature testa	monochrome	monochrome
☐ *Kernel: colour of mature uncured testa (varieties with monochrome testa only)	pink	pink
☐ *Kernel: weight per 1000 kernels	medium to high	medium to high
☐ *Kernel: dormancy period	medium	medium
☐ Resistance to: rust	absent	absent

Prior Applications and Sales

Country	Year	Current Status	Name Applied
USA	2000	Granted	'Georgia Hi/OL'

Prior sale nil.

Description: **Grant Baker,** Peanut Company of Australia Limited, Kingaroy, QLD.

Details of Application

Application Number	2006/084
Variety Name	'Konimpa'
Genus Species	*Alstroemeria* hybrid
Common Name	Peruvian Lily
Synonym	Nil
Accepted Date	8 May 2006
Applicant	Konst Breeding B.V., Nieuwveen, The Netherlands
Agent	N/A
Qualified Person	David Nichols

Details of Comparative Trial

Overseas Testing Authority	Community Plant Variety Office (CPVO)
Overseas Data Reference Number	INC 874
Location	Overseas data was verified in Monbulk, VIC.
Descriptor	Alstroemeia (*Alstroemeria*) TG/29/6.
Period	Dec 2007.
Conditions	Comparisons of most characteristics are based on Dutch trials, which were assessed under conditions of controlled environment in glasshouses at Wageningen, the Netherlands. Detailed flower descriptions of the candidate variety are based on plants growing in soil in a multispan polyhouse at Monbulk, VIC. Flowers from these plants were cut in bud and transferred to Devon Meadows, VIC, and placed in a solution of 5% sugar and 1ml/l chlorine bleach. The flowers were assessed 3 days later. Descriptions of the comparators are derived from those published in the Plant Varieties Journal.
Trial Design	Completely randomised
Measurements	Taken from all trial plants.
RHS Chart - edition	2001.

Origin and Breeding

Controlled pollination: seed parent '8656-1' x pollen parent '8124-7', in a planned breeding program at the applicant's research station at Nieuwveens, the Netherlands. Both parents are non-commercial varieties within the breeding programme. Selection criteria: growth characteristics and bi-colour flower. Propagation: a number of mature stock plants were generated from the original seedling by tissue culture through 3 generations to confirm uniformity and stability. Breeder: J. W. Konst, Konst Breeding B.V., Nieuwveen, The Netherlands.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Flower	main colour	orange red
Outer tepal	colour of centre	red
Outer tepal	colour of edges	orange
Stem	length	long to very long

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Stanata'	PVJ 12(3)

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Pink Roma'	Outer tepal colour of edges	orange	pink	PVJ 12(2)
'Cerise Miami'	Outer tepal colour of edges	orange	Pink	PVJ 12(2)

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Konimpa'	'Stanata'
☑ *Stem: length	long	very long
☑ *Stem: thickness	thick	medium
☐ *Stem: density of foliage	medium	medium
☑ *Leaf: length	long	medium
☑ *Leaf: width	narrow	medium
☐ *Leaf: shape of blade	elliptic	narrow-elliptic
☐ *Leaf: longitudinal axis of blade	recurved	recurved
☑ *Inflorescence: number of branches in umbel	many	medium
☑ *Inflorescence: length of branches in umbel	medium	long
☑ *Inflorescence: length of pedicel	short	medium
☐ *Flower: main colour	orange red	orange red
☐ *Flower: size	medium to large	large
☐ *Flower: spread of tepals	medium	medium
☐ *Outer tepal: shape of blade	broad obovate	broad obovate
☑ *Outer tepal: depth of emargination	medium	very deep
☑ *Outer tepal: main colour of inner side of blade (RHS colour chart)	32B,46B	53B,29B
☐ *Outer tepal: stripes on inner side of blade	absent	absent
☐ *Inner tepal: shape of blade	elliptic	elliptic
☑ *Inner lateral tepal: main colour of inner side of middle zone of blade (RHS colour chart)	14A	9B
☐ Inner lateral tepal: number of stripes on inner side of blade	medium	medium
☐ *Inner lateral tepal: size of stripes on inner side of blade	medium to large	medium
☐ *Stamens: main colour of filament	pink	pink
☐ *Stamens: small spots on filament	absent	absent
☑ *Stamens: colour of anthers at the start of dehiscence	brownish	greenish
☑ Pistil: anthocyanin colouration of ovary	absent or very	weak to medium

	weak	
☐ Pistil: spots on the stigma	absent	absent

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Konimpa'	'Stanata'
☐ Inner median tepal: presence of stripes	present	present
☑ Inner median tepal: presence of yellow colour	present	absent

Prior Applications and Sales

Country	Year	Current Status	Name Applied
The Netherlands	2005	Applied	'Konimpa'

First sold in Colombia in June 2005. First Australian sale July 2005.

Description: **David Nichols,** Rye, VIC.

Details of Application

Application Number	2005/353
Variety Name	'Aus-Jubilee'
Genus Species	*Ananas comosus*
Common Name	Pineapple
Synonym	Jubilee
Accepted Date	9 Feb 2006
Applicant	State of Queensland through its Department of Primary Industries and Fisheries, Brisbane, QLD
Agent	N/A
Qualified Person	Garth Sanewski

Details of Comparative Trial

Location	Maroochy Research Station, Nambour
Descriptor	Pineapple (*Ananas comosus*) TG/PINEAP (proj. 1)
Period	Planted late Sep 2005, induced on 2 Feb 2007 and harvested from Aug to Sep 2007. Flower data collected Apr 2007.
Conditions	Plants treated according to standard commercial practices with the addition of trickle irrigation. Planting density of approximately 50,000 plants/ha used.
Trial Design	Randomised Complete Block of 5 blocks and 10 plants per variety per block.
Measurements	Reference leaf data and flower data collected on 2 plants/plot (total 10 samples/variety). All plant data collected on 10 plants/plot (total of 50 plants/variety). Fruit data collected on all harvested fruit. Eye dimensions collected on 3 eyes/fruit to give a fruit sample mean. Fruit firmness data results of 3 measurements/fruit to give a fruit sample mean.
RHS Chart - edition	Third edition, 1995.

Origin and Breeding

Controlled pollination: seed parent 'Smooth Cayenne' x pollen parent '73-50' in a planned breeding program on Maroochy Research Station (MRS) at Nambour, Queensland, in 1993 using conventional hand pollination techniques. The seed was extracted and germinated in a glasshouse on MRS in 1994. The seedlings were planted on MRS in Dec 1995. The original seedling, designated 10-2594, was harvested on 12 Sep 1997. The seed parent is characterised by high yield and good plant vigour. The pollen parent is characterised by high flesh aroma, moderate high sugar content, low acidity and yellow flesh. Selection criteria: characters used in the selection included piping leaf margin, high total soluble solids, moderate acidity, good flavour, yellow flesh, and improved resistance to natural flower initiation and translucency. Propagation: the vegetative shoots on the original seedling were collected and planted on MRS in 1997. Replantings using the same method were made approximately every 2 years. In addition approximately 500 plants were produced through meristem culture at MRS using standard protocols for pineapple. Plants considered not similar to the original were discarded at each planting.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	margin type	piping
Fruit/flesh	colour	yellow

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'73-50'	Pollen parent and standard commercial fresh market cultivar.

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Smooth Cayenne'	Leaf presence of anthocyanins (upper surface)	Absent	present	
'Smooth Cayenne'	Leaf leaf margin	Piping	spiny tip	Seed parent to 'Aus-Jubilee'
'73-114'	Leaf leaf margin	Piping	spiny tip	Similar dark green leaf as 'Aus-Jubilee'

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Aus-Jubilee'	'73-50'
☐ *Plant: foliage attitude	semi-erect to spreading	semi-erect to spreading
☐ Plant: leaf emission rate (number of leaves produced from 4 months after planting to forcing)	quick to very quick	medium
☐ Reference leaf: length	short to medium	medium
☐ Reference leaf: maximum width	narrow to medium	medium
☐ Reference leaf: weight	low to medium	medium
☑ *Leaf: predominant colour (on upper face)	dark green	green
☑ *Leaf: presence of anthocyanins (on upper surface)	absent	present
☐ *Leaf: leaf edges aspect	piping	piping
☐ *Plant: fruit habit when ripe	upright	bending to upright
☐ *Peduncle: length	medium	medium to long
☐ *Suckers: mean number of underground suckers per plant	few	few
☐ *Suckers on peduncle: mean number of aerial suckers per plant	medium	few
☐ *Suckers on peduncle: size of aerial suckers at fruit harvest	small	medium
☐ *Slips: presence/absence	present	present
☐ *Slips: number of slips	few	medium
☐ *Crown: height	medium	high
☐ Crown: weight	medium	medium to large

*Fruit: breaking from peduncle	easy	easy
*Fruit: shape when ripe	ovoid	ovoid
*Fruit: predominant skin colour when ripe	golden yellow	yellow
*Fruit: colour uniformity when ripe	with a gradient	with a gradient
*Fruit: height (without neck)	short to medium	medium
*Fruit: diameter at the middle	small to medium	medium
*Fruit: weight (without crown)	low to medium	medium
Fruit: eyes number	medium	small to medium
*Fruit: eye relative surface	medium	medium to large
*Fruit: eye profile	flat	flat
*Fruit/flesh: colour	pale yellow	yellow
*Fruit/flesh: visual appraisal of density or pulp density	strong	medium
Fruit/flesh: firmness	medium to strong or firm	medium
*Fruit/flesh: texture	fibrous	smooth
Fruit/flesh: fibrousness	medium	low to medium
Fruit/flesh: aroma	medium	high
*Fruit/flesh: sugar taste	high	medium to high
*Fruit/flesh: acidic taste	low to medium	low
*Fruit/flesh: juiciness	medium	medium to high
*Fruit/juice: sugar content (using refractometer)	high	medium to high

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Aus-Jubilee'	'73-50'
Peduncle bract: presence of anthocyanin on upper bract surface	slight	strong
Peduncle bract: colour of anthocyanin on upper face	48D	48D
fruit: extent of flesh translucency midway up fruit	slight	moderate
Fruit: eye height at middle of fruit	medium	large
Fruit: eye width at middle of fruit	medium	large

Statistical Table

Organ/Plant Part: Context	'Aus-Jubilee'	'73-50'
Plant: plant height to apex of flowering syncarp (cm)		
Mean	56.50	58.75
Std. Deviation	4.93	5.06
LSD/sig	7.80	ns
Slips: number of slips		
Mean	0.27	1.20
Std. Deviation	0.60	1.39
LSD/sig	0.56	P≤0.01

☐ Sucker: length of longest sucker (cm)		
Mean	44.84	68.29
Std. Deviation	27.85	31.18
LSD/sig	15.87	P≤0.01
☑ Fruit: eye height (mm)		
Mean	24.75	29.04
Std. Deviation	1.65	2.10
LSD/sig	1.63	P≤0.01
☑ Fruit: eye width (mm)		
Mean	23.55	27.91
Std. Deviation	1.22	1.29
LSD/sig	1.06	P≤0.01
☑ Fruit: number of eyes		
Mean	118.65	94.27
Std. Deviation	19.75	11.50
LSD/sig	14.25	P≤0.01
☐ Fruit: ratio eye width to fruit diameter		
Mean	5.22	4.78
Std. Deviation	0.37	0.35
LSD/sig	0.28	P≤0.01
☐ Fruit: diameter at the middle (mm)		
Mean	123.00	133.07
Std. Deviation	10.11	7.86
LSD/sig	7.40	P≤0.01
☐ Fruit: weight without crown (g)		
Mean	1435.0	1765.00
Std. Deviation	381.00	336.00
LSD/sig	303.20	P≤0.01
☐ Fruit/flesh: juiciness (%)		
Mean	42.03	48.86
Std. Deviation	4.90	2.91
LSD/sig	3.83	P≤0.01
☑ Fruit/flesh: sugar content (using refractometer) (%)		
Mean	15.95	14.67
Std. Deviation	1.45	1.23
LSD/sig	1.09	P≤0.01
☐ Crown: height (mm)		
Mean	180.00	221.50
Std. Deviation	33.10	34.20
LSD/sig	28.7	P≤0.01
☐ Crown: weight (g)		
Mean	171.8	238.80
Std. Deviation	44.70	48.20
LSD/sig	36.78	P≤0.01
☐ Fruit: diameter of peduncle scar (mm)		
Mean	34.74	29.07

Std. Deviation	4.68	4.79
LSD/sig	3.80	P≤0.01
Sucker: number of aerial suckers		
Mean	0.99	0.80
Std. Deviation	0.63	0.64
LSD/sig	0.34	ns
Peduncle: length of peduncle (cm)		
Mean	24.18	25.04
Std. Deviation	3.36	11.44
LSD/sig	4.56	ns
Peduncle: width of peduncle (mm)		
Mean	27.04	23.31
Std. Deviation	4.51	4.03
LSD/sig	2.27	P≤0.01

Prior Applications and Sales

Nil prior applications. First sold in July, 2007. Approximately 11 pallet of fruit (7,250 fruit) test marketed as 1 consignment through Brisbane wholesale markets.

Description: **Garth Sanewski,** Maroochy Research Station, Nambour, QLD.

Details of Application

Application Number	2007/036
Variety Name	'Aus-Carnival'
Genus Species	*Ananas comosus*
Common Name	Pineapple
Synonym	Nil
Accepted Date	26 Feb 2007
Applicant	State of Queensland through its Department of Primary Industries and Fisheries, Brisbane, QLD
Agent	N/A
Qualified Person	Garth Sanewski

Details of Comparative Trial

Location	Maroochy Research Station, Nambour, QLD.
Descriptor	Pineapple (*Ananas comosus*) TG/PINEAP (proj. 1).
Period	Jun 2006 – Feb 2007.
Conditions	Plants treated according to standard commercial practices with the addition of trickle irrigation. Planting density of approximately 50,000 plants/ha.
Trial Design	Randomised Complete Block Design with 5 blocks of 10 plants per variety per block.
Measurements	Reference leaf and flower data collected on 2 plants/plot (total 10 samples/variety). All plant data collected on 10 plants/plot (total of 50 plants/variety). Fruit data collected on all harvested fruit. Eye dimensions collected on 3 eyes/fruit. Fruit firmness data collected with 3 measurements/fruit.
RHS Chart - edition	Third edition, 1995.

Origin and Breeding

Controlled pollination: 'Aus-Carnival' was developed using the seed parent '73-50' and pollen parent '53-116'. The pollinations were made in 1993 using conventional hand pollination protocols. The seed was extracted and germinated at Maroochy Research Station, Nambour in 1993. The seedlings were field-planted at Maroochy Research Station in 1995. The original seedling, designated '7-1627', was harvested on 2 June 1997. The seed parent is characterised by high flesh aroma, moderate high sugar, low acidity and yellow flesh. The pollen parent is characterised by low acidity and high resistance to internal browning. Selection criteria: characters used in the selection include piping leaf margin, good fruit weight, high total soluble solids, low acidity, good flavour, yellow flesh and high ratoon yield. Propagation: the vegetative shoots on the original seedling were collected and planted on MRS in 1997. Replantings using the same method were made approximately every 2 years. In addition approximately 500 plants were produced through meristem culture at MRS using standard protocols for pineapple. Plants considered not similar to the original were discarded at each planting.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	anthocyanin pigmentation	present
Leaf	margin type	piping

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'53-116'	Pollen parent to 'Aus-Carnival'.
'73-50'	Seed parent to 'Aus-Carnival'.

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Smooth Cayenne'	leaf	margin type	piping	spiny tip
'73-114'	leaf	anthocyanin pigmentation	present	absent
'73-114'	leaf	margin type	piping	spiny tip

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Aus-Carnival'	'53-116'	'73-50'
☐ *Plant: foliage attitude	semi-erect	semi-erect to spreading	spreading
☐ Reference leaf: length	long	short to medium	medium
☐ Reference leaf: maximum width	medium	medium	narrow to medium
☐ Reference leaf: weight	medium	low	medium
☐ *Leaf: predominant colour (on upper face)	green	green	green
☐ *Leaf: presence of anthocyanins (on upper surface)	present	present	present
☐ *Leaf: level of expression of anthocyanins	weak to medium	weak to medium	medium
☑ Leaf: cross distribution of anthocyanins	mainly on margins	uniform on margins and in the groove	mainly on margins
☑ Leaf: distribution of anthocyanins lengthwise	mainly towards the base	mainly towards the apex	mainly towards the apex
☐ *Leaf: leaf edges aspect	piping	piping	piping
☑ *Peduncle: colour of ventral upper face of bract leaves	34A	39A	34D
☐ Inflorescence: area of petal without colouration	medium	small	small to medium
☐ *Plant: fruit habit when ripe	bending to upright	bending to upright	bending to upright
☐ *Peduncle: length	medium to long	medium	medium

☐ *Suckers: mean number of underground suckers per plant	none or very few	none or very few	none or very few to few
☐ *Suckers on peduncle: mean number of aerial suckers per plant	medium to many	none or very few to few	medium
☐ *Suckers on peduncle: size of aerial suckers at fruit harvest	large to very large	small to medium	medium to large
☐ *Slips: presence/absence	present	present	present
☐ *Slips: number of slips	very few	medium	medium
☐ Crown: proportion of plants with multiple crowns	none or very low	low	none or very low
☐ *Crown: height	medium to high	medium	high
☐ Crown: weight	medium	small	medium
☐ *Fruit: breaking from peduncle	easy	easy	easy
☐ *Fruit: shape when ripe	ovoid	globular	ovoid
☐ *Fruit: predominant colour when ripe	green and yellow	green and yellow	yellow
☐ *Fruit: colour uniformity when ripe	with a gradient	with a gradient	with a gradient
☐ *Fruit: height (without neck)	medium	medium	medium
☐ *Fruit: diameter at the middle	medium	medium	medium
☐ *Fruit: weight (without crown)	medium	medium	medium
☐ Fruit: eyes number	medium	medium to large	medium
☐ *Fruit: eye relative surface	medium	small to medium	medium
☐ *Fruit: eye profile	flat	flat	flat
☐ *Fruit/flesh: colour	yellow	pale yellow	yellow
☐ Fruit/flesh: core diameter	small	small to medium	medium
☐ Fruit/flesh: eye depth	weak	weak to medium	medium
☐ *Fruit/flesh: visual appraisal of density or pulp density	medium	medium to strong	weak to medium
☐ *Fruit/flesh: texture	smooth	fibrous	smooth
☐ Fruit/flesh: fibrousness	medium	medium to high	low to medium
☐ Fruit/flesh: aroma	medium	low	high
☐ *Fruit/flesh: sugar taste	high	low to medium	medium to high
☐ *Fruit/flesh: acidic taste	medium	low	low to medium
☐ *Fruit/flesh: juiciness	medium	medium to high	medium to high
☐ *Fruit/juice: sugar content (using refractometer)	high	low to medium	medium to high

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	**'Aus-Carnival'**	**'53-116'**	**'73-50'**
☑ Crown: presence of anthocyanin on leaf tips 1 month after anthesis	moderate	slight	strong

☑ Crown: predominate colour of leaf tips 1 month after anthesis	dull red	red/brown	pink/red
☐ Fruit: extent of flesh translucency midway up fruit	very slight	moderate	very slight
☐ Reference leaf: cross-section midway along leaf	slightly concave to concave	flat to slightly concave	slightly concave to concave
☐ Crown: diameter of crown base	medium	small to medium	large
☑ Peduncle bract: presence of anthocyanin on upper bract surface	medium	strong	medium
☐ Plant: plant height to apex of flowering syncarp	tall	short	medium
☐ Fruit: eye height at middle of fruit	large	medium	medium
☐ Fruit: eye width at middle of fruit	large	medium	large

Statistical Table

Organ/Plant Part: Context	**'Aus-Carnival'**	**'53-116'**	**'73-50'**
☐ Plant: plant height to apex of flowering syncarp			
Mean	53.6	43.0	47.5
Std. Deviation	4.03	5.70	4.70
LSD/sig	4.8	P≤0.01	P≤0.01
☐ Slips: number of slips			
Mean	0.40	1.40	1.60
Std. Deviation	0.88	1.13	1.59
LSD/sig	0.61	P≤0.01	P≤0.01
☑ suckers: number of aerial suckers			
Mean	1.22	0.50	0.78
Std. Deviation	0.74	0.61	0.62
LSD/sig	0.33	P≤0.01	P≤0.01
☐ Suckers: sucker length (cm)			
Mean	63.4	22.2	50.3
Std. Deviation	37.7	22.7	31.2
LSD/sig	23.01	P≤0.01	P≤0.01
☐ Reference leaf: maximum width (mm)			
Mean	56.5	57.7	50.9
Std. Deviation	4.12	3.30	2.40
LSD/sig	3.66	ns	P≤0.01
☑ Reference leaf: leaf area (cm^2)			
Mean	520.3	417.3	423.2
Std. Deviation	93.1	39.6	66.2
LSD/sig	74.40	P≤0.01	P≤0.01
☐ Leaves: number of leaves			
Mean	40.8	37.2	38.8
Std. Deviation	5.40	3.60	5.70
LSD/sig	5.5	ns	ns
☐ Reference leaf: leaf weight (g)			

Mean	98.0	64.5	90.7
Std. Deviation	13.9	7.30	11.20
LSD/sig	12.48	P≤0.01	ns
☐ Fruit: weight of fruit (g)			
Mean	1454	1515	1505
Std. Deviation	391.0	388.8	252.0
LSD/sig	184.5	ns	ns
☐ Fruit: diameter at the middle (mm)			
Mean	122.9	124.6	125.3
Std. Deviation	11.2	8.4	8.2
LSD/sig	5.07	ns	ns
☐ Fruit: height (without neck) (mm)			
Mean	159.9	160.1	162.0
Std. Deviation	18.9	18.5	10.7
LSD/sig	8.68	ns	ns
☐ Fruit: eye height at the middle (mm)			
Mean	29.5	27.7	28.3
Std. Deviation	1.6	2.0	2.0
LSD/sig	1.05	P≤0.01	P≤0.01
☐ Fruit: diameter of peduncle scar (mm)			
Mean	24.7	23.3	25.4
Std. Deviation	4.1	3.4	2.8
LSD/sig	1.89	ns	ns
☐ Crown: diameter of crown base (mm)			
Mean	22.7	21.2	27.4
Std. Deviation	2.03	2.78	2.46
LSD/sig	1.39	P≤0.01	P≤0.01
☑ Fruit/flesh: core diameter (mm)			
Mean	13.3	17.1	19.3
Std. Deviation	2.2	3.4	2.0
LSD/sig	1.37	P≤0.01	P≤0.01
☐ Fruit/flesh: juiciness (%)			
Mean	43.0	49.4	48.7
Std. Deviation	5.70	6.8	5.1
LSD/sig	3.56	P≤0.01	P≤0.01
☐ Fruit/flesh: firmness (kg/ 0.5cm^2)			
Mean	8.8	9.5	8.2
Std. Deviation	0.7	1.4	0.9
LSD/sig	0.58	P≤0.01	P≤0.01
☑ Reference leaf: length (mm)			
Mean	1274.7	945.0	1093.4
Std. Deviation	79.7	50.5	80.2
LSD/sig	85.5	P≤0.01	P≤0.01
☑ Fruit: eye number			
Mean	95.9	111.4	97.3
Std. Deviation	17.5	16.2	10.3

LSD/sig	8.24	P≤0.01	ns
Fruit: eye width at the middle (mm)			
Mean	24.9	23.8	25.6
Std. Deviation	1.9	1.5	1.60
LSD/sig	0.89	P≤0.01	ns
Crown: height (mm)			
Mean	290.4	186.2	321.0
Std. Deviation	30.4	40.1	42.0
LSD/sig	21.3	P≤0.01	P≤0.01
Crown: weight (g)			
Mean	301.2	110.0	285.1
Std. Deviation	64.2	36.5	57.6
LSD/sig	30.7	P≤0.01	P≤0.01
Fruit/flesh: sugar content (using refractometer) (%)			
Mean	19.2	15.4	16.5
Std. Deviation	1.4	2.1	1.6
LSD/sig	0.89	P≤0.01	P≤0.01

Prior Applications and Sales

Nil.

Description: **Garth Sanewski,** Maroochy Research Station, Nambour, QLD.

Details of Application

Application Number	2003/080
Variety Name	'Emeraldstar'
Genus Species	*Pittosporum tenuifolium*
Common Name	Pittosporum
Synonym	Nil
Accepted Date	15 May 2003
Applicant	Grant Farmer McKechnie
Agent	Greenhills Propagation Nursery Pty Ltd
Qualified Person	Mark Lunghusen

Details of Comparative Trial

Location	Greenhills Propagation Nursery, Tynong, VIC.
Descriptor	Pittosporum (*Pittosporum*) PBR PITT.
Period	Spring/summer 2007.
Conditions	Plants were grown in 14cm pots in full sun in commercial pine bark based potting mix with controlled release fertiliser. Plants were grown on benches with overhead watering.
Trial Design	10 plants in block design.
Measurements	Leaf measurements taken from largest leaves.
RHS Chart - edition	2005.

Origin and Breeding

Seedling selection: a short seedling was selected from a batch of seedlings of *Pittosporum tenuifolium* in 1990. The seed parent is characterised by tall plant height. Cuttings were taken from this seedling, established, and then another generation of cuttings were taken from the young plants. This was repeated to determine distinctness, uniformity and stability. To date, the plant has been grown through three generations with no off-types being recorded. Selection criteria: leaf colour. Propagation: vegetative. Breeder: Grant McKechnie, Albany New Zealand.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	height	very short

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Green Pillar'	Closest very short variety

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Tom Thumb'	Leaf	colour	green	purple
'Shorty'	Plant	height	very short	medium

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Emeraldstar'	'Green Pillar'
☐ Plant: type	shrub	shrub
☑ Plant: density	very dense	medium
☐ Plant: attitude of distal part of branches	semi erect	erect
☑ New shoot: colour of stem	reddish	greenish
☐ New shoot: main colour of leaves (RHS Colour Chart)	yellow green N144A	yellow green N144A
☐ New shoot: main colour of midrib on leaves	greenish	greenish
☐ Stem: colour (RHS Colour Chart)	200C	Brown 200C
☐ Petiole: length	short	short
☐ Leaf blade: shape	elliptic	oblong
☐ Leaf blade: shape of apex	acute	acute
☐ Leaf blade: shape of base	obtuse	obtuse
☑ Leaf blade: undulation of margin	medium to strong	very weak to weak
☐ Leaf blade: shape of margin	entire	entire
☐ Leaf blade: shape in cross section	concave	concave
☐ Leaf blade: curvature of longitudinal axis	weak	weak
☐ Leaf blade: number of colours on upper side	one	one
☑ Leaf blade: main colour on upper side (RHS Colour Chart)	green 146A	green 144A
☑ Leaf blade: main colour of lower side (RHS Colour Chart)	green 146B	yellow green 145A
☑ Leaf blade: glossiness	weak	medium
☐ Leaf blade: anthocyanin colouration	absent of very weak	absent of very weak
☐ Leaf blade: hairiness on lower side	absent or very weak	absent or very weak

Statistical Table

Organ/Plant Part: Context	'Emeraldstar'	'Green Pillar'
☐ Plant: width (mm)		
Mean	266.00	188.00
Std. Deviation	10.75	13.98
LSD/sig	87.95	ns
☐ Leaf: length (mm)		
Mean	21.68	28.73
Std. Deviation	1.54	2.92
LSD/sig	8.36	ns
☐ Leaf: length to width ratio (mm)		
Mean	1.85	1.74
Std. Deviation	0.70	0.12
LSD/sig	0.11	ns

☑ Plant: height (mm)		
Mean	199.00	353.00
Std. Deviation	11.97	29.74
LSD/sig	144.02	P≤0.01
☐ Leaf: width (mm)		
Mean	11.72	16.47
Std. Deviation	0.94	1.01
LSD/sig	5.08	ns

Prior Applications and Sales

Country	Year	Current Status	Name Applied
New Zealand	2004	Granted	'Emeraldstar'

First sold in Australia in March 2003 under the name McKechnie.

Description: **Mark Lunghusen,** World Select Plants, Cranbourne, VIC.

Details of Application

Application Number	2006/213
Variety Name	'Golf Ball'
Genus Species	*Pittosporum tenuifolium*
Common Name	Pittosporum
Synonym	Nil
Accepted Date	26 Oct 2006
Applicant	M & R Fyfe, Hastings, New Zealand
Agent	Greenhills Propagation Nursery Pty Ltd, Tynong, Vic
Qualified Person	Mark Lunghusen

Details of Comparative Trial

Location	Greenhills Propagation Nursery, Tynong, Vic
Descriptor	Pittosporum (*Pittosporum*) PBR PITT.
Period	Spring/summer 2007
Conditions	Plants were grown in 14cm pots in full sun in commercial pine bark based potting mix with controlled release fertiliser. Plants were grown on benches with overhead watering.
Trial Design	10 plants in block design.
Measurements	Leaf measurements taken from largest leaves.
RHS Chart - edition	2005.

Origin and Breeding

Seedling selection: a short seedling was selected from a batch of seedlings of *Pittosporum tenuifolium* in 1997. The seed parent is characterised by tall plant height. Cuttings were taken from this seedling, established, and then another generation of cuttings were taken from the young plants. This was repeated to determine distinctness, uniformity and stability. To date, the plant has been grown through three generations with no off-types being recorded. Selection criteria: leaf colour. Propagation: vegetative. Breeder: Mark Fyfe, Hastings, New Zealand.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	height	very short

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Emeraldstar'	Closest very short variety

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Tom Thumb'	Leaf	colour	green	purple
'Shorty'	Plant	height	very short	medium

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Golf Ball'	'Emeraldstar'
☐ Plant: type	shrub	shrub
☐ Plant: height	very short	very short
☐ Plant: density	sparse	very dense
☐ Plant: attitude of distal part of branches	semi erect	semi erect
☐ New shoot: colour of stem	reddish	reddish
☐ New shoot: main colour of leaves (RHS Colour Chart)	yellow green N144A	yellow green N144A
☐ New shoot: main colour of midrib on leaves	greenish	greenish
☐ Stem: colour (RHS Colour Chart)	Brown 200C	Brown 200C
☐ Petiole: length	short	short
☐ Leaf blade: shape	elliptic	elliptic
☐ Leaf blade: shape of apex	acute	acute
☐ Leaf blade: shape of base	obtuse	obtuse
☑ Leaf blade: undulation of margin	very weak	medium to strong
☐ Leaf blade: shape of margin	entire	entire
☐ Leaf blade: shape in cross section	flat	flat
☐ Leaf blade: curvature of longitudinal axis	weak	weak
☐ Leaf blade: number of colours on upper side	one	one
☐ Leaf blade: main colour on upper side (RHS Colour Chart)	green 144A	green 146A
☑ Leaf blade: main colour of lower side (RHS Colour Chart)	green 145A	green 146B
☑ Leaf blade: glossiness	medium	weak
☐ Leaf blade: anthocyanin colouration	absent of very weak	absent of very weak
☐ Leaf blade: hairiness on lower side	absent or very weak	absent or very weak

Statistical Table

Organ/Plant Part: Context	'Golf Ball'	'Emeraldstar'
☑ Leaf: length (mm)		
Mean	22.21	21.68
Std. Deviation	1.88	1.54
LSD/sig	0.45	P≤0.01
☐ Leaf: width (mm)		
Mean	11.74	11.72
Std. Deviation	1.16	0.94
LSD/sig	0.02	ns
☐ Leaf: length to width ratio (mm)		
Mean	1.89	1.85
Std. Deviation	0.13	0.07
LSD/sig	0.05	ns

☑ Plant: height (mm)		
Mean	223.00	199.00
Std. Deviation	14.94	11.97
LSD/sig	19.77	P≤0.01
☐ Plant: width (mm)		
Mean	231.00	266.00
Std. Deviation	15.95	10.75
LSD/sig	36.64	ns

Prior Applications and Sales

Country	Year	Current Status	Name Applied
New Zealand	2002	Granted	'Golf Ball'
EU	2005	Applied	'Golf Ball'
USA	2003	Granted	'Golf Ball'

First sold in New Zealand in Dec 2002 under the name 'Golf Ball'

Description: **Mark Lunghusen,** World Select Plants, Cranbourne, VIC.

Details of Application

Application Number	2006/087
Variety Name	'Whitepol'
Genus Species	*Polygala* x*dalmaisiana*
Common Name	Polygala
Synonym	Nil
Accepted Date	1 Aug 2006
Applicant	Chris Cristou, Werribee South, VIC
Agent	N/A
Qualified Person	Mark Lunghusen

Details of Comparative Trial

Location	Cranbourne, VIC
Descriptor	General Descriptor (PBR GEN DES).
Period	Winter to summer 2007.
Conditions	Plants were grown in 14cm pots in full sun in commercial pine bark based potting mix with controlled release fertiliser. Plants were grown on benches with overhead watering.
Trial Design	10 plants in block design.
Measurements	Leaf measurements taken from largest leaves.
RHS Chart - edition	1995.

Origin and Breeding

Seedling selection: a seedling was selected from a garden planting of *Polygala dalmasiana* 'Dazzler' in 2002. Cuttings were taken from this seedling, established, and then another generation of cuttings were taken from the young plants. This was repeated to determine distinctness, uniformity and stability. To date, the plant has been grown through three generations with no off-types being recorded. Selection criteria: leaf colour. propagation: vegetative. Breeder: Chris Christou, Werribee South, VIC.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	size	small

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Dazzler'	This is the closest variety based on plant size. Also the parent

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Whitepol'	'Dazzler'
☐ Plant: type	shrub	shrub
☐ Plant: growth habit	bushy	bushy
☐ Plant: size	small	small
☑ Plant: height	very short	short to medium
☐ Plant: width	narrow to medium	narrow to medium
☐ Plant: time of beginning of flowering	early	early
☐ Stem: degree of hairiness	low	low
☑ Stem: presence of anthocyanin in new growth	absent	present
☐ Leaf: size	small	small
☐ Leaf: attitude	semi-erect	semi-erect
☑ Leaf: arrangement	alternate	opposite
☐ Leaf: length of blade	short	short
☐ Leaf: width of blade	narrow	narrow
☐ Leaf: length of petiole	very short	very short
☐ Leaf: shape	oval	ovate
☑ Leaf: shape of apex	rounded	acute
☑ Leaf: shape of base	cuneate	obtuse
☐ Leaf: incision of margin	absent	absent
☑ Leaf: undulation of the margin	absent	medium
☐ Leaf: shape of cross-section	flat	flat
☐ Leaf: curvature of longitudinal axis	slightly recurved	straight
☐ Leaf: glossiness of upper side	very weak	very weak
☑ Leaf: green colour	very light	light
☐ Leaf: presence of variegation	absent	absent
☑ Leaf: primary colour (RHS colour chart)	144A	146B
☐ Flower: type	single	single
☐ Flower: diameter	medium	medium
☐ Flower: fragrance	absent	absent
☐ Flower: pedicel length	medium	medium
☑ Petal: predominant colour of upper side (RHS colour chart)	155A	N80A
☑ Petal: predominant colour of lower side (RHS colour chart)	N144A	N81B
☐ Petal: eye zone (basal spot upper side)	absent	absent
☐ Petal: reflexing of margin	absent or very weak	absent or very weak
☐ Petal: incision	absent or very weak	absent or very weak

Petal: undulation	absent or very weak	absent or very weak
Petal: shape	cordate	cordate

Prior Applications and Sales

Nil.

Description: **Mark Lunghusen,** World Select Plants, Cranbourne, VIC.

Details of Application

Application Number	2003/339
Variety Name	'Cardinal'
Genus Species	*Rubus idaeus*
Common Name	Raspberry
Synonym	Nil
Accepted Date	5 Mar 2004
Applicant	Driscoll Strawberry Associates, Inc, Watsonville, CA, USA
Agent	Phillips Ormonde & Fitzpatrick, Melbourne, VIC
Qualified Person	Margaret Zorin

Details of Comparative Trial

Overseas Testing Authority	US Patent and Trademark Office (USPTO)
Overseas Data Reference Number	PP14,903
Location	Watsonville, Monterey County, California, USA. Verified at Stanthorpe QLD, Australia.
Descriptor	Raspberry (*Rubus idaeus* L.) TG/43/7.
Period	1995-2003,
Conditions	Traditional cultural practices employ rooted cuttings planted into raised ridges of soil in winter, the plants are then trellised and primocaane harvest commences approximately 7 months later in summer and autumn. At the end of the primocane fruit harvest the plants are pruned and the floricane harvest commences in spring. Test plots for verification were planted in Sep 2006 at Stanthorpe, QLD and verified in 2007.
Trial Design	Comparative trial was conducted in open fields in full sunlight in Watsonville, California in 2001 and 2002. Plants were evaluated as both primocanes and floricanes. Rooted cuttings were planted in rows adjacent to test varieties including 'Heritage', an unpatented variety grown worldwide. All plants were subjected to standard growing conditions typical of commercial raspberry production in southern California.
Measurements	Measurements of plant, flower and fruit characteristics were made approximately seven months after planting for primocane production and approximately seventeen months after planting for floricane production. All measurements were made in accordance with the UPOV Technical Guidelines and colours are described and the most similar colour designations are provided from the Royal Horticultural Society (RHS) Colour Chart.
RHS Chart - edition	2001.

Origin and Breeding

Controlled pollination: the new variety of raspberry was developed from the hybridisation of the selection 'M48.9' (an unpatented variety) as the seed parent with the selection 'Gloria' (US Plant Patent PP11,067) as the pollen parent. The parents were crossed in 1994 and seedling selection was made in 1995 in a field planting at Carpenteria, California, USA. The new variety 'Cardinal' was asexually propagated by in vitro shoot tip culture and root sucker division and root cuttings and has been shown to maintain the desired characteristics after several generations. Plant breeders: Carlos D Fear (Aptos, CA, USA), Richard E Harrison (Aptos, CA, USA), Fred M Cook (Aptos, CA, USA) and Gavin Sills (Watsonville, CA, USA), all employees of Driscoll Strawberry Associates Inc Watsonville, CA, USA.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	habit	erect
Spines	presence	absent
Leaf	green colour of upper side	dark green 147A
Very young shoot	anthocyanin colouration of apex during rapid growth	present
Fruit	general shape in lateral view	circular
Fruit	colour	medium red
Fruit	main bearing type	both previous year's cone in summer & current year's cone in autumn
Fruit	adherence to plug	medium
Time of	beginning of flowering on current seasons cane	early to medium

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Heritage'	'Heritage' is an unpatented variety grown Worldwide and used as standard comparator
'Dulcita'	'Dulcita'US Plant patent PP14,904

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Cardinal'	'Dulcita'	'Heritage'
☐ Plant: habit	upright	upright	upright
☑ *Plant: number of current season's canes	many	medium	medium
☐ *Very young shoot: anthocyanin colouration of apex during rapid growth	present	present	present
☑ *Very young shoot: intensity of anthocyanin colouration of apex during rapid growth	very weak	medium	medium
☑ Current season's cane: bloom	absent or very weak	weak	weak
☑ Current season's cane: anthocyanin colouration	absent or very weak		medium
☑ *Current season's cane: length (varieties which fruit on current season's cane in autumn)	medium to long		short to medium
☑ *Dormant cane: colour (varieties which fruit on previous season's cane in summer)	purplish brown	brown	brownish purple
☐ *Spines: presence	absent	absent	absent
☐ *Leaf: green colour of upper side	dark	dark	dark
☑ *Leaf: predominant number of leaflets	equally three and five	five	equally three and five
☐ Leaf: profile of leaflets in cross section	straight		
☑ *Leaf: rugosity	very weak	medium	medium
☑ Leaf: relative position of lateral leaflets	touching	overlapping	free
☑ Terminal leaflet: length	medium	medium to long	long
☑ Terminal leaflet: width	narrow	medium	narrow to medium
☑ Flower: size	small to medium	large	small to medium
☑ Fruiting lateral: attitude (varieties which fruit on previous year's cane in summer)	semi-erect	erect	horizontal to drooping

☑ *Fruiting lateral: length (varieties which fruit on previous year's cane in summer)	long to very long	long	short
☑ *Fruit: length	medium to long	long	short to medium
☑ *Fruit: width	medium to broad	broad	narrow to medium
☑ *Fruit: ratio length/width	medium	small	small to medium
☐ *Fruit: general shape in lateral view	circular	circular	circular
☑ Fruit: size of single drupe	large	large	small
☐ *Fruit: colour	medium red	medium red	medium red
☑ Fruit: glossiness	weak	weak	medium
☐ *Fruit: firmness	firm	medium to firm	firm
☐ Fruit: adherence to plug	medium	medium	medium
☐ *Fruit: main bearing type	both previous year's cone in summer & current year's cone in autumn	both previous year's cone in summer & current year's cone in autumn	both previous year's cone in summer & current year's cone in autumn
☑ *Plant: time of vegetative bud burst (varieties which fruit on previous year's cane in summer)	early to medium	medium	medium to late
☑ *Time of: cane emergence (varieties which fruit on current year's cane in autumn)	very early to early	early	medium to late
☐ *Time of: beginning of flowering on previous year's cane (varieties which fruit on previous year's cane in summer)	medium to late	medium	medium
☐ *Time of: beginning of flowering on current season's cane (varieties which fruit on current year's cane in autumn)	early to medium	early to medium	early to medium
☑ *Time of: beginning of fruit ripening on previous year's cane (varieties which fruit of previous year's cane in summer)	early	early to medium	medium
☐ *Time of: beginning of fruit ripening on current year's cane (varieties which fruit on current year's cane in autumn)	early	early	early to medium
☑ Length of: fruiting period on previous year's cane (varieties which fruit on previous year's cane in summer)	medium to long	short to medium	short to medium
☑ Length of: fruiting period on current year's cane (varieties which fruit on current year's cane in autumn)	long	medium to long	long to very long

Prior Applications and Sales

Country	Year	Current Status	Name Applied
Canada	2006	Applied	'Cardinal'
EU	2003	Applied	'Driscoll Cardinal'
USA	2002	Granted	'Driscoll Cardinal'

Prior sale nil.

Description: **Margaret Zorin,** 167 Collingwood Road, Birkdale, Qld 4159.

Details of Application

Application Number	2003/338
Variety Name	'Maravilla'
Genus Species	*Rubus idaeus*
Common Name	Raspberry
Synonym	Nil
Accepted Date	5 Mar 2004
Applicant	Driscoll Strawberry Associates, Inc, Watsonville, CA, USA
Agent	Phillips Ormonde & Fitzpatrick, Melbourne, VIC
Qualified Person	Margaret Zorin

Details of Comparative Trial

Overseas Testing Authority	US Patent and Trademark Office (USPTO)
Overseas Data Reference Number	PP14,804
Location	Watsonville, Monterey County, California USA Verified at Stanthorpe, Qld, Australia
Descriptor	Raspberry (*Rubus idaeus* L.) TG/43/7
Period	1998-2002
Conditions	Traditional cultural practices are employed where rooted cuttings are planted into raised ridges of soil in winter, the plants are then trellised and primocane harvest commences 7-8 months later in summer and autumn. At the end of the primocane harvest plants are pruned and the floricane harvest commences in early spring. Test plots for verification were planted in September 2006 at Stanthorpe, QLD and verified 2007.
Trial Design	Comparative trial was conducted in open fields in full sunlight and evaluated as both primocanes and floricanes in Watsonville, California between 2001 and 2002. Seedlings of 'Maravilla' were planted in rows and compared with the unpatented variety 'Heritage' and the nearest other available variety 'Francesca'. All plants were subject to standard growing conditions typical of commercial raspberry production in southern California USA.
Measurements	Measurements of plant, flower and fruit characteristics were made approximateley 7 months after planting for primocane production and seventeen months after planting for floricane production. All measurements were made in accordance with the UPOV Technical Guidelines and colours are described and most similar colour designations are provided from the Royal Horticultural Society (RHS) Colour Charts.
RHS Chart - edition	2001

Origin and Breeding

Controlled pollination: the new variety of raspberry plant 'Maravilla' was developed from the hybridisation of the selection 'Q491.1' (an unpatented variety) as the seed parent with the selection 'Q480.3' (an unpatented variety) as the pollen parent. Seedlings from this cross were planted in 1996 and final selection was made in 1998. The new variety 'Maravilla' has been asexually propagated by in vitro shoot tip culture, root sucker division and root cuttings and has been shown to maintain the desired distinguishing characteristics after propagation over several generations. Breeder: Carlos D Fear (Aptos, CA, USA), Richard E Harrison (Aptos, CA, USA), Fred M Cook (Aptos, CA, USA) and Gavin Sills (Watsonville, CA, USA), all employees of Driscoll Strawberry Associates Inc Watsonville, CA, USA.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	spines	absent
Leaf	colour of upper side	dark green
Plant	dormant cane colour	brownish purple
Fruit	colour	medium red
Fruit	adherence to plug	medium
Fruit	main bearing type	both
Plant	very young shoot anthocyanin colour at apex	present

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Heritage'	Unpatented variety in most common use throughout the World.
'Francesca'	US PP14,860 closest available commercial variety grown in California USA.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Maravilla'	'Francesca'	'Heritage'
☑ Plant: habit	semi-upright	semi-upright	upright
☐ *Plant: number of current season's canes	medium	medium	medium
☐ *Very young shoot: anthocyanin colouration of apex during rapid growth	present	present	present
☑ *Very young shoot: intensity of anthocyanin colouration of apex during rapid growth	medium	very weak to weak	medium
☑ Current season's cane: bloom	weak	strong	weak
☑ Current season's cane: anthocyanin colouration	weak	medium	medium
☑ Current season's cane: length of internode	long	short to medium	
☐ *Dormant cane: colour (varieties which fruit on previous season's cane in summer)	brownish purple	brownish purple	brownish purple
☐ *Spines: presence	absent	absent	absent
☐ *Leaf: green colour of upper side	dark	dark	dark
☑ *Leaf: predominant number of leaflets	five	equally three and five	equally three and five
☑ *Leaf: rugosity	medium	weak	medium
☑ Leaf: relative position of lateral leaflets	overlapping	free	free
☑ Terminal leaflet: length	short to medium	medium	long
☑ Terminal leaflet: width	medium to broad	medium to broad	narrow to medium
☑ Flower: size	small	medium	small to medium
☑ Fruiting lateral: attitude (varieties which fruit on previous year's cane in summer)	semi-erect	semi-erect	horizontal to drooping
☐ *Fruiting lateral: length (varieties which fruit on previous year's cane in summer)	long to very long	long	
☑ *Fruit: length	long	long	short to medium
☑ *Fruit: width	broad to very	medium to broad	narrow to medium

	broad		
☐ *Fruit: ratio length/width	small to medium	medium	small to medium
☑ *Fruit: general shape in lateral view	broad conical	broad conical	circular
☑ Fruit: size of single drupe	large	medium to large	small
☐ *Fruit: colour	medium red	medium red	medium red
☑ Fruit: glossiness	medium	weak	medium
☑ *Fruit: firmness	firm	medium	firm
☐ Fruit: adherence to plug	medium	medium	medium
☐ *Fruit: main bearing type	both previous year's cone in summer & current year's cone in autumn	both previous year's cone in summer & current year's cone in autumn	both previous year's cone in summer & current year's cone in autumn
☑ *Plant: time of vegetative bud burst (varieties which fruit on previous year's cane in summer)	early	early	medium to late
☑ *Time of: cane emergence (varieties which fruit on current year's cane in autumn)	early	early	medium to late
☐ *Time of: beginning of flowering on previous year's cane (varieties which fruit on previous year's cane in summer)	medium to late	medium	medium
☐ *Time of: beginning of flowering on current season's cane (varieties which fruit on current year's cane in autumn)	early to medium	early	early to medium
☐ *Time of: beginning of fruit ripening on previous year's cane (varieties which fruit of previous year's cane in summer)	medium to late	medium to late	medium
☑ *Time of: beginning of fruit ripening on current year's cane (varieties which fruit on current year's cane in autumn)	medium to late	early	early to medium
☑ Length of: fruiting period on previous year's cane (varieties which fruit on previous year's cane in summer)	long	medium	medium
☑ Length of: fruiting period on current year's cane (varieties which fruit on current year's cane in autumn)	long	short to medium	long to very long

Prior Applications and Sales

Country	Year	Current Status	Name Applied
Canada	2006	Applied	'Maravilla'
Chile	2006	Granted	'Driscoll Maravilla'
EU	2003	Granted	'Driscoll Maravilla'
USA	2002	Granted	'Driscoll Maravilla'
South Africa	2003	Applied	'Maravilla'

Prior sale nil.

Description: **Margaret Zorin,** 167 Collingwood Road, Birkdale, Qld 4159.

Details of Application

Application Number	2006/115
Variety Name	'Grandtang'
Genus Species	*Rosa* hybrid
Common Name	Rose
Synonym	Nil
Accepted Date	30 May 2006
Applicant	Mr H Schreuders
Agent	Grandiflora Nurseries Pty Ltd, Skye, VIC
Qualified Person	Christopher Prescott

Details of Comparative Trial

Location	145 Moores Road, Clyde, VIC (Latitude 38º09' South, elevation 16m).
Descriptor	Rose (new) TG/11/8.
Period	2007.
Conditions	Trial conducted in a controlled environment polyhouse with shade, temperature ranged between 15 and 36 degrees Celsius within the 6 weeks prior to examination (1 growth cycle) with plants on their own roots planted into 330mm (3 plants per pot) and in an open polyhouse without shade, temperature ranged between 12 and 38 degrees Celsius within the 6 weeks prior to examination (1 growth cycle) with plants on their own roots planted into 210mm (1 plant per pot) pots filled with co-co coir, nutrition was maintained as part of a commercial hydroponic system, pest and disease treatments applied as required.
Trial Design	160 plants of 'Grandtang' on benches two plants deep, arranged in rows as part of commercial flower growing operation and 6 plants of 'Tan01693' on benches two or three plants deep, arranged in blocks within the centralised testing centre for roses.
Measurements	From 6 plants at random. One sample per plant stem.
RHS Chart - edition	1995.

Origin and Breeding

Controlled pollination: 'Grandtang' was a resultant seedling from a cross between an unnamed seedling 'S11' (seed parent) from the breeding program of Harry Schreuders at his property in Skye, VIC, and 'Korblekaf' (pollen parent), between Aug and Nov 2002. The initial selection took place in Sep 2003 and went through subsequent selections in 2004, 2005 and finally in Jan 2006. All work was conducted by or under the supervision of Mr Harry Schreuders, Managing Director of Grandiflora Nurseries Pty Ltd.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Flower	colour group	orange blend
Plant	growth type	bed

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Tan01693'	

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Korblekaf'	Flower petal number	many	medium

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Grandtang'	'Tan01693'
☐ *Plant: growth type	bed	bed
☐ *Plant: growth habit (excluding varieties with growth type climber)	upright	upright
☐ Plant: height	tall to very tall	tall
☐ Young shoot: anthocyanin colouration	present	present
☑ Young shoot: intensity of anthocyanin colouration	medium	strong
☐ Stem: number of prickles	medium	medium
☐ Prickles: predominant colour	reddish	reddish
☐ Leaf: size	large	large to very large
☑ Leaf: intensity of green colour	medium	dark to very dark
☐ Leaf: anthocyanin colouration	present	present
☑ *Leaf: glossiness of upper side	weak	medium to strong
☐ *Leaflet: undulation of margin	weak	weak
☐ *Terminal leaflet: shape of blade	ovate	ovate
☑ Terminal leaflet: shape of base of blade	obtuse	rounded
☐ Terminal leaflet: shape of apex of blade	obtuse	obtuse
☐ Flowering shoot: flowering laterals	present	present
☑ Flowering shoot: number of flowering laterals	very few	few
☐ Flowering shoot: number of flowers per lateral (varieties with flowering laterals only)	very few	very few
☐ Flower bud: shape in longitudinal section	broad ovate	broad ovate
☐ *Flower: type	double	double
☑ *Flower: number of petals	many	many to very many
☐ *Flower: colour group	orange blend	orange blend
☐ Flower: colour of the centre	orange	orange
☑ Flower: density of petals	medium	dense
☐ *Flower: diameter	large to very large	large to very large
☐ *Flower: shape	irregularly rounded	irregularly rounded
☐ Flower: profile of upper part	flattened convex	flattened convex
☐ *Flower: profile of lower part	flat	flat

☑ Flower: fragrance	medium	absent or weak
☐ *Sepal: extensions	weak to medium	weak
☐ Petals: reflexing of petals one-by-one	present	present
☐ *Petal: shape	obcordate	obcordate
☑ Petal: incisions	absent or very weak	medium to strong
☐ Petal: reflexing of margin	medium to strong	medium to strong
☐ Petal: undulation	absent or very weak	absent or very weak
☐ *Petal: size	large	medium to large
☐ *Petal: length	medium	medium
☐ *Petal: width	medium to broad	medium to broad
☐ *Petal: number of colours on inner side	one	one
☑ *Petal: intensity of colour	lighter towards the top	even
☑ *Petal: main colour on the inner side (RHS Colour Chart)	23A	20A
☐ *Petal: basal spot on the inner side	present	present
☑ *Petal: size of basal spot on inner side	medium	small
☐ *Petal: colour of basal spot on inner side	medium yellow	medium yellow
☐ *Petal: main colour on the outer side (RHS Colour Chart)	31A	31A
☐ Outer stamen: predominant colour of filament	medium yellow	medium yellow
☑ Seed vessel: size	small	medium
☑ Hip: shape in longitudinal section	funnel-shaped	pitcher-shaped

Statistical Table

Organ/Plant Part: Context	**‘Grandtang’**	**‘Tan01693’**
☑ Flower: number of petals		
Mean	58.40	81.20
Std. Deviation	2.07	12.83
LSD/sig	21.78	P≤0.01

Prior Applications and Sales

Nil.

Description: **Christopher Prescott,** Prescott Roses Pty Ltd, Clyde, VIC**.**

Details of Application

Application Number	2004/012
Variety Name	'Kribigpea'
Genus Species	*Rosa* hybrid
Common Name	Rose
Synonym	Nil
Accepted Date	3 Mar 2004
Applicant	Lux Riviera S.r.l., Ventimiglia, Italy.
Agent	Grandiflora Nurseries Pty Ltd, Skye, VIC
Qualified Person	Christopher Prescott

Details of Comparative Trial

Location	145 Moores Road, Clyde, VIC (Latitude 38°09′ South, elevation 16m).
Descriptor	Rose (New) (*Rosa*) TG/11/8.
Period	2005-2007.
Conditions	Trial conducted in an open polyhouse without shade, temperature ranged between 12 and 38 degrees Celsius within the 6 weeks prior to examination (1 growth cycle) with plants on their own roots planted into 210mm (1 plant per pot) pots filled with co-co coir, nutrition was maintained as part of a commercial hydroponic system, pest and disease treatments applied as required.
Trial Design	8 plants of 'Kribigpea' and 8 plants of the comparator 'Pannaran' on benches two plants deep, arranged in blocks within the centralised testing centre for roses.
Measurements	From 6 plants at random. One sample per plant stem.
RHS Chart - edition	1995.

Origin and Breeding

Controlled pollination: 'Kribigpea' was the resultant seedling from a cross between 'Korlimlt' (seed parent and 'Rouge Antibes' (pollen parent) in 1992. The variety had gone through the next four years of subsequent selection with each year a new generation was propagated from the previous generation but in greater numbers. In 1996 'Kribigpea' was finally selected as having and maintaining the characteristics deemed necessary to fulfil the requirements of a commercial cut rose variety and was then assigned to Lux Riviera srl. Breeder: Madame Michel Kriloff, Antibes, France.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Flower	colour	orange blend
Plant	growth type	bed

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Pannaran'	

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Krivagold'	flower colour	orange blend	yellow blend

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Kribigpea'	'Pannaran'
☐ *Plant: growth type	bed	bed
☐ *Plant: growth habit (excluding varieties with growth type climber)	upright	upright
☐ Plant: height	tall	medium to tall
☐ Young shoot: anthocyanin colouration	present	present
☑ Young shoot: intensity of anthocyanin colouration	medium	weak
☑ Stem: number of prickles	very few to few	medium to many
☐ Prickles: predominant colour	reddish	reddish
☐ Leaf: size	large to very large	large to very large
☐ Leaf: intensity of green colour	medium	medium
☐ Leaf: anthocyanin colouration	present	present
☐ *Leaf: glossiness of upper side	weak to medium	weak to medium
☑ *Leaflet: undulation of margin	weak	medium
☐ *Terminal leaflet: shape of blade	ovate	ovate
☐ Terminal leaflet: shape of base of blade	rounded	rounded
☑ Terminal leaflet: shape of apex of blade	obtuse	acute
☐ Flowering shoot: flowering laterals	present	present
☐ Flowering shoot: number of flowering laterals	few to medium	medium
☐ Flowering shoot: number of flowers per lateral (varieties with flowering laterals only)	few to medium	few
☑ Flower bud: shape in longitudinal section	medium ovate	broad ovate
☐ *Flower: type	double	double
☐ *Flower: number of petals	medium	medium to many
☐ *Flower: colour group	orange blend	orange blend
☐ Flower: colour of the centre	orange	orange
☐ Flower: density of petals	loose	loose to medium
☐ *Flower: diameter	large	large
☐ *Flower: shape	irregularly rounded	irregularly rounded
☐ Flower: profile of upper part	flattened convex	flattened convex
☑ *Flower: profile of lower part	flattened convex	flat
☑ Flower: fragrance	absent or weak	medium
☐ *Sepal: extensions	medium to strong	medium

Petals: reflexing of petals one-by-one	present	present
*Petal: shape	obcordate	rounded
Petal: incisions	absent or very weak	absent or very weak
Petal: reflexing of margin	weak to medium	weak to medium
Petal: undulation	weak to medium	weak
*Petal: size	medium to large	medium
*Petal: length	medium to long	medium
*Petal: width	medium to broad	medium
*Petal: number of colours on inner side	two	two
*Petal: intensity of colour	lighter towards the base	even
*Petal: main colour on the inner side (RHS Colour Chart)	35C	35B
*Petal: secondary colour (varieties with two or more colours on inner side of petal only) (RHS Colour Chart)	11A	31B
*Petal: distribution of secondary colour on inner side (varieties with two or more colours on inner side of petal)	at base	at apex
*Petal: basal spot on the inner side	present	present
*Petal: size of basal spot on inner side	very small	small
*Petal: colour of basal spot on inner side	medium yellow	medium yellow
*Petal: main colour on the outer side (RHS Colour Chart)	4D	29C
Outer stamen: predominant colour of filament	medium yellow	orange
Seed vessel: size	medium	small to medium
Hip: shape in longitudinal section	pitcher-shaped	pitcher-shaped

Prior Applications and Sales

Country	Year	Current Status	Name Applied
Colombia	2004	Applied	‘Kribigpea’
South Korea	2002	Granted	‘Kribigpea’

First sold in Italy in Aug 2002.

Description: **Christopher Prescott,** Prescott Roses Pty Ltd, Clyde, VIC**.**

Details of Application

Application Number	2007/059
Variety Name	'Heatwave Blaze'
Genus Species	*Salvia* hybrid
Common Name	Sage
Synonym	Nil
Accepted Date	9 Mar 2007
Applicant	Plant Growers Australia Pty. Ltd., Wonga Park, VIC
Agent	Plants Management Australia Pty. Ltd., Wonga Park, VIC
Qualified Person	Steve Eggleton

Details of Comparative Trial

Location	Wonga Park, VIC
Descriptor	Salvia (New) (*Salvia*) PBR SALV 2
Period	Jul 2007 to Nov 2007
Conditions	Trial conducted in the open, plants propagated from cuttings during Jul 2007, transferred from plugs to 140mm pots in Sep 2007. Pots filled with soilless, pinebark based mix with controlled release fertilizers. Appropriate pest and disease treatments were applied as required.
Trial Design	Twelve pots of each variety in a completely randomised design.
Measurements	From ten plants randomly selected.
RHS Chart - edition	1995

Origin and Breeding

Controlled pollination: occurred between Jan and Mar 2003 at Wonga Park, VIC, Australia. This was part of a breeding program designed to hybridize forms of *Salvia greggii* with *Salvia microphylla* with the aim of producing plants being more robust as garden specimens and in a range of flower colours. An assorted range of *greggii* forms were all pollinated with *S. microphylla* 'San Carlos Festival'. This seed was collected, sown and raised. When the seedlings reached flowering maturity a selection was made on the basis of plant habit, medium and petal colour red-purple. The selection was made over a period of months from Oct 2003. From this selection cuttings were taken and further plants grown to maturity. During 2005 further plants were grown in a small production trial and once selection was approved for commercialisation these were used as mother stock. Propagation: will continue to be cuttings. Breeder: Plant Growers Australia Pty. Ltd., Wonga Park, VIC

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	density	medium
Corolla	predominant colour of lower lip	red or red-purple
Leaf	shape of apex	acute
Leaf	presence of variegation	absent
Leaf	incision of margin	present

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Heatwave Sizzle'	from same parentage
'San Carlos Festival'	parent

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
S. greggii red	leaf	incision of margin	present	absent
S. greggii red	inflorescence	number of flowers per node	1,2 or more	1 or 2 only

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Heatwave Blaze'	'Heatwave Sizzle'	'San Carlos Festival'
☐ *Plant: growth habit	bushy to spreading	bushy	bushy
☐ *Plant: density	medium	medium	medium
☑ Stem: anthocyanin colouration	medium to strong	very weak to weak	weak
☐ Leaf: shape	ovate	ovate	ovate
☐ Leaf: shape of apex	acute	acute	acute
☑ Leaf: shape of base	cuneate	cuneate	obtuse
☐ Leaf: incision of margin	present	present	present
☑ Leaf: depth of incision	shallow to medium	shallow to medium	medium to deep
☐ Leaf: type of incision	crenate	crenate	crenate
☑ Leaf: undulation of the margin	weak	weak	medium
☑ Leaf: prominence of venation	medium	medium	strong
☑ Leaf: glossiness of upper side	medium	medium	weak
☐ Leaf: presence of variegation	absent	absent	absent
☐ Leaf: predominant colour of upper side (RHS colour chart)	yellow-green 146B	yellow-green 146B	yellow-green 146B
☐ Inflorescence: number of flowers per node	1, 2 or more	1, 2 or more	1, 2 or more
☑ Caylx: anthocyanin colouration	strong to very strong	strong	weak to medium
☑ Corolla: predominant colour of lower lip (RHS colour chart)	red-purple 61A	red - purple 57A	red - purple 66A

Prior Applications and Sales

First sold in Australia in Mar 2006 under the name 'Heatwave Blaze'.

Description: **Steve Eggleton,** Plant Growers Australia Pty. Ltd., Wonga Park, VIC.

Details of Application

Application Number	2007/060
Variety Name	'Heatwave Sizzle'
Genus Species	*Salvia* hybrid
Common Name	Sage
Synonym	
Accepted Date	21 Mar 2007
Applicant	Plant Growers Australia Pty. Ltd., Wonga Park, VIC
Agent	Plants Management Australia Pty. Ltd., Wonga Park, VIC
Qualified Person	Steve Eggleton

Details of Comparative Trial

Location	Wonga Park VIC.
Descriptor	Salvia (new) (*Salvia*) PBR SALV 2
Period	Jul 07 to Nov 07.
Conditions	Trial conducted in the open, plants propagated from cuttings during Jul 07, transferred from plugs to 140mm pots in Sep 2007. Pots filled with soilless, pinebark based mix with controlled release fertilizers. Appropriate pest and disease treatments were applied as required.
Trial Design	Twelve pots of each variety in a completely randomised design.
Measurements	From ten plants randomly selected.
RHS Chart - edition	1995.

Origin and Breeding

Controlled pollination: occurred between Jan and Mar 2003 at Wonga Park, VIC, Australia. This was part of a breeding program designed to hybridize forms of *Salvia greggii* with *Salvia microphylla* with the aim of producing plants being more robust as garden specimens and in a range of flower colours. An assorted range of *S. greggii* forms were all pollinated with *S. microphylla* 'San Carlos Festival'. This seed was collected, sown and raised. When the seedlings reached flowering maturity a selection was made on the basis of plant habit, medium and petal colour red-purple. The selection was made over a period of months from Oct 2003. From this selection cuttings were taken and further plants grown to maturity. During 2005 further plants were grown in a small production trial and once selection was approved for commercialisation these were used as mother stock. Propagation: will continue to be cuttings. Breeder: Plant Growers Australia Pty. Ltd., Wonga Park, VIC.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	density	medium
Leaf	shape of apex	acute
Leaf	presence of variegation	absent
Leaf	incision of margin	present

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Heatwave Blaze'	from same parentage
'San Carlos Festival'	parent

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
S. greggii red	leaf	incision of margin	present	absent
S. greggii red	inflorescence	number of flowers per node	1, 2 or more	1 or 2 only

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Heatwave Sizzle'	'Heatwave Blaze'	'San Carlos Festival'
☐ *Plant: growth habit	bushy	bushy to spreading	bushy
☐ *Plant: density	medium	medium	medium
☑ Stem: anthocyanin colouration	very weak to weak	medium to strong	weak
☐ Leaf: shape	ovate	ovate	ovate
☐ Leaf: shape of apex	acute	acute	acute
☑ Leaf: shape of base	cuneate	cuneate	obtuse
☐ Leaf: incision of margin	present	present	present
☑ Leaf: depth of incision	shallow to medium	shallow to medium	medium to deep
☐ Leaf: type of incision	crenate	crenate	crenate
☑ Leaf: undulation of the margin	weak	weak	medium
☑ Leaf: prominence of venation	medium	medium	strong
☑ Leaf: glossiness of upper side	medium	medium	weak
☐ Leaf: presence of variegation	absent	absent	absent
☐ Leaf: predominant colour of upper side (RHS colour chart)	yellow-green 146B	yellow-green 146B	yellow -green 146B
☐ Inflorescence: number of flowers per node	1, 2 or more	1, 2 or more	1, 2 or more
☑ Caylx: anthocyanin colouration	strong	strong to very strong	weak to medium
☑ Corolla: predominant colour of lower lip (RHS colour chart)	red - purple 57A	red - purple 61A	red - purple 66A

Prior Applications and Sales

First sold in Australia in Mar 2006 under the name 'Heatwave Sizzle'.

Description: **Steve Eggleton,** Plant Growers Australia Pty. Ltd., Wonga Park, VIC.

Details of Application

Application Number	2006/297
Variety Name	'Cherry Surprise'
Genus Species	*Syzygium smithii*
Common Name	Small Leaf Lilly Pilly
Synonym	Nil
Accepted Date	16 Mar 2007
Applicant	Wirreanda Nursery, Ingleside, NSW
Agent	N/A
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Ingleside, NSW.
Descriptor	Lilly Pilly (*Acmena smithii*/*Syzygium* sp) PBR LILL
Period	Summer 2007 to late autumn 2007.
Conditions	Trial conducted in open beds, plants originally propagated by cuttings, potted to 200mm containers filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random.
RHS Chart - edition	1995

Origin and Breeding

Seedling selection: *Syzygium smithii*. The parent is characterised by a medium intensity of colour of new growth flush and medium plant width. Selection took place in Ingleside, NSW. Selection criteria: colour of new growth, shape of plant and strong growth vigour. Propagation: vegetative cuttings were found to be uniform and stable. Breeders: Mark Cruickshank and Bill Douglass, Ingleside, NSW.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Immature leaf	colour	greyed red to greyed purple
Leaf	variegation	absent

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Hot Flush'	
'Allyn Magic'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Cherry Surprise'	'Allyn Magic'	'Hot Flush'
☐ Plant: growth habit	upright	upright	upright
☑ Plant: branch density	dense to very dense	very dense	medium to dense
☐ Stem: branch angle	acute	acute	acute
☑ Stem: internode length	medium	short	medium
☑ Stem: colour of new growth (RHS colour chart)	187B	183A	185A
☑ Leaf: blade length	long	medium to long	medium
☑ Leaf: blade width	medium	broad	medium
☐ Leaf: petiole length	short	short	short
☑ Leaf: shape of blade	narrow elliptic	broad elliptic	elliptic
☐ Leaf: shape of apex	cuspidate	cuspidate	cuspidate
☐ Leaf: shape of base	acute	acute	acute
☐ Leaf: glossiness	medium	medium	medium
☐ Leaf: shape of cross section	flat	flat to concave	flat to concave
☑ Leaf: shape of longitudinal section	flat	flat to concave	convex to flat
☐ Leaf: stiffness	medium to strong	medium	medium
☐ Mature leaf: primary colour of upper side (RHS colour chart)	147A	147A	147A
☐ Mature leaf: primary colour of lower side (RHS colour chart)	146B	146B	146B
☑ Partly mature leaf: primary colour of upper side (RHS colour chart)	ca 177A	176A	175A
☑ Partly mature leaf: primary colour of lower side (RHS colour chart)	ca 165A	177B	ca 177B
☑ Newly emerged: upper side (RHS colour chart)	187B	183A	178B
☐ Leaf: variegation	absent	absent	absent

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Cherry Surprise'	'Allyn Magic'	'Hot Flush'
☑ Newly emerged leaf: size	medium	small	medium

Prior Applications and Sales

Nil.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW.

Details of Application

Application Number	2006/298
Variety Name	'Sunrise'
Genus Species	*Syzygium smithii*
Common Name	Small Leaf Lilly Pilly
Synonym	NIl
Accepted Date	16 Mar 2007
Applicant	Wirreanda Nursery, Ingleside, NSW
Agent	N/A
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Ingleside, NSW.
Descriptor	Lilly Pilly (*Acmena smithii*/*Syzygium* sp) PBR LILL.
Period	Summer 2007 to late autumn 2007.
Conditions	Trial conducted in open beds, plants originally propagated by cuttings, potted to 200mm containers filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random.
RHS Chart - edition	1995.

Origin and Breeding

Seedling selection: *Syzygium smithii*. The parent is characterised by a medium intensity of colour of new growth flush and medium plant width. Selection took place in Ingleside, NSW. Selection criteria: colour of new growth, shape of plant and strong growth vigour. Propagation: vegetative cuttings were found to be uniform and stable. Breeders: Mark Cruickshank and Bill Douglass, Ingleside, NSW.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Immature leaf	colour	greyed red to greyed purple
Leaf	variegation	absent

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Cherry Surprise'	new variety from the same breeding program
'Hot Flush'	
'Allyn Magic'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Sunrise'	'Cherry Surprise'	'Allyn Magic'	'Hot Flush
☐ Plant: growth habit	upright	upright	upright	upright
☑ Plant: branch density	dense	dense to very dense	very dense	medium to dense
☐ Stem: branch angle	acute	acute	acute	acute
☑ Stem: internode length	medium	medium	short	medium
☑ Stem: colour of new growth (RHS colour chart)	187C	187B	183A	185A
☐ Leaf: blade length	medium to long	long	medium to long	medium
☑ Leaf: blade width	medium	medium	broad	medium
☐ Leaf: petiole length	short	short	short	short
☑ Leaf: shape of blade	elliptic	narrow elliptic	broad elliptic	elliptic
☑ Leaf: shape of apex	apiculate	cuspidate	cuspidate	cuspidate
☐ Leaf: shape of base	acute	acute	acute	acute
☐ Leaf: glossiness	strong to medium	medium	medium	medium
☐ Leaf: shape of cross section	flat to concave	flat	flat to concave	flat to concave
☐ Leaf: shape of longitudinal section	flat to concave	flat	flat to concave	flat to concave
☐ Leaf: stiffness	medium	medium to strong	medium	medium
☐ Leaf: prominence of midrib on lower surface	not prominent	n/a	not prominent	not prominent
☐ Mature leaf: primary colour of upper side (RHS colour chart)	147A	147A	147A	147A
☐ Mature leaf: primary colour of lower side (RHS colour chart)	146B	146B	146B	146B
☑ Partly mature leaf: primary colour of upper side (RHS colour chart)	175A	ca 177A	176A	175A
☑ Partly mature leaf: primary colour of lower side (RHS colour chart)	ca 177B	ca 165A	177B	ca 177B
☑ Newly emerged: upper side (RHS colour chart)	187C	187B	183A	178B
☐ Leaf: variegation	absent	absent	absent	absent

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Sunrise'	'Cherry Surprise'	'Allyn Magic'	'Hot Flush
☑ Newly emerged leaf: size	small	medium	small	medium

Prior Applications and Sales

Nil.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW.

Details of Application

Application Number	2001/229
Variety Name	'Street Snow'
Genus Species	*Mimusops elengi*
Common Name	Spanish Cherry
Synonym	Nil
Accepted Date	04 Sep 2001
Applicant	Darwin Plant Wholesalers, Winnellie, NT
Agent	N/A
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Lambells Lagoon, NT.
Descriptor	Spanish Cherry (*Mimusops elengai*) PBR MIMU.
Period	Spring 2006-summer 2007.
Conditions	Trial conducted in a opens beds, plants originally propagated by cuttings, mature trees in 150L containers filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Ten pots of each variety arranged in a completely randomised design.
Measurements	From five plants at random. Two leaf samples per plant.
RHS Chart - edition	2001.

Origin and Breeding

Spontaneous mutation: spontaneous mutation from a mass growing of *Mimusops elengai* of Indonesian origin in Benara Nursery, near Jakarta, Indonesia. Material being grown for commercial use, having non-variegated leaves. Selection criteria: variegated leaf. Propagation: vegetative cuttings were taken from the original variegated plant and propagated for several generations to confirm the uniformity and stability of the selction. Breeder: Darryl South, Darwin Plant Wholesalers, Winnellie, NT.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	presence of variegation	present

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Street Elegance'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Street Snow'	'Street Elegance'
☐ Plant: growth habit	upright	upright
☐ Plant: vigour	weak to medium	medium
☑ Plant: density	medium	very dense
☐ Plant: inner angle of lateral shoots to main stem	broad acute	broad acute
☐ Plant: length of internodes	medium	medium
☐ Plant: colour of young stem	brownish green	brownish green
☑ Plant: glaucosity of young stem	medium	strong
☐ Plant: colour of older stem	light greyish brown	light greyish brown
☐ Petiole: length	short to medium	short
☐ Petiole: colour	light green	light green
☑ Leaf blade: length	medium to long	medium
☑ Leaf blade: width	broad	medium
☑ Leaf blade: shape	broad elliptic	elliptic
☑ Leaf blade: shape of apex	broad-acuminate	acuminate
☑ Leaf blade: shape of base	obtuse	cuneate
☐ Leaf bade: undulation of margin	strong	medium to strong
☐ Leaf blade: cross-section	concave	concave
☐ Leaf blade: curvature of longitudinal section	recurved	recurved
☐ Leaf blade: variegation	present	present
☐ Leaf blade: border between colours	not clearly defined	not clearly defined
☐ Leaf blade: regularity of colour patches	irregular	irregular
☑ Leaf blade: ground colour (RHS colour chart)	147A	147B
☑ Leaf blade: secondary colour (RHS colour chart)	155A	147C
☑ Leaf blade: tertiary colour (RHS colour chart)	n/a	11B
☑ Leaf blade: area of ground colour compared to other colours	small	large
☐ Leaf blade: glossiness	weak to medium	weak

Statistical Table

Organ/Plant Part: Context	'Street Snow'	'Street Elegance'
☑ Leaf blade: length (mm)		
Mean	91.10	82.40
Std. Deviation	4.40	8.60
LSD/sig	8.82	P≤0.01
☑ Leaf blade : width (mm)		
Mean	52.30	36.80
Std. Deviation	4.60	2.80
LSD/sig	4.91	P≤0.01

Prior Applications and Sales

Nil.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW.

Details of Application

Application Number	2007/146
Variety Name	'Ocean'
Genus Species	*Chlorophytum comosum*
Common Name	Spider Plant
Synonym	Nil
Accepted Date	11 Jul 2007
Applicant	Koning Smit IPR S.A., Aalsmeer, The Netherlands
Agent	Ramm Botanicals Pty Ltd, Tuggerah, NSW
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	MacMaster's Beach, NSW
Descriptor	Spider plant (*Chlorophytum comosum*) PBR CHLO
Period	Autumn-spring 2007
Conditions	Trial conducted in a fibreglass covered greenhouse, plants propagated by division, tubestock planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release fertilisers and overhead irrigated, no pest or disease treatments were required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random. One sample per plant.
RHS Chart - edition	2001

Origin and Breeding

Spontaneous mutation: parent *Chlorophytum comosum* 'Variegatum'. The parent is characterised by a yellow green leaf margin with a white central stripe along the middle of the blade. Selection took place in Sappemeeer, the Netherlands. Selection criteria: distinctive leaf colour, strong growth vigour and ease of reproduction. Propagation: vegetatively reproduced plants from micropropagation, cuttings and divisions are found to be uniform and stable. Breeder: Lammert Koning, The Netherlands.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf blade	curvature	straight

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Variegatum'	common form and parent

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Bonnie'	Leaf blade curvature	straight	strongly curved

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Ocean'	'Variegatum'
☐ Plant: growth habit	erect to semi-erect	erect to semi-erect
☑ Plant: height	tall	medium
☐ Plant: density of shoots	medium to dense	medium to dense
☑ Plant: vigour	very strong	medium
☑ Stolon: colour	green	yellow
☐ Leaf: twisting	absent	absent
☐ Leaf: arching	medium to strong	medium
☐ Leaf: length	medium to long	medium
☑ Leaf: width	wide	medium
☐ Leaf: variegation	present	present
☐ Leaf: primary colour of upper side (RHS colour chart)	147A	147A
☐ Leaf: primary colour of lower side (RHS colour chart)	147B	146A
☑ Leaf: secondary colour of upper side (variegated leaves only) (RHS colour chart)	191A	146B
☑ Leaf: secondary colour of lower side (variegated leaves only) (RHS colour chart)	198A	191A
☐ Leaf: tertiary colour of upper side (variegated leaves only) (RHS colour chart)	155A	155A
☐ Leaf: shape of blade	ensiform	ensiform
☐ Leaf: shape of apex	acute	acute
☐ Corolla: shape	rotate	rotate
☐ Corolla: colour	white	white

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Ocean'	'Variegatum'
☐ Leaf: cross-section	concave to flat	concave to flat
☑ Leaf: colour of margin	white	green
☐ Leaf blade: curvature	straight	straight

Statistical Table

Organ/Plant Part: Context	'Ocean'	'Variegatum'
☐ Leaf: length (mm)		
Mean	391.20	350.50
Std. Deviation	34.70	33.60
LSD/sig	39.0	ns
☑ Leaf: width (mm)		
Mean	29.80	22.00
Std. Deviation	4.60	2.10
LSD/sig	4.08	P≤0.01

Prior Applications and Sales

Country	Year	Current Status	Name Applied
EU	2004	Granted	‘Ocean’

First sold in The Netherlands in Mar 2004. First Australian sale Jan 2007.

Description: **Ian Paananen, Crop & Nursery Services, Central Coast, NSW**

Details of Application

Application Number	2006/088
Variety Name	'LHCOM'
Genus Species	*Lomandra hystrix*
Common Name	Spiny Headed Mat Rush
Synonym	Nil
Accepted Date	30 May 2006
Applicant	Ozbreed Pty Ltd, Clarendon, NSW
Agent	N/A
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Clarendon, NSW
Descriptor	Lomandra (*Lomandra*) PBR LOMA
Period	Autumn 2007 - spring 2007
Conditions	Trial conducted in open beds, plants propagated from cuttings, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random.
RHS Chart - edition	1995

Origin and Breeding

Seedling Selection: seed parent *L. hystrix*. The seed parent is characterised by a tall plant height, upright plant growth habit and a broad leaf width. In 2001 open-pollinated seedlings of *L. hystrix* were grown in an open bed. There were approximately 5000 plants grown in viro tubes. In 2002 approximately 200 plants were selected due to their smaller shoot and leaf sizes These were grown on for further observation. In 2003, these were reduced to 10 selections based on the same criteria. Finally, in late 2004 a single plant was identified as having narrower leaf width combined with a compact, dense growth habit with a shorter plant height than the parent form. Selection took place in Clarendon, NSW. Selection criteria: narrow leaf width, short plant height, compact habit with more horizontal basal shoot attitude. Propagation: vegetative, micropropagation is found to be uniform and stable. Breeder: Todd Layt, Clarendon, NSW.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf blade	presence of variegation	absent
Plant	sex expression	male

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
L.hystrix common male form 'LHBYF'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'LHCOM'	*L.hystrix* common male form	'LHBYF'
☑ Plant: growth habit	semi-upright	upright	semi-upright
☑ Plant: height	medium	very tall	tall
☑ Plant: density	dense	medium	medium
☐ Leaf: texture	medium	medium	medium
☐ Leaf: glaucosity	weak	weak	weak
☑ Leaf: rigidity	weak	medium	medium
☑ Leaf: length of blade	short	medium	medium
☑ Leaf: width of blade	narrow	medium	medium
☐ Leaf: cross section	flat	flat	flat
☐ Leaf: variegation	absent	absent	absent
☐ Leaf: colour (RHS colour chart)	146A	146A	146B
☐ Basal sheath: margin shredding	very weak	very weak	very weak
☐ Basal sheath: colour	medium brown	medium brown	medium brown
☐ Inflorescence: degree of branching	strong	strong	strong
☑ Inflorescence: length of floral axis	short	medium	long
☑ Inflorescence: length of peduncle	medium	long	medium
☑ Inflorescence: length of bract	medium	medium	long
☑ Inflorescence: position in relation foliage	level	below	above
☑ Inflorescence: colour of peduncle (RHS colour chart)	145B	145A	145C
☑ Flower: colour of calyx (RHS colour chart)	145B	145A	145C
☑ Flower: colour of perianth (RHS colour chart)	22A	11C	23A

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'LHCOM'	*L.hystrix* common male form	'LHBYF'
☑ Plant: duration of flowering	medium	medium	long

Statistical Table

Organ/Plant Part: Context	'LHCOM'	*L.hystrix* common male form	'LHBYF'

☑ Plant: height (mm)			
Mean	475.00	658.00	615.50
Std. Deviation	68.70	73.00	39.30
LSD/sig	70.96	P≤0.01	P≤0.01
☑ Leaf: length (mm)			
Mean	452.60	642.50	600.70
Std. Deviation	83.10	61.20	28.80
LSD/sig	70.60	P≤0.01	P≤0.01
☑ Leaf: width (mm)			
Mean	10.50	13.20	13.10
Std. Deviation	1.20	0.90	0.70
LSD/sig	1.10	P≤0.01	P≤0.01
☑ Inflorescence: length of floral axis (mm)			
Mean	233.40	341.20	411.70
Std. Deviation	44.40	63.30	50.60
LSD/sig	60.93	P≤0.01	P≤0.01
☑ Inflorescence: width (mm)			
Mean	114.80	97.20	154.00
Std. Deviation	10.10	19.70	13.30
LSD/sig	16.98	ns	P≤0.01
☑ Inflorescence: length of bract (mm)			
Mean	62.80	46.40	93.30
Std. Deviation	12.20	26.70	30.30
LSD/sig	27.79	ns	P≤0.01
☑ Peduncle: length (mm)			
Mean	123.40	184.00	141.50
Std. Deviation	29.20	28.90	33.20
LSD/sig	34.81	P≤0.01	ns

Prior Applications and Sales

Nil.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW.

Details of Application

Application Number	2006/270
Variety Name	'LHBYF'
Genus Species	*Lomandra hystrix*
Common Name	Spiny Headed Mat Rush
Synonym	Nil
Accepted Date	26 Oct 2006
Applicant	Ozbreed Pty Ltd, Clarendon, NSW
Agent	N/A
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Clarendon, NSW.
Descriptor	Lomandra (*Lomandra*) PBR LOMA
Period	Autumn 2007 - spring 2007.
Conditions	Trial conducted in open beds, plants propagated from cuttings, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random.
RHS Chart - edition	1995.

Origin and Breeding

Seedling selection: seed parent *L. hystrix*. The seed parent is characterised by a medium inflorescence width and a medium number of flowers. Approximately 3000 seedlings were grown in 1997 and originally 50 were selected as having the best growth vigour. These were grown on in pots and 3 seedlings were selected due to their male flowering habits. These were grown on as garden plants until 2003 when finally one of these plants were selected due to its prolific large yellow flowers combined with vigorous growth. Selection took place in Clarendon, NSW. Selection criteria: prolific flowering and large inflorescence size. Propagation: vegetative, micropropagation is found to be uniform and stable. Breeder: Todd Layt, Clarendon, NSW.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf blade	presence of variegation	absent
Plant	sex expression	male

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'LHCOM'	
L.hystrix common male form	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'LHBYF'	*L.hystrix* common male form	'LHCOM'
☑ Plant: growth habit	semi-upright	upright	semi-upright
☑ Plant: height	tall	very tall	medium
☑ Plant: density	medium	medium	dense
☐ Leaf: texture	medium	medium	medium
☐ Leaf: glaucosity	weak	weak	weak
☐ Leaf: rigidity	medium	medium	weak
☑ Leaf: length of blade	medium	medium	short
☑ Leaf: width of blade	medium	medium	narrow
☐ Leaf: cross section	flat	flat	flat
☐ Leaf: variegation	absent	absent	absent
☐ Leaf: colour (RHS colour chart)	146B	146A	146A
☐ Basal sheath: margin shredding	very weak	very weak	very weak
☐ Basal sheath: colour	medium brown	medium brown	medium brown
☐ Inflorescence: degree of branching	strong	strong	strong
☑ Inflorescence: length of floral axis	long	medium	short
☑ Inflorescence: length of peduncle	medium	long	medium
☑ Inflorescence: length of bract	long	medium	medium
☑ Inflorescence: position in relation foliage	above	below	level
☑ Inflorescence: colour of peduncle (RHS colour chart)	145C	145A	145B
☑ Flower: colour of calyx (RHS colour chart)	145C	145A	145B
☑ Flower: colour of perianth (RHS colour chart)	23A	11C	22A

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'LHBYF'	*L.hystrix* common male form	'LHCOM'
☑ Plant: duration of flowering	long	medium	medium

Statistical Table

Organ/Plant Part: Context	'LHBYF'	*L.hystrix* common	'LHCOM'

		male form	
☑ Plant: height (mm)			
Mean	615.50	658.00	475.00
Std. Deviation	39.30	73.00	68.70
LSD/sig	70.96	ns	P≤0.01
☑ Leaf: length (mm)			
Mean	600.70	642.50	452.60
Std. Deviation	28.80	61.20	83.10
LSD/sig	70.60	ns	P≤0.01
☑ Leaf: width (mm)			
Mean	13.10	13.20	10.50
Std. Deviation	0.70	0.90	1.20
LSD/sig	1.10	ns	P≤0.01
☑ Inflorescence: length of floral axis (mm)			
Mean	411.70	341.20	233.40
Std. Deviation	50.60	63.30	44.40
LSD/sig	60.93	P≤0.01	P≤0.01
☑ Inflorescence: width (mm)			
Mean	154.00	97.20	114.80
Std. Deviation	13.30	19.70	10.10
LSD/sig	16.98	P≤0.01	P≤0.01
☑ Inflorescence: length of bract (mm)			
Mean	93.30	46.40	62.80
Std. Deviation	30.30	26.70	12.20
LSD/sig	27.79	P≤0.01	P≤0.01
☑ Peduncle: length (mm)			
Mean	141.50	184.00	123.40
Std. Deviation	33.20	28.90	29.20
LSD/sig	34.81	P≤0.01	ns

Prior Applications and Sales

Nil.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW.

Details of Application

Application Number	2005/340
Variety Name	'Cal Giant 5'
Genus Species	*Fragaria* x*ananassa*
Common Name	Strawberry
Synonym	Galexia
Accepted Date	22 Dec 2005
Applicant	California Giant, Inc., Watsonville, CA
Agent	State of Queensland through its Department of Primary Industries and Fisheries, Brisbane, QLD
Qualified Person	Mark Herrington

Details of Comparative Trial

Location	Redlands Research Station, Delancey St., Cleveland, QLD. (Latitude 27 South, Longitude 153 East elevation 24 m).
Descriptor	Strawberry (*Fragaria*) TG/22/9
Period	Apr to Sep 2007.
Conditions	Trial conducted in a non-fumigated field of krasnozem soil, runners from commercial sources in QLD runner growing district (Stanthorpe), black polythene mulch, double rows on beds (40 cm inter-row, 40 cm intra-row and 140 cm between bed centres), trickle irrigated and fertilised, pest and disease treatments applied as required.
Trial Design	Duplicate plots each of approx 25 plants.
Measurements	From twenty to twenty-eight plants or fruit per cultivar as individual plants or fruit randomly sampled over the duplicate plots.
RHS Chart - edition	2001 for leaf colour, 1995 for fruit colour.

Origin and Breeding

Controlled pollination: main selection criteria used to develop this variety were fruit quality, disease resistance, and productivity. During the period between Oct of 1996 and May of 1997 parent material was placed in an enclosed greenhouse and controlled hybridisation between those parents took place. Of the seed pollinated 15,000 unique varieties germinated; within that group of unique varieties during the grow-out period to Aug of 1999 the selection 65H1 showed potential due to its strong flesh and skin firmness, good colour, good interior colour, good flavour, strong propensity to produce fruit, good continuing size within the hand, as well as retention of size from hand to hand, and the tremendous disease tolerance of the variety. After three successive years of testing the variety 65H1 was determined to be worthy of plant protection. At that time the variety was designated Galexia and protection has been sought as 'CalGiant 5'. The variety was tested each successive year for three years before the decision was made to seek protection for the variety. From its inception the variety has been propagated annually, asexually, at the nursery through the growing of runners. The variety has been propagated and continues to be propagated asexually to date. To date there have been no known off-types. Breeder: David W Small, Santa Maria, California, USA.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	habit	flat globose
Leaf	blistering	absent or very weak
Inflorescence	position relative to foliage	level with
Flower	relative position of petals	overlapping
Flower	size of calyx relative to corolla	same size
Fruit	length/width ratio	slightly longer than broad
Fruit	size	medium
Fruit	predominant shape	conical
Fruit	colour	red
Fruit	colour of flesh	orange red
Fruit	hollow centre	weakly expressed
Fruit	distribution of red colour of flesh	marginal and central
Time	of flowering	medium
Time	of ripening	medium
Bearing	type	partially remontant

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Cal Giant 3'	

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Cal Giant 5'	'Cal Giant 3'
☐ Plant: habit	flat globose	flat globose
☐ Plant: density	medium	medium
☐ Plant: vigour	medium	medium
☐ Leaf: colour of upper side	medium green	medium green
☐ Leaf: shape in cross section	slightly concave	strongly concave to slightly concave
☐ *Leaf: blistering	absent or very weak	absent or very weak
☐ *Leaf: glossiness	weak	weak
☐ *Terminal leaflet: length/width ratio	as long as broad	as long as broad
☐ *Terminal leaflet: shape of base	obtuse	obtuse
☐ Terminal leaflet: shape of incisions of margin	crenate	crenate
☐ Petiole: attitude of hairs	strongly outwards	strongly outwards
☐ Stipule: anthocyanin colouration	absent or very weak	absent or very weak
☐ *Inflorescence: position relative to foliage	level with	level with
☐ Flower: size	large	large
☐ *Flower: size of calyx	same size	same size
☐ *Primary flower: relative position of petals	overlapping	overlapping

Petal: length/width ratio	as long as broad	as long as broad
*Fruit: ratio of length/width	slightly longer than broad	slightly longer than broad
*Fruit: size	medium	medium
*Fruit: predominant shape	conical	conical
Fruit: band without achenes	narrow	narrow
Fruit: unevenness of surface	absent or very weak	absent or very weak
*Fruit: colour	red	red
Fruit: evenness of colour	slightly uneven	even
Fruit: glossiness	strong	strong
*Fruit: insertion of achenes	below surface	level with surface
Fruit: insertion of calyx	with fruit level	with fruit level
Fruit: attitude of the calyx segments	spreading	reflexed
Fruit: size of calyx in relation to fruit diameter	slightly larger	same size
Fruit: adherence of calyx	medium	strong
Fruit: firmness	very firm	medium
Fruit: colour of flesh	orange red	orange red
Fruit: hollow centre	weakly expressed	weakly expressed
Fruit: distribution of red colour of flesh	marginal and central	marginal and central
*Time of: flowering	medium	medium
Time of: ripening	medium	medium
*Type of: bearing	partially remontant	partially remontant

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	**‘Cal Giant 5’**	**‘Cal Giant 3’**
Leaf: green colour upper side (RHS, 2001)	147A	147A
Fruit: colour (RHS, 1995)	45A	53A

Prior Applications and Sales

Country	**Year**	**Current Status**	**Name Applied**
USA	2002	Granted	‘Cal Giant 5’
EU	2003	Granted	‘Galexia’
South Africa	2003	Applied	‘Galexia’

First sold in USA in Oct 2002.

Description: **Mark Herrington,** Maroochy Horticultural Research Station, Nambour, QLD.

Details of Application

Application Number	2002/008
Variety Name	'Arodel'
Genus Species	*Prunus avium*
Common Name	Sweet Cherry
Synonym	Nil
Accepted Date	27 Jun 2003
Applicant	Societe Anonyme des Pepinieres et Roseraies GEORGES DELBARD, Malicorne, France
Agent	Australian Nurserymen's Fruit Improvement Company, Bathurst, NSW.
Qualified Person	Peter Kennedy

Details of Comparative Trial

Location	Young, NSW. Longitude 148°18′ E, Latitude 34°18′ S.
Descriptor	Cherry (*Prunus avium*) TG/35/6.
Period	2003-2007.
Conditions	Grown under normal conditions on a Tatura Trellis training system.
Trial Design	Six trees of the candidate variety were planted in 2003.A total of 20 trees of two comparator varieties were planted in 2001 and 2003.
Measurements	From all trial plants.
RHS Chart - edition	N/A

Origin and Breeding

Open pollination: 'Arodel' is a product of uncontrolled pollination of approximately 50 cherry varieties on the French National List. Approximately 50,000 seedlings were raised and from these 28 were selected by the breeder for further evaluation. These 28 selections were in turn passed to Delbard Nurseries in France for further evaluation. In 1991 Delbard selected 7 of the 28 lines for indexing. 'Arodel' was one of the seven selections. Breeder: Paul Argot, Rive-de-Gier, France.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Fruit	maturity	very early
Fruit	size	large
Fruit	colour of skin	red

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Early Burlat'	Time of maturity: very early
'Rivedel'	Time of maturity: very early

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Empress'	Fruit	size	large	small to medium
'Burgsdorf'	Fruit	size	large	small

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Arodel'	'Early Burlat'	'Rivedel'
☐ Tree: vigour	strong to very strong	strong	medium
☐ *Tree: habit	upright to semi-upright	upright	upright
☑ *Tree: branching	medium	weak	weak
☐ One-year-old shoot: number of lenticels	medium	medium	few to medium
☐ Young shoot: anthocyanin colouration of tip	medium	weak to medium	medium
☐ Leaf blade: green colour of upper side	light to medium	light to medium	light to medium
☐ *Petiole: nectaries	present	present	present
☐ Petiole: colour of nectaries	dark red	dark red	dark red
☐ *Fruit: size	large	large	large
☐ *Fruit: shape	reniform	reniform	reniform
☐ Fruit: pistil end	depressed	depressed	depressed
☐ *Fruit: colour of skin	red	red	red
☐ Fruit: size of lenticels on skin	small	small	small
☑ Fruit: number of lenticels on skin	many	many	few
☑ Fruit: colour of juice	pink	red	red
☐ Fruit: colour of flesh	red	dark red	dark red
☐ *Fruit: firmness	medium to firm	medium	medium
☐ Fruit: acidity	medium	medium	medium
☐ Fruit: sweetness	medium	medium	medium
☐ Fruit: juiciness	strong to very strong	strong to very strong	strong to very strong
☐ *Fruit: length of stalk	medium	medium	medium
☐ Fruit: abscission layer between stalk and fruit	present	present	present
☐ Fruit: thickness of stalk	medium	medium	
☐ *Stone: size	large	medium to large	medium to large
☐ *Stone: shape	broad elliptic	broad elliptic	broad elliptic
☐ *Time of: flowering	early	early	very early to early
☐ *Time of: fruit maturity	very early	very early	very early

Statistical Table

Organ/Plant Part: Context	'Arodel'	'Early Burlat'	'Rivedel'
☑ Leaf: length (mm)			
Mean	156.9	149.6	131.0

Std. Deviation	13.99	10.28	15.61
LSD/sig	14.91	ns	P≤0.01
☑ Leaf: width (mm)			
Mean	81.45	61.85	63.3
Std. Deviation	3.46	8.06	4.47
LSD/sig	6.28	P≤0.01	P≤0.01
☐ Petiole: length (mm)			
Mean	36.5	38.5	34.5
Std. Deviation	8.31	7.98	4.00
LSD/sig	7.79	ns	ns
☑ Fruit: diameter (mm)			
Mean	27.75	26.28	26.46
Std. Deviation	0.56	1.15	1.40
LSD/sig	1.21	P≤0.01	P≤0.01
☐ Fruit: length of stalk (mm)			
Mean	29.18	27.11	27.04
Std. Deviation	2.39	4.18	1.74
LSD/sig	3.27	ns	ns
☑ Stone: diameter (mm)			
Mean	10.58	9.66	9.89
Std. Deviation	0.54	0.55	0.65
LSD/sig	0.64	P≤0.01	P≤0.01
☐ Fruit: brix (°Bx RDS)			
Mean	13.89	14.48	14.19
Std. Deviation	1.54	1.12	1.03
LSD/sig	1.38	ns	ns

Prior Applications and Sales

Country	Year	Current Status	Name Applied
France	1993	Granted	‘Arodel’

Prior sale nil.

Description: **Peter Kennedy,** Young, NSW.

Details of Application

Application Number	2003/148
Variety Name	'Dame Nancy'
Genus Species	*Prunus avium*
Common Name	Sweet Cherry
Synonym	Nil
Accepted Date	7 Jul 2003
Applicant	Minister for Agriculture, Food and Fisheries, Adelaide, SA
Agent	Australian Nurseryman's Fruit Improvement Company Limited, Bathurst, NSW
Qualified Person	Peter Kennedy

Details of Comparative Trial

Location	Young, NSW. Longitude 148°18′ E, Latitude 34°18′ S.
Descriptor	Cherry (*Prunus avium*) TG/35/6
Period	2001-2007
Conditions	Grown under normal conditions on a Tatura trellis training system.
Trial Design	Four trees of the candidate variety and six trees of the comparator variety were planted at the trial site in 2001 on a commercial orchard.
Measurements	From all trial plants.
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: 'Dame Nancy' is the result of the controlled cross of the self fertile variety 'Stella'. 'Stella' seed parents were enclosed by shadecloth to exclude pollinating insects. Methods including flower emasculation and hand hybridisation were used to make controlled crosses. Seeds from successful hybridisations were then germinated and F_1 seedlings planted in the field. Fruit was assessed from 1991 onwards and the selection of 'Dame Nancy' was made in 1992. Original clonal material has been held at Lenswood Horticultural Centre, Lenswood SA and no off types have been observed. Breeder: Dr. Andrew Granger, Lenswood Horticultural Centre, Lenswood, SA.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Fruit	size	large to very large
Fruit	maturity	medium to late
Fruit	colour of skin	vermillion on pale yellow background

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Rainier'	'Rainier', like 'Dame Nancy', is a blush cherry that is considered the world standard in blush cherries.

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Stella'	Fruit colour	vermillion with pale yellow background	Dark red

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Dame Nancy'	'Rainier'
☐ Tree: vigour	strong	strong
☑ *Tree: habit	semi-upright	upright
☑ *Tree: branching	medium	weak
☐ One-year-old shoot: number of lenticels	many	
☑ Young shoot: anthocyanin colouration of tip	absent or very weak	weak to medium
☐ Leaf blade: green colour of upper side	medium	medium to dark
☐ *Petiole: nectaries	present	present
☐ Petiole: colour of nectaries	dark red	dark red
☐ *Fruit: size	large to very large	large
☐ *Fruit: shape	reniform	reniform
☐ Fruit: pistil end	flat	flat
☐ *Fruit: colour of skin	vermillion on pale yellow background	vermillion on pale yellow background
☐ Fruit: size of lenticels on skin	small	small
☐ Fruit: number of lenticels on skin	many	many to very many
☑ Fruit: colour of juice	cream yellow	pink
☑ Fruit: colour of flesh	yellow	cream white
☐ *Fruit: firmness	medium to firm	firm
☑ Fruit: acidity	medium	low
☑ Fruit: sweetness	medium	very high
☐ Fruit: juiciness	strong to very strong	strong to very strong
☑ *Time of: flowering	late	early
☐ *Time of: fruit maturity	medium to late	medium to late

Statistical Table

Organ/Plant Part: Context	'Dame Nancy'	'Rainier'
☑ Leaf: length (mm)		
Mean	171.5	193.1
Std. Deviation	11.34	18.96
LSD/sig	17.83	P≤0.01
☑ Leaf: width (mm)		
Mean	81.1	85.7
Std. Deviation	7.12	8.81
LSD/sig	9.17	ns

☑ Petiole: length (mm)		
Mean	31.35	40.6
Std. Deviation	5.05	6.99
LSD/sig	6.96	P≤0.01
☐ Fruit: diameter (mm)		
Mean	29.4	27.90
Std. Deviation	1.41	1.82
LSD/sig	1.86	ns
☐ Fruit: length of stalk (mm)		
Mean	35.62	30.77
Std. Deviation	5.98	2.46
LSD/sig	5.22	ns
☑ Stone: diameter (mm)		
Mean	9.44	10.12
Std. Deviation	0.70	0.42
LSD/sig	0.66	P≤0.01
☑ Fruit: brix (°Bx RDS)		
Mean	16.5	19.83
Std. Deviation	0.99	3.48
LSD/sig	2.92	P≤0.01

Prior Applications and Sales

Nil.

Description: **Peter Kennedy,** Young, NSW.

Details of Application

Application Number	2007/234
Variety Name	'Hawkeye'
Genus Species	x*Triticosecale*
Common Name	Triticale
Synonym	Nil
Accepted Date	10 Oct 2007
Applicant	Australian Grain Technologies Pty Ltd, Glen Osmond, SA
Agent	N/A
Qualified Person	Gil Hollamby

Details of Comparative Trial

Location	Mintaro, South Australia.
Descriptor	Triticale (x*Triticosecale*) TG/83/4
Period	Winter to spring 2007.
Conditions	The trial was grown in a black self mulching soil which had been pasture in 2006 and wheat in 2005. The area was sprayed with Roundup Power Max (1.2L/ha) + Goal CT (75ml/ha) on 24 May 2007 and direct drilled at 2-4cm in slightly moist conditions on 25 May at 200 plants/m2 and with 90kg/ha DAP and 80kg/ha Urea. During the winter months moisture was adequate and the trial grew well. In crop weeds were controlled with 2,4-D amine 625 (1.5l/ha) on 6 Sep. Spring was dry and some moisture stress occurred. Harvest took place on 11 Dec about two weeks earlier than normal. There were no diseases of note. A similar trial was planted at Roseworthy.
Trial Design	Randomised Block Design of 3 blocks and 16 entries consisting of comparators and potential candidates. Sown in 12 ranges of 4 plots wide, block 1 being in ranges 1 to 4 and so on. Plots were 1.25m wide (5 rows) and 3.2m long. There were approx. 1000 plants per plot.
Measurements	Heading times were recorded on the same trial planted at Roseworthy 2007, but this trial later was abandoned due to a heavy infestation of Crown Rot. All other measurements and observations were recorded on plant samples taken from the Mintaro trial. At anthesis 5 primary tillers were sampled from each plot in each replicate and flag leaf measurements made. Glaucosity and leaf angle was observed at this time. After maturity plant heights to the top of the awns were recorded at 10 random locations in replicate 2 and 3 only. Twenty heads were also sampled at random from each plot in replicates 2 and 3 for head descriptions and measurements. Measurements were performed on 10 intact heads per block. Physical quality data was measured on the grain harvested from the plots. Statistical analyses were completed using GENSTAT software.
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: The cross ISR499-61/TX93-19-2 was made by Dr Kath V Cooper in the glasshouse at Waite Campus, The University of Adelaide in the spring of 1996. The female parent, ISR499-61 (NSW accession number of an imported CIMMYT line POPP1_2), had a broad head type. The male parent was a sib of 'Tickit' and 'Speedee' and had a shorter stature, resistance to cereal cyst nematode and resistance to the stem rust pathotype 34-2,12,13. F_1 generation seed was harvested in Jan 1997, and allocated the number TX97-41. Fl seed was immediately sown in pots in the glasshouse to produce F_2 generation seed, harvested May 1997, and sown as a single plot at Callington, SA. Single heads from plants showing desired agronomic type were taken in Dec and sown as head hills in the Waite Campus birdcage, under irrigation. F_3 generation head hills were harvested in May 1998 and sown as (F_4 generation) single plots at Callington. A line having desirable plant type, cereal cyst nematode resistance (SARDI test) and stem rust resistance (NRCP test), was selected and designated TX97-41-1. TX97 -41-1 was assessed for grain yield, plant type and grain conformation as F_5 replicated field trials in 1999 (2 sites), as F_6 in 2000 and as F_7 in 2001 (4 sites). Sites used were Callington, Lameroo, Cleve and Birdwood all in SA. Re-selections were taken from TX97-41-1 at the Callington site in 2001 to improve uniformity. These F_8 heads were sown as head hills at Waite Campus, in the birdcage under irrigation, in Dec 2001, harvested May 2002 and resown at Birdwood, Jun 2002. One of these reselections, designated TX97-41-1-2 after yield testing in replicated trials at 3 sites during winter 2003 was transferred by Dr. Cooper to Australian Grain Technologies under a licensing agreement where its trialling was continued by Jason Reinheimer. TX97-41-1-2 was assessed for yield, physical grain quality, disease resistance and plant type at 11 sites across Australia in 2004 as well as CCN resistance in the laboratory. In 2004, 50 single head selections were taken from a single plot of TX97-41-1-2 and were grown over summer at Roseworthy Campus, University of Adelaide. In 2005 these single selections were assessed individually for plant type, rust resistance and CCN resistance with the resistant individuals that were similar in plant type formed a bulk designated TSA0108. This line was assessed for yield, rust resistance, CCN resistance and physical grain quality at 19 sites by AGT and 15 sites by the National Variety Trial system across Australia in 2006 and again in 2007. Breeders: Dr Kath Cooper, The University of Adelaide and Mr. Jason Reinheimer, Australian Grain Technologies.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	time of ear emergence	250 to 255 Julian days
Plant	height	105 to 120cm
Flag leaf	length of blade	>180mm
Ear	degree of awning	fully awned
Ear	colour	white

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Tahara'	common variety grown in the area of adaptation.
'Tickit'	related variety.
'Kosciuszko'	visually similar in the field.

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Speedee'	Plant	time of ear emergence	253.8 Julian days	247.0 Julian days
'Jackie'	Plant	time of ear emergence	253.8 Julian days	271.0 Julian days
'Abacus'	Plant	time of ear emergence	253.8 Julian days	262.7 Julian days
'Jackie'	Flag leaf	length	205.0mm	137.9mm
'Jackie'	Flag leaf	width	17.00mm	14.30mm
'Treat'	Plant	time of ear emergence	253.8 Julian days	251.7 Julian days

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Hawkeye'	'Kosciuszko'	'Tahara'	'Tickit'
☐ *Ploidy:	hexaploid	hexaploid	hexaploid	hexaploid
☑ Flag leaf: anthocyanin colouration of auricles	medium	absent or very weak	weak to medium	medium
☐ Awn: anthocyanin colouration	absent or very weak		absent or very weak	absent or very weak
☐ Ear: glaucosity	medium		medium	medium
☐ *Stem: density of hairiness of neck	strong	strong to very strong	strong to very strong	strong
☐ *Ear: distribution of awns	fully awned	fully awned	fully awned	fully awned
☐ *Awns above the tip of ear: length	short to medium	short	short to medium	short to medium
☐ *Lower glume: length of first beak	short to medium	medium	short	medium
☐ Lower glume: size of second beak	absent or very small	absent or very small	absent or very small	absent or very small
☑ *Lower glume: hairiness on external surface	present	absent	absent	absent
☐ Straw: pith in cross section	thin to medium	medium	thin to medium	thin
☐ Ear: colour	white	white	white	white
☐ *Grain: colouration with phenol	very dark	dark	very dark	very dark
☐ *Seasonal type:	spring type	spring type	spring type	spring type

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Hawkeye'	'Kosciuszko'	'Tahara'	'Tickit'
☐ Leaves: reaction to stripe rust pathotype 110E143A+	resistant	resistant	resistant	resistant
☑ Leaves: reaction to stripe rust pathotype 134E16A+	resistant	moderately susceptible	resistant	moderately resistant
☑ Leaves: reaction to stripe rust pathotype 134E16A+J+	resistant	susceptible	moderately resistant	moderately resistant
☐ Ear: attitude at maturity	mixed erect to semi-recurved	mixed erect to semi-recurved	mixed erect to semi-recurved	mixed erect to semi-recurved
☑ Roots: reaction to high Boron levels	moderately intolerant			moderately tolerant

☑ Roots: reaction to Cereal Cyst Nematode	resistant	susceptible	resistant

Statistical Table

Organ/Plant Part: Context	'Hawkeye'	'Kosciuszko'	'Tahara'	'Tickit'
☐ Flag leaf blade: length (mm)				
Mean	205.00	182.40	206.90	203.30
Std. Deviation	28.70	33.00	29.20	21.70
LSD/sig	39.3	ns	ns	ns
☐ Flag leaf blade: width (mm)				
Mean	17.00	16.30	16.60	17.20
Std. Deviation	1.85	1.50	1.24	1.81
LSD/sig	1.7	ns	ns	ns
☑ Ear: length without awns (mm)				
Mean	100.50	128.00	107.40	107.40
Std. Deviation	6.70	11.10	9.50	8.30
LSD/sig	13.5	P≤0.01	ns	ns
☑ Ear: rachis internode length (mm)				
Mean	3.38	4.18	3.66	3.66
Std. Deviation	0.22	0.30	0.25	0.20
LSD/sig	0.38	P≤0.01	ns	ns
☐ Plant: height including awns (cm)				
Mean	111.20	118.20	115.60	109.70
Std. Deviation	4.10	5.20	3.20	3.47
LSD/sig	8.3	ns	ns	ns
☐ Plant: time of ear emergence from boot (Julian days)				
Mean	253.80	252.00	254.70	254.30
Std. Deviation	0.75	0	0.60	1.15
LSD/sig	1.9	ns	ns	ns
☑ Ear: width (mm)				
Mean	13.15	12.75	11.45	11.00
Std. Deviation	0.91	0.97	0.25	0.91
LSD/sig	1.87	ns	P≤0.01	P≤0.01
☑ Grain: test weight (kg/hl)				
Mean	79.93	78.93	76.27	76.93
Std. Deviation	0.61	0.46	0.90	0.83
LSD/sig	1.14	ns	P≤0.01	P≤0.01
☐ Grain: screenings, grain through 2mm sieve (%)				
Mean	1.69	4.65	7.05	4.96
Std. Deviation	0.30	0.74	3.85	0.39
LSD/sig	4.6	ns	P≤0.01	ns

Prior Applications and Sales

Nil.

Description: **Gil Hollamby,** Williamstown, SA.

Details of Application

Application Number	2007/235
Variety Name	'Jaywick'
Genus Species	x*Triticosecale*
Common Name	Triticale
Synonym	Nil
Accepted Date	10 Oct 2007
Applicant	Australian Grain Technologies Pty Ltd, Glen Osmond, SA
Agent	N/A
Qualified Person	Gil Hollamby

Details of Comparative Trial

Location	Mintaro, South Australia.
Descriptor	Triticale (x*Triticosecale*) TG/83/4.
Period	2007.
Conditions	The trial was grown in a black self mulching soil which had been pasture in 2006 and wheat in 2005. The area was sprayed with Roundup Power Max (1.2L/ha) + Goal CT (75ml/ha) on 24 May 2007 and direct drilled at 2-4cm in slightly moist conditions on 25 May at 200 plants/m2 and with 90kg/ha DAP and 80kg/ha Urea. During the winter months moisture was adequate and the trial grew well. In crop weeds were controlled with 2,4-D amine 625 (1.5l/ha) on 6 Sep. Spring was dry and some moisture stress occurred. Harvest took place on 11 Dec about two weeks earlier than normal. There were no diseases of note. A similar trial was planted at Roseworthy.
Trial Design	Randomised Block Design of 3 blocks and 16 entries consisting of comparators and potential candidates. Sown in 12 ranges of 4 plots wide, block 1 being in ranges 1 to 4 and so on. Plots were 1.25m wide (5 rows) and 3.2m long. There were approx. 1000 plants per plot.
Measurements	Heading times were recorded on the same trial planted at Roseworthy 2007, but this trial later was abandoned due to a heavy infestation of Crown Rot. All other measurements and observations were recorded on plant samples taken from the Mintaro trial. At anthesis 5 primary tillers were sampled from each plot in each replicate and flag leaf measurements made. Glaucosity and leaf angle was observed at this time. After maturity plant heights to the top of the awns were recorded at 10 random locations in replicate 2 and 3 only. Twenty heads were also sampled at random from each plot in replicates 2 and 3 for head descriptions and measurements. Measurements were performed on 10 intact heads. Grain quality was measured on the grain harvested from each plot. Statistical analyses were completed using GENSTAT software.
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: The female parent was a CIMMYT line accessioned in NSW as ISR499-62. Its pedigree is BGLB/2*RHINO_3. This line had a broad head type. This was crossed in the spring 1996 by Dr Kath V Cooper with TX93-19-2, a line bred by her and a sib to 'Tickit' and 'Speedee'. It was chosen for its shorter stature, stiff straw, high tillering habit, resistance to cereal cyst nematode, and resistance to triticale stem rust pathotype 34-2,12,13. The F_1 generation seed was allocated the number TX97-44. F_1 seed was immediately sown in pots in the glasshouse to produce F_2 generation seed, harvested May 1997, and sown as a single plot at Callington, South Australia. Single heads from plants showing the desired agronomic type were taken in Dec and sown as head hills in the Waite Campus birdcage, under irrigation. F_3 generation head hills were harvested in May 1998 and sown as single plots at Callington. One particular plot having desirable plant type, cereal cyst nematode resistance (tested by SARDI) and stem rust resistance (tested by NRCP) was given the line number TX97-44-7. TX97-44-7 was assessed in replicated trials for grain yield and plant type as an F_5 in 1999 (2 sites), F_6 in 2000 and F_7 in 2001 (4 sites). Sites used were Callington, Lameroo, Cleve and Birdwood, SA. In 2001 F_8 generation heads were selected from TX97-44-7 and sown as head hills at Waite Campus, in the birdcage under irrigation, in Dec 2001. These head hills were harvested in May 2002 and sown as F_9 generation as single plots at Birdwood in Jun 2002. One of these reselections was designated TX97-44-7-1. In May-Jun 2003, F_{10} generation seed of TX97-44-7-1 was sown in replicated yield trials at 3 sites, and harvested in Dec 2003-Jan 2004. Seed of the F_{11} generation was transferred to Australian Grain Technologies in March 2004, by means of a licensing agreement from Adelaide Research and Innovation. Jason Reinheimer continued with testing and reselecting TX97-44-7-1. In 2004 yield tests were carried out at 11 sites, and 50 single head selections were taken from a single plot of TX97-44-7-1. These selections were grown over summer at Roseworthy Campus, University of Adelaide. In 2005 these single selections were assessed individually for plant type, rust resistance and CCN resistance. Selections surviving were bulked as TSA0124. This line was assessed for yield, rust resistance, CCN resistance and physical grain quality at 19 sites by AGT and 15 sites by the National Variety Trial system across Australia in 2006 and 2007. Breeders: Dr Kath Cooper, The University of Adelaide and Mr. Jason Reinheimer, Australian Grain Technologies.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Ear	distribution of awns	fully awned
Ear	colour	white
Plant	time of ear emergence	250 to 255 Julian days
Plant	height	105 to 120cm
Flag leaf	length of blade	180 to 230mm

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Tickit'	related variety.
'Tahara'	same adaptation.
'Kosciuszko'	Similar adaptation.

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Speedee'	flag leaf blade length	193.2mm	233.1 mm	LSD=39.3(P=1%)
'Speedee'	flag leaf width	15.5mm	18.5mm	LSD=1.7(P=1%)
'Speedee'	Plant time of ear emergence	251.2 Julian days	247.0 Julian days	LSD=1.9 days(P=1%)
'Jackie'	Plant time of ear emergence	251.2 Julian days	271.0 Julian days	LSD=1.9 days(P=1%)

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Jaywick'	'Kosciuszko'	'Tahara'	'Tickit'
☐ *Ploidy:	hexaploid	hexaploid	hexaploid	hexaploid
☑ Flag leaf: anthocyanin colouration of auricles	absent or very weak	absent or very weak	weak to medium	medium
☐ *Stem: density of hairiness of neck	strong to very strong	strong to very strong	strong to very strong	strong
☐ *Ear: distribution of awns	fully awned	fully awned	fully awned	fully awned
☐ *Awns above the tip of ear: length	short to medium	short	short to medium	short to medium
☐ *Lower glume: length of first beak	medium	medium	short	medium
☐ Lower glume: size of second beak	absent or very small	absent or very small	absent or very small	absent or very small
☑ *Lower glume: hairiness on external surface	present	absent	absent	absent
☐ Straw: pith in cross section	thin to medium	medium	thin to medium	thin
☐ Ear: colour	white	white	white	white
☐ *Grain: colouration with phenol	dark to very dark	dark	very dark	very dark
☐ *Seasonal type:	spring type	spring type	spring type	spring type

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Jaywick'	'Kosciuszko'	'Tahara'	'Tickit'
☑ Roots: reaction to cereal cyst nematode	Resistant	susceptible	resistant	resistant
☑ Roots: reaction to high boron levels	Intolerant			moderately tolerant
☐ Leaves: reaction to stripe rust pathotype 110E143A+	resistant	resistant	resistant	resistant
☑ Leaves: reaction to stripe rust pathotype 134E16A+	resistant	moderately susceptible	resistant	moderately resistant
☐ Leaves: reaction to stripe rust	Resistant			
☑ Leaves: reaction to stripe rust pathotype 134E16A+J+	resistant	susceptible	moderately resistant	mod susceptible to mod resistant
☑ Ear: attitude at maturity	recurved	mixed erect to semi-recurved	mixed erect to semi-recurved	mixed erect to semi-recurved

Statistical Table

Organ/Plant Part: Context	'Jaywick'	'Kosciuszko'	'Tahara'	'Tickit'
☐ Flag leaf: blade length (mm)				
Mean	193.20	182.40	206.90	203.30
Std. Deviation	36.10	33.00	29.10	21.70
LSD/sig	39.3	ns	ns	ns
☐ Flag leaf: blade width (mm)				
Mean	15.50	16.30	16.60	17.20
Std. Deviation	1.80	1.50	1.20	1.80
LSD/sig	1.7	ns	ns	ns
☑ Flag leaf: sheath length (mm)				
Mean	156.00	181.90	178.00	184.10
Std. Deviation	10.40	16.25	12.50	9.70
LSD/sig	15.8	P≤0.01	P≤0.01	P≤0.01
☑ Ear: length without awns (mm)				
Mean	99.70	127.90	107.40	105.70
Std. Deviation	5.60	11.10	9.50	8.30
LSD/sig	13.5	P≤0.01	ns	ns
☑ Ear: width (mm)				
Mean	13.60	12.75	11.45	11.00
Std. Deviation	0.80	0.97	0.76	0.91
LSD/sig	1.87	ns	P≤0.01	P≤0.01
☑ Ear: rachis internode length (mm)				
Mean	3.68	4.18	3.66	3.55
Std. Deviation	0.21	0.30	0.25	0.20
LSD/sig	0.38	P≤0.01	ns	ns
☑ Plant: height (cm)				
Mean	108.10	118.20	115.60	109.70
Std. Deviation	3.90	5.20	3.20	3.50
LSD/sig	8.29	P≤0.01	ns	ns
☑ Plant: time of ear emergence from boot (Julian days)				
Mean	251.20	252.00	254.70	254.30
Std. Deviation	0.29	0.00	0.60	1.10
LSD/sig	1.9	ns	P≤0.01	P≤0.01
☑ Grain: test weight (kg/hl)				
Mean	79.40	78.93	76.27	76.93
Std. Deviation	0.40	0.46	0.90	0.83
LSD/sig	1.14	ns	P≤0.01	P≤0.01
☐ Grain: screenings, grain through a 2mm sieve (%)				
Mean	2.41	4.65	7.05	4.96
Std. Deviation	0.36	0.74	3.85	0.39
LSD/sig	4.6	ns	P≤0.01	ns

Prior Applications and Sales

Nil.

Description: **Gil Hollamby,** Williamstown, SA.

Details of Application

Application Number	2007/001
Variety Name	'LS005A01'
Genus Species	*Leucospermum cuneiforme*
Common Name	Wart-stemmed Pincushion
Synonym	Nil
Accepted Date	25 Jan 2007
Applicant	Proteaflora Enterprises Pty Ltd, Monbulk, VIC
Agent	N/A
Qualified Person	Paul Armitage

Details of Comparative Trial

Location	Monbulk, VIC.
Descriptor	Leucospermum (*Leucospermum*) TG/128/3
Period	Feb 2006- Nov 2007.
Conditions	Plants propagated by cuttings, potted to 14cm pots with soiless media. Fed by CRF fertilisers. Grown in outdoor nursery conditions. Plants pinched in Dec 2006.
Trial Design	15 plants of each variety arranged in randomised design.
Measurements	From 10 plants selected at random from each variety.
RHS Chart - edition	2001.

Origin and Breeding

Open pollination of *Leucospermum cuneiforme* 'Goldie'. The putative pollen parent is *Leucospermum cuneiforme* 'Mardi Gras Petite', plants of which were adjacent to the maternal parent. The seed parent is characterised by erect to spreading growth habit, yellow flowers and late flowering season. The putative pollen parent is characterised by erect to spreading habit, orange flowers and an early flowering season. 'LS005A01' was selected from 4 seedlings originating from the 'Goldie' cross. The candidate was selected on the basis of its erect to spreading habit, medium to late flowering season, high flower number and yellow-orange flower colour. Breeder: Sue Mathews, Proteaflora Enterprises Pty Ltd, Monbulk, VIC.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	erect to spreading
Plant	lignotuber	present
Leaf	position of broadest part	above middle
Leaf	shape of base	acute
Leaf	petiole	absent
Flowering branch	clustering of fully developed flower heads	sometimes present
Flower head	texture of involucral bract	cartilaginous
Floret	colour of apex of bud	greyish
Floret	attitude of basal part of style in relation to receptacle	oblique
Floret	colour of middle part of style	yellow
Floret	shape of pollen presenter in lateral view	triangular

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Mardi Gras Petite'	Putative pollen parent. Orange flowered early-mid season *L.cuneiflorme* variety.
'Goldie'	Seed parent. Late flowering variety with yellow inflorescences.

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'LS005A01'	'Goldie'	'Mardi Gras Petite'
☐ *Plant: growth habit	erect to spreading	erect to spreading	erect to spreading
☐ Plant: height	medium	medium	medium
☐ Plant: diameter	medium	medium	medium
☐ Plant: density of foliage	medium to dense	medium	medium
☐ *Plant: lignotuber	present	present	present
☐ Leaf: blade always upright	absent	absent	absent
☐ Leaf: predominant attitude in relation to branch	oblique	oblique	oblique
☑ Leaf: length	short to medium	medium to long	short to medium
☐ Leaf: width	narrow to medium	narrow	narrow to medium
☐ *Leaf: position of broadest part	above middle	above middle	above middle
☑ *Leaf: shape of apex	acute	truncate	obtuse
☐ *Leaf: shape of base	acute	acute	acute
☐ Leaf: shape in cross section	more or less straight	more or less straight	more or less straight
☐ Leaf: colour	green	green	green
☐ Leaf: pubescence of blade	inconspicuous	inconspicuous	inconspicuous
☐ *Leaf: incisions on distal part	present	present	present
☑ *Leaf: number of incisions on distal part	very few to few	few	medium to many
☑ *Leaf: depth of incisions on distal part	shallow	deep	medium
☐ Leaf: undulation of margin	absent	absent	absent
☐ Leaf: conspicuous colour of margin	greenish	greenish	greenish
☐ Leaf: fringe on margin	absent	absent	absent
☐ *Leaf: petiole	absent	absent	absent
☐ Flowering branch: length	short to medium	short to medium	short to medium
☐ Flowering branch: thickness	thin to medium	thin to medium	thin to medium
☐ Flowering branch: rigidity	medium to strong	medium to strong	medium to strong
☐ Flowering branch: pubescence	conspicuous	conspicuous	conspicuous

*Flowering branch: clustering of fully developed flower heads	sometimes present	sometimes present	sometimes present
Flowering branch: number of fully developed flower heads per cluster	2 to 3	2 to 3	2 to 3
Flower head: length of narrowed basal part	medium	medium	short
*Flower head: length	short to medium	medium to long	short to medium
*Flower head: diameter	small to medium	medium	small to medium
*Flower head: predominant colour	yellow	yellow	orange
*Flower head: texture of involucral bract	cartilaginous	cartilaginous	cartilaginous
Flower head: pubescence of involucral bract	conspicuous	conspicuous	conspicuous
Flower head: length of floret bract	medium	medium	medium
Flower head: width of floret bract	medium	medium	medium
Flower head: colour of apical part of floret bract	reddish	greenish	reddish
Flower head: fringe on apical margin of floret bract	present	present	present
*Flower head: diameter of perianth mass	small to medium	small to medium	small
Floret: length of perianth	medium	medium	medium
Floret: pubescence on apex of bud	conspicuous	conspicuous	conspicuous
*Floret: colour of apex of bud	greyish	greyish	greyish
*Floret: colour of perianth below apex of bud	orange	yellow	orange
*Floret: colour of rolled up perianth segments	orange red	yellow	red
Floret: intensity of colour of rolled up perianth segments	medium	medium	medium
Floret: length of style	medium	medium	medium
Floret: degree of curvature of style	weak	weak	weak
Floret: thickness of style	medium	medium to thick	medium
*Floret: attitude of basal part of style in relation to receptacle	oblique	oblique	oblique

☐ *Floret: colour of middle part of style	yellow	yellow	yellow
☐ Floret: intensity of colour of middle part of style	medium	medium	medium
☐ Floret: length of pollen presenter	medium	medium	medium to long
☐ *Floret: shape of pollen presenter in lateral view	triangular	triangular	triangular
☑ Floret: colour of pollen presenter	orange red	orange	orange
☐ Floret: intensity of colour of pollen presenter	light to medium	light to medium	light to medium
☑ *Time of: flowering	medium to late	late	early to medium

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	**'LS005A01'**	**'Goldie'**	**'Mardi Gras Petite'**
☑ Floret: colour of rolled perianth	RHS34A	RHS17A	RHS45A
☑ Floret: colour of middle of style	RHS22A	RHS21B	RHS32A
☑ Floret: colour of pollen presenter	RHS32B	RHS21C	RHS42A
☑ Leaf: shape of blade	ovate	lanceolate	cuneate

Statistical Table

Organ/Plant Part: Context	**'LS005A01'**	**'Goldie'**	**'Mardi Gras Petite'**
☑ Leaf: length (mm)			
Mean	65.40	80.90	66.20
Std. Deviation	10.13	7.81	6.94
LSD/sig	10.41	P≤0.01	ns
☑ Plant: number of inflorescences			
Mean	6.50	3.60	4.90
Std. Deviation	0.71	1.26	1.29
LSD/sig	1.39	P≤0.01	P≤0.01

Prior Applications and Sales

Nil.

Description: **Paul Armitage,** Proteaflora Enterprises Pty Ltd, Monbulk, VIC.

Details of Application

Application Number	2005/289
Variety Name	'DOW20'
Genus Species	*Waterhousea floribunda*
Common Name	Weeping Lilly Pilly
Synonym	Nil
Accepted Date	29 Apr 2006
Applicant	Downes Wholesale Nursery Pty Ltd, Rossmore, NSW
Agent	Ozbreed Pty Ltd, Clarendon, NSW
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Theresa Park, NSW.
Descriptor	Lilly Pilly (*Acmena smithii*/*Syzygium* sp) PBR LILL.
Period	Summer - autumn 2007.
Conditions	Trial conducted in open beds, plants propagated from cuttings, planted into 300mm pots filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random.
RHS Chart - edition	1995.

Origin and Breeding

Seedling selection: seed parent *Waterhousea floribunda*. The seed parent is characterised by a medium weeping growth habit, brown new stem colour, reddish immature leaf colour and weak undulation of the leaf margin. In 2000, 2 to 3 thousand seedlings arising from open-pollianted seed of *W. floribunda* were grown in an open bed. In 2001, 4 seedlings were selected due to their new growth, rippled leaf margin, green bark colour and strongly weeping habits. Finally, in 2001 a single seedling was selected due to its most extreme differences to the parent form. Selection took place in Tuckombil, NSW in 2001. Selection criteria: compact strongly weeping plant growth habit, green new stem colour, green immature leaf colour and strong undulation of the leaf margin. Propagation: vegetative cuttings were found to be uniform and stable. Breeders: Greg Hellyar and Stuart Nolan , Tuckombil, NSW.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Leaf	variegation	absent
Plant	growth habit	spreading

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
W. floribunda	parent variety used as DOW20 is the first variety of the species

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'DOW20'	*W. floribunda*
☐ Plant: growth habit	spreading	spreading
☐ Plant: height	medium to tall	tall
☐ Plant: branch density	medium	medium
☑ Stem: branch angle	horizontal	broad acute
☐ Stem: internode length	medium	medium
☐ Stem: basal diameter	medium	medium
☑ Stem: colour of mature stem (RHS colour chart)	199D	199D and 200A (scars)
☑ Stem: colour of new growth (RHS colour chart)	144B	177A
☐ Leaf: blade length	medium to long	medium
☐ Leaf: blade width	medium	medium
☐ Leaf: petiole length	medium	medium
☑ Leaf: shape of blade	narrow elliptic	elliptic
☐ Leaf: shape of apex	acuminate	acuminate
☐ Leaf: shape of base	cuneate	cuneate
☐ Leaf: glossiness	medium	medium
☐ Leaf: shape of cross section	concave	flat to concave
☐ Leaf: shape of longitudinal section	flat	flat
☐ Leaf: stiffness	very weak to weak	very weak to weak
☐ Leaf: prominence of midrib on lower surface	prominent	prominent
☐ Mature leaf: primary colour of upper side (RHS colour chart)	147A	147A
☑ Mature leaf: primary colour of lower side (RHS colour chart)	ca 147A	ca 146A
☑ Partly mature leaf: primary colour of upper side (RHS colour chart)	144A	ca 146A
☑ Partly mature leaf: primary colour of lower side (RHS colour chart)	144A	146C
☑ Newly emerged: upper side (RHS colour chart)	144A	174A
☐ Leaf: variegation	absent	absent
☑ Leaf: petiole colour (RHS colour chart)	153D	174B

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'DOW20'	*W. floribunda*
☑ Plant: degree of weeping	strong	medium
☑ Leaf: undulation of margin	strong	weak

☐ Mature leaf: colour of midrib (RHS)	154D	154D
☑ Leaf: anthocyanin coloration of midrib on lower side	absent	present

Statistical Table

Organ/Plant Part: Context	**'DOW20'**	***W. floribunda***
☐ Leaf: length (mm)		
Mean	86.70	80.90
Std. Deviation	14.20	19.70
LSD/sig	19.62	ns
☐ Leaf: width (mm)		
Mean	20.40	20.50
Std. Deviation	2.90	3.90
LSD/sig	3.92	ns

Prior Applications and Sales

Nil.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW

Details of Application

Application Number	2007/117
Variety Name	'Axe'
Genus Species	*Triticum aestivum*
Common Name	Wheat
Synonym	Nil
Accepted Date	18 May 2007
Applicant	Australian Grain Technologies Pty Ltd, Glen Osmond, SA
Agent	N/A
Qualified Person	Gil Hollamby

Details of Comparative Trial

Location	Mintaro, South Australia.
Descriptor	Wheat (*Triticum aestivum*) TG/3/11.
Period	2007.
Conditions	The trial was grown in a redbrown earth soil which had been pasture in 2006 and wheat in 2005. The area was sprayed with Roundup Power Max (1.2L/ha) + Goal CT (75ml/ha) on 24 May 2007 and direct drilled at 2-4cm in slightly moist conditions on 25 May at 200 plants/m2 and with 90kg/ha DAP and 80kg/ha Urea. During the winter months moisture was adequate and the trial grew well. In crop weeds were controlled with 2,4-D amine 625(1.5l/ha) on 6 Sep. Spring was dry and some moisture stress occurred so varieties were shorter in stature than expected. Harvest took place on 14 December about two weeks earlier than normal. There were no diseases of note. A similar trial was planted at Roseworthy.
Trial Design	Randomised Block Design of 3 blocks and 56 entries consisting of comparators and potential candidates. Sown in 12 ranges of 14 plots wide, block 1 being in ranges 1 to 4 and so on. Plots were 1.25m wide (5 rows) and 3.2m long. There were approx. 1000 plants per plot.
Measurements	Heading times were recorded on the same trial planted at Roseworthy 2007, but this trial later was abandoned due to a heavy infestation of Crown Rot. All other measurements and observations were recorded on plant samples taken from the Mintaro trial. At anthesis 5 primary tillers were sampled from each plot in each replicate and flag leaf measurements made. Glaucosity and leaf angle was observed at this time. After maturity plant heights to the top of the awns were recorded at 10 random locations in replicate 2 and 3 only. Twenty heads were also sampled at random from each plot in replicates 2 and 3 for head descriptions and measurements. Measurements were performed on 10 intact heads per block. Statistical analyses were completed using GENSTAT software.
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: The cross that produced RAC1192, coded CO5641, was completed in 1999. Two F_1 plants, each with the pedigree RAC875//Excalibur/Kukri, were intercrossed. In total 59 doubled haploids were produced from the resultant F_1s. Seed was multiplied over winter at Roseworthy Campus, Roseworthy, in 2000. This, and all subsequent seed was multiplied by self pollination. Doubled haploids were grown in 3 field nurseries in southern Australia in 2001 and 8 in 2002. The lines were assessed for rust resistance, plant type, heading date, end use quality and grain yield. Each of the lines were also assessed at Cobbitty (NSW) and Horsham (Vic) for rust resistance. One elite doubled haploid, CO5641-AH00 was identified and renamed RAC1192. RAC1192 was included in the Stage 3 and 4 testing regimes of Australian Grain Technologies in 2003-2006. The disease resistance, abiotic stress tolerance, end use quality and grain yield of RAC1192 was assessed in WA, SA, Vic, NSW and QLD as part of its inclusion in Stage 3 and 4 trials. Samples were submitted to AWB for quality testing, and RAC1192 received an AH classification. RAC1192 was included in the Stage 4 testing of AGT and the NVT system in 2007. Breeder: Mr Haydn Kuchel, Dr Stephen Jefferies and Mr Gil Hollamby, Australian Grain Technologies.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Ear	distribution of awns	fully awned
Ear	colour	white
Plant	time of ear emergence	≤ 254 Julian days

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Kukri'	early, grown in area of adaptation
'Silverstar'	very early
'Excalibur'	early variety in the area of adaptation

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety	State of Expression in Comparator Variety	Comments
'Young'	Flag leaf blade width	very wide (21.2 mm)	narrow(14.6mm)	LSD=2.2 (P=1%)
'Wyalkatchem'	Flag leaf blade width (2006)	very wide(17.6mm)	medium to narrow(13.2)	LSD=1.4 (P=1%)
'H45'	Leaves post anthesis stripe rust reaction	Moderately resistant	very susceptible	
'H45'	Flag leaf blade width (2006 trial)	very wide(18.0mm)	narrow(13.2mm)	LSD=1.4 (p=1%)

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Axe'	'Excalibur'	'Kukri'	'Silverstar'
☐ *Plant: growth habit	semi-erect to intermediate	intermediate	semi-erect	semi-erect
☑ Flag leaf: anthocyanin colouration of auricles	absent or very weak	absent or very weak	strong	absent or very weak
☑ Plant: frequency of plants with recurved flag leaves	very low to low	high	medium to high	high
☑ *Flag leaf: glaucosity of sheath	medium		weak	weak
☑ *Ear: glaucosity	medium		weak to medium	weak
☐ Culm: glaucosity of neck	medium		weak to medium	weak
☐ *Straw: pith in cross section	thin	very thin to thin	thin	thin
☐ *Ear: shape in profile	tapering	tapering	tapering	tapering
☑ *Ear: density	medium	medium	lax	lax to medium
☐ *Awns or scurs: presence	awns present	awns present	awns present	awns present
☐ *Awns of scurs at tip of ear: length	medium	medium	medium	medium
☐ *Ear: colour	white	white	white	white
☑ Lower glume: shoulder width	medium	broad	medium	narrow
☑ Lower glume: shoulder shape	straight to elevated	straight to elevated	elevated	sloping
☑ Lower glume: beak length	short to medium	medium	long	medium to long
☐ Lower glume: beak shape	straight to slightly curved	straight	moderately curved	slightly curved
☐ Lower glume: extent of internal hair	very weak	weak	weak to medium	medium
☐ Lowest lemma: beak shape	slightly curved to moderately curved	straight to slightly curved	slightly curved	straight
☐ *Grain: colour	white	white	white	white
☑ Grain: colouration with phenol	none or very light to light	dark	medium	very dark
☐ *Seasonal type:	spring type	spring type	spring type	spring type

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Axe'	'Excalibur'	'Kukri'	'Silverstar'
☐ Whole plant post anthesis: Stem rust reaction	moderately susceptible	Susceptible	resistant	
☐ Whole plant post anthesis: Stripe rust	moderately	susceptible	resistant	moderately

reaction	resistant			susceptible
☑ Leaves post anthesis: Leaf rust reaction (Lr37 virulent race)	moderately resistant	susceptible	susceptible	
☐ Glutenin composition: allele expression at GluA1	a	mixed a & b	a	a
☑ Glutenin composition: allele expression at GluB1	i	i	al	i
☑ Glutenin composition: allele expression at GluD1	d	a	d	mixed a & d
☐ Glutenin composition: allele expression at GluA3	c	mixed b & c	d	mixed b & c
☑ Glutenin composition: allele expression at GluB3	b	b	h	h
☑ Glutenin composition: allele expression at GluD3	b	a	b	b

Statistical Table

Organ/Plant Part: Context	**'Axe'**	**'Excalibur'**	**'Kukri'**	**'Silverstar'**
☑ Flag leaf: length (mm)				
Mean	179.10	n/a	198.90	246.10
Std. Deviation	43.30	n/a	35.50	31.30
LSD/sig	41.3	n/a	ns	P≤0.01
☐ Flag leaf: blade width (mm)				
Mean	21.20	n/a	17.40	16.80
Std. Deviation	2.10	n/a	2.60	1.20
LSD/sig	2.2	n/a	P≤0.01	P≤0.01
☑ Flag leaf: sheath length (mm)				
Mean	161.30	n/a	183.40	177.80
Std. Deviation	10.60	n/a	8.00	7.00
LSD/sig	15.1	n/a	P≤0.01	P≤0.01
☑ Plant: time of ear emergence (Julian days)				
Mean	248.10	253.30	254.00	248.70
Std. Deviation	2.00	0.58	1.00	2.30
LSD/sig	2.1	P≤0.01	P≤0.01	ns
☑ Whole plant: height (cm)				
Mean	82.00	83.90	91.50	88.00
Std. Deviation	3.20	2.35	3.10	3.40
LSD/sig	6.6	ns	P≤0.01	ns
☑ Ear: length without awns (mm)				
Mean	93.40	100.00	106.40	106.10
Std. Deviation	5.60	5.31	10.56	7.61
LSD/sig	8.4	P≤0.01	P≤0.01	P≤0.01
☑ Ear: rachis internode (mm)				
Mean	4.24	4.19	4.54	4.73

Std. Deviation	0.25	0.18	0.32	0.28
LSD/sig	0.35	ns	ns	P≤0.01

Prior Applications and Sales

Nil.

Description: **Gil Hollamby,** Williamstown, SA.

Details of Application

Application Number	2006/302
Variety Name	'Gladius'
Genus Species	*Triticum aestivum*
Common Name	Wheat
Synonym	Nil
Accepted Date	17 Jan 2007
Applicant	Australian Grain Technologies Pty Ltd, Glen Osmond, SA
Agent	N/A
Qualified Person	Gil Hollamby

Details of Comparative Trial

Location	Mintaro, South Australia.
Descriptor	Wheat (*Triticum aestivum*) TG/3/11.
Period	2007.
Conditions	The trial was grown in a redbrown earth soil which had been pasture in 2006 and wheat in 2005. The area was sprayed with Roundup Power Max (1.2L/ha) + Goal CT (75ml/ha) on 24 May 2007 and direct drilled at 2-4cm in slightly moist conditions on 25 May at 200 plants/m2 and with 90kg/ha DAP and 80kg/ha Urea. During the winter months moisture was adequate and the trial grew well. In crop weeds were controlled with 2,4-D amine 625(1.5l/ha) on 6 Sep. Spring was dry and some moisture stress occurred so varieties were shorter in stature than expected. Harvest took place on 14 Dec about two weeks earlier than normal. There were no diseases of note. A similar trial was planted at Roseworthy.
Trial Design	Randomised Block Design of 3 blocks and 56 entries consisting of comparators and potential candidates. Sown in 12 ranges of 14 plots wide, block 1 being in ranges 1 to 4 and so on. Plots were 1.25m wide (5 rows) and 3.2m long. There were approx. 1000 plants per plot.
Measurements	Heading times were recorded on the same trial planted at Roseworthy 2007, but this trial later was abandoned due to a heavy infestation of Crown Rot. All other measurements and observations were recorded on plant samples taken from the Mintaro trial. At anthesis 5 primary tillers were sampled from each plot in each replicate and flag leaf measurements made. Glaucosity and leaf angle was observed at this time. After maturity plant heights to the top of the awns were recorded at 10 random locations in replicate 2 and 3 only. Twenty heads were also sampled at random from each plot in replicates 2 and 3 for head descriptions and measurements. Measurements were performed on 10 intact heads per block. Statistical analyses were completed using GENSTAT software.
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: A complex crossing strategy involving the parents of RAC1262 was completed in 2001. The final cross, coded CO5693, was between an F_2 plant with the pedigree RAC875/Krichauff//Excalibur/Kukri/3/RAC875/Krichauff and a doubled haploid with pedigree RAC875//Excalibur/Kukri. In total 181 doubled haploids were produced from this cross. Seed was multiplied over summer at Roseworthy Campus, Roseworthy in 2002/3. This, and all subsequent generations were multiplied by self pollination. Doubled haploids were grown in five field nurseries in South Australia and assessed for rust resistance, plant type, heading date and grain yield. An elite doubled haploid, CO5693-E002 was identified and renamed RAC1262. RAC1262 was included in the Stage 3 testing regime of Australian Grain Technologies, undergoing grain yield evaluation at 16 locations across Australia. The disease resistance, abiotic stress tolerance, and end use quality of RAC1262 was also assessed as part of its inclusion in Stage 3 trials. RAC1262 was then included in AGT Stage 4 trials in 2005, and its grain yield evaluated at 37 sites across Australia. Samples were submitted to AWB for quality testing, and RAC1262 has received a preliminary APW classification. RAC1262 was included in the Stage 4 testing of AGT and the NVT system in 2006 and 2007. Breeders: Mr Haydn Kuchel, Dr Stephen Jefferies, Mr Gil Hollamby (Australian Grain Technologies) in collaboration with Dr Neil Howes (formerly SARDI).

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Ear	distribution of awns	fully awned
Ear	colour	white
Plant	time of ear emergence	250 to 258 Julian days

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
‘Correll’	similar adaptation.
‘Yitpi’	widely grown hard wheat in the area.
‘Kukri’	grown in the area of adaptation.

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics	State of Expression in Candidate Variety		State of Expression in Comparator Variety	Comments
‘Krichauff’	flag leaf	blade width (2006 data)	narrow(12.0mm)	wide(16.3mm)	LSD=1.4mm (P=1%)
‘Excalibur’	grain	protein composition	pure	mixed	
‘Young’	flag leaf	blade width	narrow(14.6mm)	wide(18.0mm)	LSD=2.2(P=1%)

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Gladius'	'Correll'	'Kukri'	'Yitpi'
☐ *Plant: growth habit	semi-erect	intermediate	semi-erect	intermediate
☑ Flag leaf: anthocyanin colouration of auricles	absent or very weak	absent or very weak	strong	absent or very weak
☑ Plant: frequency of plants with recurved flag leaves	low	medium	medium to high	high to very high
☑ *Flag leaf: glaucosity of sheath	very strong	strong	weak	medium
☑ *Ear: glaucosity	medium	strong	weak to medium	weak to medium
☑ Culm: glaucosity of neck	strong	strong	weak to medium	medium to strong
☐ *Straw: pith in cross section	thin	thin	thin	thin
☐ *Ear: shape in profile	parallel sided	parallel sided	tapering	parallel sided
☐ *Ear: density	lax to medium	medium	lax	medium
☐ *Awns or scurs: presence	awns present	awns present	awns present	awns present
☐ *Awns of scurs at tip of ear: length	medium	medium to long	medium	medium
☐ *Ear: colour	white	white	white	white
☑ Lower glume: shoulder width	broad	broad	medium	broad
☑ Lower glume: shoulder shape	slightly sloping to straight	straight	elevated	slightly sloping
☑ Lower glume: beak length	short	short	long	medium
☐ Lower glume: beak shape	straight	straight	moderately curved	straight
☐ Lower glume: extent of internal hair	weak	medium	weak to medium	very weak to weak
☐ Lowest lemma: beak shape	slightly curved	straight to slightly curved	slightly curved	straight
☐ *Grain: colour	white	white	white	white
☐ Grain: colouration with phenol	dark	dark to very dark	medium	very dark
☐ *Seasonal type:	spring type	spring type	spring type	spring type

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Gladius'	'Correll'	'Kukri'	'Yitpi'
☐ Whole plant post anthesis: Stripe rust reaction	moderately resistant	mod. susceptible to mod. resistant		mod. susceptible to mod. resistant
☐ Whole plant post anthesis: stem rust reaction	moderately resistant	mod susceptible to mod resistant	resistant	Susceptible
☐ Leaves post anthesis: Leaf rust reaction (Lr37 virulent race)	moderately susceptible	moderately susceptible	resistant	moderately susceptible
☐ Glutenin composition: allele expression at GluA1	a	mixed a & b	a	a

☑ Glutenin composition: allele expression at GluB1	u	u	al	u
☐ Glutenin composition: allele expression at GluD1	d	d	d	d
☐ Glutenin composition: allele expression at GluA3	c	mixed c & d	d	c
☐ Glutenin composition: allele expression at GluB3	b	mixed b & h	h	h
☑ Glutenin composition: allele expression at GluD3	a	mixed b & c	b	c

Statistical Table

Organ/Plant Part: Context	'Gladius'	'Correll'	'Kukri'	'Yitpi'
☑ Flag leaf: blade length (mm)				
Mean	166.20	172.10	198.90	197.90
Std. Deviation	28.90	21.70	35.50	36.10
LSD/sig	41.4	ns	P≤0.01	ns
☐ Flag leaf: blade width (mm)				
Mean	18.00	16.90	17.40	16.80
Std. Deviation	1.60	1.40	2.70	1.20
Lsd/sig	2.2	ns	ns	ns
☑ Flag leaf: sheath length (mm)				
Mean	155.50	170.30	183.40	189.20
Std. Deviation	6.30	8.10	8.00	10.70
LSD/sig	15.1	ns	P≤0.01	P≤0.01
☐ Plant: time of ear emergence (Julian days)				
Mean	254.00	255.70	254.00	256.00
Std. Deviation	0.00	0.60	1.00	0.00
LSD/sig	2.1	ns	ns	ns
☐ Whole plant: height (cm)				
Mean	82.00	85.60	91.50	89.20
Std. Deviation	4.10	3.30	3.10	2.50
LSD/sig	6.6	ns	P≤0.01	P≤0.01
☑ Ear: length without awns (mm)				
Mean	90.90	98.30	106.40	94.50
Std. Deviation	4.60	5.30	10.56	6.21
LSD/sig	8.4	ns	P≤0.01	ns
☐ Ear: rachis internode (mm)				
Mean	4.30	4.27	4.54	4.18
Std. Deviation	0.21	0.18	0.32	0.22
LSD/sig	0.35	ns	ns	ns

Prior Applications and Sales

Nil.

Description: **Gil Hollamby,** Williamstown, SA.

Details of Application

Application Number	2007/322
Variety Name	'Espada'
Genus Species	*Triticum aestivum*
Common Name	Wheat
Synonym	Nil
Accepted Date	17 Jan 2008
Applicant	Australian Grain Technologies Pty Ltd, Glen Osmond, SA
Agent	N/A
Qualified Person	Gil Hollamby

Details of Comparative Trial

Location	Mintaro, South Australia.
Descriptor	Wheat (*Triticum aestivum*) TG/3/11.
Period	2007.
Conditions	The trial was grown in a redbrown earth soil which had been pasture in 2006 and wheat in 2005. The area was sprayed with Roundup Power Max (1.2L/ha) + Goal CT (75ml/ha) on 24 May 2007 and direct drilled at 2-4cm in slightly moist conditions on 25 May at 200 plants/m2 and with 90kg/ha DAP and 80kg/ha Urea. During the winter months moisture was adequate and the trial grew well. In crop weeds were controlled with 2,4-D amine 625(1.5l/ha) on 6 Sep. Spring was dry and some moisture stress occurred so varieties were shorter in stature than expected. Harvest took place on 14 Dec about two weeks earlier than normal. There were no diseases of note. A similar trial was planted at Roseworthy.
Trial Design	Randomised Block Design of 3 blocks and 56 entries consisting of comparators and potential candidates. Sown in 12 ranges of 14 plots wide, block 1 being in ranges 1 to 4 and so on. Plots were 1.25m wide (5 rows) and 3.2m long. There were approx. 1000 plants per plot.
Measurements	Heading times were recorded on the same trial planted at Roseworthy 2007, but this trial later was abandoned due to a heavy infestation of Crown Rot. All other measurements and observations were recorded on plant samples taken from the Mintaro trial. At anthesis 5 primary tillers were sampled from each plot in each replicate and flag leaf measurements made. Glaucosity and leaf angle was observed at this time. After maturity plant heights to the top of the awns were recorded at 10 random locations in replicate 2 and 3 only. Twenty heads were also sampled at random from each plot in replicates 2 and 3 for head descriptions and measurements. Measurements were performed on 10 intact heads per block. Statistical analyses were completed using GENSTAT software.
RHS Chart - edition	N/A

Origin and Breeding

Controlled pollination: a complex crossing strategy involving the parents of RAC1263 was completed in 2001. The final cross, coded CO5693, was between an F_2 plant with pedigree RAC875/Krichauff//Excalibur/Kukri/3/RAC875/Kricauff and a doubled haploid with pedigree RAC875//Excalibur/Kukri. In total 181 doubled haploids were produced from this cross. Seed was multiplied over summer at Roseworthy Campus, Roseworthy, in 2002/3. This and all subsequent seed was multiplied by self pollination. Doubled haploids were grown in five field nurseries in South Australia and assessed for rust resistance, plant type, heading date and grain yield. An elite doubled haploid, CO5693-E010 was identified and renamed RAC1263. RAC1263 was included in the Stage 3 testing regime of Australian Grain Technologies, undergoing grain yield evaluation at 16 locations across Australia. The disease resistance, abiotic stress tolerance, and end use quality of RAC1263 was also assessed as part of its inclusion in Stage 3 trials. RAC1263 was then included in AGT Stage 4 trials in 2005, and its grain yield evaluated at 37 sites across Australia. Samples were submitted to AWB for quality testing, and RAC1263 has received an APW classification. RAC1263 was included in the Stage 4 testing of AGT and the NVT system in 2007. Breeders: Mr Haydn Kuchel, Dr Stephen Jefferies, Mr Gil Hollamby (Australian Grain Technologies) in collaboration with Dr Neil Howes (formerly SARDI).

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	time of ear emergence	252 to 258 Julian days
Plant	post anthesis glaucosity	strong to very strong
Ear	distribution of awns	fully awned
Ear	colour	white

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
‘Correll’	Similar area of adaptation.
‘Gladius’	Sibling.

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
‘Yitpi’	Plant	post anthesis glaucosity	very strong	medium
‘Wyalkatchem’	Plant	post anthesis glaucosity	very strong	medium
‘Kukri’	Plant	post anthesis glaucosity	very strong	weak to medium

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Espada'	'Correll'	'Gladius'
☐ *Plant: growth habit	intermediate to semi-prostrate	intermediate	semi-erect
☐ Flag leaf: anthocyanin colouration of auricles	absent or very weak	absent or very weak	absent or very weak
☐ Plant: frequency of plants with recurved flag leaves	absent or very low	medium	low
☑ *Flag leaf: glaucosity of sheath	very strong	strong	very strong
☑ *Ear: glaucosity	medium	strong	medium
☑ Culm: glaucosity of neck	very strong	strong	strong
☑ *Straw: pith in cross section	medium	thin	thin
☐ *Ear: shape in profile	parallel sided	parallel sided	parallel sided
☐ *Ear: density	medium to dense	medium	lax to medium
☐ *Awns or scurs: presence	awns present	awns present	awns present
☐ *Awns of scurs at tip of ear: length	medium	medium to long	medium
☐ *Ear: colour	white	white	white
☑ Lower glume: shoulder width	medium	broad	broad
☐ Lower glume: shoulder shape	slightly sloping to straight	straight	slightly sloping to straight
☐ Lower glume: beak length	very short to short	short	short
☐ Lower glume: beak shape	straight	straight	straight
☐ Lower glume: extent of internal hair	very weak	medium	weak
☐ Lowest lemma: beak shape	straight	straight to slightly curved	slightly curved
☐ *Grain: colour	white	white	white
☐ Grain: colouration with phenol	dark	dark to very dark	dark
☐ *Seasonal type:	spring type	spring type	spring type

Characteristics Additional to the Descriptor/TG

Organ/Plant Part: Context	'Espada'	'Correll'	'Gladius'
☑ Leaves: Reaction to stripe rust pathotype 134E16A+	resistant	moderately susceptible	
☑ Glutenin composition : allele expression at GluA1	a	Mixed a & b	a
☐ Leaves post anthesis: Leaf rust reaction (Lr37 virulent race)	moderately susceptible	moderately susceptible	moderately susceptible
☐ Glutenin composition: allele expression at GluB1	u	u	u
☐ Glutenin composition: allele expression at GluD1	d	d	d
☐ Glutenin composition: allele expression at GluA3	d	mixed c & d	c
☐ Glutenin composition: allele expression	b	mixed b & h	b

at GluB3			
☑ Glutenin composition: allele expression at GluD3	b	mixed b & c	a
☐ Whole plant post anthesis: stem rust reaction	moderately resistant		moderately resistant
☐ Whole plant post anthesis: Stripe rust reaction	moderately resistant	mod. susceptible to mod. resistant	moderately resistant

Statistical Table

Organ/Plant Part: Context	**'Espada'**	**'Correll'**	**'Gladius'**
☐ Flag leaf: blade length (mm)			
Mean	153.00	172.10	166.20
Std. Deviation	21.20	21.70	28.90
LSD/sig	41.3	ns	ns
☐ Flag leaf: blade width (mm)			
Mean	18.00	16.90	18.00
Std. Deviation	1.10	2.20	1.60
LSD/sig	2.2	ns	ns
☑ Flag leaf: sheath length (mm)			
Mean	149.70	170.30	155.50
Std. Deviation	7.40	14.10	6.30
LSD/sig	15.1	P≤0.01	ns
☐ Plant: time of ear emergence (Julian days)			
Mean	255.50	255.70	254.00
Std. Deviation	0.60	0.60	0.00
LSD/sig	2.1	ns	ns
☑ Plant: height (cm)			
Mean	78.90	85.60	82.00
Std. Deviation	3.40	3.30	4.10
LSD/sig	6.6	P≤0.01	ns
☑ Ear: length without awns (mm)			
Mean	88.50	98.30	90.90
Std. Deviation	6.40	5.30	4.60
LSD/sig	8.4	P≤0.01	ns
☑ Ear: rachis internode length (mm)			
Mean	3.92	4.27	4.30
Std. Deviation	0.22	0.18	0.21
LSD/sig	0.35	P≤0.01	P≤0.01

Prior Applications and Sales

Nil.

Description: **Gil Hollamby,** Williamstown, SA.

Details of Application

Application Number	2006/222
Variety Name	'Jedda's Dream'
Genus Species	*Agonis flexuosa*
Common Name	Willow Myrtle
Synonym	Nil
Accepted Date	15 Aug 2006
Applicant	James F Koppman and Jaqueline A Koppman, Falls Creek, NSW
Agent	N/A
Qualified Person	Ian Paananen

Details of Comparative Trial

Location	Falls Creek , NSW.
Descriptor	Willow Peppermint (*Agonis flexuosa*) PBR AGON
Period	Summer 2006-autumn 2007.
Conditions	Trial conducted in open beds, plants propagated from cuttings, planted into 200mm pots filled with soilless potting mix, nutrition maintained with slow release and liquid fertilisers, irrigation by overhead watering, pest and disease treatments not required.
Trial Design	Fifteen pots of each variety arranged in a completely randomised design.
Measurements	From ten plants at random.
RHS Chart - edition	2001.

Origin and Breeding

Spontaneous mutation: 'Jervis Bay Afterdark'. The parent is characterised by a tall plant height, upright to weeping growth habit and medium basal branching. Selection took place in Tumbi Umbi, NSW, and Falls Creek, NSW in 1999. Selection criteria: short plant height, bushy plant growth habit. Propagation: vegetative cuttings and micropropagation were found to be uniform and stable. Breeders: James and Jacquie Koppman, Falls Creek , NSW.

Choice of Comparators Characteristics used for grouping varieties to identify the most similar Variety of Common Knowledge

Organ/Plant Part	Context	State of Expression in Group of Varieties
Plant	growth habit	upright
Leaf blade	variegation	absent
Leaf blade	colour of mature leaf	greyed-purple

Most Similar Varieties of Common Knowledge identified (VCK)

Name	Comments
'Jervis Bay Afterdark'	parent variety

Varieties of Common Knowledge identified and subsequently excluded

Variety	Distinguishing Characteristics		State of Expression in Candidate Variety	State of Expression in Comparator Variety
'Burgundy'	leaf	predominant colour	greyed purple	green
'Burgundy'	growth habit	weeping of branches	absent	present

Variety Description and Distinctness - Characteristics which distinguish the candidate from one or more of the comparators are marked with a tick.

Organ/Plant Part: Context	'Jedda's Dream'	'Jervis Bay Afterdark'
☐ Plant: growth habit	upright	upright
☐ Plant: vigour	medium	medium
☑ Plant: height	short	medium
☑ Plant: density	dense	medium
☐ Stem: inner angle of lateral shoots to main stem	acute	acute
☑ Stem: length of longest primary branch	short	medium
☐ Stem: colour of young stem (RHS colour chart)	187A	187A
☐ Stem: colour of mature stem (RHS colour chart)	165B	165B
☑ Stem: degree of basal branching	strong	weak
☐ Stem: diameter	medium	medium
☑ Leaf blade: length	short	medium
☐ Leaf blade: width	medium	medium
☐ Leaf blade: shape	lanceolate	lanceolate
☐ Leaf blade: shape of apex	acute	acute
☐ Leaf blade: shape of base	cuneate	cuneate
☐ Leaf bade: undulation of margin	absent or very weak	absent or very weak
☐ Leaf blade: cross-section	concave to flat	concave
☐ Leaf blade: curvature of longitudinal section	straight to recurved	straight to recurved
☐ Leaf blade: variegation	absent	absent
☑ Leaf blade: colour of immature leaf (RHS colour chart)	146A with 187A in the margin	ca 187A
☐ Leaf blade: colour of mature leaf (RHS colour chart)	N200A	N200A
☐ Leaf blade: glossiness	medium	medium

Statistical Table

Organ/Plant Part: Context	'Jedda's Dream'	'Jervis Bay Afterdark'
☑ Plant: height (cm)		
Mean	37.20	88.30
Std. Deviation	4.40	3.10
LSD/sig	4.94	P≤0.01
☑ Branch: length (cm)		
Mean	30.90	48.60
Std. Deviation	2.60	8.80
LSD/sig	8.32	P≤0.01
☑ Stem: diameter (mm)		
Mean	5.10	6.60
Std. Deviation	0.95	0.55
LSD/sig	1.00	P≤0.01

☑ Leaf: length (mm)		
Mean	59.50	79.70
Std. Deviation	7.40	7.50
LSD/sig	9.57	P≤0.01
☑ Leaf: width (mm)		
Mean	12.00	13.80
Std. Deviation	1.20	0.90
LSD/sig	1.33	P≤0.01

Prior Applications and Sales

Nil.

Description: **Ian Paananen**, Crop & Nursery Services, Central Coast, NSW

GRANTS

Alstroemeria hybrid

PERUVIAN LILY

‘Koncalga’[Φ]

Application No: 2006/082 Grantee: **Konst Breeding B.V.**.
Certificate No: 3416 Expiry Date: 14 December, 2027.
Agent: **David Nichols - postal address for service of notice on the applicant Konst Breeding BV**, Devon Meadows, VIC.

‘Konsacram’[Φ]

Application No: 2006/083 Grantee: **Konst Breeding B.V.**.
Certificate No: 3417 Expiry Date: 14 December, 2027.
Agent: **David Nichols - postal address for service of notice on the applicant Konst Breeding BV**, Devon Meadows, VIC.

‘Konsirak’[Φ]

Application No: 2006/080 Grantee: **Konst Breeding B.V.**.
Certificate No: 3414 Expiry Date: 14 December, 2027.
Agent: **David Nichols - postal address for service of notice on the applicant Konst Breeding BV**, Devon Meadows, VIC.

‘Konzifer’[Φ]

Application No: 2006/081 Grantee: **Konst Breeding B.V.**.
Certificate No: 3415 Expiry Date: 14 December, 2027.
Agent: **David Nichols - postal address for service of notice on the applicant Konst Breeding BV**, Devon Meadows, VIC.

‘Zalsanyx’[Φ] syn Onyx[Φ]

Application No: 2006/057 Grantee: **Van Zanten Plants B.V.**.
Certificate No: 3418 Expiry Date: 14 December, 2027.
Agent: **Ramm Botanicals Holdings Pty Ltd**, Tuggerah, NSW.

‘Zaprifabi’[Φ] syn Fabiana[Φ]

Application No: 2006/058 Grantee: **Van Zanten Plants B.V.**.
Certificate No: 3419 Expiry Date: 14 December, 2027.
Agent: **Ramm Botanicals Holdings Pty Ltd**, Tuggerah, NSW.

‘Zapriteres’[Φ] syn Theresa[Φ]

Application No: 2006/059 Grantee: **Van Zanten Plants B.V.**.
Certificate No: 3420 Expiry Date: 14 December, 2027.
Agent: **Ramm Botanicals Holdings Pty Ltd**, Tuggerah, NSW.

Arctotis fastuosa

AFRICAN DAISY

‘Archise’Φ

Application No: 2005/324 Grantee: **NuFlora International Pty Ltd**, Macquarie Fields, NSW.
Certificate No: 3401 Expiry Date: 12 October, 2027.

Avena sativa

OATS

‘Graza 51’Φ

Application No: 2004/302 Grantee: **Agriculture and Agri-Food Canada**.
Certificate No: 3403 Expiry Date: 21 November, 2027.
Agent: **Pioneer Hi-Bred Australia Pty Ltd**, Toowoomba, QLD.

‘Graza 80’Φ

Application No: 2004/301 Grantee: **Agriculture and Agri-Food Canada**.
Certificate No: 3402 Expiry Date: 21 November, 2027.
Agent: **Pioneer Hi-Bred Australia Pty Ltd**, Toowoomba, QLD.

Citrus limon

LEMON

‘3 ELS 0’Φ

Application No: 2003/278 Grantee: **Craig Robert Pressler**, Emerald, QLD.
Certificate No: 3409 Expiry Date: 25 November, 2032.

‘7 ELS 1’Φ

Application No: 2003/279 Grantee: **Craig Robert Pressler**, Emerald, QLD.
Certificate No: 3410 Expiry Date: 25 November, 2032.

‘7 ELS C3’Φ

Application No: 2003/280 Grantee: **Craig Robert Pressler**, Emerald, QLD.
Certificate No: 3411 Expiry Date: 25 November, 2032.

‘Code 3X97’Φ

Application No: 2001/172 Grantee: **Craig Robert Pressler**, Emerald, QLD.
Certificate No: 3407 Expiry Date: 25 November, 2032.

‘Code 7B97’Φ

Application No: 2001/173 Grantee: **Craig Robert Pressler**, Emerald, QLD.

Certificate No: 3408 Expiry Date: 25 November, 2032.

Clematis hybrid

CLEMATIS

‘Piilu’Φ syn Little Duckling Φ

Application No: 2004/102 Grantee: **Aili Kivistik**.
Certificate No: 3413 Expiry Date: 10 December, 2027.
Agent: **Plants Management Australia Pty Ltd**, Wonga Park, VIC.

Coprosma hybrid

MIRROR BUSH

‘Fire Burst’Φ

Application No: 2005/073 Grantee: **Richard Graeme Ware**.
Certificate No: 3422 Expiry Date: 14 December, 2027.
Agent: **Greenhills Propagation Nursery Pty Ltd**, Tynong, VIC.

Fragaria x*ananassa*

STRAWBERRY

‘Driscoll El Dorado’Φ

Application No: 2006/072 Grantee: **Driscoll Strawberry Associates, Inc**.
Certificate No: 3405 Expiry Date: 21 November, 2027.
Agent: **Phillips Ormonde & Fitzpatrick**, Melbourne, VIC.

‘Driscoll Ojai’Φ

Application No: 2006/074 Grantee: **Driscoll Strawberry Associates, Inc**.
Certificate No: 3406 Expiry Date: 21 November, 2027.
Agent: **Phillips Ormonde & Fitzpatrick**, Melbourne, VIC.

Hebe diosmifolia

HEBE

‘Ohakea’Φ

Application No: 2002/253 Grantee: **Plantlife Partnership**.
Certificate No: 3429 Expiry Date: 18 December, 2027.
Agent: **Greenhills Propagation Nursery Pty Ltd**, Tynong, VIC.

Hedysarum coronorium

SULLA

'Flamenco'[Φ]

Application No: 2006/178 Grantee: **Western Australian Agriculture Authority, University of Western Australia, Rural Industries Research and Development Corporation**.
Certificate No: 3427 Expiry Date: 18 December, 2027.
Agent: **Western Australian Agriculture Authority**, Bentley Delivery Centre, WA.

Hordeum vulgare

BARLEY

'Buloke'[Φ]

Application No: 2005/206 Grantee: **Parties of the Malting Barley Quality Improvement Program**.
Certificate No: 3458 Expiry Date: 13 November, 2027.
Agent: **Agriculture Victoria Services Pty Ltd**, Attwood, VIC.

'Fitzroy'[Φ]

Application No: 2005/207 Grantee: **Parties of the Malting Barley Quality Improvement Program**.
Certificate No: 3459 Expiry Date: 13 November, 2027.
Agent: **Agriculture Victoria Services Pty Ltd**, Attwood, VIC.

'Yarra'[Φ]

Application No: 2005/208 Grantee: **Parties of the Malting Barley Quality Improvement Program**.
Certificate No: 3460 Expiry Date: 13 November, 2027.
Agent: **Agriculture Victoria Services Pty Ltd**, Attwood, VIC.

'Hindmarsh'[Φ]

Application No: 2006/290 Grantee: **Parties of the Malting Barley Quality Improvement Program**.
Certificate No: 3404 Expiry Date: 21 November, 2027.
Agent: **Agriculture Victoria Services Pty Ltd**, Attwood, VIC.

Libertia ixiodies

NEW ZEALAND IRIS

'Goldfinger'[Φ]

Application No: 2004/209 Grantee: **Naturally Native New Zealand Plants Ltd**.
Certificate No: 3421 Expiry Date: 14 December, 2027.
Agent: **Greenhills Propagation Nursery Pty Ltd**, Tynong, VIC.

Petunia hybrid

PETUNIA

‘Conblue’Φ syn Blueberry FrostΦ

Application No: 2005/109 Grantee: **Plant 21 LLC**.
Certificate No: 3426 Expiry Date: 18 December, 2027.
Agent: **Aussie Winners Pty Ltd**, Redland Bay, QLD.

‘Constraw’Φ syn Strawberry FrostΦ

Application No: 2005/108 Grantee: **Plant 21 LLC**.
Certificate No: 3425 Expiry Date: 18 December, 2027.
Agent: **Aussie Winners Pty Ltd**, Redland Bay, QLD.

Phormium tenax

NEW ZEALAND FLAX

‘Merlot’Φ

Application No: 2002/252 Grantee: **Lyndale Nurseries Auckland Ltd**.
Certificate No: 3428 Expiry Date: 18 December, 2027.
Agent: **Greenhills Propagation Nursery Pty Ltd**, Tynong, VIC.

Pittosporum tenuifolium

PITTOSPORUM, KOHUHU, TAWHIWHI

‘Screen Between’Φ

Application No: 2005/062 Grantee: **Hayden & Jeanette Heyme**.
Certificate No: 3423 Expiry Date: 13 December, 2032.
Agent: **Southern Advanced Plants Pty Ltd**, Dromana, VIC.

Protea cynaroides

GIANT PROTEA

‘Madiba’Φ

Application No: 2004/225 Grantee: **Agricultural Research Council.**
Certificate No: 3431 Expiry Date: 20 December, 2032.
Agent: **Proteaflora Enterprises Pty Ltd**, Monbulk, VIC.

Prunus persica

PEACH

‘Coconut Ice’Φ

Application No: 2003/314 Grantee: **The Horticulture and Food Research Institute of New Zealand Limited**.
Certificate No: 3412 Expiry Date: 9 December, 2032.
Agent: **A J Park**, Canberra, ACT.

Saccharum hybrid

SUGARCANE

‘KQ228’Φ

Application No: 2005/351 Grantee: **BSES Limited and CSR Ltd**, Mackay Mail Centre, QLD.
Certificate No: 3424 Expiry Date: 14 December, 2027.

Stenotaphrum secundatum

BUFFALO GRASS, ST AUGUSTINE GRASS

‘Kings Pride’Φ

Application No: 2005/341 Grantee: **J and S Gardiner Investments Pty Ltd**.
Certificate No: 3430 Expiry Date: 20 December, 2027.
Agent: **Peter McMaugh**, Carlingford, NSW.

Triticum aestivum

WHEAT

‘Odiel’Φ

Application No: 2005/112 Grantee: **Svalof Weibull AB**.
Certificate No: 3400 Expiry Date: 12 October, 2027.
Agent: **Access Genetics Pty Ltd**, Laverton North, VIC.

x*Triticosecale*

TRITICALE

‘Kosciuszko’Φ

Application No: 2002/318 Grantee: **University of New England and QAF Feeds Pty Ltd**.
Certificate No: 3399 Expiry Date: 12 October, 2027.
Agent: **Robin Jessop**, Armidale, NSW.

Denomination Changed

Application No.	Genus	Species	Common Name	Denomination Changed From	Denomination Changed To
2003/251	*Citrus*	hybrid	Mandarin	Dalahaye	Bella
2007/268	*Fragaria*	x *ananassa*	Strawberry	JUMBUK	AMELIA
2007/245	*Stenotaphrum*	*secundatum*	Buffalo Grass	Turf Force One	TF01
2007/202	*Syzygium*	*australe*	Lilly Pilly	LITTLE MISS ELEGANCE	Little Miss-Elegance
2007/241	*Avena*	*sativa*	Oats	PO 808	Dawson

Synonym Changed

Application No.	Genus	Species	Variety	Common Name	Synonym Changed From	Synonym Changed To
2006/249	*Solanum*	*tuberosum*	SUMMER DELIGHT	Potato	Crop 17	Golden Cream
2007/275	*Zoysia*	*macrantha*	MAC03	Prickly Couch	Ozgreen	Nara

Applicant's Name Amended

Application No.	Genus	Species	Variety	Changed From	Changed To
2003/272	*Phaseolus*	*vulgaris*	BN 155	Syngenta Seeds Inc.	Syngenta Crop Protection AG
2004/016	*Vitrullus*	*lanatus*	SP-1	Syngenta Seeds Inc.	Syngenta Crop Protection AG
2004/017	*Citrullus*	*lanatus*	90-4194	Syngenta Seeds Inc.	Syngenta Crop Protection AG
2007/190	*Lactuca*	*sativa*	Curletta	Syngenta Seeds Pty Ltd	Syngenta Crop Protection AG
2007/191	*Lactuca*	*sativa*	Winny	Syngenta Seeds Pty Ltd	Syngenta Crop Protection AG
2007/192	*Lactuca*	*sativa*	Robinio	Syngenta Seeds Pty Ltd	Syngenta Crop Protection AG

Change of Agent

Application No.	Genus	Species	Variety	Changed From	Changed To
2001/013	*Anthurium*	hybrid	Antinkeles	W & E Sieverding Wholesale Nursery	Sprint Horticulture
2001/351	*Euphorbia*	*characias*	Wilcott	Fleming's Nurseries Pty Ltd	Plants Management Australia Pty Ltd
2001/352	*Euphorbia*	*characias*	Charam	Fleming's Nurseries Pty Ltd	Plants Management Australia Pty Ltd
2007/233	*Citrullus*	*lanatus*	SP-4		Syngenta Seeds Pty Ltd
2007/190	*Lactuca*	*sativa*	Curletta		Syngenta Seeds Pty Ltd
2007/191	*Lactuca*	*sativa*	Winny		Syngenta Seeds Pty Ltd
2007/192	*Lactuca*	*sativa*	Robinio		Syngenta Seeds Pty Ltd

Assignment of Rights

Application No.	Genus	Species	Variety	Changed From	Changed To
1995/200	*Metrosideros*	*excelsus*	DALESE	Neil Perrott and Robert Donato	Robert Donato

The following varieties were assigned:
From:
State of Western Australia represented by the Chief Executive Officer, State of Queensland through its Department of Primary Industries and Fisheries, Department of Primary Industries for and on behalf of the State of New South Wales, Grains Research and Development Corporation
To:
State of Queensland through its Department of Primary Industries and Fisheries, Department of Primary Industries for and on behalf of the State of New South Wales, Grains Research and Development Corporation

Triticum aestivum 2002/236 'EGA Bellaroi'
Triticum aestivum 2004/218 'EGA Wentworth'
Triticum aestivum 2004/216 'EGA Wylie'
Triticum aestivum 2004/217 'EGA Gregory'
Triticum aestivum 2002/288 'EGA Wedgetail'

The following varieties were assigned:
From:
State of Western Australia represented by the Chief Executive Officer, State of Queensland through its Department of Primary Industries and Fisheries, Department of Primary Industries for and on behalf of the State of New South Wales, Grains Research and Development Corporation

To:
State of Queensland through its Department of Primary Industries and Fisheries, Department of Primary Industries for and on behalf of the State of New South Wales, Grains Research and Development Corporation

Followed by an assignment:
To:
State of Western Australia represented by the Chief Executive Officer, Grains Research and Development Corporation

Followed by an assignment:
To:
InterGrain Pty Ltd

Triticum aestivum 2003/254 'EGA Jitarning'
Triticum aestivum 2004/197 'EGA Eagle Rock'
Triticum aestivum 2003/160 'EGA 2248'
Triticum aestivum 2003/252 'EGA Blanco'
Triticum aestivum 2003/161 'EGA Bonnie Rock'
Triticum aestivum 2003/253 'EGA Castle Rock'

The following varieties were assigned:
From:
State of Western Australia through its Department of Agriculture and Food, Grains Research and Development Corporation
To:
InterGrain Pty Ltd

Triticum aestivum 1999/226 ‘Karlgarin’
Triticum aestivum 2001/221 ‘Wyalkatchem’
Triticum aestivum 2001/222 ‘Harrismith’
Triticum aestivum 2005/016 ‘Tammarin Rock’
Triticum aestivum 2005/346 ‘Bullaring’
Triticum aestivum 2006/257 ‘Binnu’

Transfer of Rights

Application No.	Genus	Species	Variety	Right Transferred From	Rights Transferred To
1998/249	*Chamelaucium*	*uncinatum*	Dancing Queen	Western Flora	Floriscape Pty Ltd
1998/250	*Chamelaucium*	hybrid	My Sweet Sixteen	Western Flora	Floriscape Pty Ltd

Surrendered - the following varieties are no longer under PBR protection

Application No.	Genus	Species	Variety	Synonym	Common Name
1989/081	*Acalypha*	hybrid	PINK CANDLES		Chenille Plant
2002/219	*Bougainvillea*	*glabra*	Purple Patch		Bougainvillea
1999/318	*Bracteantha*	*bracteata*	NN-9812AE		Everlasting Daisy
1994/051	*Brassica*	*napus*	RAINBOW		Canola
1997/046	*Brassica*	*napus*	TI1 PINNACLE		Canola
2001/309	*Brassica*	*napus var. oleifera*	ATR-EYRE		Canola
2003/154	*Calibrachoa*	hybrid	KLEC01058	Selecta White	Calibrachoa
2001/319	*Cordyline*	*fruticosa*	Gan01		Cordyline
1991/056	*Cupressus*	*glabra*	LIMELIGHT		Arizona Cypress
1992/063	*Desmanthus*	*virgatus*	BAYAMO		Desmanthus
1992/064	*Desmanthus*	*virgatus*	UMAN		Desmanthus
2002/006	*Freesia*	*hybrid*	Varafoc	Focus	Freesia
1998/022	*Gypsophila*	*paniculata*	Dangysha	Yukinko	Baby's Breath
2001/350	*Impatiens*	*hawkeri*	Balcebchro		New Guinea Impatiens
2000/070	*Impatiens*	*hawkeri*	Balcelavgo	Celebration Lavender Glow	New Guinea Impatiens
2000/072	*Impatiens*	*hawkeri*	Balcelisow	Celebration Salmon II	New Guinea Impatiens
2000/274	*Impatiens*	*hawkeri*	BFP-796	Apricot Celebration	New Guinea Impatiens
1994/008	*Impatiens*	*walleriana*	GOLDEN SURPRISE		Busy Lizzie
1997/290	*Kalanchoe*	spp.	Elves Bells		Kalanchoe
2003/263	*Lilium*	hybrid	Loire		Lily
2002/045	*Lilium*	hybrid	WINDSOR	VLETWIN	Lily
1994/139	*Rhododendron*	hybrid	PRINCESS BARBARA		Azalea
2003/071	*Rhododendron*	*simsii*	Davidel		Azalea
1995/156	*Rhododendron*	*simsii*	HEIDE HANISCH		Azalea
1999/132	*Rosa*	hybrid	Fairy Queen		Rose
1998/265	*Rosa*	hybrid	Ruiconti	Yellow Unique	Rose
1998/264	*Rosa*	hybrid	Ruioran	Orange Unique	Rose
1992/163	*Rosa*	hybrid	TANAKINOM	MONICA	Rose
1991/078	*Rosa*	hybrid	TENNESSEE		Rose
2000/009	*Solanum*	*tuberosum*	Rioja		Potato
1996/210	*Solanum*	*tuberosum*	SAXON		Potato
1991/096	*Vitis*	*vinifera*	KING HUSAINY	JADE SEEDLESS	Grape

Withdrawn- the following varieties are no longer under PBR provisional protection

Application No.	Genus	Species	Common Name	Variety
2006/280	*Acacia*	*cognata*	Bower Wattle	BW 06
2006/309	*Brassica*	*oleracea* convar. *botrytis* var. *cymosa*	Broccoli	BRM 51-1045
2001/235	*Malus*	*domestica*	Apple	MJ 806.06
2006/293	*Rosa*	hybrid	Rose	SPEfeys
2005/304	*Rosa*	hybrid	Rose	TAN94488
2005/037	*Schlumbergera*	*truncata*	Christmas Cactus	Moonlightfantasy
2006/111	*Sedum*	hybrid	Sedum	Chocolate Sauce
2001/262	*Syzygium*	*australe*	Lilly Pilly	Yuruga No. 1
2001/261	*Syzygium*	*australe*	Lilly Pilly	Yuruga No. 2
2001/260	*Syzygium*	*australe*	Lilly Pilly	Yuruga No. 3
2001/258	*Syzygium*	*australe*	Lilly Pilly	Yuruga No. 5

CORRIGENDA

Detailed descriptions of the following varieties were published in *Plant Varieties Journal* vol 15, issue 4. The first date of sale in the EU was incorrectly given as April 1999. They should be given as:

2001/311 Osteospermum **'Seidacre'**
first date of sale in the EU 1 May 1999

2001/312 Osteospermum **'Seimora'**
first date of sale in the EU August 2000

2001/313 Osteospermum **'Seikilrem'**
first date of sale in Japan and the EU 1 May 1999

'LMF500'

Application No: 2004/249. Detailed Description published in PVJ 19.4

The correct botanical name of **'LMF500'** should be: *Lomandra filiformis* subsp. *coriacea.* The excluded variety 'Mondra' should be listed as *Lomandra filiformis* subsp. *filiformis.*

'RK19'

Application No: 2007/130. Detailed Description published in PVJ 20.3

The plant heights of the two cultivars 'RK19' and 'Whittet' are not significantly different, and so the following information should not be ticked as a key difference.

Organ/Plant Part: Context	'RK19'	'Whittet'
Plant: mean height 79 days after planting (cm)		
Mean	206.10	202.00
Std. Deviation	34.10	41.40
LSD/sig	27.4	ns

'Southern Belle' and 'Emerald'

Application no: 2005/078 and 2005/079. Detailed Description published in PVJ 20.3

The agent for the following applications was incorrectly published as BerryExchange. The correct name of the agent is BerryExchange (a division of CostaExchange Ltd).

Australian Government
IP Australia

Part 3 Appendices

The appendices to *Plant Varieties Journal* **(Vol. 20 Issue 4)** are listed below:

- Home
- Appendix 1 - Fees
- Appendix 2 - Plant Breeder's Rights Advisory Committee
- Appendix 3 - Index of Accredited Consultant 'Qualified Persons'
- Appendix 4 - Index of Accredited Non-Consultant 'Qualified Persons'
- Appendix 5 - Addresses of UPOV and Member States
- Appendix 6 - Centralised Testing Centres
- Appendix 7 - List of Plant Classes for Denomination Purposes
- Appendix 8 - Register of Plant Varieties

APPENDIX 1

FEES

Two fee structures exist as a result of the transition from Plant Variety Rights to Plant Breeders Rights. For new applications (those lodged on or after 11 November 1994) the PBR fees apply. For older applications lodged before 11 November 1994 and not finally disposed of (Granted, Withdrawn, Refused etc.) the PVR fees in force at the time apply.

The Treasurer has determined that all statutory fees under PBR regulations will be exempted from GST.

Payment of Fees

All cheques for fees should be made payable and sent to:

Collector of Public Monies
C/-Plant Breeders Rights Office, IP Australia
GPO Box 200
Woden, ACT 2606

The **application fee** ($300) must accompany the application at the time of lodgement.

Consequences of not paying fees when due

Application fee
Should an application not be accompanied by the prescribed application fee the application will be deemed to be 'non-valid' and neither assigned an application number nor examined for acceptance pending the payment of the fee.

Examination fee
Non-payment of the examination fee of an application will automatically result, at the end of 12 months from the date of acceptance[1], in a refusal of the application. The consequences of refusal are the same as for applications deemed to be inactive (see 'inactive applications' below).

Consideration of a request for an extension of the period of provisional protection from the initial 12-month period may require the prior payment of the examination fee.

Certificate fee
Following the successful completion of the examination, including the public notice period, the applicant will be required and invoiced to pay the certification fee. Payment of the certification fee is a prerequisite to granting PBR and issuing the official certificate by the PBR office. Failure to pay the fee may result in a refusal to grant PBR.

Annual fee
Should an annual renewal fee not be paid within 30 days after the due date, the grant of PBR will be revoked under Section 50 of the PBR Act. To assist grantees, the PBR office will invoice grantees or their Australian agents for renewal fees.

Inactive applications
An application will be deemed inactive if, after 24 months of provisional protection (or 12 months in the case of non-payment of the examination fee) the PBR Office has not received a completed application or has not been advised to proceed with the examination or an extension of provisional protection has not been requested or not granted or a certificate fee has not been paid. Inactive applications will be examined and, should they not fully comply with Section 44 of the PBR Act 1994, they will be refused. As a result provisional protection will lapse, priority claims on that variety will be

[1] The time limit to pay examination fees on imported varieties can be deferred for a maximum of 12 months after the variety has been released from quarantine. Contact the PBR Office for further details.

lost and should the variety have been sold, it will be ineligible for plant breeders rights on reapplication. Continued use of labels or any other means to falsely imply that a variety is protected after the application has been refused is an offence under Section 75 of the Act.

FEES

Basic Fees	**Schedule**			
	A	**B**	**C**	**D**
	$			
Application	300	300	400	300
Examination - per application	1400	1200	1400	800
Certificate	300	300	250	300
Total Basic Fees	2000	1800	2050	1400
Annual Renewal - all applications	300			

Schedule

A Single applications and applications based on an official overseas test reports.

B Applicable when two or more Part 2 Applications are lodged simultaneously and the varieties are of the same genus and the examinations can be completed at one location at the same time.

C Applications lodged under PVR (prior to 10th Nov 1994)

D Applicable to 5 or more applications examined at an Accredited Centralised Testing Centre

Other Fees	
Variation to application(s) - per hour or part thereof	75
Change of Assignment - per application	100
Copy of an application (Part1 and/or Part2) , an objection or a detailed description	50
Copy of an entry in the Register	50
Lodging an objection	100
Annual subscription to Plant Varieties Journal	40
Back issues of Plant Varieties Journal	14
Administration - Other work relevant to PBR - per hour or part thereof	75
Application for declaration of essential derivation	800
Application for (a) revocation of a PBR	500
(b) revocation of a declaration of essential derivation	500
Compulsory licence	500
Request under subsection 19(11) for exemption from public access - varieties with no direct use as a consumer	100

APPENDIX 2

Plant Breeders Rights Advisory Committee (PBRAC)

(Members of the PBRAC hold office in accordance with Section 85 of the *Plant Breeder's Rights Act 1994*.)

Committee Members

Member Representing Plant Breeders Dr Paul Brennan Rock Valley Post Office via Lismore 1201 Cawongla Rd LARNOOK NSW 2480	**Member Representing Plant Breeders** Dr Glenn Dale Saltgrow PO Box 575 ASHGROVE QLD 4060
Member Representing Users Mr Robert Hansen Peanut Company of Australia PO Box 26 KINGAROY QLD 4610	**Member Representing Consumers** Ms Anne Pye PO Box 1538 MT BARKER SA 5251
Member Representing Conservation Interests Mr Bruce Lloyd Fairley downs 5250 Barmah-Shepparton Road TALLYGAROOPNA VIC 3634	**Member Representing Indigenous Interests** Mr Mark Porter 26 Callicarpa Street REEDY CREEK QLD 4227
Member with Appropriate Qualifications Mr Benny Browne Griffith Hack 509 St Kilda Road MELBOURNE VIC 3004	**Member with Appropriate Qualifications** Professor Brad Sherman TC Beirne School of Law The University of Queensland ST LUCIA QLD 4072
Registrar (Chair) Mr Doug Waterhouse IP Australia PO Box 200 Woden ACT 2606	

APPENDIX 3 - INDEX OF ACCREDITED CONSULTANT 'QUALIFIED PERSONS'

The following persons have been accredited by the PBR office based on information provided by these persons. From the information provided by the applicants, the PBR office believes that these people can fulfil the role of 'qualified person' in the application for plant breeder's rights. Neither accreditation nor publication of a name in the list of persons is an implicit recommendation of the person so listed. The PBR office cannot be held liable for damages that may arise from the omission or inclusion of a person's name in the list nor does it assume any responsibility for losses or damages arising from agreements entered into between applicants and any person in the list of accredited persons. Qualified persons charge a fee for services rendered.

A guide to the use of the index of consultants:

- locate in the left column of Table 1 the plant group for which you are applying;
- listed in the right column are the names of accredited qualified persons from which you can choose a consultant;
- in Table 2 find that consultant's name, telephone number and area in which they are willing to consult (they may consult outside the nominated area);
- using the "Nomination of Qualified Person" form as a guide, agree provisionally on the scope and terms of the consultancy; complete the form and attach it to Part 1 of the application form;
- when you are notified that your nomination of a consultant qualified person is acceptable in the letter of acceptance of your application for PBR you should again consult the qualified person when planning the rest of the application for PBR.

TABLE 1

PLANT GROUP/SPECIES/FAMILY	CONSULTANT'S NAME (TELEPHONE AND AREA IN TABLE 2)
Actinidia	Lye, Colin Paananen, Ian Richards, Graeme
Agapanthus	Paananen, Ian
Almonds	Granger, Andrew Swinburn, Garth
Alstroemeria	Paananen, Ian
Ajuga	Paananen, Ian
Apple	Cramond, Gregory Darmody, Liz Engel, Richard Fleming, Graham Langford, Garry Mackay, Alastair Malone, Michael Mitchell, Leslie Portman, Anthony Scholefield, Peter Tancred, Stephen Valentine, Bruce

Anigozanthos	Paananen, Ian Kirby, Greg Smith, Daniel
Anthurium	Paananen, Ian
Aroid	Harrison, Peter
Avocado	Lye, Colin Edwards, Arthur MacGregor, Alison Owen-Turner, John Parr, Wayne Swinburn, Garth Whiley, Tony
Azalea	Barrett, Mike Hempel, Maciej Paananen, Ian
Barley (Common)	Bhatti, Muhammad Collins, David Downes, Ross Khan, Akram Platz, Greg Rhodes, Phil Saunders, James
Berry Fruit	Darmody, Liz Fleming, Graham Greer, Neil Scholefield, Peter Zorin, Margaret
Blackberry (*Rubus* sp)	Paananen, Ian
Blandfordia	Treverrow, Florence
Blueberry	Paananen, Ian Zorin, Margaret
Bougainvillea	Iredell, Janet Willa Prince, John
Brachyscome	Paananen, Ian

Brassica	Bannan, Nathaniel Bhatti, Muhammad Chequer, Robert Cooper, Kath Downes, Ross Easton, Andrew Fennell, John Gororo, Nelson Johnston, Evan Kadkol, Gururaj Laker, Richard Light, Kate McMichael, Prue Rhodes, Phil Rudolph, Paul Sanders, Milton Saunders, James Scholefield, Peter Mouwen, Heidi Zadow, Diane
Brunia	Dunstone, Bob
Buddleia	Robb, John Paananen, Ian
Buffalo Grass	Paananen, Ian
Calibrachoa	Paananen, Ian
Camellia	Paananen, Ian Robb, John
Cannabis	Calabria, Patrick
Carnation/Dianthus	Paananen, Ian

Cereals	Bhatti, Muhammad Bullen, Kenneth Collins, David Cook, Bruce Cooper, Kath Downes, Ross Fennell, John Hare, Raymond Harrison, Peter Henry, Robert J Johnston, Evan Khan, Akram Mitchell, Leslie Moore, Stephen Oates, John Platz, Greg Porter, Richard Poulsen, David Rhodes, Phil Roake, Jeremy Rose, John Saunders, James Scattini, Walter John Siedel, John Wilson, Frances
Cherry	Cramond, Gregory Darmody, Liz Fleming, Graham Granger, Andrew Mackay, Alastair Mitchell, Leslie Pumpa, Lucy Scholefield, Peter
Chickpeas	Bhatti, Muhammad Downes,Ross Collins, David Goulden, David Rhodes, Phil Saunders, James
Chrysanthemum	Paananen, Ian
Citrus	Calabria, Patrick Edwards, Arthur Lee, Slade MacGregor, Alison Mitchell, Leslie Owen-Turner, John Parr, Wayne Scholefield, Peter Swinburn, Garth Sykes, Stephen Topp, Bruce
Clivia	Smith, Kenneth

Clover	Bannan, Nathaniel Downes, Ross James, Jennifer Johnston, Evan Lake, Andrew Miller, Jeff Mitchell, Leslie Nichols, Phillip Porter, Richard Rhodes, Phil Saunders, James
Cotton	Khan, Akram Leske, Richard
Cucurbits	Herrington, Mark McMichael, Prue Rhodes, Phil Scholefield, Peter Sykes, Stephen
Dianella	Paananen, Ian
Dogwood	Darmody, Liz Fleming, Graham
Echinacea	Paananen, Ian
Eucalyptus	Paananen, Ian
Euphorbia	Paananen, Ian
Feijoa	Parr, Wayne Scholefield, Peter
Fibre Crops	Gillespie, David Khan, Akram
Fig	Darmody, Liz Fleming, Graham Parr, Wayne
Flower Bulbs	Verdegaal, John
Forage Brassicas	Goulden, David Rhodes, Phil Saunders, James
Forage Grasses	Bannan, Nathaniel Downes, Ross Fennell, John Harrison, Peter Johnston, Evan Kirby, Greg Mitchell, Leslie Rhodes, Phil Smith, Kevin

Forage Legumes	Downes, Ross Fennell, John Foster, Kevin Harrison, Peter Hill, Jeff James, Jennifer Lake, Andrew Miller, Jeff Porter, Richard Rhodes, Phil Saunders, James Siedel, John
Fruit	Cramond, Gregory Darmody, Liz Delaporte, Kate Fleming, Graham Gillespie, David Granger, Andrew Kennedy, Peter Lenoir, Roland McCarthy, Alec Mitchell, Leslie Parr, Wayne Portman, Sian Pumpa, Lucy Schapel, Amanda Scholefield, Peter
Fuchsia	Paananen, Ian
Gerbera	Paananen, Ian
Ginger	Smith, Mike Whiley, Tony
Grapes	Burne, Peter Darmody, Liz Delaporte, Kate Farquhar, Wayne Fleming, Graham Lee, Slade Lye, Colin MacGregor, Alison Mitchell, Leslie Paananen, Ian Parr, Wayne Porter, Richard Pumpa, Lucy Schapel, Amanda Scholefield, Peter Smith, Daniel Swinburn, Garth Sykes, Stephen
Grevillea	Dunstone, Bob Herrington, Mark Paananen, Ian

Gypsophila	Paananen, Ian
Hardenbergia	Dunstone, Bob
Hops (*Humulus* sp)	Paananen, Ian
Hydrangea	Hanger, Brian Paananen, Ian
Impatiens	Paananen, Ian
Jojoba	Dunstone, Bob
Kalanchoe	Paananen, Ian
Lavender	Paananen, Ian
Legumes	Aberdeen, Ian Collins, David Cook, Bruce Cruickshank, Alan Downes, Ross Foster, Kevin Harrison, Peter Imrie, Bruce Kirby, Greg Khan, Akram Knights, Edmund Lake, Andrew Loch, Don Mitchell, Leslie Rhodes, Phil Rose, John Saunders, James Siedel, John
Lentils	Collins, David Downes, Ross Goulden, David Khan, Akram Porter, Richard Rhodes, Phil Saunders, James
Lilium	Paananen, Ian
Liriope	Paananen, Ian
Lomandra	Paananen, Ian
Lucerne	Bannan, Nathaniel Downes, Ross Johnston, Evan Lake, Andrew Mitchell, Leslie Nichols, Phillip Porter, Richard Rhodes, Phil Saunders, James

Lupin	Bhatti, Muhammad Collins, David Sanders, Milton Rhodes, Phil Saunders, James
Magnolia	Paananen, Ian
Mandevilla	Paananen, Ian
Mango	Lye, Colin Owen-Turner, John Mitchell, Leslie Parr, Wayne Whiley, Tony
Myrtaceae	Dunstone, Bob
Native grasses	Paananen, Ian Quinn, Patrick
Oat	Bhatti, Muhammad Collins, David Downes, Ross Khan, Akram Platz, Greg Rhodes, Phil Saunders, James
Oilseed crops	Downes, Ross Poulsen, David Siedel, John Rhodes, Phil Saunders, James
Olives	Bazzani, Mr Luigi Granger, Andrew
Onions	Bannan, Nathaniel Fennell, John Khan, Akram Laker, Richard McMichael, Prue Scholefield, Peter Rhodes, Phil

Ornamentals - Exotic	Abell, Peter Armitage, Paul Angus, Tim Barth, Gail Collins, Ian Cunneen, Thomas Darmody, Liz Delaporte, Kate Eggleton, Steve Fisk, Anne Marie Fleming, Graham Guy, Gareme Harrison, Peter Hempel, Maciej Johnston, Margaret Khan, Akram Kulkarni, Vinod Lamont, Greg Larkman, Clive Lenoir, Roland Lowe, Greg Lunghusen, Mark Marcsik, Doris McMichael, Prue Milne, Carolynn Mitchell, Hamish Mitchell, Leslie Nichols, David Oates, John O'Brien, Shaun Paananen, Ian Prescott, Chris Prince, John Robb, John Pumpa, Lucy Schapel, Amanda Scholefield, Peter Singh, Deo Smith, Daniel Stewart, Angus Van der Staay, Rosemaree Anne Watkins, Phillip Watkinson, Andrew

Ornamentals - Indigenous	Abell, Peter Allen, Paul Angus, Tim Barrett, Mike Barth, Gail Cunneen, Thomas Delaporte, Kate Downes, Ross Eggleton, Steve Granger, Andrew Harrison, Peter Henry, Robert J Hockings, David Jack, Brian Johnston, Margaret Kirby, Greg Khan, Akram Lenoir, Roland Lowe, Greg Lullfitz, Robert Lunghusen, Mark McMichael, Prue Milne, Carolynn Mitchell, Hamish Molyneux, W M Nichols, David Oates, John O'Brien, Shaun Paananen, Ian Prince, John Pumpa, Lucy Schapel, Amanda Scholefield, Peter Singh, Deo Slater, Tony Smith, Daniel Tan, Beng Watkins, Phillip
Ornithopus	Foster, Kevin Nichols, Phillip
Osmanthus	Paananen, Ian Robb, John
Osteospermum	Paananen, Ian

Pastures & Turf	Anderson, Malcolm Avery, Angela Bannan, Nathaniel Bhatti, Muhammad Cameron, Stephen Cook, Bruce Downes, Ross Harrison, Peter Kemp, Stuart Kirby, Greg James, Jennifer Loch, Don McMaugh, Peter Miller, Jeff Mitchell, Leslie Neylan, John Paananen, Ian Porter, Richard Rhodes, Phil Rose, John Saunders, James Smith, Raymond Scattini, Walter John Smith, Kevin Wilkes, Gregory Wilson, Frances Zorin, Margaret
Peanut	Cruickshank, Alan George, Doug
Pear	Cramond, Gregory Darmody, Liz Engel, Richard Fleming, Graham Langford, Garry Mackay, Alastair Malone, Michael Paananen, Ian Portman, Anthony Scholefield, Peter Tancred, Stephen Valentine, Bruce
Pelargonium	Paananen, Ian
Persimmon	Parr, Wayne Swinburn, Garth
Petunia	Paananen, Ian Nichols, David
Philodendron	Paananen, Ian
Philotheca	Dunstone, Bob
Phormium	Paananen, Ian
Photinia	Robb, John

Pistacia	Richardson, Clive Sykes, Stephen
Pisum	Bhatti, Muhammad Downes, Ross Goulden, David McMichael, Prue Rhodes, Phil Sanders, Milton Saunders, James
Potatoes	Delaporte, Kate Fennell, John Friemond, Terry Guertsen, Paul Hill, Jim Johnston, Evan McMichael, Prue Pumpa, Lucy Rhodes, Phil Saunders, James Schapel, Amanda Scholefield, Peter Slater, Tony Smith, Daniel Wilson, Graeme
Proteaceae	Barth, Gail Kirby, Neil Paananen, Ian Robb, John Scholefield, Peter Smith, Daniel
Prunus	Calabria, Patrick Cramond, Gregory Darmody, Liz Engel, Richard Fleming, Graham Granger, Andrew Kennedy, Peter Mackay, Alastair Malone, Michael Portman, Anthony Richards, Graeme Topp, Bruce Wilkes, Gregory Witherspoon, Jennifer
Pulse Crops	Collins, David Downes, Ross Graetz, Darren Oates, John Porter, Richard Poulsen, David Rhodes, Phil Saunders, James

Raspberry	Darmody, Liz Fleming, Graham Herrington, Mark Scholefield, Peter Zorin, Margaret
Rhododendron	Barrett, Mike Paananen, Ian
Rose	Barrett, Mike Darmody, Liz Delaporte, Kate Fleming, Graham Hanger, Brian Lee, Peter McKirdy, Simon Paananen, Ian Prescott, Chris Pumpa, Lucy Schapel, Amanda Scholefield, Peter Smith, Daniel Swane, Geoff Syrus, A Kim
Scaevola	Paananen, Ian
Sesame	Bennett, Malcolm Harrison, Peter Imrie, Bruce
Sorghum	Khan, Akram
Soybean	Harrison, Peter James, Andrew
Spathiphylum	Paananen, Ian
Spices and Medicinal Plants	Khan, Akram
Stone Fruit	Barrett, Mike Cramond, Gregory Darmody, Liz Fleming, Graham Granger, Andrew Kennedy, Peter MacGregor, Alison Mackay, Alistair Malone, Michael Scholefield, Peter Swinburn, Garth Valentine, Bruce
Strawberry	Herrington, Mark Mitchell, Leslie Morrison, Bruce Scholefield, Peter Zorin, Margaret

Sugarcane	Cox, Mike Piperidis, George
Sunflower	George, Doug
Tomato	Herrington, Mark Khan, Akram Laker, Richard McMichael, Prue Rhodes, Phil Scholefield, Peter Smith, Daniel
Tree Crops	McRae, Tony
Triticale	Bhatti, Muhammad Downes, Ross Collins, David Cooper, Kath Rhodes, Phil Saunders, James
Tropical/Sub-Tropical Crops	Harrison, Peter Kulkarni, Vinod Parr, Wayne Scholefield, Peter Whiley, Tony
Umbrella Tree	Paananen, Ian
Vegetables	Bannan, Nathaniel Delaporte, Kate Fennell, John Frkovic, Edward Gillespie, David Harrison, Peter Khan, Akram Laker, Richard Lenoir, Roland MacGregor, Alison McMichael, Prue Oates, John O'Connor, Lauren Pearson, Craig Pumpa, Lucy Rhodes, Phil Schapel, Amanda Scholefield, Peter Smith, Daniel Westra Van Holthe, Jan
Verbena	Paananen, Ian
Walnut	Mitchell, Leslie

Wheat (Aestivum & Durum Groups)	Bhatti, Muhammad Collins, David Downes, Ross Kadkol, Gururaj Khan, Akram Platz, Greg Rhodes, Phil Saunders, James Sanders, Milton
Zantedeschia	Paananen, Ian

TABLE 2

NAME	TELEPHONE	AREA OF OPERATION
Abell, Peter	0438 392 837 mobile	Australia
Aberdeen, Ian	03 5782 1029 03 5782 2073 fax	SE Australia
Allen, Paul	07 3824 0263 ph/fax	SE QLD, Northern NSW
Anderson, Malcolm	03 5573 0900 03 5571 1523 fax 017 870 252 mobile	Victoria
Angus, Tim	(64 4) 568 3878 ph/fax 001164211871076 mobile plantatim@zip.co.nz	Australia and New Zealand
Armitage, Paul	03 9756 7233 03 9756 6948 fax	Victoria
Avery, Angela	02 6030 4500 02 6030 4600 fax	South Eastern Australia
Bannan, Nathaniel	03 8318 9019 03 8318 9002 fax 0429 720 013 mobile	Australia
Barrett, Mike	02 9875 3087 02 9980 1662 fax 0407 062 494 mobile	NSW/ACT
Barth, Gail	08 8389 7479	SA and Victoria
Bazzani, Luigi	08 9772 1207 08 9772 1333 fax	Western Australia
Bennett, Malcolm	08 8973 9733 08 8973 9777 fax	NT, QLD, NSW, WA
Bhatti, Muhammad	08 9671 1322 ph 08 9671 1352 fax	Western Australia
Burne, Peter	08 8582 0338 ph 08 8583 2104 fax 0418 834 102 mobile	South Australia
Calabria, Patrick	02 6963 6360 0438 636 219 mobile	Riverina area of NSW
Chequer, Robert	03 5382 1269 0419 145 262 mobile	Victoria
Collins, David	08 9623 2343 ph/fax 0154 42694 mobile	Central Western Wheatbelt of Western Australia
Cooper, Kath	08 8339 3049 0429 191 848 mobile	South Australia
Cox, Mike	07 4132 5200 07 4132 5253 fax	Queensland and NSW
Cramond, Gregory	08 8390 0299 08 8390 0033 fax 0417 842 558 mobile	Australia
Cruickshank, Alan	07 4160 0722 07 4162 3238 fax	QLD
Cunneen, Thomas	02 4889 8647 02 4889 8657 fax	Sydney Region
Darmody, Liz	03 9756 6105 03 9752 0005 fax	Australia
Delaporte, Kate	08 8373 2488 08 8373 2442 fax 0427 394 240 mobile	South Australia
Downes, Ross	02 4474 0456 ph 02 4474 0476 fax 0402472601 mobile	ACT, South East Australia

Dunstone, Bob	02 6281 1754 ph/fax	South East NSW
Easton, Andrew	07 4690 2666 07 4630 1063 fax	QLD and NSW
Edwards, Arthur	08 8586 1232 08 8595 1394 fax 0409 609 300 mobile	SE Australia
Eggleton, Steve	03 9876 1097 03 9876 1696 fax	Melbourne Region
Engel, Richard	08 9397 5941 08 9397 5941 fax	WA
Fennell, John	03 5334 7871 03 5334 7892 fax 0419 881 887	Australia
Farquhar, Wayne	08 85657000 08 85657011 fax	South Australia
Fleming, Graham	03 9756 6105 03 9752 0005 fax	Australia
Friemond, Terry	08 9203 6720 08 9203 6720 fax 0438 915 811 mobile	Western Australia
Foster, Kevin	08 9368 3804 08 9474 2840 fax	Mediterranean areas of Australia
Frkovic, Edward	02 6962 7333 02 6964 1311 fax	Australia
George, Doug	07 5460 1308 07 5460 1112 fax	Australia
Gillespie, David	07 4155 6344 07 4155 6656 fax	Wide Bay Burnett District, QLD
Gororo, Nelson	03 5382 5911 03 5382 5755 fax 0428 534 770 mobile	Mediterranean areas of Australia
Goulden, David	64 3 325 6400 64 3 325 2074 fax	New Zealand
Graetz, Darren	08 8303 9362 08 8303 9424 fax	South Australia
Granger, Andrew	08 8389 8809 08 8389 8899 fax	South Australia
Greer, Neil	07 5441 1118 07 5476 0098 fax 0418 881 755 mobile	Australia
Guertsen, Paul	02 6845 3789 02 6845 3382 fax 0407 658 105 mobile	NSW, VIC, SE QLD
Hanger, Brian	03 9837 5547 ph/fax 0418 598106 mobile	Victoria
Hare, Ray	02 6763 1232 02 6763 1222 fax	QLD, NSW VIC & SA
Harrison, Peter	08 8948 1894 ph 08 8948 3894 fax 0407 034 083 mobile	Tropical/Sub-tropical Australia, including NT and NW of WA and tropical arid areas
Hempel, Maciej	02 4628 0376 02 4625 2293 fax	NSW, QLD, VIC, SA
Henry, Robert J	02 6620 3010 02 6622 2080 fax	Australia
Herrington, Mark	07 5441 2211 07 5441 2235 fax	Southern Queensland
Hill, Jeff	08 8303 9487 08 8303 9607 fax	South Australia

Hill, Jim	03 6428 2519 03 6428 2049 fax 0428 262 765 mobile	Australia
Hockings, David	07 5494 3385 ph/fax	Southern Queensland
Imrie, Bruce	02 4474 0951 02 4474 0952 imriecsc@sci.net.au	SE Australia
Iredell, Janet Willa	07 3202 6351 ph/fax	SE Queensland
Jack, Brian	08 9952 5040 08 9952 5053 fax	South West WA
James, Andrew	07 3214 2278 07 3214 2272 fax	Australia
James, Jennifer	+64 6 3518214	Manawatu Region, New Zealand
Johnston, Evan	64 3358 1745 0214 417 13 mobile	Canterbury, New Zealand
Johnston, Margaret	07 5460 1240 07 5460 1455 fax	SE Queensland
Kadkol, Gururaj	03 5382 1269 03 5381 1210 fax	North Western Victoria
Kemp, Stuart	03 8390 8150 0437 278 873 mobile	SE Australia
Kennedy, Peter	02 6382 7600 02 6382 2228 fax	New South Wales
Khan, Akram	02 9351 8821 02 9351 8875 fax	New South Wales
Kirby, Greg	08 8201 2176 08 8201 3015 fax	South Australia
Kirby, Neil	02 4754 2637 02 4754 2640 fax	New South Wales
Knights, Edmund	02 6763 1100 02 6763 1222 fax	North Western NSW
Kulkarni, Vinod	08 9992 2221 08 9992 2049 fax	Australia
Lake, Andrew	08 8177 0558 0418 818 798 mobile lake@arcom.com.au	SE Australia
Laker, Richard	08 87258987 08 8723 0142 fax 0417 855 592 mobile	Australia
Lamont, Greg	02 8778 5388 02 9734 9866 fax	Sydney region
Langford, Garry	03 6266 4344 03 6266 4023 fax 0418 312 910 mobile	Australia
Larkman, Clive	03 9735 3831 03 9739 6370 larkman@tpgi.com.au	Victoria
Lee, Peter	03 6330 1147 03 6330 1927 fax	SE Australia
Lee, Slade	02 6620 3410 02 6622 2080 fax	Queensland/Northern New South Wales
Lenoir, Roland	02 6231 9063 ph/fax	Australia
Leske, Richard	07 4671 3136 07 4671 3113 fax	Cotton growing regions of QLD & NSW
Light, Kate	03 5362 2175 0419 145 768 mobile	Victoria
Loch, Don	07 3286 1488 07 3286 3094 fax	Queensland

Lowe, Greg	02 4389 8750 02 4389 4958 fax 0411 327390 mobile	Sydney, Central Coast NSW
Lullfitz, Robert	08 9447 6360	South West WA
Lunghusen, Mark	03 5998 2083 03 5998 2089fax 0407 050 133 mobile	Melbourne & environs
Lye, Colin	07 4671 0044 07 4671 0066 fax 0427 786 668 mobile	NT, QLD and NSW
MacGregor, Alison	03 5023 4644 0419 229 713 mobile	Southern Australia – Murray Valley Region
Mackay, Alastair	08 9310 5342 ph/fax 0159 87221 mobile	Western Australia
McMaugh, Peter	02 9872 7833 02 9872 7855 fax	Australia
Malone, Michael	+64 6 877 8196 +64 6 877 4761 fax	New Zealand
Marcsik, Doris	08 8999 2017 08 8999 2049	Northern Territory and Queensland
McCarthy, Alec	08 9780 6273 08 9780 6136 fax	South West WA
McKirdy, Simon	042 163 8229 mobile	Australia
McMichael, Prue	08 8373 2488 08 8373 2442 fax	SE Australia
McRae, Tony	08 8723 0688 08 8723 0660 fax	Australia
Miller, Jeff	64 6 356 8019 extn 8027 64 3 351 8142 fax	Manawatu region, New Zealand
Milne, Carolynn	07 3206 3509	QLD
Mitchell, Hamish	03 9737 9568 03 9737 9899 fax	Victoria
Mitchell, Leslie	03 5821 2021 03 5831 1592 fax	VIC, Southern NSW
Molyneux, William	03 5965 2011 03 5965 2033 fax	Victoria
Moore, Stephen	02 6799 2230 02 6799 2239 fax	NSW
Morrison, Bruce	03 9210 9251 03 9800 3521 fax	East of Melbourne
Mouwen, Heidi	07 4690 2666 07 4630 1063	QLD, NSW
Neylan, John	03 9886 6200 0413 620 256 mobile	VIC, NSW, SA
Nichols, David	03 5977 4755 03 5977 4921 fax	SE Melbourne, Mornington Peninsula and Dandenong Ranges, Victoria
Nichols, Phillip	08 9387 7442 08 9383 9907 fax	Western Australia
Oates, John	02 4473 8465	Sydney region, Eastern Australia
O'Brien, Shaun	07 5442 3055 07 5442 3044 fax 0407 584 417 mobile	SE Queensland
O'Connor, Lauren	07 3359 3113 0418 510 480 mobile	Australia
Owen-Turner, John	07 4129 5217 07 4129 5511 fax	Burnett region, Central Queensland region

Paananen, Ian	02 4381 0051 02 8569 1896 fax 0412 826 589 mobile	Australia (based in Sydney) and New Zealand
Parr, Wayne	07 4129 4147 07 4129 4463 fax	QLD, Northern NSW
Piperidis, George	07 3331 3373 07 3871 0383 fax	QLD, Northern NSW
Platz, Greg	07 4639 8817 07 4639 8800 fax	QLD, Northern NSW
Porter, Richard	08 8431 5396 08 8431 5396 fax 0413 270 670 mobile	Adelaide region, South Australia
Portman, Anthony	08 9274 5355 08 9250 1859 fax	South-west Western Australia
Portman, Sian	08 9725 0660 0421 606 651 mobile	Western Australia
Poulsen, David	07 4661 2944 07 4661 5257 fax	SE QLD, Northern NSW
Prescott, Chris	03 5998 5100 03 5998 5333 0417 340 558 mobile	Victoria
Prince, John	07 5533 0211 07 5533 0488 fax	SE QLD
Pumpa, Lucy	08 8373 2488 08 8373 2422 fax 0400 041 881 mobile	South Australia
Quinn, Patrick	03 5427 0485	SE Australia
Richards, Graeme	02 4570 1358 02 4570 1314 fax 0405 178 211 mobile	Australia
Richardson, Clive	03 51550255	Victoria
Rhodes, Phil	64 3322 5405 0211 862 422 mobile phil@epr.co.nz	New Zealand
Roake, Jeremy	02 9351 8830 02 9351 8875 fax	Sydney Region
Robb, John	02 4376 1330 02 4376 1271 fax 0199 19252 mobile	Sydney, Central Coast NSW
Rose, John	07 4661 2944 07 4661 5257 fax	SE Queensland
Rudolph, Paul	03 5381 2168 03 5381 1210 fax 0438 083 840 mobile	Victoria
Saunders, James	03 8318 9016 03 8318 9002 fax 0408 037 801 mobile	Australia
Sanders, Milton	08 9825 8087 08 9387 4388 fax 0427 031 951 mobile	Southern Australia: WA,Vic, NSW, SA
Scattini, Walter	07 3356 0863 ph/fax	Tropical and sub-tropical Australia
Schapel, Amanda	08 8373 2488 0408 344 843 mobile	South Australia
Scholefield, Peter	08 8373 2488 08 8373 2442 fax 018 082022 mobile	SE Australia
Singh, Deo	0418 880787 mobile 07 3207 5998 fax	Brisbane

Slater, Tony	03 9210 9222 03 9800 3521 fax 0408 656 021 mobile	SE Australia
Smith, Daniel	08 8373 2488 08 8373 2442 fax	South Australia
Smith, Kenneth	02 4570 9069	Australia
Smith, Kevin	03 5573 0900 03 5571 1523 fax	SE Australia
Smith, Mike	07 5444 9630	SE Queensland
Smith, Stuart	03 6336 5234 03 6334 4961 fax	SE Australia
Stewart, Angus	02 4385 9788ph/fax 0419 632 123 mobile	Sydney, Gosford
Swane, Geoff	02 6889 1545 02 6889 2533 fax 0419 841580 mobile	Central western NSW
Swinburn, Garth	03 5023 4644 03 5023 5814 fax	Murray Valley Region - from Swan Hill (Vic) to Waikere (SA)
Sykes, Stephen	03 5051 3100 03 5051 3111 fax	Victoria
Syrus, A Kim	03 8556 2555 03 8556 2955 fax	Adelaide
Tan, Beng	08 9266 7168 08 9266 2495	Perth & environs
Tancred, Stephen	07 4681 2931 07 4681 4274 fax 0157 62888 mobile	QLD, NSW
Treverrow, Florence	02 6629 3359	Australia
Topp, Bruce	07 4681 1255 07 4681 1769 fax	SE QLD, Northern NSW
Valentine, Bruce	02 6361 3919 02 6361 3573 fax	New South Wales
Van der Staay, Rosemaree Anne	03 6248 6863 03 6248 7402 fax	Tasmania
Verdegaal, John	03 6458 3581 03 6458 3581 fax	Australia and New Zealand
Watkins, Phillip	08 9537 1811 08 9537 3589 fax 0416 191 472 mobile	Perth Region
Watkinson, Andrew	07 5445 6654 0409 065 266 mobile	Northern NSW and Southern QLD
Westra Van Holthe, Jan	03 9706 3033 03 9706 3182 fax	Australia
Whiley, Tony	07 5441 5441	QLD
Wilkes, Gregory	02 4570 1358 02 4570 1314 fax 0418 642 359 mobile	Sydney region
Wilson, Frances	64 3 318 8514 64 3 318 8549 fax	Canterbury, New Zealand
Wilson, Graeme	03 5957 1200 03 5957 1210 fax	SE Australia
Zadow, Diane	03 5382 1269 03 5381 1210 fax 0419 145 763 mobile	Victoria
Zorin, Margaret	07 3207 4306 0418 984 555	Eastern Australia

Appendix 4 Index of Accredited Non-Consultant Qualified Persons

Name	Name
Ali, S	Lowe, Russell
Allen, Antony	Luckett, David
Armour, David	Mack, Ian
Baelde, Arie	Mann, Dorham
Baker, Grant	Mansfield, Daniel
Bally, Ian	Mason, Lloyd
Barr, Andrew	Matic, Rade
Bell, David	Matthews, Michael
Bernuetz, Andrew	McCallum, Lesley
Birmingham, Erika	McDonald, David
Box, Amanda	Mendham, Neville
Brennan, Paul	Menzies, Kim
Brewer, Lester	Miller, Kylie
Brindley, Tony	Moody, David
Brindle, Sean	Moss, Ian
Buchanan, Peter	Mullins, Kathleen
Bunker, John	Mungall, Neil
Bunker, Kerry	Neilson, Peter
Burton, Wayne	Newman, Allen
Cameron, Nick	Noone, Brian
Cant, Russell	Norriss, Michael
Chesher, Wayne	Oakes, John
Chivers, Ian	Offord, Cathy
Clayton-Greene, Kevin	O'Brien, Tim
Constable, Greg	O'Sullivan, Robert
Cook, Esther	Paull, Jeff
Corcoran, Lisa	Pearce, Bob
Coventry, Stewart	Potter, Trent
Craig, Andrew	Pressler, Craig
Craigie, Gail	Reeve, Christopher
Culvenor, Richard	Reid, Peter
Dawson, Iain	Reinke, Russell
Crowhurst, Max	Roberts, Sean
De Betue, Remco	Roche, Matthew
de Koning, Carolyn	Rose, Ian
Dear, Brian	Sanders, Milton
Delaporte, Kate	Sandral, Graeme
Done, Anthony	Sanewski, Garth
Donnelly, Peter	Schilg, Karl
Downe, Graeme	Schreuders, Harry
Dryden, Susan	Scott, Ralph
Eastwood, Russell	Senior, Michael
Eglinton, Jason	Siemon, Fran
Eisemann, Robert	Smith, Chris
Elliott, Philip	Smith, Raymond
Evans, Pedro	Smith, Malcolm
Fitzgibbon, John	Smith, Susan
Flett, Peter	Snelling, Cath
Geary, Judith	Snowball, Richard
Gibbons, Philip	Stiller, Warwick

Gillies, Leanne	Stuart, Peter
Glover, Russell	Sturgess, Eric
Granger, Andrew	Sutton, John
Gurciullo, Gaetano	Tonks, John
Haire, Chris	Trimboli, Daniel
Harden, Patrick	Taylor, Kerry
Hollamby, Gil	Trigg, Pamela
Hoppo, Suzanne	Urwin, Nigel
Howie, Jake	Van der Spek, Folke
Hoxha, Adriana	Vater, Daniel
Hunt, Melissa	Vaughan, Peter
Hurst, Andrea	Venkatanagappa, Shoba
Irwin, John	Venn, Neil
Janhsen, Joanne	Warner, Bradley
Johnson, Peter	Warren, Andrew
Jupp, Noel	Watson, Brigid
Kaehne, Ian	Weatherly, Lilia
Katelaris, Andrew	Wei, Xianming
Kebblewhite, Tony	Whalley, RDB
Kempff, Stefan	Williams, Rex
Kennedy, Chris	Wilson, Stephen
Kobelt, Eric	Wilson, Rob
Lacey, Kevin	Winter, Bruce
Lawson, Marion	Wirthensohn, Michelle
Lee, Kathryn	Wright, Gary
Leighton, A	Yan, Guijun
Leonforte, Antonio	Zeppa, Aldo
Lewin, Laurence	
Lewis, Hartley	
Loi, Angelo	

APPENDIX 5

ADDRESSES OF UPOV AND MEMBER STATES

International Union for the Protection of New Varieties of Plants (UPOV):

International Union for the Protection of New Varieties of Plants (UPOV)
34, Chemin des Colombettes
CH-1211
Geneva 20
SWITZERLAND

Phone: (41-22) 338 9111
Fax: (41-22) 733 0336
Web site: http://www.upov.int

List of Addresses of Plant Variety Protection Offices in UPOV Member States

Status of Ratification in UPOV member States is available from UPOV website.

APPENDIX 6

CENTRALISED TESTING CENTRES

Under Plant Breeder's Rights Regulations introduced in 1996, establishments may be officially authorised by the PBR office to conduct test growings. An authorised establishment will be known as Centralised Test Centre (CTC).

Usually, the implementation of PBR in Australia relies on a 'breeder testing' system in which the applicant, in conjunction with a nominated Qualified Person (QP), establishes, conducts and reports a comparative trial. More often than not, trials by several breeders are being conducted concurrently at different sites. This makes valid comparisons difficult and often results in costly duplication.

While the current system is and will remain satisfactory, other optional testing methods are now available which will add flexibility to the PBR process.

Centralised Testing is one such optional system. It is based upon the authorisation of private or public establishments to test one or more genera of plants. Applicants can choose to submit their varieties for testing by a CTC or continue to do the test themselves. Remember, using a CTC to test your variety is voluntary.

The use of CTCs recognises the advantages of testing a larger number of candidate varieties (with a larger number of comparators) in a single comprehensive trial. Not only is there an increase in scientific rigour but also there are substantial economies of scale and commensurate cost savings. A CTC will establish, conduct and report each trial on behalf of the applicant.

The PBR office has amended its fees so that cost savings can be passed to applicants who choose to test their varieties in a CTC. Accordingly, when 5 or more candidate varieties of the same genus are tested simultaneously, each will qualify for the CTC examination fee of $800. This is a saving of nearly 40% over the normal fee of $1400.

Trials containing less than 5 candidate varieties capable of being examined simultaneously will not be considered as Centralised test trials regardless of the authorisation of the facility. Candidate varieties in non-qualifying small trials will not qualify for CTC reduction of examination fees.

Establishments wishing to be authorised as a CTC may apply in writing to the PBR office outlining their claims against the selection criteria. Initially, only one CTC will be authorised for each genus. Exemptions to this rule can be claimed due to special circumstances, industry needs and quarantine regulations. Authorisations will be reviewed periodically.

Authorisation of CTCs is not aimed solely at large research institutions. Smaller establishments with appropriate facilities and experience can also apply for CTC status. There is no cost for authorisation as a CTC.

APPLICATIONS FOR AUTHORISATION AS A 'CENTRALISED TESTING CENTRE'

Establishments interested in gaining authorisation as a Centralised Testing Centre should apply in writing addressing each of the Conditions and Selection Criteria outlined below.

Conditions and Selection Criteria

To be authorised as a CTC, the following conditions and criteria will need to be met:

Appropriate facilities
While in part determined by the genera being tested, all establishments must have facilities that allow the conduct and completion of moderate to large-scale scientific experiments without undue environmental influences. Again dependent on genera, a range of complementary testing and propagation facilities (e.g. outdoor, glasshouse, shadehouse, tissue culture stations) is desirable.

Experienced staff
Adequately trained staff, and access to appropriately accredited Qualified Persons, with a history of successful PVR/PBR applications will need to be available for all stages of the trial from planting to the presentation of the

analysed data. These staff will require the authority to ensure timely maintenance of the trial. Where provided by the PBR office, the protocol and technical guidelines for the conduct of the trial must be followed.

Substantial industry support
Normally the establishment will be recognised by a state or national industry society or association. This may include/be replaced by a written commitment from major nurseries or other applicants, who have a history of regularly making applications for PBR in Australia, to use the facility.

Capability for long-term storage of genetic material
Depending upon the genus, a CTC must be in a position to make a long-term commitment to collect and maintain, at minimal cost, genetic resources of vegetatively propagated species as a source of comparative varieties. Applicants indicating a willingness to act as a national genetic resource centre in perpetuity will be favoured.

Contract testing for 3rd Parties
Unless exempted in writing by the PBR office operators of a CTC must be prepared to test varieties submitted by a third party.

Relationship between CTC and 3rd Parties
A formal arrangement between the CTC and any third party including fees for service will need to be prepared and signed before the commencement of the trial. It will include among other things: how the plant material will be delivered (e.g. date, stage of development plant, condition etc); allow the applicant and/or their agent and QP access to the site during normal working hours; and release the use of all trial data to the owners of the varieties included in the trial.

One trial at a time
Unless exempted in writing by the PBR office, all candidates and comparators should be tested in a single trial.

One CTC per genus
Normally only one CTC will be authorised to test a genus. Special circumstances may exist (environmental factors, quarantine etc) to allow more than one CTC per genus, though a special case will need to be made to the PBR office. More than one CTC maybe allowed for roses.

One CTC may be authorised to test more than one genus.
Authorisations for each genus will be reviewed periodically.

Authorised Centralised Test Centres (CTCs)

Following publication of applications for accreditation and ensuing public comment, the following organisations/individuals are authorised to act as CTCs. Any special conditions are also listed.

Name	Location	Approved Genera	Facilities	Name of QP	Date of accreditation
Agriculture Victoria, National Potato Improvement Centre	Toolangi, VIC	Potato	Outdoor, field, greenhouse, tissue culture laboratory	R Kirkham	31/3/97
Bureau of Sugar Experiment Stations	Cairns, Tully, Ingham, Ayr, Mackay, Bundaberg, Brisbane QLD	*Saccharum*	Field, glasshouse, tissue culture, pathology	G Piperidis	30/6/97
Ag-Seed Research	Horsham and other sites	Canola	Field, glasshouse, shadehouse, laboratory and biochemical analyses	P Rudolph	30/6/97
Agriculture Western Australia	Northam WA	Wheat	Field, laboratory	D Collins	30/6/97
University of Sydney, Plant Breeding Institute	Camden, NSW	*Argyranthemum, Diascia, Mandevilla*	Outdoor, field, irrigation, greenhouses with controlled micro-climates, controlled environment rooms,	J Oates	30/6/97

			tissue culture, molecular genetics and cytology lab.		
Boulters Nurseries Monbulk Pty Ltd	Monbulk, VIC	Clematis	Outdoor, shadehouse, greenhouse	M Lunghusen	30/9/97
Geranium Cottage Nursery	Galston, NSW	Pelargonium	Field, controlled environment house	I Paananen	30/11/97
Agriculture Victoria	Hamilton, VIC	*Perennial ryegrass, tall fescue, tall wheat grass, white clover, Persian clover*	Field, shadehouse, glasshouse, growth chambers. Irrigation. Pathology and tissue culture. Access to DNA and molecular marker technology. Cold storage.	M Anderson	30/6/98
Koala Blooms	Monbulk, VIC	*Bracteantha*	Outdoor, irrigation	M Lunghusen	30/6/98
Redlands Nursery	Redland Bay, QLD	*Aglaonema*	Outdoor, shadehouse, glasshouse and indoor facilities	K Bunker	30/6/98
Protected Plant Promotions	Macquarie Fields , NSW	New Guinea Impatiens including *Impatiens hawkeri* and its hybrids	Glasshouse	I Paananen	30/9/98
University of Queensland, Gatton College	Lawes, QLD	Some tropical pastures	Field, irrigation, glasshouse, small phytotron, plant nursery & propagation, tissue culture, seed and chemical lab, cool storage	To be advised	30/9/98
Jan and Peter Iredell	Moggill, QLD	Bougainvillea	Outdoor, shadehouse	J Iredell	30/9/98
Protected Plant Promotions	Macquarie Fields, NSW	*Verbena*	Glasshouse	I Paananen	31/12/98
Avondale Nurseries Ltd	Glenorie, NSW	*Agapanthus*	Greenhouse, tissue culture with commercial partnership	I Paananen	31/12/98
Paradise Plants	Kulnura, NSW	*Camellia, Lavandula, Osmanthus, Ceratopetalum*	Field, glasshouse, shadehouse, irrigation, tissue culture lab	J Robb	31/12/98
Prescott Roses	Berwick, VIC	*Rosa*	Field, controlled environment greenhouses	C Prescott	31/12/98
F & I Baguley Flower and Plant Growers	Clayton South, VIC	*Euphorbia*	Controlled glasshouses, quarantine facilities, tissue culture	G Guy	31/3/99
Paradise Plants	Kulnura, NSW	*Limonium, Raphiolepis, Eriostemon, Lonicera Jasminum*	Field, glasshouse, shadehouse, irrigation, tissue culture lab	J Robb	30/6/00
Ramm Pty Ltd	Macquarie Fields, NSW	*Angelonia*	Glasshouse	I Paananen	30/6/00
Carol's Propagation	Alexandra Hills, QLD	*Cuphea, Anthurium*	Field beds, wide range of comparative varieties	C Milne D Singh	30/6/00
Queensland Department of Primary Industries, Redlands Research Station	Cleveland, QLD	*Cynodon, Zoysia* and other selected warm season-season turf and amenity species	Field, glasshouse, irrigation, tissue culture lab	D Loch	30/9/00

Luff Partnership	Kulnura, NSW	*Bracteantha*	Field beds, irrigation, shade house, propagation house, cool rooms,	I Dawson	31/12/00
Ramm Pty Ltd	Macquarie Fields, NSW	*Petunia, Calibrachoa*	Glasshouse	I Paananen J Oates	31/12/00
NSW Agriculture	Temora	*Triticum, Hordeum, Avena*	Field, irrigation, glasshouse, climate controlled areas	P Breust	31/3/01
Bywong Nursery	Bungendore NSW	*Leptospermum*	Field, shadehouse, greenhouse	P Ollerenshaw	31/3/01
S J Saperstein	Mullumbimby NSW	*Rhododendron* (vireya types)	Field and propagation facilities	S Saperstein	31/12/01
Redlands Nursery	Redland Bay, QLD	*Osteospermum, Rhododendron*	Outdoor, shadehouse, glasshouse and indoor facilities	K Bunker	31/3/02
Ramm Pty Ltd	Macquarie Fields, NSW	*Euphorbia*	Glasshouse	I Paananen	31/3/02
Oasis Horticulture Pty Ltd	Springwood,	*Impatiens, Euphorbia*	AQIS accredited quarantine facilities; glasshouse, shadehouse, field, tissue culture	B Sidebottom A Bernuetz M Hunt N Derera T Angus	30/9/02
Carol's Propagation	Alexandra Hills, QLD	*Dahlia*	Field beds, wide range of comparative varieties	C Milne D Singh	31/12/03
Carol's Propagation	Brookfield, QLD	*Anubias*	Glasshouse specifically designed for aquatic plants	C Milne D Singh	31/3/04
Queensland Department of Primary Industries, Maroochy Research Station	Nambour, QLD	*Ananas*	Field, plots, pots, shadehouse, temperature controlled glasshouse and tissue culture lab	G. Sanewski	31/3/04
Abulk Pty Ltd	Clarendon, NSW	*Dianella*	Normal nursery facilities with access to micro propagation.	I Paananen	31/3/04
Proteaflora Nursery Pty Ltd	Monbulk, VIC	*Plectranthus*	Fogged propagation house, greenhouses and irrigated outdoor facilities	Paul Armitage	30/6/04
Berrimah Agricultural Research Centre	Darwin	*Zingiber*	Irrigated shadehouse, outdoor facilities, cool storage, high level post entry quarantine facility, tissue culture lab, pathology and entomology diagnostic services	D Marcsik	30/9/04
Ball Australia	Keysborough, VIC	*Impatiens, Verbena*	Controlled climate glasshouse and environment rooms, germination chamber, quarantine house, cool storage, irrigation and outdoor facilities.	D. Nichols	30/9/04
Floreta Pty Ltd	Redland Bay QLD	*Bracteantha*	Purpose built, secure greenhouse, access to fog house, registered quarantine facility on site.	K Bunker	31/12/04
Boulevarde Nurseries Mildura Pty Ltd	Irymple VIC	*Zantedeschia*	Glasshouse, shade house, propagation facilities, field areas, irrigation, cool rooms, tissue culture lab, hydroponics,	K Mullins	31/12/04

			quarantine facilities		
Buchanan's Nursery	Hodgsonvale, QLD	*Prunus*	Outdoor facilities including a collection of 90 varieties of common knowledge.	P Buchanan	31/12/04
Ball Australia	Keysborough, VIC	*Calibrachoa, Osteospermum*	Controlled climate glasshouse and environment rooms, germination chamber, quarantine house, cool storage, irrigation and outdoor facilities.	D. Nichols	30/9/05
Queensland Department of Primary Industries, Southedge Research Centre	Mareeba, QLD	*Mangifera*	Glasshouse, shadehouse, laboratory complex including biotech, propagation , outdoor facilities	I Bally	30/09/05
Blueberry Farms of Australia	Corindi Beach NSW and optional sites Tumbarumba NSW and Tasmania	*Vaccinium*	Extensive irrigated growing beds. Birds, hail and frost protection. Post harvest facilities including cool rooms. Access to tissue culture laboratories.	I Paananen	15/10/07

The following applications are pending:

Name	**Location**	**Genera applied for**	**Facilities**	**Name of QP**
Ball Australia	Keysborough, VIC	*Kalanchoe*	Controlled climate glasshouse and environment rooms, germination chamber, quarantine house, cool storage, irrigation and outdoor facilities.	D. Nichols
Yates Botanical Pty Ltd	Somersby and Tuggerah, NSW	*Rosa*	Tissue culture lab, glasshouse, quarantine and nursery facilities	I Paananen
Aussie Winners Pty Ltd	Redland Bay, QLD	*Fuchsia*	Comprehensive growing facilities	I Paananen
Schreurs Australia Pty Ltd	Leppington, NSW	*Rosa*	Comprehensive growing facilities	I Paananen

Comments (both for or against) either the continued accreditation of a CTC or applications to become a CTC are invited. Written comments are confidential and should be addressed to:

The Registrar
Plant Breeder's Rights Office
IP Australia
PO Box 200
Woden, ACT 2606
Fax (02) 6283 7999

Closing date for comment: 30 April 2008.

APPENDIX 7 List of Classes for Variety Denomination Purposes

UPOV Variety Denomination Classes: (UPOV/INF/12/1: ANNEX I)

A Variety Denomination Should not be Used More than Once in the Same Class

For the purposes of providing guidance on the third and fourth sentences of paragraph 2 of Article 20 of the 1991 Act and of Article 13 of the 1978 Act and the 1961 Convention, variety denomination classes have been developed. A variety denomination should not be used more than once in the same class. The classes have been developed such that the botanical taxa within the same class are considered to be closely related and/or liable to mislead or to cause confusion concerning the identity of the variety.

The variety denomination classes are as follows:

(a) General Rule (one genus / one class): for genera and species not covered by the List of Classes in this Annex, a genus is considered to be a class;

(b) Exceptions to the General Rule (list of classes):

(i) classes within a genus: List of classes in this Annex: Part I;

(ii) classes encompassing more than one genus: List of classes in this Annex: Part II.

LIST OF CLASSES

Part I

Classes within a genus

	Botanical names	UPOV codes
Class 1.1	Brassica oleracea	BRASS_OLE
Class 1.2	Brassica other than Brassica oleracea	other than BRASS_OLE
Class 2.1	Beta vulgaris L. var. alba DC., Beta vulgaris L. var. altissima	BETAA_VUL_GVA; BETAA_VUL_GVS
Class 2.2	Beta vulgaris ssp. vulgaris var. conditiva Alef. (syn.: B. vulgaris L. var. rubra L.), B. vulgaris L. var. cicla L., B. vulgaris L. ssp. vulgaris var. vulgaris	BETAA_VUL_GVC; BETAA_VUL_GVF
Class 2.3	Beta other than classes 2.1 and 2.2.	other than classes 2.1 and 2.2
Class 3.1	Cucumis sativus	CUCUM_SAT
Class 3.2	Cucumis melo	CUCUM_MEL
Class 3.3	Cucumis other than classes 3.1 and 3.2	other than classes 3.1 and 3.2
Class 4.1	Solanum tuberosum L.	SOLAN_TUB
Class 4.2	Solanum other than class 4.1	other than class 4.1

LIST OF CLASSES (Continuation)

Part II

Classes encompassing more than one genus

	Botanical names	UPOV codes
Class 201	Secale, Triticale, Triticum	SECAL; TRITL; TRITI
Class 202	Panicum, Setaria	PANIC; SETAR
Class 203*	Agrostis, Dactylis, Festuca, Festulolium, Lolium, Phalaris, Phleum and Poa	AGROS; DCTLS; FESTU; FESTL; LOLIU; PHALR; PHLEU; POAAA
Class 204*	Lotus, Medicago, Ornithopus, Onobrychis, Trifolium	LOTUS; MEDIC; ORNTP; ONOBR; TRFOL
Class 205	Cichorium, Lactuca	CICHO; LACTU
Class 206	Petunia and Calibrachoa	PETUN; CALIB
Class 207	Chrysanthemum and Ajania	CHRYS; AJANI
Class 208	(Statice) Goniolimon, Limonium, Psylliostachys	GONIO; LIMON; PSYLL_
Class 209	(Waxflower) Chamelaucium, Verticordia	CHMLC; VERTI; VECHM
Class 210	Jamesbrittania and Sutera	JAMES; SUTER
Class 211	Edible Mushrooms	
	Agaricus bisporus	AGARI_BIS
	Agaricus blazei	AGARI_BLA
	Agrocybe cylindracea	AGROC_CYL
	Auricularia auricura	AURIC_AUR
	Auricularia polytricha (Mont.) Sscc.	AURIC_POL
	Dictyophora indusiata (Ventenat:Persoon) Fischer	DICTP_IND
	Flammulina velutipes	FLAMM_VEL
	Ganoderma lucidum (Leyss:Fries) Karsten	GANOD_LUC
	Grifola frondosa	GRIFO_FRO
	Hericium erinaceum	HERIC_ERI
	Hypsizigus marmoreus	HYPSI_MAR
	Hypsizigus ulmarius	HYPSI_ULM
	Lentinula edodes	LENTI_ELO
	Lepista nuda (Bulliard:Fries) Cooke	LEPIS_NUD
	Lepista sordida (Schumacher:Fries) Singer	LEPIS_SOR
	Lyophyllum decastes	LYOPH_DEC
	Lyophyllum shimeji (Kawamura) Hongo	LYOPH_SHI
	Meripilus giganteus (Persoon:Fries) Karten	MERIP_GIG
	Mycoleptodonoides aitchisonii (Berkeley) Maas Geesteranus	MYCOL_AIT
	Naematoloma sublateritium	NAEMA_SUB
	Panellus serotinus	PANEL_SER
	Pholiota adiposa	PHLIO_ADI
	Pholiota nameko	PHLIO_NAM
	Pleurotus cornucopiae var.citrinooileatus	PLEUR_COR
	Pleurotus cystidiosus	PLEUR_CYS
	Pleurotus cystidiosus subsp. Abalonus	PLEUR_CYS_ABA
	Pleurotus eryngii	PLEUR_ERY
	Pleurotus ostreatus	PLEUR_OST
	Pleurotus pulmonarius	PLEUR_PUL
	Polyporus tuberaster (Jacquin ex Persoon) Fries	POLYO_TUB
	Sparassis crispa (Wulfen) Fries	SPARA_CRI
	Tricholoma giganteum Massee	MACRO_GIG

* Classes 203 and 204 are not solely established on the basis of closely related species.

APPENDIX 8

REGISTER OF PLANT VARIETIES

Register of Plant Varieties contains the legal description of the varieties granted Plant Breeder's Rights. A person may inspect the Register at any reasonable time. Following are the contact details for Registers (1988-2000) kept in each state and territories*

South Australia

Ms Lisa Halskov
AQIS
8 Butler Street
PORT ADELAIDE SA 5000
Phone 08 8305 9706

New South Wales

Mr. Alex Jabs
General Services
AQIS
2 Hayes Road
ROSEBERY NSW 2018
Phone 02 9364 7293

Victoria and Tasmania

Mr. Colin Hall
AQIS
Building D, 2nd Floor
World Trade Centre
Flinders Street
MELBOURNE VIC 3005
Phone 03 9246 6810

Queensland

Mr. Ian Haseler
AQIS
2nd Floor
433 Boundary Street
SPRING HILL QLD 4000
Phone 07 3246 8755

Australian Capital Territory, Northern Territory and Western Australia
ACT and NT Registers are kept
in the Library of PBR Office in Canberra
Phone (02) 6283 2999

* In accordance with an amendment to section 61 of Plant Breeder's Rights Act, from 2002 the Register of Plant Varieties will be available from the Library of PBR Office in Canberra. The Register is also electronically available from the PBR website at http://pbr.ipaustralia.plantbreeders.gov.au/

Plant Varieties Journal Mailing List

The Plant Varieties Journal mailing list informs subscribers whenever the new journal is posted on the IP Australia web site.

- Home