

Subgroup	N	Exposed (FIS)	Deaths	HR	95% CI	p-value	p-interaction	
Age group (years)								
55–75	1124	15	254	1.14	0.41...3.15	0.804		
<65	1496	38	202	0.96	0.38...2.41	0.924		
>75	464	14	147	0.79	0.25...2.49	0.688		
LVEF category (%)								
30–35	1321	26	200	0.71	0.25...2.01	0.518		
<30	1763	41	403	1.10	0.55...2.19	0.783		
NYHA class							0.236	
III...IV	928	21	251	0.61	0.22...1.68	0.337		
I...II	2156	46	352	1.37	0.68...2.73	0.376		
Atrial fibrillation/flutter							0.845	
Yes	1143	34	206	0.96	0.38...2.37	0.921		
Diabetes mellitus								0.576
Yes	1839	44	318	1.15	0.55...2.41	0.706		
Sex								0.258
Female	402	8	74	0.35	0.05...2.66	0.309		
Male	2682	59	529	1.18	0.65...2.14	0.585		
HR = Hazard Ratio; CI = Confidence Interval; FIS = patients with inappropriate ICD shock; Deaths = all-cause deaths in subgroup. Each HR reflects the within-subgroup association between inappropriate ICD shock and all-cause mortality. p-interaction tests whether the effect of shock differs across subgroup levels (Step 9); reported once per subgroup variable.								