

Generalization Bounds via Validation

Thomas Walker

1 NOTATION

- $x \in \mathcal{X}$, input.
- $y \in \mathcal{Y}$, target.
- \mathcal{L} , loss function.
- $\mathcal{R}[f] = \mathbb{E}_{x,y \sim \mathbb{P}_{(\mathcal{X}, \mathcal{Y})}} (\mathcal{L}(f(x), y))$, expected risk of a function f . Where $\mathbb{P}_{(\mathcal{X}, \mathcal{Y})}$ is the true distribution.
- $f_{\mathcal{A}(S)} : \mathcal{X} \rightarrow \mathcal{Y}$, function learnt by a learning algorithm \mathcal{A} on training set $S := S_m := \{(x_1, y_1), \dots, (x_m, y_m)\}$
- \mathcal{F} , a characterized set of functions known as the hypothesis space.
- $\mathcal{L}_{\mathcal{F}} := \{g : g \in \mathcal{F}, g(x, y) = \mathcal{L}(f(x), y)\}$, the family of loss functions associated to \mathcal{F} .
- For vector v its dimension is d_v .

2 SETUP

Machine learning aims to minimize $\mathcal{R}(f_{\mathcal{A}(S)})$. However, this is **non-computable** as $\mathbb{P}_{(\mathcal{X}, \mathcal{Y})}$ is unknown. Therefore, one minimizes the empirical risk

$$\mathcal{R}_S(f_{\mathcal{A}(S)}) = \frac{1}{|S|} \sum_{(x,y) \in S} \mathcal{L}(f_{\mathcal{A}(S)}(x), y),$$

where the generalization gap is given by $\mathcal{R}(f_{\mathcal{A}(S)}) - \mathcal{R}_S(f_{\mathcal{A}(S)})$.

3 GENERALIZATION BOUNDS VIA VALIDATION

The training-validation paradigm involves holding out a validation set to optimize model architecture. Giving rise to the hypothesis that *deep neural networks can obtain good generalization error by performing a model search on the validation set*.

AFFILIATION Imperial College London

CORRESPONDENCE thomas.walker21@imperial.ac.uk

DATE July 2023

Proposition 3.1 *Let $S_{m_{\text{val}}}^{(\text{val})}$ be a held-out validation set, where $|S_{m_{\text{val}}}^{(\text{val})}| = m_{(\text{val})}$. Assume that $m_{(\text{val})}$ is an i.i.d sample from $\mathbb{P}_{(X,Y)}$. Let $\kappa_{f,i} = \mathcal{R}(f) - \mathcal{L}(f(x_i), y_i)$ for $(x_i, y_i) \in S_{m_{\text{val}}}^{(\text{val})}$. Suppose that $\mathbb{E}(\kappa_{f,i}^2) \leq \gamma^2$ and $|\kappa_{f,i}| \leq C$ almost surely for all $(f, i) \in \mathcal{F}_{\text{val}} \times \{1, \dots, m_{\text{val}}\}$. Then, for $\delta \in (0, 1]$, with probability $1 - \delta$*

$$\mathcal{R}(f) \leq \mathcal{R}_{S_{m_{\text{val}}}^{(\text{val})}}(f) + \frac{2C \log\left(\frac{|\mathcal{F}_{\text{val}}|}{\delta}\right)}{3m_{\text{val}}} + \sqrt{\frac{2\gamma^2 \log\left(\frac{|\mathcal{F}_{\text{val}}|}{\delta}\right)}{m_{\text{val}}}}$$

holds for all $f \in \mathcal{F}_{\text{val}}$.

Remarks 3.2

- \mathcal{F}_{val} is independent of $S_{m_{\text{val}}}^{(\text{val})}$.
- The bound is only dependent on the validation error on $S_{m_{\text{val}}}^{(\text{val})}$.

Implementation 3.3