

CS 613 - Machine Learning

Assignment 2 - Logistic Regression

Robert Thompson

1 Theory

1. For the function $J = (x_1w_1 - 5x_2w_2 - 2)^2$, where $w = [w_1, w_2]$ are our weights to learn:

(a) What are the partial gradients, $\frac{\partial J}{\partial w_1}$ and $\frac{\partial J}{\partial w_2}$?

i. Partial Gradient of $\frac{\partial J}{\partial w_1}$:

A. Chain Rule

$$\frac{\partial J}{\partial w_1} = 2(x_1w_1 - 5x_2w_2 - 2) \frac{\partial J}{\partial w_1} (x_1w_1 - 5x_2w_2 - 2)$$

B. Sum/Difference Rule

$$\frac{\partial J}{\partial w_1} (x_1w_1 - 5x_2w_2 - 2) = \frac{\partial J}{\partial w_1} (x_1w_1) - \frac{\partial J}{\partial w_1} (5x_2w_2) - \frac{\partial J}{\partial w_1} (2)$$

$$\frac{\partial J}{\partial w_1} (x_1w_1) = x_1$$

$$\frac{\partial J}{\partial w_1} (5x_2w_2) = 0$$

$$\frac{\partial J}{\partial w_1} (2) = 0$$

$$= x_1 - 0 - 0$$

$$= x_1$$

C. Partial Gradient $\frac{\partial J}{\partial w_1} = 2x_1(x_1w_1 - 5x_2w_2 - 2)$

i. Partial Gradient of $\frac{\partial J}{\partial w_2}$:

A. Chain Rule

$$\frac{\partial J}{\partial w_2} = 2(x_1w_1 - 5x_2w_2 - 2) \frac{\partial J}{\partial w_2} (x_1w_1 - 5x_2w_2 - 2)$$

B. Sum/Difference Rule

$$\frac{\partial J}{\partial w_2} (x_1w_1 - 5x_2w_2 - 2) = \frac{\partial J}{\partial w_2} (x_1w_1) - \frac{\partial J}{\partial w_2} (5x_2w_2) - \frac{\partial J}{\partial w_2} (2)$$

$$\frac{\partial J}{\partial w_2} (x_1w_1) = 0$$

$$\frac{\partial J}{\partial w_2} (5x_2w_2) = 5x_2$$

$$\frac{\partial J}{\partial w_2} (2) = 0$$

$$= 0 - 5x_2 - 0$$

$$= -5x_2$$

C. Partial Gradient $\frac{\partial J}{\partial w_2} = -10x_2(x_1w_1 - 5x_2w_2 - 2)$

(b) What are the value of the partial gradients given current values of $w = [0, 0]$, $x = [1, 1]$?

i. Let $\frac{\partial J}{\partial w_1} = 2x_1(x_1w_1 - 5x_2w_2 - 2)$

A. Plug in $w = [0, 0]$, $x = [1, 1]$

$$\frac{\partial J}{\partial w_1} = 2(1)(1 * 0 - 5(1)(0) - 2)$$

B. Partial Gradient Value of $\frac{\partial J}{\partial w_1} = -4$


ii. Let $\frac{\partial J}{\partial w_2} = -10x_2(x_1w_1 - 5x_2w_2 - 2)$

- A. Plugin $w = [0, 0], x = [1, 1]$
 $\frac{\partial J}{\partial w_2} = -10(1)(1 * 0 - 5(1)(0) - 2)$
- B. Partial Gradient Value of $\frac{\partial J}{\partial w_2} = 20$

2 Spambase Logistic Regression Classifier

1. Plot of Training and Validation Log Loss as a Function of the Epoch

- Learning Rate: 0.1
- Epochs: 10,000
- Stability Constant: $10e - 7$



2. Training Statistics

- Precision: 0.9267676767676768
- Recall: 0.8900565885206144
- F-Measure: 0.9080412371134021


- Accuracy: 0.9272905119008803

3. Validation Statistics

- Precision: 0.9092495636998255
- Recall: 0.9045138888888888
- F-Measure: 0.9068755439512619
- Accuracy: 0.9302477183833116

4. Training and Validation Precision-Recall Graph

- Learning Rate: 0.1
- Epochs: 10,000
- Stability Constant: $10e - 7$



3 Logistic Regression for Multi-Class Classification

1. Validation Accuracy: 0.96
2. Validation Confusion Matrix

- $$\begin{bmatrix} 19 & 0 & 0 \\ 0 & 15 & 2 \\ 0 & 0 & 14 \end{bmatrix}$$