

STAT 345/445 Lecture 9

Differentiating under an integral sign – Section 2.4

- A technical section on when we can switch the order of limits, integrals, and sums

Moment Generating functions

- Proof of

$$\left. \frac{d}{dt} M_X(t) \right|_{t=0} = E(X)$$

goes like this:

$$\frac{d}{dt} M_X(t) = \frac{d}{dt} \int_{-\infty}^{\infty} \overbrace{e^{tx} f(x)}^{M(t)} dx = \int_{-\infty}^{\infty} \underbrace{\left(\frac{\partial}{\partial t} e^{tx} f(x) \right)}_{\text{if we can differentiate under the integral sign}} dx$$

if we can *differentiate under the integral sign*

$$\Rightarrow \frac{d}{dt} M_X(t) = \int_{-\infty}^{\infty} \overbrace{xe^{tx} f(x)}^{M'(t)} dx = E(Xe^{tx})$$

$$\Rightarrow \left. \frac{d}{dt} M_X(t) \right|_{t=0} = E(Xe^{0 \cdot X}) = E(X)$$

$e^{tx} f(x)$ is
a function of
both t and x

Differentiating under an integral sign

Theorem: (Simplified Leibnitz's Rule)

If $f(x, \theta)$ is differentiable with respect to θ and a and b are constants then

$$\frac{d}{d\theta} \overbrace{\int_a^b f(x, \theta) dx}^{\text{func. of } \theta} = \int_a^b \underbrace{\frac{\partial}{\partial \theta}}_{\uparrow} f(x, \theta) dx$$

- Finite range integral \Rightarrow can switch the order of $\frac{d}{d\theta}$ and $\int dx$
- If $a = -\infty$ and/or $b = \infty$ we have to be careful
 - Recall: A derivative is a limit
 - \Rightarrow The issue is actually: When can we change the order of limits and integration?

What is it we want to do?

$$\begin{aligned}\infty + \infty &= \infty \\ -\infty - \infty &= -\infty \\ \infty - \infty &\text{ not defined,}\end{aligned}$$

- Recall:

$$\frac{\partial}{\partial \theta} f(x, \theta) = \lim_{\delta \rightarrow 0} \frac{f(x, \theta + \delta) - f(x, \theta)}{\delta}$$

Therefore:

$$\int_{-\infty}^{\infty} \frac{\partial}{\partial \theta} f(x, \theta) dx = \int_{-\infty}^{\infty} \lim_{\delta \rightarrow 0} \frac{f(x, \theta + \delta) - f(x, \theta)}{\delta} dx$$

$$\begin{aligned}\frac{d}{d\theta} \int_{-\infty}^{\infty} f(x, \theta) dx &= \lim_{\delta \rightarrow 0} \frac{\int_{-\infty}^{\infty} f(x, \theta + \delta) dx - \int_{-\infty}^{\infty} f(x, \theta) dx}{\delta} \\ &= \lim_{\delta \rightarrow 0} \int_{-\infty}^{\infty} \frac{f(x, \theta + \delta) - f(x, \theta)}{\delta} dx\end{aligned}$$

When can we differentiate under the integral sign?

Theorem

If $f(x, \theta)$ is differentiable with respect to θ and there exists a constant $\delta_0 > 0$ and a function $g(x, \theta)$ that satisfies

$$(i) \left| \frac{\partial}{\partial \theta} f(x, \theta) \right|_{\theta=\theta'} \leq g(x, \theta)$$

for all θ' such that $|\theta' - \theta| \leq \delta_0$

$$(ii) \int_{-\infty}^{\infty} g(x, \theta) dx < \infty$$

then

$$\frac{d}{d\theta} \int_{-\infty}^{\infty} f(x, \theta) dx = \int_{-\infty}^{\infty} \frac{\partial}{\partial \theta} f(x, \theta) dx$$

derivative of $f(x, \theta)$
has to behave!
need to be
bounded by
some function

When can we differentiate under the integral sign?

- Grad: Understand that conditions (i) and (ii) basically mean:
 - The (partial) derivative $\frac{\partial}{\partial \theta} f(x, \theta)$ has to behave!
 - It has to be dominated by a function $g(x, \theta)$ that has a finite integral (w.r.t. x)
 - at least at some θ' close to θ
- UG: Just remember that changing the order of a derivative and an integral with an infinite range can't always be done

Example

- Let $X \sim \text{Expo}(\lambda)$. The pdf for X is

$$f(x) = \frac{1}{\lambda} e^{-x/\lambda} \quad \text{for } x > 0$$

Show that for an integer $n \geq 1$

$$E(X^{n+1}) = \lambda E(X^n) + \lambda^2 \frac{d}{d\lambda} E(X^n)$$

Derivatives and infinite sums

- Finite sums are no problem:

$$\frac{d}{d\theta} \sum_{x=0}^n f(x, \theta) = \sum_{x=0}^n \frac{\partial}{\partial \theta} f(x, \theta)$$

- When does the following hold?

$$\frac{d}{d\theta} \sum_{x=0}^{\infty} f(x, \theta) = \sum_{x=0}^{\infty} \frac{\partial}{\partial \theta} f(x, \theta)$$

- Basically: Both series have to converge
See Theorem 2.4.8 in the textbook

Example: Geometric distribution

- Let $X \sim \text{Geometric}(\theta)$. The pmf for X is

$$f(x) = \theta(1 - \theta)^x \quad \text{for } x = 0, 1, 2, \dots \text{ and } 0 < \theta < 1$$

- Lets see where this takes us:

$$\frac{d}{d\theta} \sum_{x=0}^{\infty} f(x)$$

- Convenient facts about the **geometric series**

$$\sum_{x=0}^{\infty} r^x = \frac{1}{1-r} \quad \text{for } |r| < 1$$

$$\sum_{x=0}^n r^x = \frac{1 - r^{n+1}}{1 - r} \quad \text{for } r \neq 1$$