


Microtubule + PD-(L)1: 2 molecules stopped early

3 stopped trials across 2 molecules: Paclitaxel and Belantamab Mafodotin. Against 5.0% for PD-(L)1 combinations without Microtubule and 4.3% across all oncology Phase 2,3 trials in the same window. Rates count trials that stopped early for efficacy, safety or benefit–risk reasons; trials that completed and missed their endpoints are not counted.

1.5% 3 of 200 closed trials (95% CI 0.5%–4.3%)	3 / 0 / 0 efficacy / safety / both (adds to 3)	29.5% of 677 trials have closed · 477 still open or unresolved	0.6% stopped for a biological reason within 3 years of starting (95% CI 0.1%–2.2%; 301 still at risk; 1.5% area-wide)
--	--	--	---

How this compares



What is in this class: 369 distinct experimental drugs across 677 trials (antibody 648, small molecule 606, adc 121, unknown 95). Most tested: PACLITAXEL (420), CARBOPLATIN (192), PEMBROLIZUMAB (172), CISPLATIN (138), DOCETAXEL ANHYDROUS (114). A class groups drugs by what they act on; check that the grouping is one you would make before reading the rate as a property of the mechanism.

How much does this rest on one decision? The 3 stops came from 3 sponsor-asset programmes across 3 sponsors; the largest (Tufts Medical Center) contributed 1; removing that programme's trials from both sides leaves 1.0%. Screened alongside 85 other segments in this area, it does not survive a 10% false-discovery correction ($q=1$, Benjamini-Yekutieli). A further 18 closed trials here were terminated with no cause recorded in the registry; if every one of them were biological the rate would be 10.5%. Read it as a screen worth checking against the underlying trials.

Bars show the share of closed trials stopped for biological reasons. Confidence intervals overlap where sample sizes are small — read the counts, not just the bars.

By start cohort

Trials started	Stops	Closed	Rate	95% CI
2015–2017	0	59	0.0%	0.0%–6.1%
2018–2020	1	79	1.3%	0.2%–6.8%
2021–2024	2	62	3.2%	0.9%–11.0%

Where the stops sit

Efficacy / futility: 3.
Sponsors with most stops: Tufts Medical Center (1), GSK (1), Novartis (1).

The stopped trials

Trial	Ph	Sponsor	Experimental drugs	Stopped	Type	Registry stop reason
NCT04856176	2	Tufts Medical Center	Granulocyte-Macrophage Colony-Sti..., PEMBROLIZUMAB, PEMETREXED, PACLITAXEL	2024-12	Efficacy	Futility
NCT06655818	2	GSK	BELANTAMAB MAFODOTIN, DOSTARLIMAB	2024-02	Efficacy	The study was terminated due to lack of efficacy
NCT03631199	3	Novartis	CANAKINUMAB, PEMBROLIZUMAB, CARBOPLATIN, CISPLATIN	2021-08	Efficacy	Results of primary analysis showed addition of canakinumab to combination treatment did not improve tumor response or overall survival; the decision to stop the trial was not due to safety concerns

Method

Denominator: ClinicalTrials.gov interventional Phase 2,3 oncology trials started 2015–2024 that have closed (completed or terminated).
Numerator: terminated trials whose registry stop reason is classified as biological (efficacy, safety or benefit–risk) by Classification V2 — held-out precision 95.5%, recall 95.3% (n=600). Drugs are linked to ChEMBL and the NCI Thesaurus; 76% of industry oncology trials in this window carry a resolved drug target, and a trial without one cannot enter a mechanism class. Intervals are Wilson 95%.

Limits

Not a failure rate: trials that completed with negative results are not counted, and programs discontinued after a completed trial do not appear. Stop reasons are sponsor-reported. This is a closed-trial proportion, not a time-to-event analysis: only 29.5% of trials in this segment have closed, and a trial that stops early enters the denominator sooner than one that runs to completion, which can inflate the rate in immature segments. Recent cohorts have fewer closed trials, so their rates are less stable. Research signals, not clinical or investment advice.

Any mechanism, sponsor or indication, updated weekly. The dataset behind this brief covers every stopped oncology trial with an efficacy or safety signal plus the full denominator universe. Free sample and licensing: clinicaltrialfailures.com/pricing

Sources: ClinicalTrials.gov (NLM); ChEMBL (EMBL-EBI, CC BY-SA 3.0); NCI Thesaurus (NCI); RxNorm/RxClass (NLM); SEC EDGAR. Classification, linkage and rates by Clinical Trial Failures. Rebuilt weekly; this brief covers trials started 2015–2024.