

Monoamine reuptake: 9 of 22 completed trials missed their primary endpoint

9 completed trials across 7 sponsors and 7 molecules ran to the end and missed their primary endpoint on the sponsor's own posted result; 11 met it. Only 3 of 68 closed trials were stopped early for efficacy or safety, which is why this class is read through its completed trials rather than a stop rate. Most completed trials post nothing readable, so the count is a floor.

9 of 22	7 / 7	4.4%	74.7%
completed trials with a readable primary result missed it · 11 met it	sponsors / molecules behind the misses	stopped early: 3 of 68 closed trials (95% CI 1.5%–12.2%)	of 91 trials have closed · 23 still open or unresolved

9 of 22


Completed and missed the primary endpoint. Of 58 completed trials in this segment, 22 posted a primary result on ClinicalTrials.gov that can be read — the sponsor's own comparison held to the sponsor's own threshold, or its statement that the endpoint was missed: 9 missed and 11 met, 2 split across co-primary endpoints. Counted separately and never in the rate above; a floor, not a rate. 12 of the 54 completed trials that finished more than a year ago have posted no results at all — a gap in what can be known, not a sign of failure.

The completed trials that missed

Trial	Ph	Sponsor	Experimental drugs	Started	What the posted result says
NCT04885257	2	VA Office of Research and Development	METHYLPHENIDATE	2022-01	Change in Modified Rankin Scale at 12 Weeks — p 0.896 against 0.05: not significant. Method: Wilcoxon (Mann-Whitney).
NCT04437485	2	Indiana University	BUPROPION	2020-10	Hemoglobin A1c at 6 Months — p 0.64 against 0.05: not significant. Method: ANCOVA.
NCT04345471	3	Mochida Pharmaceutical Company, Ltd.	DESVENLAFAXINE	2020-05	Changes in Total MADRS Score From the Baseline to Week 8 Visit During the Treatment Period — p 0.509 against the sponsor's stated threshold of 0.05: not significant. LS mean difference -0.6 (95% CI -2.5 to 1.2). Method: MMRM.
NCT03569475	3	AbbVie	LEVOMILNACIPRAN	2018-07	Change From Baseline in Children's Depression Rating Scale-Revised (CDRS-R) — p 0.2215 against 0.05: not significant. Least Square Mean Difference -1.7 (95% CI -4.49 to 1.04). Method: Mixed Model for Repeated Measures.
NCT03315793	3	Shionogi	DULOXETINE	2017-12	Change From Baseline on the Children's Depression Rating Scale-Revised (CDRS-R) Total Score — p 0.5587 against 0.05: not significant. Least squares mean difference 1.39 (98% CI -3.3 to 6.08). Method: mixed-effects model repeated measures.
NCT02600494	3	Intra-Cellular Therapies, Inc.	LUMATEPERONE	2015-12	Change From Baseline to Day 43 in Montgomery-Åsberg Depression Rating Scale (MADRS) Total Score — p 0.514 against 0.05: not significant. Least Squares Mean Difference 0.9 (95% CI -1.83 to 3.53). Method: Mixed Effects Model for Repeated Measure.

The 6 most recently started of 9 completed trials that missed. The diligence report lists every one, with the numbers or the sentence each verdict was read from.

How this compares



What is in this class: 119 distinct experimental drugs across 91 trials (small molecule 91, unknown 3). Most tested: LUMATEPERONE (17), VORTIOXETINE (14), MIDOMAFETAMINE (11), DULOXETINE (11), SERTRALINE (7). A class groups drugs by what they act on; check that the grouping is one you would make before reading the rate as a property of the mechanism.

How much does this rest on one decision? The 3 stops came from 2 sponsor-asset programmes across 2 sponsors; the largest (H. Lundbeck A/S) contributed 2; removing that programme's trials from both sides leaves 1.6%. Screened alongside 12 other segments, the rate is within what screening that many classes could produce ($q=1$); the molecules and trials that failed here are the record either way. A further 1 closed trials here were terminated with no cause recorded in the registry; if every one of them were biological the rate would be 5.9%. Read it as a screen worth checking against the underlying trials.

Bars show the share of closed trials stopped for biological reasons. Confidence intervals overlap where sample sizes are small — read the counts, not just the bars.

The stopped trials

Trial	Ph	Sponsor	Experimental drugs	Stopped	Type	Registry stop reason
NCT05611749	2	Scripps Health	DULOXETINE	2024-03	Safety	Enrollment was extremely low and there were too many adverse events.
NCT05014919	3	H. Lundbeck A/S	VORTIOXETINE	2022-03	Efficacy	The study was terminated based on new efficacy data from another study.
NCT02871297	3	H. Lundbeck A/S	VORTIOXETINE	2022-03	Efficacy	The study was terminated based on new efficacy data from another study.

Method

Denominator: ClinicalTrials.gov interventional Phase 2,3 psychiatry & mental health trials started 2015–2024 that have closed (completed or terminated). Numerator: terminated trials whose registry stop reason is classified as biological (efficacy, safety or benefit–risk) by Classification V2 — held-out precision 95.5%, recall 95.3% ($n=600$). Drugs are linked to ChEMBL and the NCI Thesaurus; 44% of industry psychiatry & mental health trials in this window carry a resolved drug target, and a trial without one cannot enter a mechanism class. Intervals are Wilson 95%.

Limits

Not a failure rate: completed trials that missed their endpoints are not in the rate. They are counted separately where the sponsor posted a primary result that can be read — most did not, so that count is a floor — and programmes discontinued after a completed trial do not appear. Stop reasons are sponsor-reported. This is a closed-trial proportion, not a time-to-event analysis: only 74.7% of trials in this segment have closed, and a trial that stops early enters the denominator sooner than one that runs to completion, which can inflate the rate in immature segments. Recent cohorts have fewer closed trials, so their rates are less stable. Research signals, not clinical or investment advice.

The whole cohort, trial by trial. The diligence report names every trial behind these figures — the ones still running and the ones that completed and missed their primary endpoint included — and places your molecule against each one that failed. €99, delivered immediately: clinicaltrialfailures.com/asset-check

By start cohort

Trials started	Stops	Closed	Rate	95% CI
2015–2017	1	29	3.4%	0.6%–17.2%
2018–2020	0	17	0.0%	0.0%–18.4%
2021–2024	2	22	9.1%	2.5%–27.8%

Where the stops sit

Efficacy / futility: 2, Safety: 1.
Sponsors with most stops: H. Lundbeck A/S (2), Scripps Health (1).