CLINICAL TRIAL FAILURES · DISCONTINUATION RATE · NEUROLOGY PHASE 2,3 · STARTS 2015–2024

GABA-A: 3 molecules stopped early, all small molecules

3 stopped trials, but 3 molecules — all small molecules: Clorazepic Acid, Darigabat and Ganaxolone. 1 of the stops were the trial's own verdict and 2 followed a decision taken elsewhere. Against 9.6% for all neurology trials with a resolved drug target and 6.5% across all neurology Phase 2,3 trials in the same window. The rate counts trials that stopped early for efficacy, safety or benefit–risk reasons. Trials that ran to completion and missed their primary endpoint are counted separately below, never in the rate.

7.9%
3 of 38 closed trials (95% CI 2.7%–20.8%)


3 / 0 / 0
efficacy / safety / both (adds to 3)

77.6%
of 49 trials have closed · 11 still open or unresolved

4.8%
stopped for a biological reason within 3 years of starting (95% CI 1.2%–17.8%; 18 still at risk; 3.4% area-wide)

2 of 3 **Completed and missed the primary endpoint.** Of 29 completed trials in this segment, 3 posted a primary result on ClinicalTrials.gov that can be read — the sponsor's own comparison held to the sponsor's own threshold, or its statement that the endpoint was missed: 2 missed and 1 met. Counted separately and never in the rate above; a floor, not a rate. 5 of the 26 completed trials that finished more than a year ago have posted no results at all — a gap in what can be known, not a sign of failure.

How this compares



What is in this class: 110 distinct experimental drugs across 49 trials (small molecule 49, unknown 6). Most tested: DIAZEPAM (9), GANAXOLONE (7), CENOBAMATE (7), PADSEVONIL (5), CLOBAZAM (4). A class groups drugs by what they act on; check that the grouping is one you would make before reading the rate as a property of the mechanism.

How much does this rest on one decision? The 3 stops came from 3 sponsor-asset programmes across 3 sponsors; the largest (Marinus Pharmaceuticals) contributed 1; removing that programme's trials from both sides leaves 6.1%. Screened alongside 27 other segments, the rate is within what screening that many classes could produce (q=1); the molecules and trials that failed here are the record either way. A further 5 closed trials here were terminated with no cause recorded in the registry; if every one of them were biological the rate would be 21.1%. Read it as a screen worth checking against the underlying trials.

Bars show the share of closed trials stopped for biological reasons. Confidence intervals overlap where sample sizes are small — read the counts, not just the bars.

By start cohort

Trials started	Stops	Closed	Rate	95% CI
2015–2017	1	17	5.9%	1.0%–27.0%
2018–2020	1	15	6.7%	1.2%–29.8%
2021–2024	1	6	16.7%	3.0%–56.4%

Where the stops sit

Efficacy / futility: 3.
Sponsors with most stops: AbbVie (1), Assistance Publique - Hôpitaux de Paris (1), Marinus Pharmaceuticals (1).

The stopped trials

Trial	Ph	Sponsor	Experimental drugs	Stopped	Type	Registry stop reason
NCT04686786	2	AbbVie ABBV	DARIGABAT	2024-12	Efficacy	Closed early after CVL-865-SZ-001 (NCT04244175) did not meet its primary objective

Trial	Ph	Sponsor	Experimental drugs	Stopped	Type	Registry stop reason
NCT04726592	3	Assistance Publique - Hôpitaux de Paris	CLORAZEPIC ACID	2024-04	Efficacy	It has been determined that continuing with enrollments in this study is no longer justified due to the lack of significant impact on the expected outcomes.
NCT02519439	3	Marinus Pharmaceuticals	GANAXOLONE	2016-12	Efficacy	Ganaxolone missed its primary endpoint in the double-blind portion of the 1042-0603 study. Due to this outcome Marinus discontinued this extension study.

Method

Denominator: ClinicalTrials.gov interventional Phase 2,3 neurology trials started 2015–2024 that have closed (completed or terminated). Numerator: terminated trials whose registry stop reason is classified as biological (efficacy, safety or benefit–risk) by Classification V2 — held-out precision 95.5%, recall 95.3% (n=600). Drugs are linked to ChEMBL and the NCI Thesaurus; 45% of industry neurology trials in this window carry a resolved drug target, and a trial without one cannot enter a mechanism class. Intervals are Wilson 95%.

Limits

Not a failure rate: completed trials that missed their endpoints are not in the rate. They are counted separately where the sponsor posted a primary result that can be read — most did not, so that count is a floor — and programmes discontinued after a completed trial do not appear. Stop reasons are sponsor-reported. This is a closed-trial proportion, not a time-to-event analysis: only 77.6% of trials in this segment have closed, and a trial that stops early enters the denominator sooner than one that runs to completion, which can inflate the rate in immature segments. Recent cohorts have fewer closed trials, so their rates are less stable. Research signals, not clinical or investment advice.

The whole cohort, trial by trial. The diligence report names every trial behind these figures — the ones still running and the ones that completed and missed their primary endpoint included — and places your molecule against each one that failed. Free during the beta: enter your work email at **clinicaltrialfailures.com/asset-check** and it opens straight away.

Sources: ClinicalTrials.gov (NLM); ChEMBL (EMBL-EBI, CC BY-SA 3.0); NCI Thesaurus (NCI); RxNorm/RxClass (NLM); SEC EDGAR. Classification, linkage and rates by Clinical Trial Failures. Rebuilt weekly; this brief covers trials started 2015–2024.