

CD20: 2 molecules stopped early, all monoclonal antibodies


6 stopped trials, but 2 molecules — all monoclonal antibodies: Obinutuzumab and Rituximab. Against 5.0% for all oncology trials with a resolved drug target and 4.3% across all oncology Phase 2,3 trials in the same window. The rate counts trials that stopped early for efficacy, safety or benefit–risk reasons. Trials that ran to completion and missed their primary endpoint are counted separately below, never in the rate.

3.6% 6 of 169 closed trials (95% CI 1.6%–7.5%)	4 / 1 / 0 efficacy / safety / both (adds to 6)	33.6% of 503 trials have closed · 334 still open or unresolved	0.7% stopped for a biological reason within 3 years of starting (95% CI 0.2%–2.2%; 272 still at risk; 1.5% area-wide)
--	--	--	---

0 of 3

Completed and missed the primary endpoint. Of 99 completed trials in this segment, 3 posted a primary superiority analysis on ClinicalTrials.gov; 0 came back non-significant and 3 significant. Counted separately and never in the rate above. The rest posted no analysis to read, so this is a floor, not a rate.

How this compares



What is in this class: 202 distinct experimental drugs across 503 trials (antibody 481, small molecule 384, unknown 52, adc 48). Most tested: RITUXIMAB (302), OBINUTUZUMAB (129), CYCLOPHOSPHAMIDE ANHYDROUS (120), PREDNISONE (90), VINCRISTINE (89). A class groups drugs by what they act on; check that the grouping is one you would make before reading the rate as a property of the mechanism.

How much does this rest on one decision? The 6 stops came from 6 sponsor-asset programmes across 4 sponsors; the largest (M.D. Anderson Cancer Center) contributed 1; removing that programme's trials from both sides leaves 3.0%. Screened alongside 85 other segments in this area, it does not survive a 10% false-discovery correction (q=1, Benjamini-Yekutieli). A further 13 closed trials here were terminated with no cause recorded in the registry; if every one of them were biological the rate would be 11.2%. Read it as a screen worth checking against the underlying trials.

Bars show the share of closed trials stopped for biological reasons. Confidence intervals overlap where sample sizes are small — read the counts, not just the bars.

By start cohort

Trials started	Stops	Closed	Rate	95% CI
2015–2017	4	88	4.5%	1.8%–11.1%
2018–2020	1	52	1.9%	0.3%–10.1%
2021–2024	1	29	3.4%	0.6%–17.2%

Where the stops sit

Efficacy / futility: 4, Benefit–risk: 1, Safety: 1.
Sponsors with most stops: Fondazione Italiana Linfomi - ETS (3), ADC Therapeutics S.A. (1), Johnson & Johnson (1), M.D. Anderson Cancer Center (1).

The stopped trials

Trial	Ph	Sponsor	Experimental drugs	Stopped	Type	Registry stop reason
NCT05144009	2	ADC Therapeutics S.A.	LONCASTUXIMAB TESIRINE, RITUXIMAB	2024-01	Benefit–risk	The benefit-risk profile does not support continuation of the LOTIS-9 trial.
NCT02703272	3	Johnson & Johnson	IBRUTINIB, RITUXIMAB, IFOSFAMIDE, CARBOPLATIN	2021-06	Efficacy	IDMC recommended that enrolment be stopped, as the EFS hazard ratio and associated p-value crossed the futility boundary specified in protocol (July 2020).

Trial	Ph	Sponsor	Experimental drugs	Stopped	Type	Registry stop reason
NCT03890289	2	Fondazione Italiana Linfomi - ETS	IDELALISIB, OBINUTUZUMAB	2021-04	Safety	Study was halted prematurely due to safety, since toxicity stopping rules have been met
NCT02420717	2	M.D. Anderson Cancer Center	CYCLOPHOSPHAMIDE ANHYDROUS, CYTARABINE, DASATINIB ANHYDROUS, DEXAMETHASONE	2021-01	Efficacy	Study was closed early due to low accrual and lack of response.
NCT02495454	2	Fondazione Italiana Linfomi - ETS	OBINUTUZUMAB	2017-03	Efficacy	Primary objective not obtained after interim analysis
NCT02670317	2	Fondazione Italiana Linfomi - ETS	OBINUTUZUMAB, IBRUTINIB, CYCLOPHOSPHAMIDE ANHYDROUS, DOXORUBICIN	2017-02	Efficacy	GA101-CHOP not advantage from rituximab-CHOP

Method

Denominator: ClinicalTrials.gov interventional Phase 2,3 oncology trials started 2015–2024 that have closed (completed or terminated).
Numerator: terminated trials whose registry stop reason is classified as biological (efficacy, safety or benefit–risk) by Classification V2 — held-out precision 95.5%, recall 95.3% (n=600). Drugs are linked to ChEMBL and the NCI Thesaurus; 76% of industry oncology trials in this window carry a resolved drug target, and a trial without one cannot enter a mechanism class. Intervals are Wilson 95%.

Limits

Not a failure rate: completed trials that missed their endpoints are not in the rate. They are counted separately where the sponsor posted a primary analysis that can be read — most did not, so that count is a floor — and programmes discontinued after a completed trial do not appear. Stop reasons are sponsor-reported. This is a closed-trial proportion, not a time-to-event analysis: only 33.6% of trials in this segment have closed, and a trial that stops early enters the denominator sooner than one that runs to completion, which can inflate the rate in immature segments. Recent cohorts have fewer closed trials, so their rates are less stable. Research signals, not clinical or investment advice.

The whole cohort, trial by trial. The evidence package names every trial behind these figures — the ones still running and the ones that completed and missed their primary endpoint included — and places your molecule against each one that failed. €99, delivered immediately: clinicaltrialfailures.com/asset-check

Sources: ClinicalTrials.gov (NLM); ChEMBL (EMBL-EBI, CC BY-SA 3.0); NCI Thesaurus (NCI); RxNorm/RxClass (NLM); SEC EDGAR. Classification, linkage and rates by Clinical Trial Failures. Rebuilt weekly; this brief covers trials started 2015–2024.