

ALK / ROS1 / RET: 6 molecules stopped early, all small molecules


7 stopped trials, but 6 molecules — all small molecules: Regorafenib, Alectinib, Ceritinib, EGFR/VEGFR/RET Inhibitor HA121-28 and 2 more. Against 5.3% for all oncology trials with a resolved drug target and 4.5% across all oncology Phase 2,3 trials in the same window. The rate counts trials that stopped early for efficacy, safety or benefit-risk reasons. Trials that ran to completion and missed their primary endpoint are counted separately below, never in the rate.

4.6%	4 / 0 / 0	37.2%	1.7%
7 of 153 closed trials (95% CI 2.2%–9.1%)	efficacy / safety / both (adds to 7)	of 411 trials have closed · 258 still open or unresolved	stopped for a biological reason within 3 years of starting (95% CI 0.7%–4.0%; 203 still at risk; 1.6% area-wide)

2 of 4

Completed and missed the primary endpoint. Of 100 completed trials in this segment, 4 posted a primary result on ClinicalTrials.gov that can be read — the sponsor's own comparison held to the sponsor's own threshold, or its statement that the endpoint was missed: 2 missed and 2 met. Counted separately and never in the rate above; a floor, not a rate. 46 of the 90 completed trials that finished more than a year ago have posted no results at all — a gap in what can be known, not a sign of failure.

How this compares



What is in this class: 239 distinct experimental drugs across 411 trials (small molecule 410, antibody 99, unknown 53, adc 13). Most tested: REGORAFENIB (108), SORAFENIB (75), ALECTINIB (37), SUNITINIB (35), CRIZOTINIB (34). A class groups drugs by what they act on; check that the grouping is one you would make before reading the rate as a property of the mechanism.

How much does this rest on one decision? The 7 stops came from 7 sponsor-asset programmes across 7 sponsors; the largest (UNICANCER) contributed 1; removing that programme's trials from both sides leaves 3.9%. Screened alongside 85 other segments, the rate is within what screening that many classes could produce (q=1); the molecules and trials that failed here are the record either way. A further 6 closed trials here were terminated with no cause recorded in the registry; if every one of them were biological the rate would be 8.5%. Read it as a screen worth checking against the underlying trials.

Bars show the share of closed trials stopped for biological reasons. Confidence intervals overlap where sample sizes are small — read the counts, not just the bars.

By start cohort

Trials started	Stops	Closed	Rate	95% CI
2015–2017	3	80	3.8%	1.3%–10.5%
2018–2020	1	43	2.3%	0.4%–12.1%
2021–2024	3	30	10.0%	3.5%–25.6%

Where the stops sit

Efficacy / futility: 4, Benefit-risk: 3.
Sponsors with most stops: Great Novel Therapeutics Biotech & Medicals Corporation (1), Bristol Myers Squibb (1), CSPC ZhongQi Pharmaceutical Technology Co., Ltd. (1), UNICANCER (1), Sun Yat-sen University (1), Dana-Farber Cancer Institute (1).

The stopped trials

Trial	Ph	Sponsor	Experimental drugs	Stopped	Type	Registry stop reason
NCT05770882	2	Great Novel Therapeutics Biotech & Medicals Corporation	TUCIDINOSTAT, REGORAFENIB	2025-01	Benefit–risk	The overall profile does not support development for Hepatocellular Carcinoma
NCT05071183	2	Bristol Myers Squibb 	REPOTRECTINIB, TRAMETINIB	2023-03	Benefit–risk	Inability to achieve and optimize a dose that could provide a positive benefit risk profile
NCT04784520	2	CSPC ZhongQi Pharmaceutical Technology Co., Ltd.	EGFR/VEGFR/RET Inhibitor HA121-28	2022-12	Efficacy	The primary endpoint did not meet expectation.
NCT03722108	2	UNICANCER	REGORAFENIB, IRINOTECAN	2022-05	Benefit–risk	Unfavorable benefit-risk balance in the experimental arm following the IDMC for intermediate efficacy and safety analysis
NCT02867280	3	Sun Yat-sen University	SORAFENIB	2019-08	Efficacy	Midterm analysis showed negative results.
NCT03131206	2	Dana-Farber Cancer Institute	ALECTINIB	2018-11	Efficacy	Study closed due to slow accrual and lack of efficacy.

The 6 most recent of 7 stopped trials. Every molecule behind all 7 is named above; the complete trial list, the trials still running and the terminations with no readable cause ship with the dataset and the diligence report.

Method

Denominator: ClinicalTrials.gov interventional Phase 2,3 oncology trials started 2015–2024 that have closed (completed or terminated).
Numerator: terminated trials whose registry stop reason is classified as biological (efficacy, safety or benefit–risk) by Classification V2 — held-out precision 95.5%, recall 95.3% (n=600). Drugs are linked to ChEMBL and the NCI Thesaurus; 76% of industry oncology trials in this window carry a resolved drug target, and a trial without one cannot enter a mechanism class. Intervals are Wilson 95%.

Limits

Not a failure rate: completed trials that missed their endpoints are not in the rate. They are counted separately where the sponsor posted a primary result that can be read — most did not, so that count is a floor — and programmes discontinued after a completed trial do not appear. Stop reasons are sponsor-reported. This is a closed-trial proportion, not a time-to-event analysis: only 37.2% of trials in this segment have closed, and a trial that stops early enters the denominator sooner than one that runs to completion, which can inflate the rate in immature segments. Recent cohorts have fewer closed trials, so their rates are less stable. Research signals, not clinical or investment advice.

The whole cohort, trial by trial. The diligence report names every trial behind these figures — the ones still running and the ones that completed and missed their primary endpoint included — and places your molecule against each one that failed. €99, delivered immediately: clinicaltrialfailures.com/asset-check

Sources: ClinicalTrials.gov (NLM); ChEMBL (EMBL-EBI, CC BY-SA 3.0); NCI Thesaurus (NCI); RxNorm/RxClass (NLM); SEC EDGAR. Classification, linkage and rates by Clinical Trial Failures. Rebuilt weekly; this brief covers trials started 2015–2024.