

CD3 T-cell engagers: 4 molecules stopped early


4 stopped trials across 4 molecules: Anti-CD3/CD7-Ricin Toxin A Immunotoxin, B7H3/CD3-targeting T-cell Engaging Molecule TAK-280, Tak-186 and Vadacabtagene Leraleucel. Against 5.3% for all oncology trials with a resolved drug target and 4.5% across all oncology Phase 2,3 trials in the same window. The rate counts trials that stopped early for efficacy, safety or benefit–risk reasons. Trials that ran to completion and missed their primary endpoint are counted separately below, never in the rate.

6.3%	2 / 2 / 0	22.3%	0.9%
4 of 63 closed trials (95% CI 2.5%–15.2%)	efficacy / safety / both (adds to 4)	of 283 trials have closed · 220 still open or unresolved	stopped for a biological reason within 3 years of starting (95% CI 0.2%–3.4%; 130 still at risk; 1.6% area-wide)

1 of 2

Completed and missed the primary endpoint. Of 31 completed trials in this segment, 2 posted a primary result on ClinicalTrials.gov that can be read — the sponsor's own comparison held to the sponsor's own threshold, or its statement that the endpoint was missed: 1 missed and 1 met. Counted separately and never in the rate above; a floor, not a rate. 14 of the 27 completed trials that finished more than a year ago have posted no results at all — a gap in what can be known, not a sign of failure.

How this compares



What is in this class: 173 distinct experimental drugs across 283 trials (antibody 240, small molecule 94, adc 28, bispecific antibody 28). Most tested: BLINATUMOMAB (70), MOSUNETUZUMAB (32), DEXAMETHASONE (30), RITUXIMAB (28), GLOFITAMAB (28). A class groups drugs by what they act on; check that the grouping is one you would make before reading the rate as a property of the mechanism.

How much does this rest on one decision? The 4 stops came from 4 sponsor-asset programmes across 3 sponsors; the largest (Xenikos) contributed 1; removing that programme's trials from both sides leaves 4.8%. Screened alongside 85 other segments, the rate is within what screening that many classes could produce (q=1); the molecules and trials that failed here are the record either way. A further 7 closed trials here were terminated with no cause recorded in the registry; if every one of them were biological the rate would be 17.5%. Read it as a screen worth checking against the underlying trials.

Bars show the share of closed trials stopped for biological reasons. Confidence intervals overlap where sample sizes are small — read the counts, not just the bars.

By start cohort

Trials started	Stops	Closed	Rate	95% CI
2015–2017	1	16	6.2%	1.1%–28.3%
2018–2020	0	18	0.0%	0.0%–17.6%
2021–2024	3	29	10.3%	3.6%–26.4%

Where the stops sit

Efficacy / futility: 2, Safety: 2.
Sponsors with most stops: Takeda (2), Xenikos (1), Bristol Myers Squibb (1).

The stopped trials

Trial	Ph	Sponsor	Experimental drugs	Stopped	Type	Registry stop reason
NCT05220098	2	Takeda TAK	B7H3/CD3-targeting T-cell Engagin...	2025-07	Efficacy	Terminated due to limited anti-cancer activity
NCT04844073	2	Takeda TAK	TAK-186	2025-06	Efficacy	Sponsor decision based on limited anti-cancer activity of TAK-186.

Trial	Ph	Sponsor	Experimental drugs	Stopped	Type	Registry stop reason
NCT04934670	3	Xenikos	Anti-CD3/CD7-Ricin Toxin A Immuno...	2022-11	Safety	The study met the protocol defined stopping boundary for Day 60 mortality when comparing mortality between the T-Guard and ruxolitinib arms
NCT02535364	2	Bristol Myers Squibb BMY	VADACABTAGENE LERALEUCEL	2017-04	Safety	Safety reasons

Method

Denominator: ClinicalTrials.gov interventional Phase 2,3 oncology trials started 2015–2024 that have closed (completed or terminated).
Numerator: terminated trials whose registry stop reason is classified as biological (efficacy, safety or benefit–risk) by Classification V2 — held-out precision 95.5%, recall 95.3% (n=600). Drugs are linked to ChEMBL and the NCI Thesaurus; 76% of industry oncology trials in this window carry a resolved drug target, and a trial without one cannot enter a mechanism class. Intervals are Wilson 95%.

Limits

Not a failure rate: completed trials that missed their endpoints are not in the rate. They are counted separately where the sponsor posted a primary result that can be read — most did not, so that count is a floor — and programmes discontinued after a completed trial do not appear. Stop reasons are sponsor-reported. This is a closed-trial proportion, not a time-to-event analysis: only 22.3% of trials in this segment have closed, and a trial that stops early enters the denominator sooner than one that runs to completion, which can inflate the rate in immature segments. Recent cohorts have fewer closed trials, so their rates are less stable. Research signals, not clinical or investment advice.

The whole cohort, trial by trial. The diligence report names every trial behind these figures — the ones still running and the ones that completed and missed their primary endpoint included — and places your molecule against each one that failed. €99, delivered immediately: clinicaltrialfailures.com/asset-check

Sources: ClinicalTrials.gov (NLM); ChEMBL (EMBL-EBI, CC BY-SA 3.0); NCI Thesaurus (NCI); RxNorm/RxClass (NLM); SEC EDGAR. Classification, linkage and rates by Clinical Trial Failures. Rebuilt weekly; this brief covers trials started 2015–2024.