

Alarmins (TSLP / IL-33): 4 of 7 completed trials missed their primary endpoint

4 completed trials across 4 sponsors and 4 molecules ran to the end and missed their primary endpoint on the sponsor's own posted result; 2 met it. Only 0 of 24 closed trials were stopped early for efficacy or safety, which is why this class is read through its completed trials rather than a stop rate. Most completed trials post nothing readable, so the count is a floor.

4 of 7

completed trials with a readable primary result missed it · 2 met it

4 / 4

sponsors / molecules behind the misses

0.0%

stopped early: 0 of 24 closed trials (95% CI 0.0%–13.8%)

77.4%

of 31 trials have closed · 7 still open or unresolved


4 of 7

Completed and missed the primary endpoint. Of 23 completed trials in this segment, 7 posted a primary result on ClinicalTrials.gov that can be read — the sponsor's own comparison held to the sponsor's own threshold, or its statement that the endpoint was missed: 4 missed and 2 met, 1 split across co-primary endpoints. Counted separately and never in the rate above; a floor, not a rate. 4 of the 20 completed trials that finished more than a year ago have posted no results at all — a gap in what can be known, not a sign of failure.

The completed trials that missed

Trial	Ph	Sponsor	Experimental drugs	Started	What the posted result says
NCT04386616	2	Roche	ASTEGOLIMAB, EFMARODOCOKIN ALFA	2020-06	Time to Recovery, Defined as the Time to a Clinical Status Score of 1 or 2 on the 7-Category Ordinal Scale (Whichever Occurs First) by Day 28 — p 0.9266 against 0.05: not significant. Hazard Ratio (HR) 1.01 (95% CI 0.75 to 1.36). Method: Unstratified Log Rank.
NCT03615040	2	University of Leicester	ASTEGOLIMAB	2018-10	Number of Moderate to Severe Exacerbation (Defined as Requiring Treatment With Systemic Corticosteroids and/or Antibiotics in the Community or Hospital or Hospitalisation). — p 0.195 against 0.05: not significant. Incidence Rate Ratio 0.78 (95% CI 0.53 to 1.14). Method: t-test, 2 sided.
NCT03546907	2	Sanofi	ITEPEKIMAB	2018-07	Annualized Rate of Moderate to Severe Acute Exacerbation Events in Chronic Obstructive Pulmonary Disease (AECOPD) Participants — p 0.1296 against 0.05: not significant. Risk Ratio (RR) 0.808 (95% CI 0.613 to 1.065). Method: Negative binomial regression model.
NCT03406078	3	AstraZeneca	TEZEPELUMAB	2018-03	Categorized Percent Reduction From Baseline in the Daily OCS Dose While Not Losing Asthma Control — p 0.434 against 0.05: not significant. Cumulative Odds Ratio 1.28 (95% CI 0.69 to 2.35). Method: Proportional odds model.

How this compares



What is in this class: 9 distinct experimental drugs across 31 trials (antibody 31, protein 1, small molecule 1). Most tested: TEZEPELUMAB (17), ITEPEKIMAB (7), ASTEGOLIMAB (6), ETOKIMAB (1), EFMARODOCOKIN ALFA (1). A class groups drugs by what they act on; check that the grouping is one you would make before reading the rate as a property of the mechanism.

How much does this rest on one decision? The 0 stops came from 0 sponsor-asset programmes across 0 sponsors. Screened alongside 13 other segments, the rate is within what screening that many classes could produce (q=1); the molecules and trials that failed here are the record either way. Read it as a screen worth checking against the underlying trials.

Bars show the share of closed trials stopped for biological reasons. Confidence intervals overlap where sample sizes are small — read the counts, not just the bars.

Method

Denominator: ClinicalTrials.gov interventional Phase 2,3 respiratory trials started 2015–2024 that have closed (completed or terminated).
Numerator: terminated trials whose registry stop reason is classified as biological (efficacy, safety or benefit–risk) by Classification V2 — held-out precision 95.5%, recall 95.3% (n=600). Drugs are linked to ChEMBL and the NCI Thesaurus; 42% of industry respiratory trials in this window carry a resolved drug target, and a trial without one cannot enter a mechanism class. Intervals are Wilson 95%.

By start cohort

Trials started	Stops	Closed	Rate	95% CI
2018–2020	0	11	0.0%	0.0%–25.9%
2021–2024	0	9	0.0%	0.0%–29.9%

Where the stops sit

Sponsors with most stops: .

Limits

Not a failure rate: completed trials that missed their endpoints are not in the rate. They are counted separately where the sponsor posted a primary result that can be read — most did not, so that count is a floor — and programmes discontinued after a completed trial do not appear. Stop reasons are sponsor-reported. This is a closed-trial proportion, not a time-to-event analysis: only 77.4% of trials in this segment have closed, and a trial that stops early enters the denominator sooner than one that runs to completion, which can inflate the rate in immature segments. Recent cohorts have fewer closed trials, so their rates are less stable. Research signals, not clinical or investment advice.

The whole cohort, trial by trial. The diligence report names every trial behind these figures — the ones still running and the ones that completed and missed their primary endpoint included — and places your molecule against each one that failed. €99, delivered immediately: clinicaltrialfailures.com/asset-check

Sources: ClinicalTrials.gov (NLM); ChEMBL (EMBL-EBI, CC BY-SA 3.0); NCI Thesaurus (NCI); RxNorm/RxClass (NLM); SEC EDGAR. Classification, linkage and rates by Clinical Trial Failures. Rebuilt weekly; this brief covers trials started 2015–2024.