


MET: 8 molecules stopped early

11 stopped trials across 8 molecules: Cabozantinib, Capmatinib, Amg-337, Amivantamab and 4 more. Against 5.0% for all oncology trials with a resolved drug target and 4.3% across all oncology Phase 2,3 trials in the same window. Rates count trials that stopped early for efficacy, safety or benefit–risk reasons; trials that completed and missed their endpoints are not counted.

9.1% 11 of 121 closed trials (95% CI 5.2%–15.5%)	6 / 4 / 0 efficacy / safety / both (adds to 11)	37.9% of 319 trials have closed · 198 still open or unresolved	2.7% stopped for a biological reason within 3 years of starting (95% CI 1.3%–5.6%; 186 still at risk; 1.5% area-wide)
--	---	--	---

How this compares



What is in this class: 213 distinct experimental drugs across 319 trials (small molecule 296, antibody 135, unknown 22, adc 12). Most tested: CABOZANTINIB (134), NIVOLUMAB (40), CRIZOTINIB (34), CAPMATINIB (29), AMIVANTAMAB (28). A class groups drugs by what they act on; check that the grouping is one you would make before reading the rate as a property of the mechanism.

How much does this rest on one decision? The 11 stops came from 11 sponsor-asset programmes across 10 sponsors; the largest (University of Nebraska) contributed 1; removing that programme's trials from both sides leaves 8.3%. Screened alongside 85 other segments in this area, it does not survive a 10% false-discovery correction ($q=1$, Benjamini-Yekutieli). A further 5 closed trials here were terminated with no cause recorded in the registry; if every one of them were biological the rate would be 13.2%. Read it as a screen worth checking against the underlying trials.

Bars show the share of closed trials stopped for biological reasons. Confidence intervals overlap where sample sizes are small — read the counts, not just the bars.

By start cohort

Trials started	Stops	Closed	Rate	95% CI
2015–2017	2	42	4.8%	1.3%–15.8%
2018–2020	7	46	15.2%	7.6%–28.2%
2021–2024	2	33	6.1%	1.7%–19.6%

Where the stops sit

Efficacy / futility: 6, Safety: 4, Benefit–risk: 1.
Sponsors with most stops: Novartis (2), Genmab (1), Johnson & Johnson (1), Jiangsu Alphamab Biopharmaceuticals Co., Ltd (1), NantPharma, LLC (1), University of Nebraska (1).

The stopped trials

Trial	Ph	Sponsor	Experimental drugs	Stopped	Type	Registry stop reason
NCT06685068	2	Genmab GMAB	Anti-EGFR/c-Met Antibody-drug Con...	2026-03	Benefit–risk	Genmab has decided to terminate the GEN1286 program due to an unfavorable benefit-risk profile not supporting further development.
NCT05653427	2	Johnson & Johnson JNJ	AMIVANTAMAB	2023-10	Efficacy	N1–15 subjects data analysis did not meet statistical consideration of ORR and study primary/secondary objectives were not met. Enrollment was terminated for futility
NCT04323436	2	Novartis NVS	SPARTALIZUMAB, CAPMATINIB	2022-12	Safety	Study termination based on sponsor decision due to lack of tolerability observed with capmatinib and spartalizumab combination treatment when compared to data from capmatinib single agent studies
NCT03838848	2	Jiangsu Alphamab Biopharmaceuticals Co., Ltd	ERFONRILIMAB, CARBOPLATIN, PEMETREXED, NINGETINIB	2022-07	Efficacy	Cohort A,B,C end enrollments. Cohort D and E were considered to have no significant clinical benefit at the SMC meeting and decided to terminate enrollment.

Trial	Ph	Sponsor	Experimental drugs	Stopped	Type	Registry stop reason
NCT04139317	2	Novartis NVS	CAPMATINIB	2022-01	Safety	Sponsor decision to terminate due to the lack of tolerability observed in patients treated with capmatinib and pembrolizumab in the combination arm as compared to patients treated with pembrolizumab alone in the pembrolizumab single agent arm
NCT03132155	2	NantPharma, LLC	AMG-337	2021-08	Efficacy	Prematurely terminated due to lack of therapeutic effect

The 6 most recent of 11 stopped trials. Every molecule behind all 11 is named above; the complete trial list, the trials still running and the terminations with no readable cause ship with the dataset and the evidence package.

Method

Denominator: ClinicalTrials.gov interventional Phase 2,3 oncology trials started 2015–2024 that have closed (completed or terminated).
Numerator: terminated trials whose registry stop reason is classified as biological (efficacy, safety or benefit–risk) by Classification V2 — held-out precision 95.5%, recall 95.3% (n=600). Drugs are linked to ChEMBL and the NCI Thesaurus; 76% of industry oncology trials in this window carry a resolved drug target, and a trial without one cannot enter a mechanism class. Intervals are Wilson 95%.

Limits

Not a failure rate: trials that completed with negative results are not counted, and programs discontinued after a completed trial do not appear. Stop reasons are sponsor-reported. This is a closed-trial proportion, not a time-to-event analysis: only 37.9% of trials in this segment have closed, and a trial that stops early enters the denominator sooner than one that runs to completion, which can inflate the rate in immature segments. Recent cohorts have fewer closed trials, so their rates are less stable. Research signals, not clinical or investment advice.

Any mechanism, sponsor or indication, updated weekly. The dataset behind this brief covers every stopped oncology trial with an efficacy or safety signal plus the full denominator universe. Free sample and licensing: clinicaltrialfailures.com/pricing

Sources: ClinicalTrials.gov (NLM); ChEMBL (EMBL-EBI, CC BY-SA 3.0); NCI Thesaurus (NCI); RxNorm/RxCClass (NLM); SEC EDGAR. Classification, linkage and rates by Clinical Trial Failures. Rebuilt weekly; this brief covers trials started 2015–2024.