CLINICAL TRIAL FAILURES · DISCONTINUATION RATE · ONCOLOGY PHASE 2,3 · STARTS 2015–2024

# CD20: 2 molecules stopped early, all monoclonal antibodies

**6 stopped trials, but 2 molecules — all monoclonal antibodies: Obinutuzumab and Rituximab.** Against 5.2% for all oncology trials with a resolved drug target and 4.5% across all oncology Phase 2,3 trials in the same window. The rate counts trials that stopped early for efficacy, safety or benefit–risk reasons. Trials that ran to completion and missed their primary endpoint are counted separately below, never in the rate.

**3.5%**
6 of 171 closed trials (95% CI 1.6%–7.4%)


**4 / 1 / 0**
efficacy / safety / both (adds to 6)

**33.9%**
of 504 trials have closed · 333 still open or unresolved

**0.7%**
stopped for a biological reason within 3 years of starting (95% CI 0.2%–2.2%; 275 still at risk; 1.5% area-wide)

**1 of 3** **Completed and missed the primary endpoint.** Of 100 completed trials in this segment, 3 posted a primary result on ClinicalTrials.gov that can be read — the sponsor's own comparison held to the sponsor's own threshold, or its statement that the endpoint was missed: 1 missed and 2 met. Counted separately and never in the rate above; a floor, not a rate. 43 of the 96 completed trials that finished more than a year ago have posted no results at all — a gap in what can be known, not a sign of failure.

## How this compares




**What is in this class:** 202 distinct experimental drugs across 504 trials (antibody 482, small molecule 385, unknown 52, adc 49). Most tested: RITUXIMAB (303), OBINUTUZUMAB (129), CYCLOPHOSPHAMIDE ANHYDROUS (120), PREDNISONE (91), VINCRISTINE (88). A class groups drugs by what they act on; check that the grouping is one you would make before reading the rate as a property of the mechanism.

**How much does this rest on one decision?** The 6 stops came from 6 sponsor-asset programmes across 4 sponsors; the largest (M.D. Anderson Cancer Center) contributed 1; removing that programme's trials from both sides leaves 2.9%. Screened alongside 85 other segments, the rate is within what screening that many classes could produce (q=1); the molecules and trials that failed here are the record either way. A further 14 closed trials here were terminated with no cause recorded in the registry; if every one of them were biological the rate would be 11.7%. Read it as a screen worth checking against the underlying trials.

Bars show the share of closed trials stopped for biological reasons. Confidence intervals overlap where sample sizes are small — read the counts, not just the bars.

## By start cohort

| Trials started | Stops | Closed | Rate | 95% CI |
|---|---|---|---|---|
| 2015–2017 | 4 | 88 | 4.5% | 1.8%–11.1% |
| 2018–2020 | 1 | 53 | 1.9% | 0.3%–9.9% |
| 2021–2024 | 1 | 30 | 3.3% | 0.6%–16.7% |

## Where the stops sit

Efficacy / futility: 4, Benefit–risk: 1, Safety: 1.
Sponsors with most stops: Fondazione Italiana Linfomi - ETS (3), ADC Therapeutics S.A. (1), Johnson & Johnson (1), M.D. Anderson Cancer Center (1).

## The stopped trials

| Trial | Ph | Sponsor | Experimental drugs | Stopped | Type | Registry stop reason |
|---|---|---|---|---|---|---|
| NCT05144009 | 2 | ADC Therapeutics S.A. | LONCASTUXIMAB TESIRINE, RITUXIMAB | 2024-01 | Benefit–risk | The benefit-risk profile does not support continuation of the LOTIS-9 trial. |

| Trial | Ph | Sponsor | Experimental drugs | Stopped | Type | Registry stop reason |
|---|---|---|---|---|---|---|
| NCT02703272 | 3 | Johnson & Johnson JNJ | IBRUTINIB, RITUXIMAB, IFOSFAMIDE, CARBOPLATIN | 2021-06 | Efficacy | IDMC recommended that enrolment be stoped, as the EFS hazard ratio and associated p-value crossed the futility boundary specified in protocol (July 2020). |
| NCT03890289 | 2 | Fondazione Italiana Linfomi - ETS | IDELALISIB, OBINUTUZUMAB | 2021-04 | Safety | Study was halted prematurely due to safety, since tocity stopping rules have been met |
| NCT02420717 | 2 | M.D. Anderson Cancer Center | CYCLOPHOSPHAMIDE ANHYDROUS, CYTARABINE, DASATINIB ANHYDROUS, DEXAMETHASONE | 2021-01 | Efficacy | Study was closed early due to low accrual and lack of response. |
| NCT02495454 | 2 | Fondazione Italiana Linfomi - ETS | OBINUTUZUMAB | 2017-03 | Efficacy | Primary objective not obtained after interim analysis |
| NCT02670317 | 2 | Fondazione Italiana Linfomi - ETS | OBINUTUZUMAB, IBRUTINIB, CYCLOPHOSPHAMIDE ANHYDROUS, DOXORUBICIN | 2017-02 | Efficacy | GA101-CHOP not advantage from rituximab-CHOP |

**Method**
Denominator: ClinicalTrials.gov interventional Phase 2,3 oncology trials started 2015–2024 that have closed (completed or terminated). Numerator: terminated trials whose registry stop reason is classified as biological (efficacy, safety or benefit–risk) by Classification V2 — held-out precision 95.5%, recall 95.3% (n=600). Drugs are linked to ChEMBL and the NCI Thesaurus; 76% of industry oncology trials in this window carry a resolved drug target, and a trial without one cannot enter a mechanism class. Intervals are Wilson 95%.

**Limits**
Not a failure rate: completed trials that missed their endpoints are not in the rate. They are counted separately where the sponsor posted a primary result that can be read — most did not, so that count is a floor — and programmes discontinued after a completed trial do not appear. Stop reasons are sponsor-reported. This is a closed-trial proportion, not a time-to-event analysis: only 33.9% of trials in this segment have closed, and a trial that stops early enters the denominator sooner than one that runs to completion, which can inflate the rate in immature segments. Recent cohorts have fewer closed trials, so their rates are less stable. Research signals, not clinical or investment advice.

**The whole cohort, trial by trial.** The diligence report names every trial behind these figures — the ones still running and the ones that completed and missed their primary endpoint included — and places your molecule against each one that failed. €99, delivered immediately: **clinicaltrialfailures.com/asset-check**

Sources: ClinicalTrials.gov (NLM); ChEMBL (EMBL-EBI, CC BY-SA 3.0); NCI Thesaurus (NCI); RxNorm/RxClass (NLM); SEC EDGAR. Classification, linkage and rates by Clinical Trial Failures. Rebuilt weekly; this brief covers trials started 2015–2024.