CLINICAL TRIAL FAILURES · DISCONTINUATION RATE · ONCOLOGY PHASE 2,3 · STARTS 2015–2024

CD38 + PD-(L)1: 2 molecules stopped early

4 stopped trials across 2 molecules: Isatuximab and Modakafusp Alfa. Against 5.1% for PD-(L)1 combinations without CD38 and 4.5% across all oncology Phase 2,3 trials in the same window. The rate counts trials that stopped early for efficacy, safety or benefit–risk reasons. Trials that ran to completion and missed their primary endpoint are counted separately below, never in the rate.

33.3%
4 of 12 closed trials (95% CI 13.8%–60.9%)

4 / 0 / 0
efficacy / safety / both (adds to 4)

70.6%
of 17 trials have closed · 5 still open or unresolved

0.0%
stopped for a biological reason within 3 years of starting (95% CI 0.0%–0.0%; 13 still at risk; 1.5% area-wide)

1 of 1 **Completed and missed the primary endpoint.** Of 6 completed trials in this segment, 1 posted a primary result on ClinicalTrials.gov that can be read — the sponsor's own comparison held to the sponsor's own threshold, or its statement that the endpoint was missed: 1 missed and 0 met. Counted separately and never in the rate above; a floor, not a rate. 1 of the 6 completed trials that finished more than a year ago have posted no results at all — a gap in what can be known, not a sign of failure.

How this compares

CD38 + PD-(L)1
33.3%
4/12 closed

PD-(L)1 combinations without CD38
5.1%
112/2196 closed

All oncology Phase 2,3
4.5%
355/7935 closed

What is in this class: 36 distinct experimental drugs across 17 trials (antibody 17, small molecule 5, protein 2, unknown 2). Most tested: DARATUMUMAB (9), ISATUXIMAB (7), NIVOLUMAB (4), CEMIPLIMAB (4), POMALIDOMIDE (3). A class groups drugs by what they act on; check that the grouping is one you would make before reading the rate as a property of the mechanism.

How much does this rest on one decision? The 4 stops came from 3 sponsor-asset programmes across 2 sponsors; the largest (Sanofi) contributed 2; removing that programme's trials from both sides leaves 20.0%. Screened alongside 85 other segments, the rate is within what screening that many classes could produce (q=0.371); the molecules and trials that failed here are the record either way. Read it as a screen worth checking against the underlying trials.

Bars show the share of closed trials stopped for biological reasons. Confidence intervals overlap where sample sizes are small — read the counts, not just the bars.

By start cohort

Trials started	Stops	Closed	Rate	95% CI
2015–2017	0	6	0.0%	0.0%–39.0%
2018–2020	4	5	80.0%	37.6%–96.4%

Where the stops sit

Efficacy / futility: 4.
Sponsors with most stops: Sanofi (3), Teva Branded Pharmaceutical Products R&D LLC (1).

The stopped trials

Trial	Ph	Sponsor	Experimental drugs	Stopped	Type	Registry stop reason
NCT04157517	2	Teva Branded Pharmaceutical Products R&D LLC	MODAKAFUSP ALFA, PEMBROLIZUMAB	2023-12	Efficacy	Study was terminated early due to futility.
NCT03769181	2	Sanofi SNY	ISATUXIMAB, CEMIPLIMAB	2022-11	Efficacy	Study was stopped after interim analysis for all 4 cohorts with results either not fulfilling the pre-planned interim analysis criteria or fulfilling the criteria but as per sponsor decision. It was not due to any safety concern

Trial	Ph	Sponsor	Experimental drugs	Stopped	Type	Registry stop reason
NCT03637764	2	Sanofi SNY	ISATUXIMAB, ATEZOLIZUMAB	2022-05	Efficacy	Study was stopped after interim analysis for 4 cohorts (HCC, SCCHN, EOC, and GBM) indicated that the results did not fulfill the preplanned interim analysis criteria to move to Phase 2 stage 2 in all 4 cohorts.
NCT03367819	2	Sanofi SNY	ISATUXIMAB, CEMIPLIMAB	2021-03	Efficacy	The study was stopped after interim analysis indicated that overall result was not sufficient to satisfy per-protocol criteria to move forward in metastatic, castration-resistant prostate cancer (mCRPC) and non-small cell lung cancer (NSCLC) cohorts.

Method

Denominator: ClinicalTrials.gov interventional Phase 2,3 oncology trials started 2015–2024 that have closed (completed or terminated). Numerator: terminated trials whose registry stop reason is classified as biological (efficacy, safety or benefit–risk) by Classification V2 — held-out precision 95.5%, recall 95.3% (n=600). Drugs are linked to ChEMBL and the NCI Thesaurus; 76% of industry oncology trials in this window carry a resolved drug target, and a trial without one cannot enter a mechanism class. Intervals are Wilson 95%.

Limits

Not a failure rate: completed trials that missed their endpoints are not in the rate. They are counted separately where the sponsor posted a primary result that can be read — most did not, so that count is a floor — and programmes discontinued after a completed trial do not appear. Stop reasons are sponsor-reported. This is a closed-trial proportion, not a time-to-event analysis: only 70.6% of trials in this segment have closed, and a trial that stops early enters the denominator sooner than one that runs to completion, which can inflate the rate in immature segments. Recent cohorts have fewer closed trials, so their rates are less stable. Research signals, not clinical or investment advice.

The whole cohort, trial by trial. The diligence report names every trial behind these figures — the ones still running and the ones that completed and missed their primary endpoint included — and places your molecule against each one that failed. Free during the beta, in exchange for a work email, delivered immediately: **clinicaltrialfailures.com/asset-check**

Sources: ClinicalTrials.gov (NLM); ChEMBL (EMBL-EBI, CC BY-SA 3.0); NCI Thesaurus (NCI); RxNorm/RxClass (NLM); SEC EDGAR. Classification, linkage and rates by Clinical Trial Failures. Rebuilt weekly; this brief covers trials started 2015–2024.