

# Data Structures and Algorithms <sup>1</sup>

BITS-Pilani K. K. Birla Goa Campus

---

<sup>1</sup>Material for the presentation taken from Cormen, Leiserson, Rivest and Stein, *Introduction to Algorithms, Third Edition*;

## Ch 4: Divide-and-Conquer

**Divide** the problem into a number of subproblems that are smaller instances of the same problem.

**Conquer** the subproblems by solving them recursively.

**Combine** the solutions to the subproblems into the solution for the original problem.

# Minimum Element

- Q. You are given an unsorted array. Find the minimum element in the array using a divide-and-conquer approach.

# Minimum Element

Q. You are given an unsorted array. Find the minimum element in the array using a divide-and-conquer approach.

```
1: procedure FINDMINIMUM(arr, left, right)  
2:   if left = right then  
3:     return arr[left]  
4:   mid  $\leftarrow \lfloor (left + right)/2 \rfloor$ 
```

# Minimum Element

Q. You are given an unsorted array. Find the minimum element in the array using a divide-and-conquer approach.

```
1: procedure FINDMINIMUM(arr, left, right)  
2:   if left = right then  
3:     return arr[left]  
4:   mid  $\leftarrow \lfloor (\textit{left} + \textit{right}) / 2 \rfloor$   
5:   minLeft  $\leftarrow$  FINDMINIMUM(arr, left, mid)  
6:   minRight  $\leftarrow$  FINDMINIMUM(arr, mid + 1, right)
```

# Minimum Element

Q. You are given an unsorted array. Find the minimum element in the array using a divide-and-conquer approach.

```
1: procedure FINDMINIMUM(arr, left, right)
2:   if left = right then
3:     return arr[left]
4:   mid  $\leftarrow \lfloor (\textit{left} + \textit{right}) / 2 \rfloor$ 
5:   minLeft  $\leftarrow$  FINDMINIMUM(arr, left, mid)
6:   minRight  $\leftarrow$  FINDMINIMUM(arr, mid + 1, right)
7:   return min(minLeft, minRight)
```

# Minimum Element

Q. You are given an unsorted array. Find the minimum element in the array using a divide-and-conquer approach.

```
1: procedure FINDMINIMUM(arr, left, right)
2:   if left = right then
3:     return arr[left]
4:   mid  $\leftarrow \lfloor (left + right)/2 \rfloor$ 
5:   minLeft  $\leftarrow$  FINDMINIMUM(arr, left, mid)
6:   minRight  $\leftarrow$  FINDMINIMUM(arr, mid + 1, right)
7:   return min(minLeft, minRight)
```

(a) Find the recurrence equation for the above algorithm?

# Minimum Element

Q. You are given an unsorted array. Find the minimum element in the array using a divide-and-conquer approach.

```
1: procedure FINDMINIMUM(arr, left, right)
2:   if left = right then
3:     return arr[left]
4:   mid  $\leftarrow \lfloor (left + right)/2 \rfloor$ 
5:   minLeft  $\leftarrow$  FINDMINIMUM(arr, left, mid)
6:   minRight  $\leftarrow$  FINDMINIMUM(arr, mid + 1, right)
7:   return min(minLeft, minRight)
```

- (a) Find the recurrence equation for the above algorithm?
- (b) What is the worst case running time of the algorithm?



# Matrix multiplication

- ▶  $C = A \cdot B$ , where  $A$ ,  $B$  and  $C$  are  $n \times n$  matrices

# Matrix multiplication

- ▶  $C = A \cdot B$ , where  $A$ ,  $B$  and  $C$  are  $n \times n$  matrices
- ▶  $c_{ij} = \sum_{k=1}^n a_{ik} \cdot b_{kj}$

# Matrix multiplication

- ▶  $C = A \cdot B$ , where  $A$ ,  $B$  and  $C$  are  $n \times n$  matrices
- ▶ 
$$c_{ij} = \sum_{k=1}^n a_{ik} \cdot b_{kj}$$

**MATRIX-MULTIPLY( $A, B, C, n$ )**

```
1  for  $i = 1$  to  $n$ 
2      for  $j = 1$  to  $n$ 
3          for  $k = 1$  to  $n$ 
4               $c_{ij} = c_{ij} + a_{ik} \cdot b_{kj}$ 
```

# Matrix multiplication

- ▶ MATRIX-MULTIPLY procedure runs in  $\Theta(n^3)$  time.

# Matrix multiplication

- ▶ MATRIX-MULTIPLY procedure runs in  $\Theta(n^3)$  time.
- ▶ Volker Strassen came up with an algorithm that takes  $O(n^{\lg 7})$  time, which is  $o(n^{2.81})$ . ( $\lg 7 = 2.8074$ )

# Matrix multiplication

- ▶ MATRIX-MULTIPLY procedure runs in  $\Theta(n^3)$  time.
- ▶ Volker Strassen came up with an algorithm that takes  $O(n^{\lg 7})$  time, which is  $o(n^{2.81})$ . ( $\lg 7 = 2.8074$ )
- ▶ Strassen's algorithm uses a divide-and-conquer approach.

# Matrix multiplication: divide-and-conquer

$$A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}, \quad B = \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}, \quad C = \begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix},$$

# Matrix multiplication: divide-and-conquer

$$A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}, \quad B = \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}, \quad C = \begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix},$$

so that we rewrite the equation  $C = A \cdot B$  as

$$\begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix} = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \cdot \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}.$$



# Matrix multiplication: divide-and-conquer

$$A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}, \quad B = \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}, \quad C = \begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix},$$

so that we rewrite the equation  $C = A \cdot B$  as

$$\begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix} = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \cdot \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}.$$

$$C_{11} = A_{11} \cdot B_{11} + A_{12} \cdot B_{21},$$

$$C_{12} = A_{11} \cdot B_{12} + A_{12} \cdot B_{22},$$

$$C_{21} = A_{21} \cdot B_{11} + A_{22} \cdot B_{21},$$

$$C_{22} = A_{21} \cdot B_{12} + A_{22} \cdot B_{22}.$$

# Matrix multiplication: divide-and-conquer

MATRIX-MULTIPLY-RECURSIVE( $A, B, C, n$ )

```
1  if  $n == 1$ 
2    // Base case.
3       $c_{11} = c_{11} + a_{11} \cdot b_{11}$ 
4      return
5    // Divide.
6    partition  $A, B$ , and  $C$  into  $n/2 \times n/2$  submatrices
       $A_{11}, A_{12}, A_{21}, A_{22}; B_{11}, B_{12}, B_{21}, B_{22};$ 
      and  $C_{11}, C_{12}, C_{21}, C_{22}$ ; respectively
7    // Conquer.
8    MATRIX-MULTIPLY-RECURSIVE( $A_{11}, B_{11}, C_{11}, n/2$ )
9    MATRIX-MULTIPLY-RECURSIVE( $A_{11}, B_{12}, C_{12}, n/2$ )
10   MATRIX-MULTIPLY-RECURSIVE( $A_{21}, B_{11}, C_{21}, n/2$ )
11   MATRIX-MULTIPLY-RECURSIVE( $A_{21}, B_{12}, C_{22}, n/2$ )
12   MATRIX-MULTIPLY-RECURSIVE( $A_{12}, B_{21}, C_{11}, n/2$ )
13   MATRIX-MULTIPLY-RECURSIVE( $A_{12}, B_{22}, C_{12}, n/2$ )
14   MATRIX-MULTIPLY-RECURSIVE( $A_{22}, B_{21}, C_{21}, n/2$ )
15   MATRIX-MULTIPLY-RECURSIVE( $A_{22}, B_{22}, C_{22}, n/2$ )
```

# Matrix multiplication: divide-and-conquer

$$\begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix} = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \cdot \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}$$

$$C_{11} = A_{11} \cdot B_{11} + A_{12} \cdot B_{21} ,$$

$$C_{12} = A_{11} \cdot B_{12} + A_{12} \cdot B_{22} ,$$

$$C_{21} = A_{21} \cdot B_{11} + A_{22} \cdot B_{21} ,$$

$$C_{22} = A_{21} \cdot B_{12} + A_{22} \cdot B_{22} .$$

# Matrix multiplication: divide-and-conquer

$$\begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix} = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \cdot \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}$$

$$C_{11} = A_{11} \cdot B_{11} + A_{12} \cdot B_{21} ,$$

$$C_{12} = A_{11} \cdot B_{12} + A_{12} \cdot B_{22} ,$$

$$C_{21} = A_{21} \cdot B_{11} + A_{22} \cdot B_{21} ,$$

$$C_{22} = A_{21} \cdot B_{12} + A_{22} \cdot B_{22} .$$

►  $T(n) = 8T(n/2) + D(n) + C(n)$

# Matrix multiplication: divide-and-conquer

$$\begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix} = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \cdot \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}$$

$$C_{11} = A_{11} \cdot B_{11} + A_{12} \cdot B_{21} ,$$

$$C_{12} = A_{11} \cdot B_{12} + A_{12} \cdot B_{22} ,$$


$$C_{21} = A_{21} \cdot B_{11} + A_{22} \cdot B_{21} ,$$

$$C_{22} = A_{21} \cdot B_{12} + A_{22} \cdot B_{22} .$$

▶  $T(n) = 8T(n/2) + D(n) + C(n)$

▶  $T(n) = 8T(n/2) + \Theta(1) + \Theta(1)$

# Recursion tree for merge sort



# Matrix multiplication: divide-and-conquer

►  $T(n) = c8^{\lg n} + \dots$

# Matrix multiplication: divide-and-conquer

►  $T(n) = c8^{\lg n} + \dots$

$$T(n) = cn^{\lg 8} + \dots$$



# Matrix multiplication: divide-and-conquer

►  $T(n) = c8^{\lg n} + \dots$

$$T(n) = cn^{\lg 8} + \dots$$

$$T(n) = \Theta(n^3)$$

# Matrix multiplication: divide-and-conquer

- ▶  $T(n) = c8^{\lg n} + \dots$   
 $T(n) = cn^{\lg 8} + \dots$   
 $T(n) = \Theta(n^3)$
- ▶ Strassen modified the divide-and-conquer approach explained above such that  $T(n) = O(n^{\lg 7})$

# Matrix multiplication: divide-and-conquer

- ▶  $T(n) = c8^{\lg n} + \dots$   
 $T(n) = cn^{\lg 8} + \dots$   
 $T(n) = \Theta(n^3)$
- ▶ Strassen modified the divide-and-conquer approach explained above such that  $T(n) = O(n^{\lg 7})$
- ▶  $T(n) = c7^{\lg n} + \dots$

# Matrix multiplication: divide-and-conquer

- ▶  $T(n) = c8^{\lg n} + \dots$   
 $T(n) = cn^{\lg 8} + \dots$   
 $T(n) = \Theta(n^3)$
- ▶ Strassen modified the divide-and-conquer approach explained above such that  $T(n) = O(n^{\lg 7})$
- ▶  $T(n) = c7^{\lg n} + \dots$   
 $T(n) = cn^{\lg 7} + \dots$

# Matrix multiplication: divide-and-conquer

- ▶  $T(n) = c8^{\lg n} + \dots$   
 $T(n) = cn^{\lg 8} + \dots$   
 $T(n) = \Theta(n^3)$
- ▶ Strassen modified the divide-and-conquer approach explained above such that  $T(n) = O(n^{\lg 7})$
- ▶  $T(n) = c7^{\lg n} + \dots$   
 $T(n) = cn^{\lg 7} + \dots$   
 $T(n) = \Theta(n^{\lg 7})$

# Matrix multiplication: divide-and-conquer

►  $T(n) = c8^{\lg n} + \dots$

$$T(n) = cn^{\lg 8} + \dots$$

$$T(n) = \Theta(n^3)$$

- Strassen modified the divide-and-conquer approach explained above such that  $T(n) = O(n^{\lg 7})$

►  $T(n) = c7^{\lg n} + \dots$

$$T(n) = cn^{\lg 7} + \dots$$

$$T(n) = \Theta(n^{\lg 7})$$

n	0	1	2	3	4	5	6
$8^n$	1	8	64	512	4096	32768	262144
$7^n$	1	7	49	343	2401	16807	117649
$2^n$	1	2	4	8	16	32	64

# Strassen's algorithm : Divide (Step 1)

Divide step involves finding ten  $n/2 \times n/2$  matrices ( $S_i$ ).

# Strassen's algorithm : Divide (Step 1)

Divide step involves finding ten  $n/2 \times n/2$  matrices ( $S_i$ ).

$$S_1 = B_{12} - B_{22} ,$$

$$S_2 = A_{11} + A_{12} ,$$

$$S_3 = A_{21} + A_{22} ,$$

$$S_4 = B_{21} - B_{11} ,$$

$$S_5 = A_{11} + A_{22} ,$$

$$S_6 = B_{11} + B_{22} ,$$

$$S_7 = A_{12} - A_{22} ,$$

$$S_8 = B_{21} + B_{22} ,$$

$$S_9 = A_{11} - A_{21} ,$$

$$S_{10} = B_{11} + B_{12} .$$



## Strassen's algorithm : Conquer (Step 2)

$$P_1 = A_{11} \cdot S_1$$

$$P_2 = S_2 \cdot B_{22}$$

$$P_3 = S_3 \cdot B_{11}$$

$$P_4 = A_{22} \cdot S_4$$

$$P_5 = S_5 \cdot S_6$$

$$P_6 = S_7 \cdot S_8$$

$$P_7 = S_9 \cdot S_{10}$$

## Strassen's algorithm : Conquer (Step 2)

$$P_1 = A_{11} \cdot S_1 = A_{11} \cdot B_{12} - A_{11} \cdot B_{22} ,$$

$$P_2 = S_2 \cdot B_{22} = A_{11} \cdot B_{22} + A_{12} \cdot B_{22} ,$$

$$P_3 = S_3 \cdot B_{11} = A_{21} \cdot B_{11} + A_{22} \cdot B_{11} ,$$

$$P_4 = A_{22} \cdot S_4 = A_{22} \cdot B_{21} - A_{22} \cdot B_{11} ,$$

$$P_5 = S_5 \cdot S_6 = A_{11} \cdot B_{11} + A_{11} \cdot B_{22} + A_{22} \cdot B_{11} + A_{22} \cdot B_{22} ,$$

$$P_6 = S_7 \cdot S_8 = A_{12} \cdot B_{21} + A_{12} \cdot B_{22} - A_{22} \cdot B_{21} - A_{22} \cdot B_{22} ,$$

$$P_7 = S_9 \cdot S_{10} = A_{11} \cdot B_{11} + A_{11} \cdot B_{12} - A_{21} \cdot B_{11} - A_{21} \cdot B_{12} .$$

# Matrix multiplication: divide-and-conquer

$$\begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix} = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \cdot \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}$$

$$C_{11} = A_{11} \cdot B_{11} + A_{12} \cdot B_{21} ,$$

$$C_{12} = A_{11} \cdot B_{12} + A_{12} \cdot B_{22} ,$$

$$C_{21} = A_{21} \cdot B_{11} + A_{22} \cdot B_{21} ,$$

$$C_{22} = A_{21} \cdot B_{12} + A_{22} \cdot B_{22} .$$

## Strassen's algorithm : Combine (Step 3)

$$C_{11} = P_5 + P_4 - P_2 + P_6 .$$

$$\begin{array}{r} A_{11} \cdot B_{11} + A_{11} \cdot B_{22} + A_{22} \cdot B_{11} + A_{22} \cdot B_{22} \\ \quad \quad \quad - A_{22} \cdot B_{11} \quad \quad \quad + A_{22} \cdot B_{21} \\ \quad \quad \quad - A_{11} \cdot B_{22} \quad \quad \quad - A_{12} \cdot B_{22} \\ \quad \quad \quad \quad \quad \quad - A_{22} \cdot B_{22} - A_{22} \cdot B_{21} + A_{12} \cdot B_{22} + A_{12} \cdot B_{21} \\ \hline A_{11} \cdot B_{11} \quad \quad \quad \quad \quad \quad \quad \quad \quad \quad \quad \quad \quad \quad \quad + A_{12} \cdot B_{21} , \end{array}$$

## Strassen's algorithm : Combine (Step 3)

$$C_{12} = P_1 + P_2 ,$$

$$\begin{array}{r} A_{11} \cdot B_{12} - A_{11} \cdot B_{22} \\ + A_{11} \cdot B_{22} + A_{12} \cdot B_{22} \\ \hline A_{11} \cdot B_{12} \qquad + A_{12} \cdot B_{22} , \end{array}$$

## Strassen's algorithm : Combine (Step 3)

$$C_{12} = P_1 + P_2 ,$$

$$\begin{array}{r} A_{11} \cdot B_{12} - A_{11} \cdot B_{22} \\ + A_{11} \cdot B_{22} + A_{12} \cdot B_{22} \\ \hline A_{11} \cdot B_{12} \qquad + A_{12} \cdot B_{22} , \end{array}$$

$$C_{21} = P_3 + P_4$$

$$\begin{array}{r} A_{21} \cdot B_{11} + A_{22} \cdot B_{11} \\ - A_{22} \cdot B_{11} + A_{22} \cdot B_{21} \\ \hline A_{21} \cdot B_{11} \qquad + A_{22} \cdot B_{21} , \end{array}$$

# Strassen's algorithm : Combine (Step 3)

$$C_{12} = P_1 + P_2 ,$$

$$C_{21} = P_3 + P_4$$

$$\begin{array}{r} A_{11} \cdot B_{12} - A_{11} \cdot B_{22} \\ + A_{11} \cdot B_{22} + A_{12} \cdot B_{22} \\ \hline A_{11} \cdot B_{12} \qquad + A_{12} \cdot B_{22} , \end{array}$$

$$\begin{array}{r} A_{21} \cdot B_{11} + A_{22} \cdot B_{11} \\ - A_{22} \cdot B_{11} + A_{22} \cdot B_{21} \\ \hline A_{21} \cdot B_{11} \qquad + A_{22} \cdot B_{21} , \end{array}$$

$$C_{22} = P_5 + P_1 - P_3 - P_7 ,$$

$$\begin{array}{r} A_{11} \cdot B_{11} + A_{11} \cdot B_{22} + A_{22} \cdot B_{11} + A_{22} \cdot B_{22} \\ - A_{11} \cdot B_{22} \qquad + A_{11} \cdot B_{12} \\ - A_{22} \cdot B_{11} \qquad - A_{21} \cdot B_{11} \\ - A_{11} \cdot B_{11} \qquad - A_{11} \cdot B_{12} + A_{21} \cdot B_{11} + A_{21} \cdot B_{12} \\ \hline A_{22} \cdot B_{22} \qquad + A_{21} \cdot B_{12} , \end{array}$$

# Strassen's algorithm

►  $T(n) = 7T(n/2) + D(n) + C(n)$



# Strassen's algorithm

- ▶  $T(n) = 7T(n/2) + D(n) + C(n)$
- ▶  $T(n) = 7T(n/2) + \Theta(n^2) + \Theta(n^2)$

# Strassen's algorithm

- ▶  $T(n) = 7T(n/2) + D(n) + C(n)$
- ▶  $T(n) = 7T(n/2) + \Theta(n^2) + \Theta(n^2)$
- ▶  $T(n) = 7^{\lg n} + \dots$

# Strassen's algorithm

- ▶  $T(n) = 7T(n/2) + D(n) + C(n)$
- ▶  $T(n) = 7T(n/2) + \Theta(n^2) + \Theta(n^2)$
- ▶  $T(n) = 7^{\lg n} + \dots$   
 $T(n) = n^{\lg 7} + \dots$

# Strassen's algorithm

- ▶  $T(n) = 7T(n/2) + D(n) + C(n)$
- ▶  $T(n) = 7T(n/2) + \Theta(n^2) + \Theta(n^2)$
- ▶  $T(n) = 7^{\lg n} + \dots$   
 $T(n) = n^{\lg 7} + \dots$   
 $T(n) = O(n^{\lg 7}) = o(n^{2.81})$

# Strassen's algorithm

- ▶  $T(n) = 7T(n/2) + D(n) + C(n)$
- ▶  $T(n) = 7T(n/2) + \Theta(n^2) + \Theta(n^2)$
- ▶  $T(n) = 7^{\lg n} + \dots$   
 $T(n) = n^{\lg 7} + \dots$   
 $T(n) = O(n^{\lg 7}) = o(n^{2.81})$
- ▶ Why should  $n$  be a power of 2?

# Strassen's algorithm

- Q. How would you modify Strassen's algorithm to multiply  $n \times n$  matrices in which  $n$  is not an exact power of 2?

# Strassen's algorithm

- Q. How would you modify Strassen's algorithm to multiply  $n \times n$  matrices in which  $n$  is not an exact power of 2?

$$n^{\lg 7} \leq m^{\lg 7} <$$

# Strassen's algorithm

- Q. How would you modify Strassen's algorithm to multiply  $n \times n$  matrices in which  $n$  is not an exact power of 2?

$$n^{\lg 7} \leq m^{\lg 7} < (2n)^{\lg 7}$$



# Strassen's algorithm

- Q. How would you modify Strassen's algorithm to multiply  $n \times n$  matrices in which  $n$  is not an exact power of 2?

$$n^{\lg 7} \leq m^{\lg 7} < (2n)^{\lg 7} = 2^{\lg 7} n^{\lg 7}$$

# Strassen's algorithm

- Q. How would you modify Strassen's algorithm to multiply  $n \times n$  matrices in which  $n$  is not an exact power of 2?

$$n^{\lg 7} \leq m^{\lg 7} < (2n)^{\lg 7} = 2^{\lg 7} n^{\lg 7} = 7n^{\lg 7}$$

# Strassen's algorithm

- Q. How would you modify Strassen's algorithm to multiply  $n \times n$  matrices in which  $n$  is not an exact power of 2?

$$n^{\lg 7} \leq m^{\lg 7} < (2n)^{\lg 7} = 2^{\lg 7} n^{\lg 7} = 7n^{\lg 7}$$

# Strassen's algorithm

- Q. How would you modify Strassen's algorithm to multiply  $n \times n$  matrices in which  $n$  is not an exact power of 2?

$$n^{\lg 7} \leq m^{\lg 7} < (2n)^{\lg 7} = 2^{\lg 7} n^{\lg 7} = 7n^{\lg 7}$$

The above implies that  $m^{\lg 7} = \Theta(n^{\lg 7})$ .

## Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

- Find an upper bound for  $T(n) = 2T(\lfloor n/2 \rfloor) + n$

## Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

- ▶ Find an upper bound for  $T(n) = 2T(\lfloor n/2 \rfloor) + n$
- ▶ We will first guess a solution:  $O(n \lg n)$  (using Recursion tree)

## Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

- ▶ Find an upper bound for  $T(n) = 2T(\lfloor n/2 \rfloor) + n$
- ▶ We will first guess a solution:  $O(n \lg n)$  (using Recursion tree)
- ▶ To show :  $T(n) = O(n \lg n)$

## Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

- ▶ Find an upper bound for  $T(n) = 2T(\lfloor n/2 \rfloor) + n$
- ▶ We will first guess a solution:  $O(n \lg n)$  (using Recursion tree)
- ▶ To show :  $T(n) = O(n \lg n)$   
 $T(n) \leq dn \lg n$



## Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

- ▶ Find an upper bound for  $T(n) = 2T(\lfloor n/2 \rfloor) + n$
- ▶ We will first guess a solution:  $O(n \lg n)$  (using Recursion tree)
- ▶ To show :  $T(n) = O(n \lg n)$   
$$T(n) \leq dn \lg n$$
- ▶ We will assume that  $T(n/2) \leq d(n/2) \lg(n/2)$ . Then show that  $T(n) \leq dn \lg n$ . (Inductive step)

# Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

Inductive step : Assume  $T(n/2) \leq d(n/2) \lg(n/2)$

# Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

Inductive step : Assume  $T(n/2) \leq d(n/2) \lg(n/2)$

$$T(\lfloor n/2 \rfloor) \leq d(\lfloor n/2 \rfloor) \lg(\lfloor n/2 \rfloor) \leq d(n/2) \lg(n/2)$$

# Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

Inductive step : Assume  $T(n/2) \leq d(n/2) \lg(n/2)$

$$T(\lfloor n/2 \rfloor) \leq d(\lfloor n/2 \rfloor) \lg(\lfloor n/2 \rfloor) \leq d(n/2) \lg(n/2)$$

$$T(n) = 2T(\lfloor n/2 \rfloor) + n$$

# Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

Inductive step : Assume  $T(n/2) \leq d(n/2) \lg(n/2)$

$$T(\lfloor n/2 \rfloor) \leq d(\lfloor n/2 \rfloor) \lg(\lfloor n/2 \rfloor) \leq d(n/2) \lg(n/2)$$

$$\begin{aligned} T(n) &= 2T(\lfloor n/2 \rfloor) + n \\ &\leq 2d(n/2) \lg(n/2) + n \end{aligned}$$

# Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

Inductive step : Assume  $T(n/2) \leq d(n/2) \lg(n/2)$

$$T(\lfloor n/2 \rfloor) \leq d(\lfloor n/2 \rfloor) \lg(\lfloor n/2 \rfloor) \leq d(n/2) \lg(n/2)$$

$$\begin{aligned} T(n) &= 2T(\lfloor n/2 \rfloor) + n \\ &\leq 2d(n/2) \lg(n/2) + n \\ &= dn(\lg n - \lg 2) + n \end{aligned}$$

# Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

Inductive step : Assume  $T(n/2) \leq d(n/2) \lg(n/2)$

$$T(\lfloor n/2 \rfloor) \leq d(\lfloor n/2 \rfloor) \lg(\lfloor n/2 \rfloor) \leq d(n/2) \lg(n/2)$$

$$\begin{aligned} T(n) &= 2T(\lfloor n/2 \rfloor) + n \\ &\leq 2d(n/2) \lg(n/2) + n \\ &= dn(\lg n - \lg 2) + n \\ &= dn(\lg n - 1) + n \end{aligned}$$

# Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

Inductive step : Assume  $T(n/2) \leq d(n/2) \lg(n/2)$

$$T(\lfloor n/2 \rfloor) \leq d(\lfloor n/2 \rfloor) \lg(\lfloor n/2 \rfloor) \leq d(n/2) \lg(n/2)$$

$$\begin{aligned} T(n) &= 2T(\lfloor n/2 \rfloor) + n \\ &\leq 2d(n/2) \lg(n/2) + n \\ &= dn(\lg n - \lg 2) + n \\ &= dn(\lg n - 1) + n \\ &= dn \lg n - dn + n \end{aligned}$$



# Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

Inductive step : Assume  $T(n/2) \leq d(n/2) \lg(n/2)$

$$T(\lfloor n/2 \rfloor) \leq d(\lfloor n/2 \rfloor) \lg(\lfloor n/2 \rfloor) \leq d(n/2) \lg(n/2)$$

$$\begin{aligned} T(n) &= 2T(\lfloor n/2 \rfloor) + n \\ &\leq 2d(n/2) \lg(n/2) + n \\ &= dn(\lg n - \lg 2) + n \\ &= dn(\lg n - 1) + n \\ &= dn \lg n - dn + n \\ &\leq dn \lg n, \text{ for } d \geq 1 \end{aligned}$$

# Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

Inductive step : Assume  $T(n/2) \leq d(n/2) \lg(n/2)$

$$T(\lfloor n/2 \rfloor) \leq d(\lfloor n/2 \rfloor) \lg(\lfloor n/2 \rfloor) \leq d(n/2) \lg(n/2)$$

$$\begin{aligned} T(n) &= 2T(\lfloor n/2 \rfloor) + n \\ &\leq 2d(n/2) \lg(n/2) + n \\ &= dn(\lg n - \lg 2) + n \\ &= dn(\lg n - 1) + n \\ &= dn \lg n - dn + n \\ &\leq dn \lg n, \text{ for } d \geq 1 \end{aligned}$$

Base case is problematic

$$T(1) = 1$$

# Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

Inductive step : Assume  $T(n/2) \leq d(n/2) \lg(n/2)$

$$T(\lfloor n/2 \rfloor) \leq d(\lfloor n/2 \rfloor) \lg(\lfloor n/2 \rfloor) \leq d(n/2) \lg(n/2)$$

$$\begin{aligned} T(n) &= 2T(\lfloor n/2 \rfloor) + n \\ &\leq 2d(n/2) \lg(n/2) + n \\ &= dn(\lg n - \lg 2) + n \\ &= dn(\lg n - 1) + n \\ &= dn \lg n - dn + n \\ &\leq dn \lg n, \text{ for } d \geq 1 \end{aligned}$$

Base case is problematic

$$T(1) = 1 \not\leq d 1 \lg 1$$

# Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

Inductive step : Assume  $T(n/2) \leq d(n/2) \lg(n/2)$

$$T(\lfloor n/2 \rfloor) \leq d(\lfloor n/2 \rfloor) \lg(\lfloor n/2 \rfloor) \leq d(n/2) \lg(n/2)$$

$$\begin{aligned} T(n) &= 2T(\lfloor n/2 \rfloor) + n \\ &\leq 2d(n/2) \lg(n/2) + n \\ &= dn(\lg n - \lg 2) + n \\ &= dn(\lg n - 1) + n \\ &= dn \lg n - dn + n \\ &\leq dn \lg n, \text{ for } d \geq 1 \end{aligned}$$

Base case is problematic

$$T(1) = 1 \not\leq d 1 \lg 1 = 0$$

# Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

►  $T(2) = 2T(\lfloor 2/2 \rfloor) + 2$

## Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

►  $T(2) = 2T(\lfloor 2/2 \rfloor) + 2 = 4$

## Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

►  $T(2) = 2T(\lfloor 2/2 \rfloor) + 2 = 4 \leq d2 \lg 2$

## Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

- ▶  $T(2) = 2T(\lfloor 2/2 \rfloor) + 2 = 4 \leq d2 \lg 2$
- ▶  $T(3) = 2T(\lfloor 3/2 \rfloor) + 3$



## Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

- ▶  $T(2) = 2T(\lfloor 2/2 \rfloor) + 2 = 4 \leq d2 \lg 2$
- ▶  $T(3) = 2T(\lfloor 3/2 \rfloor) + 3 = 5$

## Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

- ▶  $T(2) = 2T(\lfloor 2/2 \rfloor) + 2 = 4 \leq d2 \lg 2$
- ▶  $T(3) = 2T(\lfloor 3/2 \rfloor) + 3 = 5 \leq d3 \lg 3$

## Substitution method : $T(n) = 2T(\lfloor n/2 \rfloor) + n$

- ▶  $T(2) = 2T(\lfloor 2/2 \rfloor) + 2 = 4 \leq d2 \lg 2$
- ▶  $T(3) = 2T(\lfloor 3/2 \rfloor) + 3 = 5 \leq d3 \lg 3$
- ▶ We only need to show that  $T(n) \leq dn \lg n$  for  $n \geq n_0$ , where  $n_0$  need not be 1.

Substitution method :  $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

- Find a tight upper bound for the recurrence  
 $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

- ▶ Find a tight upper bound for the recurrence  
 $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$
- ▶ What can be an initial guess for the above question based on recursion tree?

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

- ▶ Find a tight upper bound for the recurrence  
 $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$
- ▶ What can be an initial guess for the above question based on recursion tree?
- ▶ We guess  $T(n) = O(n)$ .

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

- ▶ Find a tight upper bound for the recurrence  
 $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$
- ▶ What can be an initial guess for the above question based on recursion tree?
- ▶ We guess  $T(n) = O(n)$ .  
To prove :  $T(n) \leq cn \quad \forall n \geq n_0$



# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

- ▶ Find a tight upper bound for the recurrence  
 $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$
- ▶ What can be an initial guess for the above question based on recursion tree?
- ▶ We guess  $T(n) = O(n)$ .  
To prove :  $T(n) \leq cn \quad \forall n \geq n_0$   
Induction step assumption :

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

- ▶ Find a tight upper bound for the recurrence  
 $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$
- ▶ What can be an initial guess for the above question based on recursion tree?
- ▶ We guess  $T(n) = O(n)$ .

To prove :  $T(n) \leq cn \quad \forall n \geq n_0$

Induction step assumption :

$$T(\lfloor n/2 \rfloor) \leq c(\lfloor n/2 \rfloor) \quad , \quad T(\lceil n/2 \rceil) \leq c(\lceil n/2 \rceil)$$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

- ▶ Find a tight upper bound for the recurrence  
 $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$
- ▶ What can be an initial guess for the above question based on recursion tree?
- ▶ We guess  $T(n) = O(n)$ .  
To prove :  $T(n) \leq cn \quad \forall n \geq n_0$   
Induction step assumption :

$$T(\lfloor n/2 \rfloor) \leq c(\lfloor n/2 \rfloor) \quad , \quad T(\lceil n/2 \rceil) \leq c(\lceil n/2 \rceil)$$
$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

- ▶ Find a tight upper bound for the recurrence  
 $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$
- ▶ What can be an initial guess for the above question based on recursion tree?
- ▶ We guess  $T(n) = O(n)$ .  
To prove :  $T(n) \leq cn \quad \forall n \geq n_0$   
Induction step assumption :

$$T(\lfloor n/2 \rfloor) \leq c(\lfloor n/2 \rfloor) \quad , \quad T(\lceil n/2 \rceil) \leq c(\lceil n/2 \rceil)$$
$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1 \leq c\lfloor n/2 \rfloor + c\lceil n/2 \rceil + 1$$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

- ▶ Find a tight upper bound for the recurrence  
 $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$
- ▶ What can be an initial guess for the above question based on recursion tree?
- ▶ We guess  $T(n) = O(n)$ .

To prove :  $T(n) \leq cn \quad \forall n \geq n_0$

Induction step assumption :

$$T(\lfloor n/2 \rfloor) \leq c\lfloor n/2 \rfloor \quad , \quad T(\lceil n/2 \rceil) \leq c\lceil n/2 \rceil$$

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1 \leq c\lfloor n/2 \rfloor + c\lceil n/2 \rceil + 1$$

$$T(n) \leq c\lfloor n/2 \rfloor + c\lceil n/2 \rceil + 1$$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

- ▶ Find a tight upper bound for the recurrence  
 $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$
- ▶ What can be an initial guess for the above question based on recursion tree?
- ▶ We guess  $T(n) = O(n)$ .

To prove :  $T(n) \leq cn \quad \forall n \geq n_0$

Induction step assumption :

$$\begin{aligned} T(\lfloor n/2 \rfloor) &\leq c\lfloor n/2 \rfloor \quad , \quad T(\lceil n/2 \rceil) \leq c\lceil n/2 \rceil \\ T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1 &\leq c\lfloor n/2 \rfloor + c\lceil n/2 \rceil + 1 \\ T(n) &\leq c\lfloor n/2 \rfloor + c\lceil n/2 \rceil + 1 \\ &= cn + 1 \end{aligned}$$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

- ▶ Find a tight upper bound for the recurrence  
 $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$
- ▶ What can be an initial guess for the above question based on recursion tree?
- ▶ We guess  $T(n) = O(n)$ .

To prove :  $T(n) \leq cn \quad \forall n \geq n_0$

Induction step assumption :

$$\begin{aligned} T(\lfloor n/2 \rfloor) &\leq c\lfloor n/2 \rfloor \quad , \quad T(\lceil n/2 \rceil) \leq c\lceil n/2 \rceil \\ T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1 &\leq c\lfloor n/2 \rfloor + c\lceil n/2 \rceil + 1 \\ T(n) &\leq c\lfloor n/2 \rfloor + c\lceil n/2 \rceil + 1 \\ &= cn + 1 \quad (\text{cannot proceed}) \end{aligned}$$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

We need to slightly modify the initial guess for the recurrence  
 $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$ .



# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

We need to slightly modify the initial guess for the recurrence

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1.$$

To prove:  $T(n) \leq cn - d \quad \forall n \geq n_0.$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

We need to slightly modify the initial guess for the recurrence

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1.$$

To prove:  $T(n) \leq cn - d \quad \forall n \geq n_0.$

$$T(\lfloor n/2 \rfloor) \leq c(\lfloor n/2 \rfloor) - d$$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

We need to slightly modify the initial guess for the recurrence

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1.$$

To prove:  $T(n) \leq cn - d \quad \forall n \geq n_0.$

$$T(\lfloor n/2 \rfloor) \leq c(\lfloor n/2 \rfloor) - d$$

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

We need to slightly modify the initial guess for the recurrence

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1.$$

To prove:  $T(n) \leq cn - d \quad \forall n \geq n_0.$

$$T(\lfloor n/2 \rfloor) \leq c(\lfloor n/2 \rfloor) - d$$

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1 \leq c\lfloor n/2 \rfloor - d + c\lceil n/2 \rceil - d + 1$$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

We need to slightly modify the initial guess for the recurrence

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1.$$

To prove:  $T(n) \leq cn - d \quad \forall n \geq n_0.$

$$T(\lfloor n/2 \rfloor) \leq c(\lfloor n/2 \rfloor) - d$$

$$\begin{aligned} T(n) &= T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1 \leq c\lfloor n/2 \rfloor - d + c\lceil n/2 \rceil - d + 1 \\ &= cn - 2d + 1 \end{aligned}$$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

We need to slightly modify the initial guess for the recurrence

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1.$$

To prove:  $T(n) \leq cn - d \quad \forall n \geq n_0.$

$$T(\lfloor n/2 \rfloor) \leq c(\lfloor n/2 \rfloor) - d$$

$$\begin{aligned} T(n) &= T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1 \leq c\lfloor n/2 \rfloor - d + c\lceil n/2 \rceil - d + 1 \\ &= cn - 2d + 1 \\ &= cn - d - d + 1 \end{aligned}$$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

We need to slightly modify the initial guess for the recurrence

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1.$$

To prove:  $T(n) \leq cn - d \quad \forall n \geq n_0.$

$$T(\lfloor n/2 \rfloor) \leq c(\lfloor n/2 \rfloor) - d$$

$$\begin{aligned} T(n) &= T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1 \leq c\lfloor n/2 \rfloor - d + c\lceil n/2 \rceil - d + 1 \\ &= cn - 2d + 1 \\ &= cn - d - d + 1 \\ &\leq cn - d, \text{ for } d \geq 1 \end{aligned}$$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

We need to slightly modify the initial guess for the recurrence

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1.$$

To prove:  $T(n) \leq cn - d \quad \forall n \geq n_0.$

$$T(\lfloor n/2 \rfloor) \leq c(\lfloor n/2 \rfloor) - d$$

$$\begin{aligned} T(n) &= T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1 \leq c\lfloor n/2 \rfloor - d + c\lceil n/2 \rceil - d + 1 \\ &= cn - 2d + 1 \\ &= cn - d - d + 1 \\ &\leq cn - d, \text{ for } d \geq 1 \end{aligned}$$

$$T(1) = 1 \leq cn - d$$



# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

We need to slightly modify the initial guess for the recurrence

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1.$$

To prove:  $T(n) \leq cn - d \quad \forall n \geq n_0.$

$$T(\lfloor n/2 \rfloor) \leq c(\lfloor n/2 \rfloor) - d$$

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1 \leq c\lfloor n/2 \rfloor - d + c\lceil n/2 \rceil - d + 1$$

$$= cn - 2d + 1$$

$$= cn - d - d + 1$$

$$\leq cn - d, \text{ for } d \geq 1$$

$$T(1) = 1 \leq cn - d, \text{ for } d \geq 1 \text{ and } c \geq d + 1$$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

We need to slightly modify the initial guess for the recurrence

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1.$$

To prove:  $T(n) \leq cn - d \quad \forall n \geq n_0.$

$$T(\lfloor n/2 \rfloor) \leq c(\lfloor n/2 \rfloor) - d$$

$$\begin{aligned} T(n) &= T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1 \leq c\lfloor n/2 \rfloor - d + c\lceil n/2 \rceil - d + 1 \\ &= cn - 2d + 1 \\ &= cn - d - d + 1 \\ &\leq cn - d, \text{ for } d \geq 1 \end{aligned}$$

$$T(1) = 1 \leq cn - d, \text{ for } d \geq 1 \text{ and } c \geq d + 1$$

For  $c = 2, d = 1$

# Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

We need to slightly modify the initial guess for the recurrence

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1.$$

To prove:  $T(n) \leq cn - d \quad \forall n \geq n_0$ .

$$T(\lfloor n/2 \rfloor) \leq c(\lfloor n/2 \rfloor) - d$$

$$\begin{aligned} T(n) &= T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1 \leq c\lfloor n/2 \rfloor - d + c\lceil n/2 \rceil - d + 1 \\ &= cn - 2d + 1 \\ &= cn - d - d + 1 \\ &\leq cn - d, \text{ for } d \geq 1 \end{aligned}$$

$$T(1) = 1 \leq cn - d, \text{ for } d \geq 1 \text{ and } c \geq d + 1$$

$$\text{For } c = 2, d = 1 \quad T(n) \leq 2n - 1 \quad \forall n \geq 1$$

## Substitution method : $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1$

- ▶ Instead of showing  $T(n) = O(n)$  we have instead shown that  $T(n) = O(cn - d)$ , which implies  $T(n) = O(n)$

## Substitution method : $T(n) = T(n - 1) + n$

- Find a lower bound for the recurrence  $T(n) = T(n - 1) + n$

## Substitution method : $T(n) = T(n - 1) + n$

- ▶ Find a lower bound for the recurrence  $T(n) = T(n - 1) + n$
- ▶ What would be a good guess?

## Substitution method : $T(n) = T(n-1) + n$

- ▶ Find a lower bound for the recurrence  $T(n) = T(n-1) + n$
- ▶ What would be a good guess?  $T(n) = \Omega(n^2)$

## Substitution method : $T(n) = T(n-1) + n$

- ▶ Find a lower bound for the recurrence  $T(n) = T(n-1) + n$
- ▶ What would be a good guess?  $T(n) = \Omega(n^2)$

To show :  $T(n) \geq cn^2 \quad \forall n \geq n_0$



## Substitution method : $T(n) = T(n-1) + n$

- ▶ Find a lower bound for the recurrence  $T(n) = T(n-1) + n$
- ▶ What would be a good guess?  $T(n) = \Omega(n^2)$

To show :  $T(n) \geq cn^2 \quad \forall n \geq n_0$

Induction step:

$$T(n-1) \geq c(n-1)^2$$

## Substitution method : $T(n) = T(n-1) + n$

- ▶ Find a lower bound for the recurrence  $T(n) = T(n-1) + n$
- ▶ What would be a good guess?  $T(n) = \Omega(n^2)$

To show :  $T(n) \geq cn^2 \quad \forall n \geq n_0$

Induction step:

$$\begin{aligned} T(n-1) &\geq c(n-1)^2 \\ &= c(n^2 + 1 - 2n) = cn^2 + c - 2cn \end{aligned}$$

## Substitution method : $T(n) = T(n-1) + n$

- ▶ Find a lower bound for the recurrence  $T(n) = T(n-1) + n$
- ▶ What would be a good guess?  $T(n) = \Omega(n^2)$

To show :  $T(n) \geq cn^2 \quad \forall n \geq n_0$

Induction step:

$$\begin{aligned} T(n-1) &\geq c(n-1)^2 \\ &= c(n^2 + 1 - 2n) = cn^2 + c - 2cn \\ T(n) &\geq cn^2 + c - 2cn + n \end{aligned}$$

## Substitution method : $T(n) = T(n-1) + n$

- ▶ Find a lower bound for the recurrence  $T(n) = T(n-1) + n$
- ▶ What would be a good guess?  $T(n) = \Omega(n^2)$

To show :  $T(n) \geq cn^2 \quad \forall n \geq n_0$

Induction step:

$$\begin{aligned} T(n-1) &\geq c(n-1)^2 \\ &= c(n^2 + 1 - 2n) = cn^2 + c - 2cn \\ T(n) &\geq cn^2 + c - 2cn + n \quad (\text{when } c=1/2, \text{ R.H.S.} \geq cn^2) \end{aligned}$$

## Substitution method : $T(n) = T(n-1) + n$

- ▶ Find a lower bound for the recurrence  $T(n) = T(n-1) + n$
- ▶ What would be a good guess?  $T(n) = \Omega(n^2)$

To show :  $T(n) \geq cn^2 \quad \forall n \geq n_0$

Induction step:

$$\begin{aligned} T(n-1) &\geq c(n-1)^2 \\ &= c(n^2 + 1 - 2n) = cn^2 + c - 2cn \\ T(n) &\geq cn^2 + c - 2cn + n \quad (\text{when } c=1/2, \text{ R.H.S.} \geq cn^2) \\ &\geq cn^2 \quad (\text{for } c=1/2 \text{ and } n \geq 1) \end{aligned}$$

## Substitution method : $T(n) = T(n-1) + n$

- ▶ Find a lower bound for the recurrence  $T(n) = T(n-1) + n$
- ▶ What would be a good guess?  $T(n) = \Omega(n^2)$

To show :  $T(n) \geq cn^2 \quad \forall n \geq n_0$

Induction step:

$$\begin{aligned} T(n-1) &\geq c(n-1)^2 \\ &= c(n^2 + 1 - 2n) = cn^2 + c - 2cn \\ T(n) &\geq cn^2 + c - 2cn + n \quad (\text{when } c=1/2, \text{ R.H.S.} \geq cn^2) \\ &\geq cn^2 \quad (\text{for } c=1/2 \text{ and } n \geq 1) \end{aligned}$$

Base case:

## Substitution method : $T(n) = T(n-1) + n$

- ▶ Find a lower bound for the recurrence  $T(n) = T(n-1) + n$
- ▶ What would be a good guess?  $T(n) = \Omega(n^2)$

To show :  $T(n) \geq cn^2 \quad \forall n \geq n_0$

Induction step:

$$\begin{aligned} T(n-1) &\geq c(n-1)^2 \\ &= c(n^2 + 1 - 2n) = cn^2 + c - 2cn \\ T(n) &\geq cn^2 + c - 2cn + n \quad (\text{when } c=1/2, \text{ R.H.S.} \geq cn^2) \\ &\geq cn^2 \quad (\text{for } c=1/2 \text{ and } n \geq 1) \end{aligned}$$

Base case:

$$T(1) \geq cn^2$$

# Solving recurrences : Recursion tree

- ▶ Recursion tree can be used to generate a good guess for the Substitution method.




# Solving recurrences : Recursion tree

- ▶ Recursion tree can be used to generate a good guess for the Substitution method.
- ▶ Finding a good guess for  $T(n) = 3T(\lfloor n/4 \rfloor) + \Theta(n^2)$


# Solving recurrences : Recursion tree

- ▶ Recursion tree can be used to generate a good guess for the Substitution method.
- ▶ Finding a good guess for  $T(n) = 3T(\lfloor n/4 \rfloor) + \Theta(n^2)$
- ▶ We will ignore the floors and ceilings, because usually they don't affect the asymptotic order of growth.


Recursion tree :  $T(n) = 3T(n/4) + cn^2$



Recursion tree :  $T(n) = 3T(n/4) + cn^2$



# Recursion tree : $T(n) = 3T(n/4) + cn^2$



(d)

Total:  $O(n^2)$

# Recursion tree : $T(n) = 3T(n/4) + cn^2$

- ▶ Number of levels :

$$\frac{n}{4^{i-1}} = 1$$

# Recursion tree : $T(n) = 3T(n/4) + cn^2$

- ▶ Number of levels :

$$\frac{n}{4^{i-1}} = 1$$

$$i = \log_4 n + 1$$

# Recursion tree : $T(n) = 3T(n/4) + cn^2$

- ▶ Number of levels :

$$\frac{n}{4^{i-1}} = 1$$
$$i = \log_4 n + 1$$

- ▶ Running time for the non-leaf levels:

$$cn^2 \left( 1 + \frac{3}{16} + \left( \frac{3}{16} \right)^2 + \dots \right) (\log_4 n \text{ terms})$$



# Recursion tree : $T(n) = 3T(n/4) + cn^2$

- ▶ Number of levels :

$$\frac{n}{4^{i-1}} = 1$$
$$i = \log_4 n + 1$$

- ▶ Running time for the non-leaf levels:

$$cn^2 \left( 1 + \frac{3}{16} + \left( \frac{3}{16} \right)^2 + \dots \right) (\log_4 n \text{ terms})$$
$$< cn^2 \left( 1 + \frac{3}{16} + \left( \frac{3}{16} \right)^2 + \dots \right) (\text{infinite series sum})$$

# Recursion tree : $T(n) = 3T(n/4) + cn^2$

- ▶ Number of levels :

$$\frac{n}{4^{i-1}} = 1$$
$$i = \log_4 n + 1$$

- ▶ Running time for the non-leaf levels:

$$cn^2 \left( 1 + \frac{3}{16} + \left( \frac{3}{16} \right)^2 + \dots \right) (\log_4 n \text{ terms})$$
$$< cn^2 \left( 1 + \frac{3}{16} + \left( \frac{3}{16} \right)^2 + \dots \right) (\text{infinite series sum})$$
$$= \frac{16}{13} cn^2$$

# Recursion tree : $T(n) = 3T(n/4) + cn^2$

- ▶ Number of levels :

$$\frac{n}{4^{i-1}} = 1$$
$$i = \log_4 n + 1$$

- ▶ Running time for the non-leaf levels:

$$cn^2 \left( 1 + \frac{3}{16} + \left( \frac{3}{16} \right)^2 + \dots \right) (\log_4 n \text{ terms})$$
$$< cn^2 \left( 1 + \frac{3}{16} + \left( \frac{3}{16} \right)^2 + \dots \right) (\text{infinite series sum})$$
$$= \frac{16}{13} cn^2 = O(n^2)$$

# Recursion tree : $T(n) = 3T(n/4) + cn^2$

- ▶ Running time for leaf level nodes :

$$3^{\log_4 n} = n^{\log_4 3}$$

# Recursion tree : $T(n) = 3T(n/4) + cn^2$

- ▶ Running time for leaf level nodes :

$$3^{\log_4 n} = n^{\log_4 3}$$

- ▶ Total running time

# Recursion tree : $T(n) = 3T(n/4) + cn^2$

- ▶ Running time for leaf level nodes :

$$3^{\log_4 n} = n^{\log_4 3}$$

- ▶ Total running time

$$= O(n^2) + \Theta(n^{\log_4 3})$$

# Recursion tree : $T(n) = 3T(n/4) + cn^2$


- ▶ Running time for leaf level nodes :

$$3^{\log_4 n} = n^{\log_4 3}$$

- ▶ Total running time

$$\begin{aligned} &= O(n^2) + \Theta(n^{\log_4 3}) \\ &= O(n^2) \end{aligned}$$

# Recursion tree : $T(n) = 3T(n/4) + cn^2$



(d)

Total:  $O(n^2)$



## Recursion tree : $T(n) = 3T(n/4) + cn^2$

- By looking at the recursion tree we can guess  $T(n) = O(n^2)$


# Recursion tree : $T(n) = 3T(n/4) + cn^2$

- ▶ By looking at the recursion tree we can guess  $T(n) = O(n^2)$
- ▶ Then we can use the substitution method to show that our guess is correct.

# Recursion tree : $T(n) = 3T(n/4) + cn^2$

- ▶ By looking at the recursion tree we can guess  $T(n) = O(n^2)$
- ▶ Then we can use the substitution method to show that our guess is correct.
- ▶ To prove :  $T(n) \leq dn^2 \quad \forall n \geq n_0$

# Recursion tree : $T(n) = T(n/3) + T(2n/3) + O(n)$



Recursion tree :  $T(n) = T(n/3) + T(2n/3) + O(n)$

► Number of levels :

$$\frac{n}{\left(\frac{3}{2}\right)^{i-1}} = 1$$

# Recursion tree : $T(n) = T(n/3) + T(2n/3) + O(n)$

- ▶ Number of levels :

$$\frac{n}{\left(\frac{3}{2}\right)^{i-1}} = 1$$
$$i = \log_{\frac{3}{2}} n + 1$$

# Recursion tree : $T(n) = T(n/3) + T(2n/3) + O(n)$

- ▶ Number of levels :

$$\frac{n}{\left(\frac{3}{2}\right)^{i-1}} = 1$$
$$i = \log_{\frac{3}{2}} n + 1$$

- ▶ Running time :

$$T(n) = O(n \log_{\frac{3}{2}} n)$$

# Recursion tree : $T(n) = T(n/3) + T(2n/3) + O(n)$

- ▶ Number of levels :

$$\frac{n}{\left(\frac{3}{2}\right)^{i-1}} = 1$$
$$i = \log_{\frac{3}{2}} n + 1$$

- ▶ Running time :

$$T(n) = O(n \log_{\frac{3}{2}} n)$$

- ▶ Is  $\Theta(\log_{10} n) = \Theta(\log_3 n)$ ?



# Recursion tree : $T(n) = T(n/3) + T(2n/3) + O(n)$

- ▶ Number of levels :

$$\frac{n}{\left(\frac{3}{2}\right)^{i-1}} = 1$$
$$i = \log_{\frac{3}{2}} n + 1$$

- ▶ Running time :

$$T(n) = O(n \log_{\frac{3}{2}} n)$$

- ▶ Is  $\Theta(\log_{10} n) = \Theta(\log_3 n)$ ?
- ▶ Running time :

$$T(n) = O(n \log_{\frac{3}{2}} n) = O(n \lg n)$$

# Recursion tree : $T(n) = T(n/3) + T(2n/3) + O(n)$

- ▶ Number of levels :

$$\frac{n}{\left(\frac{3}{2}\right)^{i-1}} = 1$$
$$i = \log_{\frac{3}{2}} n + 1$$

- ▶ Running time :

$$T(n) = O(n \log_{\frac{3}{2}} n)$$

- ▶ Is  $\Theta(\log_{10} n) = \Theta(\log_3 n)$ ?
- ▶ Running time :

$$T(n) = O(n \log_{\frac{3}{2}} n) = O(n \lg n)$$

- ▶ We can again use the substitution method to show the upper bound.

# Polynomially larger/smaller

- For any positive  $\epsilon$  (however small), is  $n^\epsilon = \omega(\lg n)$  ?

# Polynomially larger/smaller

- ▶ For any positive  $\epsilon$  (however small), is  $n^\epsilon = \omega(\lg n)$  ?
- ▶ Using L'Hospital's rule:

$$\lim_{n \rightarrow \infty} \frac{\lg n}{n^\epsilon} = 0$$

# Polynomially larger/smaller

- ▶ For any positive  $\epsilon$  (however small), is  $n^\epsilon = \omega(\lg n)$  ?
- ▶ Using L'Hospital's rule:

$$\lim_{n \rightarrow \infty} \frac{\lg n}{n^\epsilon} = 0$$

- ▶ Is  $n^{1+\epsilon} = \omega(n \lg n)$ ?

# Polynomially larger/smaller

►  $n = o(n^2)$

# Polynomially larger/smaller

- ▶  $n = o(n^2)$  ( $n$  is asymptotically smaller)

# Polynomially larger/smaller

- ▶  $n = o(n^2)$  ( $n$  is asymptotically smaller)
- ▶  $n^{1+\epsilon} = o(n^2)$



# Polynomially larger/smaller

- ▶  $n = o(n^2)$  ( $n$  is asymptotically smaller)
- ▶  $n^{1+\epsilon} = o(n^2)$  ( $n$  is polynomially smaller because we can find a positive constant  $\epsilon$ )

# Polynomially larger/smaller

- ▶  $n = o(n^2)$  ( $n$  is asymptotically smaller)
- ▶  $n^{1+\epsilon} = o(n^2)$  ( $n$  is polynomially smaller because we can find a positive constant  $\epsilon$ )
- ▶  $n = o(n \lg n)$

# Polynomially larger/smaller

- ▶  $n = o(n^2)$  ( $n$  is asymptotically smaller)
- ▶  $n^{1+\epsilon} = o(n^2)$  ( $n$  is polynomially smaller because we can find a positive constant  $\epsilon$ )
- ▶  $n = o(n \lg n)$  ( $n$  is asymptotically smaller)


# Polynomially larger/smaller

- ▶  $n = o(n^2)$  ( $n$  is asymptotically smaller)
- ▶  $n^{1+\epsilon} = o(n^2)$  ( $n$  is polynomially smaller because we can find a positive constant  $\epsilon$ )
- ▶  $n = o(n \lg n)$  ( $n$  is asymptotically smaller)
- ▶ However,  $n^{1+\epsilon} \neq o(n \lg n)$

# Polynomially larger/smaller

- ▶  $n = o(n^2)$  ( $n$  is asymptotically smaller)
- ▶  $n^{1+\epsilon} = o(n^2)$  ( $n$  is polynomially smaller because we can find a positive constant  $\epsilon$ )
- ▶  $n = o(n \lg n)$  ( $n$  is asymptotically smaller)
- ▶ However,  $n^{1+\epsilon} \neq o(n \lg n)$  ( $n$  is **not** polynomially smaller)

# Master method : $T(n) = 3T(n/4) + cn^2$



(d)

Total:  $O(n^2)$

# Solving recurrences : Master method

## **Theorem 4.1 (Master theorem)**

Let  $a \geq 1$  and  $b > 1$  be constants, let  $f(n)$  be a function, and let  $T(n)$  be defined on the nonnegative integers by the recurrence

$$T(n) = aT(n/b) + f(n),$$

where we interpret  $n/b$  to mean either  $\lfloor n/b \rfloor$  or  $\lceil n/b \rceil$ . Then  $T(n)$  has the following asymptotic bounds:

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■



$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 9T(n/3) + n$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 9T(n/3) + n$

$$n^{\log_b a} = n^{\log_3 9}$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 9T(n/3) + n$

$$n^{\log_b a} = n^{\log_3 9}$$

$$f(n) = n =$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 9T(n/3) + n$

$$n^{\log_b a} = n^{\log_3 9}$$

$$f(n) = n = O(n^{2-\epsilon})$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 9T(n/3) + n$

$$n^{\log_b a} = n^{\log_3 9}$$

$$f(n) = n = O(n^{2-\epsilon})$$

$$T(n) = \Theta(n^2)$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = T(2n/3) + 1$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = T(2n/3) + 1$

$$n^{\log_b a} = n^{\log_{\frac{3}{2}} 1}$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = T(2n/3) + 1$

$$n^{\log_b a} = n^{\log_{\frac{3}{2}} 1} = n^0$$



$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = T(2n/3) + 1$

$$n^{\log_b a} = n^{\log_{\frac{3}{2}} 1} = n^0$$

$$f(n) = \Theta(1)$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = T(2n/3) + 1$

$$n^{\log_b a} = n^{\log_{\frac{3}{2}} 1} = n^0$$

$$f(n) = \Theta(1) = \Theta(n^{\log_b a})$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = T(2n/3) + 1$

$$n^{\log_b a} = n^{\log_{\frac{3}{2}} 1} = n^0$$

$$f(n) = \Theta(1) = \Theta(n^{\log_b a})$$

$$T(n) = \Theta(n^{\log_b a} \lg n)$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = T(2n/3) + 1$

$$n^{\log_b a} = n^{\log_{\frac{3}{2}} 1} = n^0$$

$$f(n) = \Theta(1) = \Theta(n^{\log_b a})$$

$$\begin{aligned} T(n) &= \Theta(n^{\log_b a} \lg n) \\ &= \Theta(\lg n) \end{aligned}$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 3T(n/4) + n \lg n$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 3T(n/4) + n \lg n$

$$n^{\log_b a} = n^{\log_4 3}$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 3T(n/4) + n \lg n$

$$n^{\log_b a} = n^{\log_4 3}$$

$$f(n) = n \lg n$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 3T(n/4) + n \lg n$

$$n^{\log_b a} = n^{\log_4 3}$$

$$f(n) = n \lg n = \Omega(n^{\log_4 3 + \epsilon}) ?$$



$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 3T(n/4) + n \lg n$

$$n^{\log_b a} = n^{\log_4 3}$$

$$f(n) = n \lg n = \Omega(n^{\log_4 3 + \epsilon}) ?$$

$$a(n/b) \lg(n/b) \leq cn \lg n$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 3T(n/4) + n \lg n$

$$n^{\log_b a} = n^{\log_4 3}$$

$$f(n) = n \lg n = \Omega(n^{\log_4 3 + \epsilon}) ?$$

$$a(n/b) \lg(n/b) \leq cn \lg n$$

$$3(n/4) \lg(n/4) \leq (3/4)n \lg n, \text{ for } c = 3/4$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 3T(n/4) + n \lg n$

$$n^{\log_b a} = n^{\log_4 3}$$

$$f(n) = n \lg n = \Omega(n^{\log_4 3 + \epsilon}) ?$$

$$a(n/b) \lg(n/b) \leq cn \lg n$$

$$3(n/4) \lg(n/4) \leq (3/4)n \lg n, \text{ for } c = 3/4$$

$$T(n) = \Theta(n \lg n)$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 2T(n/2) + n \lg n$

$$n^{\log_b a} = n^{\log_2 2} = n$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 2T(n/2) + n \lg n$

$$n^{\log_b a} = n^{\log_2 2} = n$$

$$f(n) = n \lg n$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 2T(n/2) + n \lg n$

$$n^{\log_b a} = n^{\log_2 2} = n$$

$$f(n) = n \lg n = \Omega(n^{1+\epsilon}) ?$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 2T(n/2) + n \lg n$

$$n^{\log_b a} = n^{\log_2 2} = n$$

$$f(n) = n \lg n = \Omega(n^{1+\epsilon}) ?$$

Master method cannot be applied.

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 7T(n/2) + \Theta(n^2)$

$$n^{\log_b a} = n^{\lg 7} \approx n^{2.81}$$



$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 7T(n/2) + \Theta(n^2)$

$$n^{\log_b a} = n^{\lg 7} \approx n^{2.81}$$

$$f(n) = \Theta(n^2)$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 7T(n/2) + \Theta(n^2)$

$$n^{\log_b a} = n^{\lg 7} \approx n^{2.81}$$

$$f(n) = \Theta(n^2) = O(n^{\log_b a - \epsilon})$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 7T(n/2) + \Theta(n^2)$

$$n^{\log_b a} = n^{\lg 7} \approx n^{2.81}$$

$$\begin{aligned} f(n) &= \Theta(n^2) = O(n^{\log_b a - \epsilon}) \\ &= O(n^{\lg 7 - \epsilon}) ? \end{aligned}$$

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 7T(n/2) + \Theta(n^2)$

$$n^{\log_b a} = n^{\lg 7} \approx n^{2.81}$$

$$\begin{aligned} f(n) &= \Theta(n^2) = O(n^{\log_b a - \epsilon}) \\ &= O(n^{\lg 7 - \epsilon}) ? \end{aligned}$$

$$T(n) = \Theta(n^{\log_b a})$$

$$T(n) = aT(n/b) + f(n) , a \geq 1, b > 1$$

1. If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$ .
3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$ . ■

►  $T(n) = 7T(n/2) + \Theta(n^2)$

$$n^{\log_b a} = n^{\lg 7} \approx n^{2.81}$$

$$\begin{aligned} f(n) &= \Theta(n^2) = O(n^{\log_b a - \epsilon}) \\ &= O(n^{\lg 7 - \epsilon}) ? \end{aligned}$$

$$\begin{aligned} T(n) &= \Theta(n^{\log_b a}) \\ &= \Theta(n^{\lg 7}) \end{aligned}$$

# Proof of the master theorem

- ▶ Section 4.6 will not be a part of our syllabus

# Part II Sorting and Order Statistics

- ▶ Record : Collection of data

# Part II Sorting and Order Statistics

- ▶ Record : Collection of data
- ▶ Key : Value to be sorted



## Part II Sorting and Order Statistics

- ▶ Record : Collection of data
- ▶ Key : Value to be sorted
- ▶ Satellite data

## Part II Sorting and Order Statistics

- ▶ Record : Collection of data
- ▶ Key : Value to be sorted
- ▶ Satellite data
- ▶ If satellite data is large, we permute an array of pointers to the records.

# Sorting algorithms

- ▶ *In place* sorting : If at any time only a constant number of elements are stored outside the array.

# Sorting algorithms

- ▶ *In place* sorting : If at any time only a constant number of elements are stored outside the array.
- ▶ MERGE procedure does not operate in place.

# Sorting algorithms

- ▶ *In place* sorting : If at any time only a constant number of elements are stored outside the array.
- ▶ MERGE procedure does not operate in place.
- ▶ Ch 6 : Heapsort that uses a data structure called heap.

# Sorting algorithms

- ▶ *In place* sorting : If at any time only a constant number of elements are stored outside the array.
- ▶ MERGE procedure does not operate in place.
- ▶ Ch 6 : Heapsort that uses a data structure called heap.
- ▶ Heapsort sorts  $n$  elements *in place* in  $O(n \lg n)$  time.

# Sorting algorithms

Algorithm	Worst-case running time	Average-case/expected running time
Insertion sort	$\Theta(n^2)$	$\Theta(n^2)$
Merge sort	$\Theta(n \lg n)$	$\Theta(n \lg n)$
Heapsort	$O(n \lg n)$	—
Quicksort	$\Theta(n^2)$	$\Theta(n \lg n)$ (expected)
Counting sort	$\Theta(k + n)$	$\Theta(k + n)$
Radix sort	$\Theta(d(n + k))$	$\Theta(d(n + k))$
Bucket sort	$\Theta(n^2)$	$\Theta(n)$ (average-case)

# Ch 6 : Heapsort

- ▶ Heapsort uses the Heap data structure.



# Ch 6 : Heapsort

- ▶ Heapsort uses the Heap data structure.
- ▶ To understand heap, we need to understand a few terminologies.

# Binary tree

- ▶ Binary tree is a tree data structure where each node can have at most two child nodes : left child node and right child node.

# Binary tree

- ▶ Binary tree is a tree data structure where each node can have at most two child nodes : left child node and right child node.

