

CONTROL ENGINEERING

1. Two types of algae evolve, in competition, in an aqueous solution. The equations describing the evolution of the two populations of algae are

$$\dot{x}_1 = x_1 \left(-x_1 + \frac{u}{1+x_2} \right), \quad \dot{x}_2 = x_2(-x_2 + u),$$

where x_1 denotes the concentration of the first type, x_2 the concentration of the second type, and u the concentration of nutrient.

- Assume $u > 0$ and constant. Determine all equilibrium points of the system. [4 marks]
- Write the linearized models of the system around each of the equilibrium points determined in part a). [8 marks]
- Using the linearized models determined in part b) determine (if possible) the stability properties of the equilibrium points computed in part a). [4 marks]
- Show that all linearized models determined in part b) are not controllable. [4 marks]

2. A cart of mass $M = 1$ has two inverted pendulums attached to it of lengths l_1 and l_2 and both of mass m . Let θ_1 and θ_2 be the angles of the pendulums with respect to a vertical axis directed upward and let f be the force on the cart.

For small values of θ_1 and θ_2 the dynamic behaviour of the pendulums is described by the differential equations

$$m(f - mg\theta_1 - mg\theta_2 + l_1\ddot{\theta}_1) = mg\theta_1, \quad m(f - mg\theta_1 - mg\theta_2 + l_2\ddot{\theta}_2) = mg\theta_2,$$

where g denotes the gravitational acceleration.

- Let $x_1 = \theta_1$, $x_2 = \theta_2$, $x_3 = \dot{\theta}_1$, $x_4 = \dot{\theta}_2$, $u = f$, $y = x_1 - x_2$ and $x = [x_1 \ x_2 \ x_3 \ x_4]'$. Write a state space representation of the considered system, i.e. determine matrices A , B and C such that

$$\dot{x} = Ax + Bu \quad y = Cx.$$

[4 marks]

- Study the controllability property of the system as a function of the physical parameters l_1 and l_2 . [6 marks]
- Study the observability property of the system as a function of the physical parameters l_1 and l_2 . [6 marks]
- Assume $l_1 = l_2$ and write a second order differential equation describing the behaviour of $\xi = \theta_1 - \theta_2$. Use this differential equation to assess the stabilizability property of the system. [4 marks]

3. Consider a herd of cattle composed of cows and calves. Let $x_1(t)$ be the number of calves in year t and $x_2(t)$ the number of cows in year t . The dynamical behaviour of the herd is described by the equation

$$x(t+1) = Ax(t) = \begin{bmatrix} \frac{1}{2} & \frac{2}{5} \\ \frac{1-k}{2} & \frac{4}{5} \end{bmatrix} x(t),$$

where $x(t) = [x_1(t), x_2(t)]'$ and $k \in [0, 1]$ denotes the portion of calves slaughtered each year.

- a) Compute the equilibrium points of the system as a function of $k \in [0, 1]$.
[4 marks]
- b) Determine for which values of k the system is stable, asymptotically stable, unstable.
[4 marks]
- c) Show that for any initial condition $x(0)$ such that $x_1(0) \geq 0$ and $x_2(0) \geq 0$, the free response $x(t)$ of the system is such that $x_1(t) \geq 0$ and $x_2(t) \geq 0$, for all $t \geq 0$.
[4 marks]
- d) Assume $k = 1/2$.

- i) Show that the free-response of the system converges to the line

$$5x_1 - 4x_2 = 0.$$


(Hint: write a difference equation for the variable $z(t) = 5x_1(t) - 4x_2(t)$ and show that $z(t)$ tends to zero as t tends to ∞ .)
[4 marks]

- ii) Suppose that for each slaughtered calf C_1 GBP are earned and that each cow costs C_2 GBP a year. The *revenue* of the herd in the year t is therefore

$$y(t) = C_1 k x_1(t) - C_2 x_2(t).$$

Determine a condition on C_1 and C_2 so that the asymptotic revenue is non-negative for each $x_1(0) \geq 0$ and $x_2(0) \geq 0$.
[4 marks]

4. The chemical reaction describing the production of water, namely



can be described by the nonlinear continuous-time system

$$\begin{aligned}\dot{H} &= -2k_1H^2O + 2k_2W, \\ \dot{O} &= -k_1H^2O + k_2W, \\ \dot{W} &= -2k_2W + 2k_1H^2O,\end{aligned}$$

where $H \geq 0$, $O \geq 0$ and $W \geq 0$ denote the concentrations of hydrogen, oxygen, and water, respectively, and $k_1 > 0$ and $k_2 > 0$ are positive constants which quantify the speed of the reaction.

To study the dynamical properties of the system consider the variables

$$x_1 = W, \quad x_2 = W + 2O, \quad x_3 = W + H.$$

- Show that the variables (x_1, x_2, x_3) define a new set of coordinates for the system and determine (H, O, W) as a function of (x_1, x_2, x_3) .
(Hint: show that there is a one-to-one relation between the variables (H, O, W) and the variables (x_1, x_2, x_3) .) [4 marks]
- Write differential equations for x_2 and x_3 . Integrate the resulting differential equations and comment on the results. [4 marks]
- Write a differential equation for x_1 and show that \dot{x}_1 can be written as a cubic polynomial in x_1 with coefficients that depend upon $x_2(0)$, $x_3(0)$, k_1 and k_2 . In particular, show that

$$\dot{x}_1 = A - Bx_1 + Cx_1^2 - Dx_1^3, \quad (\star)$$

where A , B , C and D are functions of $x_2(0)$, $x_3(0)$, k_1 and k_2 and take non-negative values. [4 marks]

- Suppose that for all $x_2(0) > 0$ and $x_3(0) > 0$ the system (\star) has only one equilibrium x_1^* .
 - Sketch \dot{x}_1 as a function of x_1 and argue that the equilibrium $x_1 = x_1^*$ is a globally asymptotically stable equilibrium for the x_1 -system. [4 marks]
 - Argue that the overall system with state (x_1, x_2, x_3) has infinitely many equilibria. Using the results of part d.i) determine the stability properties of these equilibria. [4 marks]

5. Consider a linear, time-varying, continuous-time system described by the equation

$$\dot{x} = A(t)x.$$

A common *belief* is the following.

(C) If the matrix $A(t)$ has constant eigenvalues with negative real part then the linear, time-varying, system is asymptotically stable.

To disprove the claim (C) consider the matrix

$$A(t) = \begin{bmatrix} -1 & e^{2t} \\ 0 & -1 \end{bmatrix}.$$

Let $t_0 = 0$.

- a) Show that the matrix $A(t)$ has constant eigenvalues with negative real part. [2 marks]

- b) Determine the state transition matrix of the system, i.e. the matrix $\Phi(t, 0)$ such that

$$\Phi(0, 0) = I, \quad \frac{d\Phi(t, 0)}{dt} = A(t)\Phi(t, 0).$$

(Hint: integrate the differential equations describing the system.) [8 marks]

- c) Show that for almost any selection of the initial conditions $x(0)$

$$\lim_{t \rightarrow \infty} \|x(t)\| = \infty.$$

Determine the set of initial conditions such that

$$\lim_{t \rightarrow \infty} \|x(t)\| = 0.$$

[4 marks]

- d) Using the results in part c) conclude that the considered linear, time-varying system, is not stable. [2 marks]

- e) Show that the linear, time-varying, system

$$\dot{x} = B(t)x,$$

with

$$B(t) = \begin{bmatrix} -1 & b(t) \\ 0 & -1 \end{bmatrix}$$

and $|b(t)| \leq \bar{b}$, for some \bar{b} positive, is asymptotically stable. [4 marks]

6. Consider a linear, single-input, single-output, system described by the equations

$$\sigma x = Ax + Bu, \quad y = Cx,$$

where $x(t) \in \mathbb{R}^n$ is the state, $u(t) \in \mathbb{R}$ is the input, and $y(t) \in \mathbb{R}$ is the output.

Consider the problem of studying the reachability and observability properties of the system using the PBH tests.

- a) Show, using the PBH reachability test, that the system is reachable if and only if there is no left eigenvector of A which is orthogonal to B .
(Hint: recall that a left eigenvector of A is a row vector w such that $wA = \lambda w$, for some λ which is an eigenvalue of A .) [4 marks]
- b) Show, using the PBH observability test, that the system is observable if and only if there is no right eigenvector of A which is orthogonal to C .
(Hint: recall that a right eigenvector of A is a column vector v such that $Av = \lambda v$, for some λ which is an eigenvalue of A .) [4 marks]
- c) Consider the class of linear systems described by the equations

$$\begin{aligned} \sigma x_1 &= \lambda_1 x_1 + B_1 u, \\ \sigma x_2 &= \lambda_2 x_2 + B_2 u, \\ &\vdots \\ \sigma x_n &= \lambda_n x_n + B_n u, \\ y &= C_1 x_1 + C_2 x_2 + \cdots + C_n x_n, \end{aligned}$$

with $\lambda_i \neq \lambda_j$ for $i \neq j$.

- i) Using the results in part a) determine conditions on the coefficients B_i such that the system is reachable. [4 marks]
- ii) Using the results in part b) determine conditions on the coefficients C_i such that the system is observable. [2 marks]
- d) Let

$$A = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 1 \\ 0 & 0 & \lambda \end{bmatrix}, \quad B = \begin{bmatrix} B_1 \\ B_2 \\ B_3 \end{bmatrix}.$$

Show, using the results in part a), that the system is not reachable (regardless of the values of the coefficients B_1, B_2 and B_3). [6 marks]