Central Nervous System Stabilization & Pre-Surgical Optimization Protocol

Interim Pre-Hab, Neuro-Inhibition & Sensory De-Escalation Roadmap

Comprehensive Patient Strategic Report

A Strategic Integrative Care Blueprint for Stiff Person Syndrome (SPS) and Joint Interventions

CLINICAL CONTEXT:

Condition Profile: Stiff Person Syndrome (SPS) | Upcoming Interventions: Knee, Shoulder, Ankle (12 Mo)

Document Date: 6 November 2026 | Prepared by: Personal Care & Health Strategy Advisory

Classification: Personalized Comprehensive Advisory Protocol

Contents

1. Executive Summary & Neurological Architecture . . . 3

Multi-System Coordination Matrix . . . 3

2. Low-Cost Daily Water Mineralization Blueprint . . . 4

Electrolyte Concentration Specifications (Per 1.0 Litre Water) . . . 4

Electrolyte Balancing Guidelines . . . 4

3. Low-Excitotoxin Nutritional Protocol (Zero Nuts, Zero Legumes) . . . 5

Excitotoxin Elimination Rules . . . 5

Structured Food Group Selections . . . 5

Daily Operational Meal Architecture . . . 5

4. Precision Micronutrient & Neuro-Protective Stack . . . 6

Phase 1: Neurological Inhibition & Anti-Spasm Agents . . . 6

Phase 2: Pre-Surgical Tissue & Systemic Support . . . 6

5. Sensory De-Escalation & Vagal Nerve Retraining . . . 7

Sensory Environmental Hygiene . . . 7

Vagus Nerve & Autonomic Down-Regulation Practices . . . 7

6. Restorative Sleep Architecture & Joint Ergonomics . . . 7

7. Joint-Spared Isometric 'Pre-Hab' Program . . . 8

Prescribed Daily Low-Impact Movements . . . 8

8. 12-Month Action Roadmap & Pre-Operative Checkpoints . . . 8

1. Executive Summary & Neurological Architecture

This strategic protocol provides a coordinated, proactive roadmap to stabilize your central nervous system (CNS) over the next 12 months in preparation for Stiff Person Syndrome (SPS) therapies and multi-joint orthopedic surgeries (knee, shoulder, and ankle).

In Stiff Person Syndrome, autoimmune-mediated inhibition of glutamic acid decarboxylase (GAD) impairs the synthesis of gamma-aminobutyric acid (GABA), the nervous system's chief inhibitory brake. Consequently, upper and lower motor pathways enter an unabating state of excitability. Any peripheral pain, sudden auditory or tactile stimulus, emotional tension, or physical instability cascades into debilitating muscle spasms and rigidity.

> **Clinical Mission**
> **Primary Objective:** Systematically suppress central excitotoxins (glutamate, aspartate, high sensory load) while aggressively reinforcing inhibitory GABAergic pathways, stabilizing sleep architecture, and preserving connective tissue and joint stability without aggravating damaged articular structures.

Multi-System Coordination Matrix

Managing four concurrent interventions requires tight synchronization across distinct metabolic, mechanical, and neurological domains:

- **Neuromuscular Tone:** Maximize GABA receptor activation and NMDA receptor channel blocking to elevate spasm thresholds.
- **Sensory Gating:** Prevent sympathetic autonomic surges triggered by sudden environmental inputs.
- **Nutritional Tissue Pre-Hab:** Fuel collagen synthesis and bone remodelling for joint repair without relying on nuts or legumes.
- **Electrolyte Hydration:** Maintain physiological serum sodium, potassium, and magnesium for smooth nerve firing without commercial expense.

2. Low-Cost Daily Water Mineralization Blueprint

Proper cellular hydration is critical to prevent peripheral muscle cramps from triggering generalized central spasms. Commercial electrolyte packets cost $45 to $70 monthly. You can prepare an identical, medical-grade formulation for pennies per day using pharmaceutical/food-grade bulk salts.

Electrolyte Concentration Specifications (Per 1.0 Litre Water)

Component	Commercial Grade Source	Quantity / Litre	Elemental Yield	Physiological Function
Sodium	Unrefined Grey Sea Salt or Pink Himalayan Salt	1/4 tsp (~1.5 g)	~580 mg Na	Sustains blood volume & neuromuscular membrane potential
Potassium	Potassium Chloride (NoSalt / Nu-Salt / Lite Salt)	1/8 tsp (~0.75 g)	~360 mg K	Direct cellular antagonist to resting muscular contraction
Magnesium	Magnesium Malate or Bisglycinate powder	1/4 tsp (~1.0 g)	~120 mg Mg	Endogenous NMDA receptor blocker & muscle relaxant
Alkalinizing Citrate	Fresh Squeezed Lemon or Lime Juice	15 mL (1/2 fruit)	Trace organic acids	Buffers gastric acidity and optimizes cation absorption

Table 1. Standard low-cost daily hydration and mineralization formula

Practical Mixing Tip
Batch Preparation Protocol: In a 2-litre glass pitcher, combine 1/2 tsp sea salt and 1/4 tsp potassium chloride with the juice of one fresh lemon. Keep in refrigerator. Consume across the day between meals. Keep high-dose magnesium capsules separate (at night) to avoid gastrointestinal acceleration or bitter taste.

Electrolyte Balancing Guidelines

Electrolyte equilibrium is critical for neuromuscular stability in SPS:

- **Sodium Timing:** Spread sodium intake across the daytime hours. Avoid large boluses right before lying down to prevent fluid shifts.
- **Potassium Safety:** Never exceed 1/2 tsp total potassium chloride daily unless guided by serum electrolyte blood panels (potassium is strictly regulated by renal clearance).
- **Temperature:** Consume water at room temperature or mildly warm. Ice-cold fluids can trigger visceral startle reflexes and esophageal spasms in sensitive individuals.

3. Low-Excitotoxin Nutritional Protocol (Zero Nuts, Zero Legumes)

Because you do not tolerate or enjoy nuts and legumes, standard plant-based anti-inflammatory regimens fail. The target is an anti-inflammatory, low-histamine, low-excitotoxin nutritional matrix emphasizing easily assimilated animal proteins, clean complex tubers, and nutrient-dense fruit oils.

Excitotoxin Elimination Rules

- **Free & Hydrolyzed Glutamates:** Strictly eliminate MSG (E621), autolyzed yeast extract, bouillon stock cubes, textured vegetable protein, and processed savoury chips.
- **Aged & High-Histamine Foods:** Remove aged hard cheeses, cured/charcuterie meats (salami, pepperoni), and long-fermented condiments. Histamine stimulates central wakefulness and sympathetic arousal.
- **Excitatory Sweeteners & Stimulants:** Zero aspartame (breaks down to excitatory aspartic acid), zero energy drinks. Limit caffeine to a single morning beverage or switch to organic decaf/herbal infusions.

Structured Food Group Selections

Food Category	Approved Staples (Nut & Legume Free)	Target Benefits
Primary Proteins	Pastured whole eggs, fresh chicken/turkey breast, wild salmon, cod, grass-fed beef, lamb	Tryptophan for serotonin/melatonin, high bioavailable heme iron, zero inflammatory lectins
Connective Peptides	Fresh homemade beef/poultry bone broth (simmered 3-4 hours), hydrolyzed collagen	Supplies glycine and proline directly into damaged knee, shoulder, and ankle capsules
Starch & Complex Carbs	Sweet potatoes, yams, white/jasmine rice, rolled oats, butternut squash, carrots, beets	Maintains stable glycogen without the digestive distress or anti-nutrients of legumes
Healthy Dietary Fats	Cold-pressed extra virgin olive oil, fresh avocados, avocado oil, grass-fed ghee, pure coconut oil	Replaces nut fats with stable monounsaturated and medium-chain fatty acids
Neuro-Protective Greens	Steamed spinach, Swiss chard, bok choy, zucchini, asparagus, broccoli florets	Abundant magnesium, sulforaphane, folate, and potassium with high digestive tolerance

Table 2. Approved nutrient categories replacing nuts and legumes

Daily Operational Meal Architecture

- **Breakfast (07:30 - 08:30):** 3 pasture-raised scrambled eggs cooked in grass-fed ghee, 1 cup gently wilted baby spinach, 1/2 roasted sweet potato seasoned with pink salt. Accompanied by warm chamomile or rooibos tea.
- **Lunch (12:30 - 13:30):** 150g grilled wild salmon or baked chicken thighs, 1 cup cooked jasmine rice or boiled baby potatoes, steamed zucchini and roasted carrots with 1 tbsp unheated extra virgin olive oil.
- **Afternoon Re-Fuel (16:00):** 1 sliced avocado sprinkled with sea salt and lemon juice, or fresh blueberries with 10g pure collagen stirred into warm herbal tea.
- **Dinner (18:30 - 19:30):** 150g slow-cooked tender beef chuck or turkey mince, roasted butternut squash, steamed broccoli florets, and 1 warm cup of fresh bone broth.

4. Precision Micronutrient & Neuro-Protective Stack

The following evidence-based protocol directly addresses both SPS GABAergic tone and connective tissue prep for your knee, shoulder, and ankle surgeries.

Phase 1: Neurological Inhibition & Anti-Spasm Agents

Active Compound	Optimal Dose	Administration Timing	Clinical Rationale & Mechanism
Magnesium Bisglycinate	350 - 400 mg	45 min before sleep	Glycine acts as an inhibitory neurotransmitter; magnesium blunts NMDA receptor tone.
Magnesium L-Threonate	144 mg (elemental)	Morning with meal	Proven blood-brain barrier penetration; reduces central neuro-inflammation.
L-Theanine	100 - 200 mg	Mid-morning & 15:00	Induces alpha brain wave activity; competes with glutamate at excitatory receptors.
Myo-Inositol	2,000 - 3,000 mg	Evening with water	Second-messenger modulator; down-regulates anxiety and peripheral nerve sensitivity.

Table 3. Neurological calming and anti-excitatory micronutrients

Phase 2: Pre-Surgical Tissue & Systemic Support

Active Compound	Optimal Dose	Administration Timing	Clinical Rationale & Mechanism
Vitamin D3 + K2 (MK-7)	5,000 IU D3 + 100 mcg K2	Breakfast (fat meal)	Critical immune modulator in SPS autoimmunity; directs calcium to bone matrices.
Methylated B-Complex	1 Capsule daily	Morning breakfast	Essential cofactors for GAD enzyme and peripheral myelin sheath preservation.
Omega-3 Fish Oil	2,000 - 3,000 mg (1.5g EPA)	With lunch or dinner	Suppresses systemic inflammatory cytokines (IL-1, IL-6, TNF-alpha).
Hydrolyzed Collagen + Vit C	15g Collagen + 500mg Vit C	45 min prior to pre-hab	Directly stimulates tenocyte and chondrocyte collagen synthesis in joint capsules.

Table 4. Surgical joint preparation and immunological stabilizers

Critical Surgical Precaution

MANDATORY SURGICAL PAUSE: Discontinue high-dose Omega-3 fish oil, Vitamin E, and any high-dose anti-platelet herbal extracts exactly **10 to 14 days prior** to any surgical intervention (knee, shoulder, ankle) to eliminate hemorrhage risk. Always provide this exact table to your surgical anesthesiologist.

5. Sensory De-Escalation & Vagal Nerve Retraining

In SPS, sensory startle (acoustic, tactile, or unexpected vestibular shifts) is the trigger for acute spasm storms. Creating an environmental buffer is non-negotiable.

Sensory Environmental Hygiene

- **Acoustic Buffering:** High-fidelity acoustic earplugs (such as Loop Engage or custom musician plugs) reduce ambient dB spikes while maintaining speech clarity in social/public settings.
- **Visual Shielding:** Replace sharp LED cool-white bulbs (5000K+) with soft warm lighting (2200K - 2700K). Avoid strobe effects, flickers, and blue light post-sundown.
- **Daily Zero-Stimulus Windows:** Schedule two 15-minute blocks daily: total darkness (eye mask), quiet or consistent low-frequency brown noise, lying flat in a fully supported position.

Vagus Nerve & Autonomic Down-Regulation Practices

- **The Physiological Sigh:** Two sequential nasal inhalations (one deep, one sharp 'top-up' inhale) followed by an unforced, prolonged oral exhalation. Perform 5 consecutive rounds whenever muscle tightness begins to climb.
- **Passive Body Scan (Never Progressive Muscle Relaxation):** In SPS, deliberately contracting a muscle during PMR can trigger an intractable spasm. Use purely passive, mindful observation of body weight, warmth, and surface contact.

6. Restorative Sleep Architecture & Joint Ergonomics

Transitioning from stage 1 to stage 2 sleep is the most vulnerable phase for nocturnal spasms. Maintaining physiological alignment across the knee, shoulder, and ankle prevents compression-triggered awakenings.

- **Shoulder De-Loading:** If side sleeping on the non-injured side, place a firm hug-pillow under the affected upper arm. This prevents forward slump and internal rotation torque on the shoulder rotator cuff and labrum.
- **Knee & Ankle Separation:** Position a contoured memory foam knee-to-ankle spacer between the legs. Preventing bone-on-bone contact between the knees and malleoli stops nociceptive signals from traveling up the spinal cord.
- **Back Sleeping Alternative:** Place a cylindrical wedge or rolled pillow beneath the knees to maintain 15-20 degrees of knee flexion, relaxing the hamstrings and taking tension off the lumbosacral plexus.
- **Pre-Bed Thermal Normalization:** 20 minutes before bed, submerge lower legs/feet in a warm Epsom salt bath (magnesium sulfate). Warmth counteracts cold-induced SPS rigidity while transdermal heat relaxes calf muscle spindles.

7. Joint-Spared Isometric 'Pre-Hab' Program

Because the knee, shoulder, and ankle are all undergoing procedures, conventional gym movements, aggressive stretching, and ballistic exercises are strictly contraindicated. The objective is isometric neuromuscular activation without articular shear.

Prescribed Daily Low-Impact Movements

Target Area	Exercise Protocol	Dosage & Sets	Biomechanical Objective
Knee & Quadriceps	Quad Setting / Straight-Leg Submaximal Press (Towel roll under knee, press gently into bed)	3 sets x 8 reps (5-sec hold)	Maintains vastus medialis neural mapping without patellofemoral friction.
Shoulder Complex	Scapular Retraction & Depression (Squeeze blades back and down with arms at sides)	3 sets x 10 reps (5-sec hold)	Strengthens rhomboids and lower trapezius; decompresses subacromial space.
Ankle & Foot	Non-Weight-Bearing Ankle Alphabets & Seated Toe Spreads	2 sets through alphabet (slow)	Circulates synovial fluid and maintains dorsal/plantar nerve firing safely.
Core / Pelvis	Gentle Supine Pelvic Tilts (Flatten lower back against mattress, engage transverse abdominis)	3 sets x 8 reps (4-sec hold)	Stabilizes the lumbopelvic core, taking asymmetric strain off lower extremity joints.

Table 5. Non-weight-bearing isometric pre-hab protocol

8. 12-Month Action Roadmap & Pre-Operative Checkpoints

A structured chronology ensures you arrive at both neurological treatments and joint surgeries in the strongest possible physiological state:

- **Months 1 to 3 (Foundational Stabilization):** Establish daily mineralized hydration formula; transition fully to nut-free, legume-free whole-food eating; establish bedroom sensory gating and evening magnesium routine.
- **Months 4 to 6 (Connective Tissue Loading & Pre-Hab):** Introduce daily hydrolyzed collagen + Vitamin C protocol 45 minutes prior to isometric joint pre-hab; build consistent vagal regulation habits.
- **Months 7 to 9 (Neurological Synchronization):** Coordinate with surgical team and neurologist regarding anesthesia protocols (specifying Stiff Person Syndrome, avoiding sudden GABA withdrawal, planning post-op spasm control).
- **Months 10 to 12 (Pre-Surgical Clearance):** Discontinue high-dose omega-3s and blood thinners 14 days pre-op; conduct pre-admission clinic walkthrough; prepare adaptive home environment (crutches, braces, ice machines).

Final Guidance

Summary Commitment: Consistency in low-level habits—calm environments, mineralized water, anti-inflammatory proteins, and daily gentle isometrics—will do more to safeguard your central nervous system over the next 365 days than any sporadic high-intensity intervention. Take this report to your medical specialists to align your whole team.