Name	[redacted]	VID	[redacted]
Gender	Male	Client name	WALKIN
Phone	[redacted]		
DOB	[redacted]1982		
ID number	[redacted]		
Address	53, NGUYEN DU, Sai Gon Ward, Ho Chi Minh City		

Registered **24/02/2026 12:17 PM** Collection **24/02/2026 12:27 PM** Received **24/02/2026 01:13 PM** Reported **13/03/2026 03:07 PM**

TEST NAME	RESULT	REFERENCE RANGE	UNIT
Mean Corpuscular Hemoglobin (MCH)	29.10	27 - 32	pg
Mean Corpuscular Hemoglobin Concentration (MCHC)	33.3	31.5 - 34.5	g/dL
Red Cell Distribution Width (RDW)	13.20	11.6 - 14	%
Platelets (PLT)	163.00	150 - 419	$10^3/\mu L$
Mean Platelets Volume (MPV)	11.70	9.04 - 12.79	fL
Platelets Distribution Width (PDW)	13.10	10 - 17	10 GSD
Plateletcrit (PCT)	**0.191**	0.23 - 0.24	%
ELECTROPHORESIS (HPLC)			
HbA1c [Whole blood] EDTA Whole Blood, [Tosoh] G11/G11LA			
HbA1c (NGSP)	5.1	Normal: < 5.7 Prediabetes: 5.7 - 6.4 Diabetes: ≥ 6.5	%
HbA1c (IFCC)	33.00	Normal: < 39 Prediabetes: 39 - 47 Diabetes: ≥ 48	mmol/mol
Estimated Average Glucose (eAG)	5.53 100		mmol/L mg/dL

Interpretation

1. Trends in HbA1c are a better indicator of diabetic control than a solitary test.

2. Interference of Haemoglobinopathies in HbA1c estimation:

a. If HbF is > 25%, or homozygous hemoglobinopathy is detected, Fructosamine is recommended for diabetic monitoring.

b. On Tosoh G11, HbA1c is corrected for Heterozygous HbS and HbC traits.

3. Estimated Average Glucose (eAG) is a calculated average of your blood sugar (Glucose) levels over a period of 2 to 3 months.

BIOCHEMISTRY			
Glucose Fasting [Plasma] Fasting Plasma, [Abbott] Alinity c	4.50 81	Normal: 3.9 - 5.5 Prediabetes: 5.6 - 6.9 Diabetes: ≥ 7	mmol/L mg/dL