Home

About

Problems by Topics

Linear Algebra

Gauss-Jordan Elimination

Inverse Matrix

Linear Transformation

Vector Space

Eigen Value

Cayley-Hamilton Theorem

Diagonalization

Exam Problems

By Topics

Group Theory

Abelian Group

Group Homomorphism

Sylow's Theorem

Field Theory

Module Theory

Ring Theory

LaTex/MathJax

Login/Join us

Records

Solve later Problems

My Solved Problems

Teaching

Math 2568 Linear Algebra Spring 2018

_

LINEAR ALGEBRA

FOLLOW:






Eigenvalues of a Stochastic Matrix is I'm Yu Tsumura.

LINEAR ALGEBRA PROBLEMS BY TOPICS

Always Less than or Equal to 1

BY YU · PUBLISHED 11/17/2016 · UPDATED 07/06/2017





Problem 185

Let $A = (a_{ij})$ be an $n \times n$ matrix. We say that $A = (a_{ij})$ is a **right stochastic matrix** if each entry a_{ij} is nonnegative and the sum of the entries of each row is 1. That is, we have

 $a_{ij} \ge 0$ and $a_{i1} + a_{i2} + \cdots +$ for $1 \le i, j \le n$.

Let $A = (a_{ij})$ be an $n \times n$ right stochastic matrix. Then show the following statements.

- (a) The stochastic matrix A has an eigenvalue 1.
- **(b)** The absolute value of any eigenvalue of the stochastic matrix A is less than or equal to 1.

The list of linear algebra problems is available here.




SUBSCRIBE TO BLOG VIA EMAIL

Enter your email address to subscribe to this blog and receive notifications of new posts by email.

Email Address

Subscribe

Q To search type and hit enter





Contents [hide]

Problem 185

Proof.

- (a) The stochastic matrix A has an eigenvalue 1.
- (b) The absolute value of any eigenvalue of the stochastic matrix A is less than or equal to 1.

Remark.



Proof.

(a) The stochastic matrix A has an eigenvalue 1.

We compute that

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \dots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ \vdots & \vdots & \vdots \\ 1 \end{bmatrix}$$

Here the second equality follows from the definition of a right stochastic matrix.

(Each row sums up to 1.)

This computation shows that 1 is







LINEAR ALGEBRA

- Introduction to Matrices
- Elementary Row Operations
- Gaussian-Jordan Elimination
- Solutions of Systems of Linear Equations
- Linear Combination and Linear Independence
- Nonsingular Matrices
- Inverse Matrices
- Subspaces in \mathbb{R}^n
- Bases and Dimension of Subspaces in \mathbb{R}^n

an eigenvector of A and $\begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix}$ is an

eigenvector corresponding to the eigenvalue 1.

(b) The absolute value of any eigenvalue of the stochastic matrix A is less than or equal to 1.

Let λ be an eigenvalue of the stochastic matrix A and let ${\bf v}$ be a corresponding eigenvector.

That is, we have

$$A\mathbf{v} = \lambda \mathbf{v}$$
.

Comparing the i-th row of the both sides, we obtain

$$a_{i1}v_1 + a_{i2}v_2 + \cdots + a_{in}v_n =$$

for i = 1, ..., n.

Let

$$|v_k| = \max\{|v_1|, |v_2|, \dots, |v_n|\},\$$

namely v_k is the entry of **v** that has the maximal absolute value.

Note that $|v_k| > 0$ since otherwise we have $\mathbf{v} = \mathbf{0}$ and this contradicts that an eigenvector is a nonzero vector.

Then from (*) with i = k, we have

- General Vector Spaces
- Subspaces in General Vector Spaces
- Linearly Independency of General Vectors
- Bases and Coordinate Vectors
- Dimensions of General Vector Spaces
- **–** Linear Transformation from \mathbb{R}^n to \mathbb{R}^m
- Linear Transformation Between Vector Spaces
- Orthogonal Bases
- Determinants of Matrices
- Computations of Determinants
- Introduction to Eigenvalues and Eigenvectors
- Eigenvectors and Eigenspaces
- Diagonalization of Matrices
- The Cayley-Hamilton Theorem
- Dot Products and Length of Vectors
- Eigenvalues and Eigenvectors of Linear Transformations
- Jordan Canonical Form

CATEGORIES

- Elementary Number Theory (1)
- Field Theory (26)

$$|\lambda| \cdot |v_k| = |a_{k1}v_1 + a_{k2}v_2 + \dots + \leq a_{k1}|v_1| + a_2|v_2| + \dots \leq a_{k1}|v_k| + a_2|v_k| + \dots = (a_{k1} + a_{k2} + \dots + a_{kn})$$

Since $|v_k| > 0$, it follows that

$$\lambda \leq 1$$

as required.



Remark.

A stochastic matrix is also called probability matrix, transition matrix, substitution matrix, or Markov matrix.

Click here if solved $\nleq 7$

0 Like 1

Share 1

Tweet

More from my site



Stochastic
Matrix
(Markov
Matrix) and its
Eigenvalues
and

Eigenvectors

(a) Let

$$A = \begin{bmatrix} a_{11} & a_1 \\ a_{21} & a_2 \end{bmatrix}$$

be a matrix such that $a_{11} + a_{12} = 1$ and $a_{21} + a_{22} = 1$. Namely, the sum of the entries in each

- **⇐** General (7)
- **►** Group Theory (126)
- Linear Algebra (485)
- ► Math-Magic (1)
- ► Module Theory (13)
- ► Ring theory (64)
- ► Welcome (1)

MATHJAX

Mathematical equations are created by MathJax. See How to use MathJax in WordPress if you want to write a mathematical blog.



RECENT POSTS

- If the Nullity of a Linear Transformation is Zero, then Linearly Independent Vectors are Mapped to Linearly Independent Vectors
- Find All Values of *x* such that the Matrix is Invertible
- Find All Eigenvalues and Corresponding Eigenvectors for the 3×3 matrix

Eigenvalues of a Stochastic Matrix is Always Less than or Equal to 1 – Problems in Mathematics

row is 1. (Such a matrix is called (right) stochast matrix (also termed [...]



Find the Limit of a Matrix Let

$$A = \begin{bmatrix} \frac{1}{7} & \frac{3}{7} \\ \frac{3}{7} & \frac{1}{7} \\ \frac{3}{7} & \frac{3}{7} \end{bmatrix}$$

be 3×3 matrix. Find

$$\lim_{n\to\infty}A^n$$
.

(Nagoya
University
Linear [...]



Basis with Respect to Which the Matrix for Linear **Transformation** is Diagonal Let P_1 be the vector space of all real polynomials of degree 1 or less Consider the linear transformation $T: P_1 \rightarrow P_1$ defined by

$$T(ax + b) = (1)$$

for any $ax + b \in P_1$.

(a) With respect to the basis $B = \{1, x\}$, find the matrix of the linear transformation

- Find All Values of *a* which Will Guarantee that *A* Has Eigenvalues 0, 3, and -3.
- Compute the Determinant of a Magic Square

LINEAR ALGEBRA

If A is an Idempotent Matrix, then When I - kA is an Idempotent Matrix?

LINEAR ALGEBRA

Express a Hermitian Matrix as a Sum of Real Symmetric Matrix and a Real Skew-Symmetric Matrix

LINEAR ALGEBRA

Quiz 3. Condition that Vectors are Linearly Dependent/ Orthogonal Vectors are Linearly Independent

LINEAR ALGEBRA

Trace of the Inverse Matrix of a Finite Order Matrix

LINEAR ALGEBRA

Find Matrix Representation of Linear Transformation From \mathbb{R}^2 to \mathbb{R}^2

TOP POSTS

- How to Diagonalize a Matrix. Step by Step Explanation.
- Positive definite Real Symmetric Matrixand its Eigenvalues
- The Intersection of Two Subspaces is also a Subspace
- The Matrix for the Linear Transformation of the Reflection Across a Line in the Plane

Eigenvalues of Real Skew-Symmetric Matrix are Zero or Purely Imaginary and

 $[\ldots]$



Every Vector is Eigenvector, then Matrix is a Multiple of Identity Matrix Let A be an $n \times n$ matrix. Assume that every vector **x** in \mathbb{R}^n is an eigenvector for some eigenvalue of A. Prove that there exists $\lambda \in \mathbb{R}$ such that $A = \lambda I$, where I is the $n \times n$ identity matrix. Proof. Let us write [...]



Given All Eigenvalues and Eigenspaces, Compute a Matrix Product Let C be a 4×4 matrix with all eigenvalues $\lambda = 2, -1$ and eigensapces \ [E 2=\Span\left $\$ \quad \begin{bmatrix $1 \setminus 1 \setminus 1 \setminus 1$ \end{bmatrix} \quad\right \} \text{ and } $E_{-1}=\Span \ le$ \quad\begin{br $1 \setminus 2 \setminus 1 \setminus 1$

- the Rank is Even
- 6 Express a Vector as a Linear Combination of Other Vectors
- 7 Prove a Group is Abelian if $(ab)^2 = a^2b^2$
- Eigenvalues of a Hermitian Matrix are Real Numbers
- Rotation Matrix in the Plane and its Eigenvalues and Eigenvectors
- Vector Form for the General Solution of a System of Linear Equations

SITE MAP & INDEX

Site Map

Index

[...]



Eigenvalues of

a Hermitian

Matrix are

Real Numbers

Show that

eigenvalues of

a Hermitian

matrix A are

real numbers.

(The Ohio

State

University

Linear Algebra

Exam

Problem) We

give two

proofs. These

two proofs are

essentially the

same. The

second proof

is a bit simpler

and concise

compared to

the first one.

[...]



Determine

Eigenvalues,

Eigenvectors,

Diagonalizable

From a Partial

Information of

a Matrix

Suppose the

following

information is

known about a

 3×3 matrix

 $A. \setminus$

[A\begin{bmatı

 $1 \setminus 1 \setminus 1$

\end{bmatrix}=

 $1 \setminus 1 \setminus 2 \setminus 1$

\end{bmatrix},

\quad

A\begin{bmatri

1\\-1\\1[...]



Given

Eigenvectors

and

Eigenvalues,

Compute a

Matrix

Product

(Stanford

University

Exam)

Suppose that

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix}$$
 is an

eigenvector of

a matrix A

corresponding

to the

eigenvalue 3

and that
$$\begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

is an

eigenvector of

 \boldsymbol{A}

corresponding

to the

eigenvalue

-2. Compute

 $A^2\geq \sum_{b\in A}$

4 [...]

Tags: eigenvalue eigenvector

linear algebra Markov matrix matrix

probability matrix stochastic matrix

substitution matrix transition matrix

triangle inequality vector

PREVIOUS

STORY

Eigenvalues

of Squared

Matrix and

Upper

Triangular

Matrix

NEXT

STORY

Is the

Determinant >

of a Matrix

Additive?

YOU MAY ALSO LIKE...



Finite Linearly Find the Order Depende Matrix

Matrix nt if and Represen

and its only if a tation of

Trace Vector $T(f)(x) = f(x^2)$

07/30/2016 Can be if it is a

Written Linear as a Transfor

as a Transfor Linear mation

Combina 01/21/2018

tion of

Remaini ng

03/20/2017

Vectors

LEAVE A REPLY

Comment		
Name *	Email *	
Website		
☐ Notify me	of follow-up	
comments by	email.	
☐ Notify me	of new posts by	
email.		

TAGS

abelian group augmented matrix basis basis for a vector space characteristic polynomial commutative ring determinant determinant of a matrix diagonalization diagonal matrix eigenvalue

eigenvalue
eigenvector elementary row
operations exam field theory
finite group group group
homomorphism group
theory homomorphism
ideal inverse matrix

linear algebra

invertible matrix kernel

SUBSCRIBE TO BLOG VIA EMAIL

Enter your email address to subscribe to this blog and receive notifications of new posts by email.

Join 350 other subscribers

Email Address					
Subscribe					

SEARCH MORE PROBLEMS

 ${\bf Q}$ To search type and hit ente

CREATE AN ACCOUNT

Username			
Email			
Password			
Repeat			
Password			
First Name			
Last Name			
Membership	Free		
Level			
Register			

If you are a member, Login here.

linear combination linearly

independent linear

transformation matrix

matrix representation nonsingular

matrix normal subgroup null

space Ohio State Ohio

State.LA rank ring ring

theory subgroup subspace

symmetric matrix system of linear

equations transpose Vector

vector space

Home

About

group theory

Linear Algebra

Index

Site Map



Problems in Mathematics © 2019. All Rights Reserved.



