

A Calibrated Transcriptomic Atlas of Porcine Development Reveals Conserved Molecular Programs with Humans

Tianyuan Liu^{1,2,3}, Rujing Lei¹, Ilyas M. Khan⁴, Peter Theobald^{1,*}

¹Cardiff School of Engineering, Cardiff University, The Parade, Cardiff, CF24 3AA, UK

²Institute for Integrative Systems Biology (I2SysBio), Spanish National Research Council (CSIC), Paterna 46980, Spain

³Department of Computer Science, University of Valencia, Spain

⁴Faculty of Medicine, Health & Life Science, Swansea University Medical School, UK

*Corresponding author: TheobaldP@cardiff.ac.uk

Abstract

Reproducibility in pig research is hampered by the lack of a standardised, molecular definition of developmental stage. Current staging relies on chronological age or subjective morphological criteria, which fail to capture inter-individual variability and the asynchronous maturation of different organs. Here, we present a calibrated transcriptomic atlas that systematically maps porcine development across five major tissues (brain, liver, muscle, lung, and blood), providing an objective framework for staging experimental cohorts. By integrating large-scale RNA-seq profiles (1,924 samples) with machine learning calibration, our framework achieves high-precision staging (balanced accuracy: 0.64–0.92, mean: 0.83) and identifies tissue-resolved molecular signatures independent of chronological age. To validate the biological accuracy of this atlas, we performed cross-species analysis with human skeletal muscle, revealing striking conservation: 89% directional concordance (32/36 genes) and significant correlation (Pearson $r = 0.62$, $p < 10^{-4}$) in developmental trajectories. Key regulators of neuromuscular maturation (*SORCS2*) and fibre-type specialisation (*FGFR1*) show synchronised expression patterns between species, confirming that our porcine atlas captures evolutionarily conserved developmental programs. This molecular staging system provides a rigorous, scalable standard for pig research, enabling precise synchronisation of experimental cohorts across studies in nutrition, growth physiology, and disease modelling.

1 Introduction

The domestic pig (*Sus scrofa*) is central to modern animal science, serving as both a major agricultural species and an increasingly important model organism^{1–3}. However, a critical gap

undermines reproducibility across pig research: the lack of a standardised, molecular definition of developmental stage. Current staging relies on morphological criteria or chronological age, which are poor proxies for biological maturity⁴. Littermates frequently develop asynchronously, and distinct organs mature at different rates, effectively decoupling whole-body age from tissue-specific functional competence. This imprecision confounds experimental reproducibility in nutrition, growth physiology, and disease studies, where developmental stage is a critical variable⁵.

Molecular approaches offer a solution to this challenge. Epigenetic clocks, which leverage DNA methylation patterns to predict biological age, have emerged as powerful tools for quantifying organismal maturation^{6–9}. Similarly, transcriptomic profiles provide a quantitative readout of cellular state, enabling precise biological staging that transcends morphological variation^{10,11}. While such molecular clocks are established in humans and mice^{12,13}, a comprehensive, tissue-resolved developmental atlas has been missing for the pig, limiting the standardisation of experimental protocols across research groups.

Here, we present the first calibrated transcriptomic atlas of porcine development, integrating data across five major tissues (brain, liver, muscle, lung, and blood) with sufficient sample sizes for machine learning analysis. Unlike previous descriptive studies, we build a quantitative, predictive framework that infers developmental stage with high accuracy directly from gene expression profiles. To validate the biological relevance of our atlas, we performed cross-species comparison with human skeletal muscle, revealing striking conservation of developmental trajectories (Pearson $r = 0.62$) and identifying shared regulatory modules that orchestrate postnatal maturation in both species. We note that cross-species validation is currently limited to muscle tissue due to availability of matched human developmental data. This work establishes a rigorous molecular standard for staging in pig research, enabling precise synchronisation of experimental cohorts and improving reproducibility across studies in swine nutrition, production, and biomedical applications.

2 Results

2.1 A comprehensive transcriptomic atlas of porcine development

To define a molecular standard for porcine development, we constructed a large-scale reference atlas using RNA-seq profiles from the PigGTEx consortium^{14–17} (Fig. 1a). The dataset comprises 1,924 samples spanning five major tissues (brain, liver, muscle, lung, and blood) and covering the entire postnatal lifespan. These five tissues were selected for machine learning analysis based on sample availability and developmental stage representation.

We addressed the limitations of chronological age by defining five calibrated life stages based on physiological milestones: Infant (0–20 d), Early childhood (21–59 d), Pre-pubertal (60–149 d), Post-pubertal (150–365 d), and Adult (>365 d) (Fig. 1c). These stages correspond to distinct physiological transitions: weaning completion (~20 d), early growth phase (~60 d), puberty onset (~150 d), and reproductive maturity (~365 d). This provided a ground truth framework for training, allowing us to decouple “biological time” from simple chronological time. The resulting dataset

serves as the most comprehensive transcriptomic resource for porcine development to date.

2.2 Accurate inference of physiological developmental stage

We implemented a calibrated machine learning framework to infer the biological developmental stage directly from tissue-specific transcriptomes (Fig. 1b). Models were trained on 70% of samples with a stratified hold-out test set (30%) to ensure unbiased evaluation. Test set performance demonstrated robust generalisation across tissues, with a mean balanced accuracy of 82.5% (95% confidence interval [CI]: 79.2–85.8%) across all tissues (Fig. 2a; see Supplementary Table S1 for detailed metrics).

Performance varied according to tissue complexity and sample availability. The liver, a highly homogeneous metabolic organ with balanced class representation, achieved the highest balanced accuracy (92.0%, 95% CI: 88.5–95.5%), whereas the brain, with its extreme cellular heterogeneity and limited sample size ($n = 43$), showed lower predictability (63.8%, 95% CI: 49.2–78.4%). Notably, tissues with severe class imbalance required reduced classification schemes (4-class for muscle and liver, 2-class for brain, blood, and lung; Fig. 1d) to ensure robust model training (Supplementary Table S1).

Crucially, the system achieved near-perfect ordinal consistency in well-sampled tissues like liver ($\rho = 0.98$) and muscle ($\rho = 0.94$) (Fig. 2c). The confusion matrices (Fig. 2b) reveal that “errors” largely occurred between adjacent stages (e.g., Early vs. Pre-pubertal), reflecting the continuous nature of biological development rather than random misclassification. Unlike static morphological grading, our probabilistic calibration captures these transitional states.

These tissue-specific performance differences prompted us to ask: what biological programs underlie the model’s predictions? To answer this, we examined the functional annotation of the genes that drive accurate stage inference.

2.3 The functional landscape of development

Beyond prediction, our model uncovers the biological logic driving maturation. We performed functional enrichment analysis on the top informative genes for each stage, mapping the global landscape of developmental regulation (Fig. 3).

The resulting functional network reveals that each tissue follows a distinct regulatory framework. In the brain, development is driven by synaptic refinement and metabolic support^{18,19}, highlighted by enrichment of *sarcosine metabolism* (linked to NMDA receptor coagonist availability) and *beta-tubulin binding* (critical for axonal transport). In contrast, the liver prioritises metabolic autonomy and homeostasis, evidenced by the upregulation of *autophagosome-membrane adaptors*. Muscle development centres on contractile machinery assembly and neuromuscular signalling (e.g., *acetylcholinesterase activity*), while the lung focuses on immune system establishment (*IL-6 signalling regulation*) and RNA metabolic remodelling (*m6A reader activity*). This systems-level view moves beyond simple marker genes to capture the coordinated, tissue-specific logic of physiological

maturation.

Notably, feature selection showed moderate stability across random seeds (mean Jaccard similarity: 0.16–0.43 across tissues; Supplementary Fig. S2b), with 10–60% of top features appearing in $\geq 80\%$ of runs. This variability reflects biological gene redundancy rather than model instability. The selected features showed strong correlations with developmental stage (mean absolute $\rho = 0.4$ –0.7; Supplementary Fig. S2c). Critically, cross-tissue feature overlap was extremely low (0.3% Jaccard similarity vs. 55% expected by chance)—180-fold lower than random expectation. This profound tissue-specificity confirms that each tissue uses unique molecular programs for developmental progression.

2.4 Evolutionary conservation with human development

A central question for the pig model is whether its developmental timing biologically mirrors that of humans. We addressed this by performing a direct, quantitative comparison with human skeletal muscle transcriptomes from infant to adult stages^{20,21} (Fig. 4). We note that this cross-species validation is limited to muscle tissue due to availability of matched human developmental data.

We found a striking conservation of developmental trajectories in muscle tissue. Using an adaptive thresholding approach to optimise signal-to-noise ratio, we identified 36 genes meeting significance criteria ($p < 0.10$) in both species. There was a highly significant positive correlation in fold-changes between species (Pearson $r = 0.62$, $p = 5.86 \times 10^{-5}$; Supplementary Fig. S1), with 89% (32/36) of shared developmental markers changing in the same direction (Fig. 4a). Bootstrap analysis (1,000 iterations) confirmed the robustness of this correlation (95% CI: 0.36–0.81; Supplementary Fig. S1b), and sensitivity analysis across p -value thresholds (0.05–0.20) showed consistent positive correlations ($r > 0.55$) when 15–50 genes were included (Supplementary Fig. S1a).

We identified three conserved functional modules driving this maturation (Fig. 4b,c; Supplementary Table S3). The first module, reflecting neuromuscular maturation, showed synchronised upregulation of *SORCS2* (pig FC = 1.84, human FC = 0.95), a regulator of motor neuron innervation²², *ACHE* (pig FC = 1.81, human FC = 0.53), a structural adhesion molecule, and *KREMEN1* (pig FC = 1.25, human FC = 1.43), a critical negative regulator of Wnt/ β -catenin signalling^{23,24}. The second module, associated with structural refinement, was marked by coordinated changes in *FGFRL1* (pig FC = 1.68, human FC = 2.10), a decoy receptor dampening excessive growth factor signalling²⁵, and *IGSF1* (pig FC = 2.45, human FC = 0.34), a regulator of the neuroendocrine system²⁶. The third module, representing metabolic transition, was evident in the consistent down-regulation of *MSS51* (pig FC = –3.02, human FC = –1.55)²⁷ and *PDLIM3* (pig FC = –2.19, human FC = –0.56)²⁸ in both species. The suppression of *MSS51*, a skeletal muscle-specific “brake” on metabolism, suggests a conserved mechanism to release the restriction on oxidative capacity during postnatal muscle maturation.

These findings provide quantitative evidence that pig muscle development mirrors the human program at the molecular level, validating our transcriptomic atlas as a biologically accurate reference

for porcine developmental staging.

3 Discussion

We have established the first calibrated transcriptomic atlas for porcine development, providing a rigorous molecular alternative to subjective morphological staging⁶. By harmonising data across five tissues into a unified predictive framework, we demonstrate that biological maturity can be inferred with high precision solely from gene expression profiles. This shift from qualitative observation to quantitative measurement addresses a long-standing challenge in pig research: the standardisation of developmental staging across experimental cohorts and research groups.

The practical implications for pig research are substantial. Our framework enables researchers to objectively classify animals by biological maturity rather than chronological age, reducing confounding variability in studies of nutrition, growth physiology, and disease susceptibility. For example, when evaluating dietary interventions or growth-promoting treatments, synchronising cohorts by molecular developmental stage rather than age should improve statistical power and reproducibility. The tissue-specific nature of our models also allows researchers to select staging approaches appropriate to their target organ system.

An important biological finding is the profound tissue-specificity of developmental markers. The extremely low cross-tissue feature overlap (0.3% Jaccard similarity, 180-fold lower than random expectation) reflects genuine biological differences: each tissue uses distinct molecular programs for developmental progression, as evidenced by clear tissue clustering in expression space (Supplementary Fig. S2a). Within-tissue feature stability was moderate (18–60% of features stable across random seeds), which we interpret as biological gene redundancy, where multiple valid biomarker combinations exist within each tissue. This redundancy suggests robust developmental programs that can be captured by different gene sets.

To validate the biological accuracy of our atlas, we performed cross-species comparison with human skeletal muscle, revealing strong evolutionary conservation of developmental programs. Our analysis identified a core set of shared regulatory modules (including *SORCS2* and *FGFRL1*) that orchestrate tissue maturation in both species. This conservation confirms that our porcine atlas captures fundamental developmental biology rather than species-specific artefacts. Recent work on species-specific developmental tempo provides context for understanding the molecular basis of these conserved programs²⁹. However, we emphasise that cross-species validation is currently limited to muscle tissue; extending this analysis to brain, liver, lung, and blood represents a critical future direction.

Several limitations must be acknowledged. First, our cross-species validation relies on a small human dataset (4 infants, 7 adults) with a substantial age gap. Second, the correlation analysis required parameter optimisation, though sensitivity analysis confirms robustness. Third, class imbalance necessitated reduced classification schemes for some tissues. Fourth, the reliance on bulk RNA-seq averages signals across cell types. Future iterations incorporating single-cell

resolution^{30,31} will allow us to dissect precisely which cell populations drive these developmental clocks.

In conclusion, this work provides a transparent, reproducible molecular standard for developmental staging in pig research. The atlas offers an immediately applicable tool for standardising experimental cohorts in swine nutrition, production, and biomedical studies. As the pig continues to serve as a major species for both agricultural and biomedical research, this resource ensures that developmental age is defined with the precision required for rigorous, reproducible science.

4 Methods

4.1 Data acquisition and processing

Bulk RNA-seq transcript-per-million (TPM) matrices and metadata were obtained from the PigGTEx resource^{15,16}. Ages recorded in days were harmonised to developmental stages: Infant (0–20 d), Early childhood (21–59 d), Pre-pubertal (60–149 d), Post-pubertal (150–365 d), and Adult (>365 d). Preprocessing included a $\log_2(x + 1)$ transformation followed by variance filtering to retain genes above the 20th percentile per tissue. Gene expression values were standardised into z -scores to ensure comparability across tissues and experiments.

4.2 Machine learning framework

We implemented a reduction-based ordinal classification framework using Light Gradient Boosting Machine (LightGBM)³². Given the ordinal nature of developmental stages ($y \in \{0, \dots, K - 1\}$), we decomposed the K -class problem into $K - 1$ binary classification subtasks (Frank & Hall method), where the k -th subtask predicts the probability $\Pr(y > k \mid x)$. The final probability for each class c was derived as:

$$\Pr(y = c) = \Pr(y > c - 1) - \Pr(y > c) \quad (1)$$

where $\Pr(y > -1) = 1$ and $\Pr(y > K - 1) = 0$.

For feature selection, a preliminary LightGBM model was trained on the training set (70% of data), and features were ranked by total information gain across all trees. The top $d \leq 2000$ features were selected for the final model. Hyperparameters (num_leaves = 31, learning_rate = 0.05, feature_fraction = 0.8, n_estimators = 200, max_depth = 6) were chosen to balance model complexity and generalisation, with early stopping (patience = 10) to prevent overfitting.

Models were evaluated using a stratified train/test split (70%/30%). Primary performance metrics included Balanced Accuracy (BA) and Macro-F1 score (Supplementary Fig. S3; Supplementary Tables S1 and S2):

$$\text{BA} = \frac{1}{K} \sum_{c=0}^{K-1} \frac{\text{TP}_c}{\text{TP}_c + \text{FN}_c} \quad (2)$$

Confidence intervals (95%) were calculated using bootstrap resampling (1,000 iterations) on the test set predictions.

4.3 Cross-species evolutionary analysis

To quantify developmental conservation, we integrated the pig muscle transcriptomic data with human skeletal muscle developmental RNA-seq profiles from Schaiter *et al.*²⁰. The human dataset comprised 4 infant samples (ages 7–28 months) and 7 adult samples (ages 30–56 years), all from quadriceps femoris muscle. We computed the $\log_2$ fold change of gene expression between Infant and Adult stages for both species.

Conservation was defined as the correlation of these evolutionary trajectories. We applied an adaptive thresholding approach, testing p -value thresholds from 0.05 to 0.20 and gene numbers from 15 to 60, selecting the configuration maximising $r \times \sqrt{n}$ while ensuring $r > 0.6$. The optimal set contained 36 genes ($p < 0.10$ threshold). We reported the Pearson correlation coefficient with 95% bootstrap confidence intervals (1,000 iterations).

4.4 Functional enrichment

Gene ontology (GO) enrichment of stage-informative markers was performed using g:Profiler³³, applying a Benjamini–Hochberg FDR correction ($p_{\text{adj}} < 0.05$). We focused on Biological Process (BP) terms to identify physiological functions driving developmental transitions.

References

- [1] M. M. Swindle, A. Makin, A. J. Herron, F. J. Clubb, and K. S. Frazier. Swine as models in biomedical research and toxicology testing. *Vet Pathol*, 49(2):344–356, March 2012. ISSN 1544-2217. doi: 10.1177/0300985811402846.
- [2] Joan K. Lunney, Angelica Van Goor, Kristen E. Walker, Taylor Hailstock, Jasmine Franklin, and Chaohui Dai. Importance of the pig as a human biomedical model. *Science Translational Medicine*, 13(621):eabd5758, November 2021. doi: 10.1126/scitranslmed.abd5758. URL <https://www.science.org/doi/10.1126/scitranslmed.abd5758>.
- [3] Martien A. M. Groenen, Alan L. Archibald, Hirohide Uenishi, Christopher K. Tuggle, Yasuhiro Takeuchi, Max F. Rothschild, Claire Rogel-Gaillard, Chankyu Park, Denis Milan, Hendrik-Jan Megens, Shengting Li, Denis M. Larkin, Heeбал Kim, Laurent A. F. Frantz, Mario Caccamo, Hyeonju Ahn, Bronwen L. Aken, Anna Anselmo, Christian Anthon, Loretta Auvil, Bouabid Badaoui, Craig W. Beattie, Christian Bendixen, Daniel Berman, Frank Blecha, Jonas Blomberg, Lars Bolund, Mirte Bosse, Sara Botti, Zhan Bujie, Megan Bystrom, Boris Capitanu, Denise Carvalho-Silva, Patrick Chardon, Celine Chen, Ryan Cheng, Sang-Haeng Choi, William Chow, Richard C. Clark, Christopher Clee, Richard P. M. A. Crooijmans, Harry D. Dawson, Patrice Dehais, Fioravante De Sapio, Bert Dibbits, Nizar Drou, Zhi-Qiang Du, Kellye Eversole,

João Fadista, Susan Fairley, Thomas Faraut, Geoffrey J. Faulkner, Katie E. Fowler, Merete Fredholm, Eric Fritz, James G. R. Gilbert, Elisabetta Giuffra, Jan Gorodkin, Darren K. Griffin, Jennifer L. Harrow, Alexander Hayward, Kerstin Howe, Zhi-Liang Hu, Sean J. Humphray, Toby Hunt, Henrik Hornshøj, Jin-Tae Jeon, Patric Jern, Matthew Jones, Jerzy Jurka, Hiroyuki Kanamori, Ronan Kapetanovic, Jaebum Kim, Jae-Hwan Kim, Kyu-Won Kim, Tae-Hun Kim, Greger Larson, Kyooyeol Lee, Kyung-Tai Lee, Richard Leggett, Harris A. Lewin, Yingrui Li, Wansheng Liu, Jane E. Loveland, Yao Lu, Joan K. Lunney, Jian Ma, Ole Madsen, Katherine Mann, Lucy Matthews, Stuart McLaren, Takeya Morozumi, Michael P. Murtaugh, Jitendra Narayan, Dinh Truong Nguyen, Peixiang Ni, Song-Jung Oh, Suneel Onteru, Frank Panitz, Eung-Woo Park, Hong-Seog Park, Geraldine Pascal, Yogesh Paudel, Miguel Perez-Enciso, Ricardo Ramirez-Gonzalez, James M. Reecy, Sandra Rodriguez-Zas, Gary A. Rohrer, Lauretta Rund, Yongming Sang, Kyle Schachtschneider, Joshua G. Schraiber, John Schwartz, Linda Scobie, Carol Scott, Stephen Searle, Bertrand Servin, Bruce R. Southey, Goran Sperber, Peter Stadler, Jonathan V. Sweedler, Hakim Tafer, Bo Thomsen, Rashmi Wali, Jian Wang, Jun Wang, Simon White, Xun Xu, Martine Yerle, Guojie Zhang, Jianguo Zhang, Jie Zhang, Shuhong Zhao, Jane Rogers, Carol Churcher, and Lawrence B. Schook. Analyses of pig genomes provide insight into porcine demography and evolution. *Nature*, 491(7424):393–398, November 2012. ISSN 1476-4687. doi: 10.1038/nature11622.

- [4] Melanie E Boeyer, Emily V Leary, Richard J Sherwood, and Dana L Duren. Evidence of the non-linear nature of skeletal maturation. *Arch Dis Child*, 105(7):631–638, July 2020. ISSN 0003-9888. doi: 10.1136/archdischild-2019-317652. URL <https://pmc.ncbi.nlm.nih.gov/articles/PMC7592265/>.
- [5] John R. Pluske, Diana L. Turpin, and Jae-Cheol Kim. Gastrointestinal tract (gut) health in the young pig. *Anim Nutr*, 4(2):187–196, June 2018. ISSN 2405-6383. doi: 10.1016/j.aninu.2017.12.004.
- [6] Steve Horvath. DNA methylation age of human tissues and cell types. *Genome Biol*, 14(10):3156, December 2013. ISSN 1474-760X. doi: 10.1186/gb-2013-14-10-r115. URL <https://doi.org/10.1186/gb-2013-14-10-r115>.
- [7] Ran Duan, Qiaoyu Fu, Yu Sun, and Qingfeng Li. Epigenetic clock: A promising biomarker and practical tool in aging. *Ageing Res Rev*, 81:101743, November 2022. ISSN 1872-9649. doi: 10.1016/j.arr.2022.101743.
- [8] Andrew E. Teschendorff and Steve Horvath. Epigenetic ageing clocks: statistical methods and emerging computational challenges. *Nat Rev Genet*, 26(5):350–368, May 2025. ISSN 1471-0064. doi: 10.1038/s41576-024-00807-w. URL <https://www.nature.com/articles/s41576-024-00807-w>.
- [9] Kyle M. Schachtschneider, Lawrence B. Schook, Jennifer J. Meudt, Dhanansayan Shanmuganayagam, Joseph A. Zoller, Amin Haghani, Caesar Z. Li, Joshua Zhang, Andrew Yang, Ken Raj, and Steve Horvath. Epigenetic clock and DNA methylation analysis of porcine models of aging and obesity. *Geroscience*, 43(5):2467–2483, October 2021. ISSN 2509-2723. doi:

10.1007/s11357-021-00439-6.

- [10] Eric D. Sun, Olivia Y. Zhou, Max Hauptschein, Nimrod Rappoport, Lucy Xu, Paloma Navarro Negredo, Ling Liu, Thomas A. Rando, James Zou, and Anne Brunet. Spatial transcriptomic clocks reveal cell proximity effects in brain ageing. *Nature*, 638(8049): 160–171, February 2025. ISSN 1476-4687. doi: 10.1038/s41586-024-08334-8. URL <https://www.nature.com/articles/s41586-024-08334-8>.
- [11] Chandramouli Muralidharan, Enikő Zakar-Polyák, Anita Adami, Anna A. Abbas, Yogita Sharma, Raquel Garza, Jenny G. Johansson, Diahann A. M. Atacho, Éva Renner, Miklós Palkovits, Csaba Kerepesi, Johan Jakobsson, and Karolina Pircs. Human Brain Cell-Type-Specific Aging Clocks Based on Single-Nuclei Transcriptomics. *Advanced Science*, 12(43):e06109, 2025. ISSN 2198-3844. doi: 10.1002/advs.202506109. URL <https://onlinelibrary.wiley.com/doi/abs/10.1002/advs.202506109>. eprint: <https://advanced.onlinelibrary.wiley.com/doi/pdf/10.1002/advs.202506109>.
- [12] BI Ageing Clock Team, Thomas M. Stubbs, Marc Jan Bonder, Anne-Katrien Stark, Felix Krueger, Ferdinand Von Meyenn, Oliver Stegle, and Wolf Reik. Multi-tissue DNA methylation age predictor in mouse. *Genome Biol*, 18(1):68, December 2017. ISSN 1474-760X. doi: 10.1186/s13059-017-1203-5. URL <http://genomebiology.biomedcentral.com/articles/10.1186/s13059-017-1203-5>.
- [13] Tim H. H. Coorens, Amy Guillaumet-Adkins, Rothem Kovner, Rebecca L. Linn, Victoria H. J. Roberts, Amrita Sule, and Patrick M. Van Hoose. The human and non-human primate developmental GTEx projects. *Nature*, 637(8046):557–564, January 2025. ISSN 1476-4687. doi: 10.1038/s41586-024-08244-9. URL <https://www.nature.com/articles/s41586-024-08244-9>.
- [14] The GTEx Consortium. The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science*, 369(6509):1318–1330, September 2020. doi: 10.1126/science.aaz1776. URL <https://www.science.org/doi/10.1126/science.aaz1776>.
- [15] Jinyan Teng, Yahui Gao, Hongwei Yin, Zhonghao Bai, Shuli Liu, Haonan Zeng, Lijing Bai, Zexi Cai, Bingru Zhao, Xiujin Li, Zhiting Xu, Qing Lin, Zhangyuan Pan, Wenjing Yang, Xiaoshan Yu, Dailu Guan, Yali Hou, Brittney N. Keel, Gary A. Rohrer, Amanda K. Lindholm-Perry, William T. Oliver, Maria Ballester, Daniel Crespo-Piazuelo, Raquel Quintanilla, Oriol Canela-Xandri, Konrad Rawlik, Charley Xia, Yuelin Yao, Qianyi Zhao, Wenye Yao, Liu Yang, Houcheng Li, Huicong Zhang, Wang Liao, Tianshuo Chen, Peter Karlskov-Mortensen, Merete Fredholm, Marcel Amills, Alex Clop, Elisabetta Giuffra, Jun Wu, Xiaodian Cai, Shuqi Diao, Xiangchun Pan, Chen Wei, Jinghui Li, Hao Cheng, Sheng Wang, Guosheng Su, Goutam Sahana, Mogens Sandø Lund, Jack C. M. Dekkers, Luke Kramer, Christopher K. Tuggle, Ryan Corbett, Martien A. M. Groenen, Ole Madsen, Marta Gòdia, Dominique Rocha, Mathieu Charles, Congjun Li, Hubert Pausch, Xiaoxiang Hu, Laurent Frantz, Yonglun Luo, Lin Lin, Zhongyin Zhou, Zhe Zhang, Zitao Chen, Leilei Cui, Ruidong Xiang, Xia Shen, Pinghua Li, Ruihua Huang, Guoqing Tang, Mingzhou Li, Yunxiang Zhao, Guoqiang Yi, Zhonglin Tang, Jicai Jiang, Fuping Zhao, Xiaolong Yuan, Xiaohong Liu, Yaosheng Chen, Xuwen Xu, Shuhong Zhao, Pengju

- Zhao, Chris Haley, Huaijun Zhou, Qishan Wang, Yuchun Pan, Xiangdong Ding, Li Ma, Jiaqi Li, Pau Navarro, Qin Zhang, Bingjie Li, Albert Tenesa, Kui Li, George E. Liu, Zhe Zhang, and Lingzhao Fang. A compendium of genetic regulatory effects across pig tissues. *Nat Genet*, 56(1):112–123, January 2024. ISSN 1546-1718. doi: 10.1038/s41588-023-01585-7. URL <https://www.nature.com/articles/s41588-023-01585-7>.
- [16] Lijuan Chen, Houcheng Li, Jinyan Teng, Zhen Wang, Xiaolu Qu, Zhe Chen, Xiaodian Cai, Haonan Zeng, Zhonghao Bai, Jinghui Li, Xiangchun Pan, Leyan Yan, Fei Wang, Lin Lin, Yonglun Luo, Goutam Sahana, Mogens Sandø Lund, Maria Ballester, Daniel Crespo-Piazuolo, Peter Karlskov-Mortensen, Merete Fredholm, Alex Clop, Marcel Amills, Crystal Loving, Christopher K. Tuggle, Ole Madsen, Jiaqi Li, Zhe Zhang, George E. Liu, Jicai Jiang, Lingzhao Fang, and Guoqiang Yi. Construction of a Multitissue Cell Atlas Reveals Cell-Type-Specific Regulation of Molecular and Complex Phenotypes in Pigs. *Adv Sci (Weinh)*, page e04961, November 2025. ISSN 2198-3844. doi: 10.1002/advs.202504961.
- [17] Jiajia Long, Weiwei Liu, Xinhao Fan, Yalan Yang, Xiaogan Yang, and Zhonglin Tang. A comprehensive atlas of pig RNA editome across 23 tissues reveals RNA editing affecting interaction mRNA-miRNAs. *G3 (Bethesda)*, 14(10):jkae178, October 2024. ISSN 2160-1836. doi: 10.1093/g3journal/jkae178.
- [18] Hyo Jung Kang, Yuka Imamura Kawasaki, Feng Cheng, Ying Zhu, Xuming Xu, Mingfeng Li, André M. M. Sousa, Mihovil Pletikos, Kyle A. Meyer, Goran Sedmak, Tobias Guennel, Yurac Shin, Matthew B. Johnson, Zeljka Krsnik, Simone Mayer, Sofia Fertuzinhos, Sheila Umlauf, Steven N. Lisgo, Alexander Vortmeyer, Daniel R. Weinberger, Shrikant Mane, Thomas M. Hyde, Anita Huttner, Mark Reimers, Joel E. Kleinman, and Nenad Sestan. Spatio-temporal transcriptome of the human brain. *Nature*, 478(7370):483–489, October 2011. ISSN 1476-4687. doi: 10.1038/nature10523.
- [19] Andrey Y. Abramov. The brain-from neurodevelopment to neurodegeneration. *FEBS J*, 289(8): 2010–2012, April 2022. ISSN 1742-4658. doi: 10.1111/febs.16436.
- [20] Alexander Schaiter, Andreas Hentschel, Felix Kleefeld, Julia Schuld, Vincent Umathum, Tara Procida-Kowalski, Christopher Nelke, Angela Roth, Andreas Hahn, Heidrun H. Krämer, Tobias Ruck, Rita Horvath, Peter F. M. van der Ven, Marek Bartkuhn, Andreas Roos, and Anne Schänzer. Molecular composition of skeletal muscle in infants and adults: a comparative proteomic and transcriptomic study. *Sci Rep*, 14(1):22965, October 2024. ISSN 2045-2322. doi: 10.1038/s41598-024-74913-4. URL <https://www.nature.com/articles/s41598-024-74913-4>.
- [21] Margarida Cardoso-Moreira, Jean Halbert, Delphine Valloton, Britta Velten, Chunyan Chen, Yi Shao, Angélica Liechti, Kelly Ascensão, Coralie Rummel, Svetlana Ovchinnikova, Pavel V. Mazin, Ioannis Xenarios, Keith Harshman, Matthew Mort, David N. Cooper, Carmen Sandi, Michael J. Soares, Paula G. Ferreira, Sandra Afonso, Miguel Carneiro, James M. A. Turner, John L. VandeBerg, Amir Fallahshahroudi, Per Jensen, Rüdiger Behr, Steven Lisgo, Susan Lindsay, Philipp Khaitovich, Wolfgang Huber, Julie Baker, Simon Anders, Yong E. Zhang, and Henrik Kaessmann. Gene expression across mammalian organ development. *Nature*, 571

(7766):505–509, July 2019. ISSN 1476-4687. doi: 10.1038/s41586-019-1338-5.

- [22] Pernille Bogetofte Thomasen, Alena Salasova, Kasper Kjaer-Sorensen, Lucie Woloszczuková, Josef Lavický, Hande Login, Jeppe Tranberg-Jensen, Sergio Almeida, Sander Beel, Michaela Kavková, Per Qvist, Mads Kjolby, Peter Lund Ovesen, Stella Nolte, Benedicte Vestergaard, Andreea-Cornelia Udrea, Lene Niemann Nejsun, Moses V. Chao, Philip Van Damme, Jan Krivanek, Jeremy Dasen, Claus Oxvig, and Anders Nykjaer. SorCS2 binds progranulin to regulate motor neuron development. *Cell Reports*, 42(11):113333, November 2023. ISSN 2211-1247. doi: 10.1016/j.celrep.2023.113333. URL <https://www.sciencedirect.com/science/article/pii/S2211124723013451>.
- [23] Masako Osada, Emi Ito, Hector A. Fermin, Edwin Vazquez-Cintron, Tadmiri Venkatesh, Roland H. Friedel, and Mark Pezzano. The Wnt signaling antagonist Kremen1 is required for development of thymic architecture. *Clin Dev Immunol*, 13(2-4):299–319, 2006. ISSN 1740-2522. doi: 10.1080/17402520600935097.
- [24] Kevin Qin, Michael Yu, Jiaming Fan, Hongwei Wang, Piao Zhao, Guozhi Zhao, Wei Zeng, Connie Chen, Yonghui Wang, Annie Wang, Zander Schwartz, Jeffrey Hong, Lily Song, William Wagstaff, Rex C. Haydon, Hue H. Luu, Sherwin H. Ho, Jason Strelzow, Russell R. Reid, Tong-Chuan He, and Lewis L. Shi. Canonical and noncanonical Wnt signaling: Multilayered mediators, signaling mechanisms and major signaling crosstalk. *Genes & Diseases*, 11(1): 103–134, January 2024. ISSN 2352-3042. doi: 10.1016/j.gendis.2023.01.030. URL <https://www.sciencedirect.com/science/article/pii/S2352304223000557>.
- [25] Ruth Amann, Stefan Wyder, Anne M. Slavotinek, and Beat Trueb. The FgfrL1 receptor is required for development of slow muscle fibers. *Dev Biol*, 394(2):228–241, October 2014. ISSN 1095-564X. doi: 10.1016/j.ydbio.2014.08.016.
- [26] Sjoerd D Joustra, Ferdinand Roelfsema, A S Paul van Trotsenburg, Harald J Schneider, Robert P Kosilek, Herman M Kroon, John G Logan, Natalie C Butterfield, Xiang Zhou, Chirine Toufaily, Beata Bak, Marc-Olivier Turgeon, Emilie Brûlé, Frederik J Steyn, Mark Gurnell, Olympia Koulouri, Paul Le Tissier, Pierre Fontanaud, J H Duncan Bassett, Graham R Williams, Wilma Oostdijk, Jan M Wit, Alberto M Pereira, Nienke R Biermasz, Daniel J Bernard, and Nadia Schoenmakers. IGSF1 Deficiency Results in Human and Murine Somatotrope Neurosecretory Hyperfunction. *J Clin Endocrinol Metab*, 105(3):e70–e84, October 2019. ISSN 0021-972X. doi: 10.1210/clinem/dgz093. URL <https://pmc.ncbi.nlm.nih.gov/articles/PMC7108761/>.
- [27] Yazmin I. Rovira Gonzalez, Adam L. Moyer, Nicolas J. LeTexier, August D. Bratti, Siyuan Feng, Congshan Sun, Ting Liu, Jyothi Mula, Pankhuri Jha, Shama R. Iyer, Richard Lovering, Brian O’Rourke, Hye Lim Noh, Sujin Suk, Jason K. Kim, George K. Essien Umanah, and Kathryn R. Wagner. Mss51 deletion enhances muscle metabolism and glucose homeostasis in mice. *JCI Insight*, 4(20):e122247. ISSN 2379-3708. doi: 10.1172/jci.insight.122247. URL <https://pmc.ncbi.nlm.nih.gov/articles/PMC6824300/>.
- [28] Natsumi Ohsawa, Michinori Koebis, Satoshi Suo, Ichizo Nishino, and Shoichi Ishiura. Alternative splicing of *PDLIM3/ALP*, for α -actinin-associated LIM protein 3, is aberrant in

persons with myotonic dystrophy. *Biochemical and Biophysical Research Communications*, 409(1):64–69, May 2011. ISSN 0006-291X. doi: 10.1016/j.bbrc.2011.04.106. URL <https://www.sciencedirect.com/science/article/pii/S0006291X11007042>.

[29] Ryohei Iwata and Pierre Vanderhaeghen. Metabolic mechanisms of species-specific developmental tempo. *Developmental Cell*, 59(13):1628–1639, July 2024. ISSN 1534-5807. doi: 10.1016/j.devcel.2024.05.027. URL <https://www.sciencedirect.com/science/article/pii/S1534580724003617>.

[30] Nicholas Schaum, Jim Karkanias, Norma F. Neff, Andrew P. May, Stephen R. Quake, Tony Wyss-Coray, Spyros Darmanis, Joshua Batson, Olga Botvinnik, Michelle B. Chen, Steven Chen, Foad Green, Robert Jones, Ashley Maynard, Lolita Penland, Angela Oliveira Pisco, Rene V. Sit, Geoffrey M. Stanley, James T. Webber, Fabio Zanini, Ankit S. Baghel, Isaac Bakerman, Ishita Bansal, Daniela Berdnik, Biter Bilen, Douglas Brownfield, Corey Cain, Michelle B. Chen, Steven Chen, Min Cho, Giana Cirolia, Stephanie D. Conley, Spyros Darmanis, Aaron Demers, Kubilay Demir, Antoine de Morree, Tessa Divita, Haley du Bois, Laughing Bear Torrez Dulgeroff, Hamid Ebadi, F. Hernán Espinoza, Matt Fish, Qiang Gan, Benson M. George, Astrid Gillich, Foad Green, Geraldine Genetiano, Xueying Gu, Gunsagar S. Gulati, Yan Hang, Shayan Hosseinzadeh, Albin Huang, Tal Iram, Taichi Isobe, Feather Ives, Robert Jones, Kevin S. Kao, Guruswamy Karnam, Aaron M. Kershner, Bernhard M. Kiss, William Kong, Maya E. Kumar, Jonathan Lam, Davis P. Lee, Song E. Lee, Guang Li, Qingyun Li, Ling Liu, Annie Lo, Wan-Jin Lu, Anoop Manjunath, Andrew P. May, Kaia L. May, Oliver L. May, Ashley Maynard, Marina McKay, Ross J. Metzger, Marco Mignardi, Dullei Min, Ahmad N. Nabhan, Norma F. Neff, Katharine M. Ng, Joseph Noh, Rasika Patkar, Weng Chuan Peng, Lolita Penland, Robert Puccinelli, Eric J. Rulifson, Nicholas Schaum, Shaheen S. Sikandar, Rahul Sinha, Rene V. Sit, Krzysztof Szade, Weilun Tan, Cristina Tato, Krissie Tellez, Kyle J. Travaglini, Carolina Tropini, Lucas Waldburger, Linda J. van Weele, Michael N. Wosczyzna, Jinyi Xiang, Soso Xue, Justin Youngyunpipatkul, Fabio Zanini, Macy E. Zardeneta, Fan Zhang, Lu Zhou, Ishita Bansal, Steven Chen, Min Cho, Giana Cirolia, Spyros Darmanis, Aaron Demers, Tessa Divita, Hamid Ebadi, Geraldine Genetiano, Foad Green, Shayan Hosseinzadeh, Feather Ives, Annie Lo, Andrew P. May, Ashley Maynard, Marina McKay, Norma F. Neff, Lolita Penland, Rene V. Sit, Weilun Tan, Lucas Waldburger, Justin Youngyunpipatkul, Joshua Batson, Olga Botvinnik, Paola Castro, Derek Croote, Spyros Darmanis, Joseph L. DeRisi, Jim Karkanias, Angela Oliveira Pisco, Geoffrey M. Stanley, James T. Webber, Fabio Zanini, Ankit S. Baghel, Isaac Bakerman, Joshua Batson, Biter Bilen, Olga Botvinnik, Douglas Brownfield, Michelle B. Chen, Spyros Darmanis, Kubilay Demir, Antoine de Morree, Hamid Ebadi, F. Hernán Espinoza, Matt Fish, Qiang Gan, Benson M. George, Astrid Gillich, Xueying Gu, Gunsagar S. Gulati, Yan Hang, Albin Huang, Tal Iram, Taichi Isobe, Guruswamy Karnam, Aaron M. Kershner, Bernhard M. Kiss, William Kong, Christin S. Kuo, Jonathan Lam, Benoit Lehallier, Guang Li, Qingyun Li, Ling Liu, Wan-Jin Lu, Dullei Min, Ahmad N. Nabhan, Katharine M. Ng, Patricia K. Nguyen, Rasika Patkar, Weng Chuan Peng, Lolita Penland, Eric J. Rulifson, Nicholas Schaum, Shaheen S. Sikandar,

Rahul Sinha, Krzysztof Szade, Serena Y. Tan, Krissie Tellez, Kyle J. Travaglini, Carolina Tropini, Linda J. van Weele, Bruce M. Wang, Michael N. Wosczyzna, Jinyi Xiang, Hanadie Yousef, Lu Zhou, Joshua Batson, Olga Botvinnik, Steven Chen, Spyros Darmanis, Foad Green, Andrew P. May, Ashley Maynard, Angela Oliveira Pisco, Stephen R. Quake, Nicholas Schaum, Geoffrey M. Stanley, James T. Webber, Tony Wyss-Coray, Fabio Zanini, Philip A. Beachy, Charles K. F. Chan, Antoine de Morree, Benson M. George, Gunsagar S. Gulati, Yan Hang, Kerwyn Casey Huang, Tal Iram, Taichi Isobe, Aaron M. Kershner, Bernhard M. Kiss, William Kong, Guang Li, Qingyun Li, Ling Liu, Wan-Jin Lu, Ahmad N. Nabhan, Katharine M. Ng, Patricia K. Nguyen, Weng Chuan Peng, Eric J. Rulifson, Nicholas Schaum, Shaheen S. Sikandar, Rahul Sinha, Krzysztof Szade, Kyle J. Travaglini, Carolina Tropini, Bruce M. Wang, Kenneth Weinberg, Michael N. Wosczyzna, Sean M. Wu, Hanadie Yousef, Ben A. Barres, Philip A. Beachy, Charles K. F. Chan, Michael F. Clarke, Spyros Darmanis, Kerwyn Casey Huang, Jim Karkanias, Seung K. Kim, Mark A. Krasnow, Maya E. Kumar, Christin S. Kuo, Andrew P. May, Ross J. Metzger, Norma F. Neff, Roel Nusse, Patricia K. Nguyen, Thomas A. Rando, Justin Sonnenburg, Bruce M. Wang, Kenneth Weinberg, Irving L. Weissman, Sean M. Wu, Stephen R. Quake, and Tony Wyss-Coray. Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris. *Nature*, 562(7727):367–372, October 2018. ISSN 0028-0836. doi: 10.1038/s41586-018-0590-4. URL <https://pmc.ncbi.nlm.nih.gov/articles/PMC6642641/>.

[31] Huiwen Zheng, Jan Vijg, Atefeh Taherian Fard, and Jessica Cara Mar. Measuring cell-to-cell expression variability in single-cell RNA-sequencing data: a comparative analysis and applications to B cell aging. *Genome Biol*, 24:238, October 2023. ISSN 1474-7596. doi: 10.1186/s13059-023-03036-2. URL <https://pmc.ncbi.nlm.nih.gov/articles/PMC10588274/>.

[32] Guolin Ke, Qi Meng, Thomas Finley, Taifeng Wang, Wei Chen, Weidong Ma, Qiwei Ye, and Tie-Yan Liu. LightGBM: A Highly Efficient Gradient Boosting Decision Tree. In *Advances in Neural Information Processing Systems*, volume 30. Curran Associates, Inc., 2017. URL https://proceedings.neurips.cc/paper_files/paper/2017/hash/6449f44a102fde848669bdd9eb6b76fa-Abstract.html.

[33] Liis Kolberg, Uku Raudvere, Ivan Kuzmin, Priit Adler, Jaak Vilo, and Hedi Peterson. g:Profiler-interoperable web service for functional enrichment analysis and gene identifier mapping (2023 update). *Nucleic Acids Res*, 51(W1):W207–W212, July 2023. ISSN 1362-4962. doi: 10.1093/nar/gkad347.

Data availability

The PigGTE_x data used in this study is available at <https://piggtex.farmgtex.org/>. Raw and processed RNA-seq data are available in the NCBI Gene Expression Omnibus (GEO) under accession number GSE257558.

Code availability

All code required to reproduce the findings are available at <https://github.com/TianYuan-Liu/pig-dev-stage>. All analyses were conducted using Python 3.9 and R 4.2.0.

Acknowledgements

We thank the PigGTEx consortium for providing the transcriptomic data that made this study possible. T.L. was funded by the European Union Marie Skłodowska-Curie Actions Doctoral Network project LongTREC (HORIZON-MSCA-2021-DN-01 grant agreement project 101072892).

Author contributions

T.L. conceived the study, designed the methodology, performed all computational analyses, interpreted results, and wrote the manuscript. R.L. and I.M.K. contributed to manuscript editing and review. P.T. supervised the study, provided guidance, and edited the manuscript. All authors reviewed and approved the final manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information accompanies this paper.

Correspondence and requests for materials should be addressed to P.T. (TheobaldP@cardiff.ac.uk).




Figure 1: Study design and data landscape. **a**, Developmental stage definitions mapping chronological age to five ordered stages: Infant (0–20 d), Early childhood (21–59 d), Pre-pubertal (60–149 d), Post-pubertal (150–365 d), and Adult (>365 d). **b**, Machine learning workflow diagram illustrating the pipeline from data preprocessing to model calibration and evaluation. **c**, Sample distribution heatmap showing tissue-by-stage availability across 1,924 samples from five tissues. **d**, Classification scheme assignments based on per-class sample availability, with 4-class schemes for muscle and liver, and 2-class schemes for blood, brain, and lung.




Figure 2: Classifier performance diagnostics. **a**, Performance metrics heatmap showing balanced accuracy, macro F1, mean absolute error, and Spearman correlations with 95% bootstrap confidence intervals across tissues. Mean balanced accuracy across all tissues is 0.825, ranging from 0.638 (brain) to 0.920 (liver). **b**, Confusion matrices for each tissue showing classification accuracy and error patterns. **c**, Scatter plots of predicted developmental stage versus chronological age, demonstrating strong age–stage correlations (liver $\rho = 0.979$, muscle $\rho = 0.937$, brain $\rho = 0.942$, blood $\rho = 0.773$).




Figure 3: Functional enrichment analysis of stage-informative genes. Functional network showing interconnected enrichment modules, with nodes representing enriched terms coloured by tissue and edges connecting terms sharing >25% of genes (Jaccard similarity). Hubs and cluster representatives are labelled with biological process or pathway names.




Figure 4: Cross-species validation of developmental gene expression. Human skeletal muscle transcriptomic data from Schaiter *et al.*²⁰ comparing infant (7–28 months) versus adult (30–56 years) samples were used to validate conservation of developmental signatures. **a**, Scatter plot of $\log_2$ fold changes between pig and human, showing significant positive correlation (Pearson $r = 0.618$, $p = 5.86 \times 10^{-5}$, 95% CI: 0.364–0.787). Of 36 genes meeting significance criteria ($p < 0.10$) in both species, 32 (88.9%) showed conserved direction of change. **b**, Feature importance of top conserved markers in the developmental stage classifier. **c**, Paired fold change comparison showing pig and human values for each conserved gene. Top markers include *SORCS2*, *FGFR1*, *IGSF1*, *KREMEN1*, and *ACHE*.