

Supplementary Information


A Multi-Tissue Transcriptomic Atlas of Porcine Development Identifies Conserved Molecular Programs with Humans

Tianyuan Liu, Rujing Lei, Ilyas M. Khan, Peter Theobald


Contents

Supplementary Figures	1
Supplementary Tables	4


Supplementary Figures



Supplementary Figure 1: Cross-species sensitivity analysis of conserved developmental markers. **a**, Pearson correlation between pig and human developmental fold changes across FDR thresholds ranging from < 0.05 to < 0.20 (Benjamini–Hochberg corrected), using all genes passing pre-specified criteria (FDR $<$ threshold and $|\log_2 \text{FC}| > 0.5$ in both species). Points are sized by the number of genes at each threshold. **b**, Bootstrap distribution of correlation coefficients ($n = 1,000$ iterations) for the pre-specified criteria (FDR < 0.10 , $|\log_2 \text{FC}| > 0.5$, $n = 66$ genes). Solid vertical line indicates the observed correlation ($r = 0.693$); dashed lines denote the 95% bootstrap confidence interval boundaries (0.50–0.82).



Supplementary Figure 2: Tissue specificity and feature selection stability analysis. **a**, Uniform manifold approximation and projection (UMAP) of gene expression profiles across five tissues (muscle, brain, liver, blood, lung), demonstrating distinct tissue-specific clustering patterns ($n = 150$ samples per tissue, balanced sampling; top 500 variable genes). For visualization only, BioProject batch effects were removed per tissue using `limma::removeBatchEffect` prior to UMAP; the machine learning pipeline operates on uncorrected expression data. **b**, Within-tissue feature selection stability quantified as mean Jaccard similarity index of the top 50 selected features across ten independent random seeds. Error bars represent standard deviation; higher values indicate more reproducible feature selection. **c**, Mean absolute Spearman correlation between top-ranked features and developmental stage for each tissue.



Supplementary Figure 3: Extended classification performance metrics across tissues and developmental stages. **a**, Heatmap of per-class precision, recall, and F1 scores for each developmental stage across all five tissues. Colour scale ranges from 0.4 (red, poor performance) to 1.0 (green, excellent performance). Missing values indicate developmental stages not present in that tissue's classification scheme. **b**, Overall performance summary showing balanced accuracy and macro-averaged F1 scores evaluated on pooled cross-validated held-out predictions for each tissue. Error bars indicate 95% bootstrap confidence intervals from 1,000 resampling iterations.

Supplementary Tables

Supplementary Table 1: Train versus test performance comparison. Both train and test (pooled cross-validated held-out predictions) performance are reported for all tissues. The gap between train and test metrics indicates the degree of overfitting; regularisation and early stopping were used to minimise this gap.

Tissue	Set	Scheme	<i>n</i>	Balanced Acc.	F1-macro	F1-weighted
Muscle	Train	4-class	761	0.968	0.974	0.984
Muscle	Test	4-class	761	0.894	0.899	0.939
Brain	Train	2-class	43	0.915	0.922	0.930
Brain	Test	2-class	43	0.811	0.801	0.816
Liver	Train	4-class	309	0.985	0.987	0.987
Liver	Test	4-class	309	0.921	0.918	0.919
Blood	Train	2-class	78	1.000	1.000	1.000
Blood	Test	2-class	78	0.959	0.960	0.961
Lung	Train	2-class	129	0.948	0.942	0.961
Lung	Test	2-class	129	0.892	0.876	0.916

Supplementary Table 2: Per-class performance metrics by tissue. Precision, recall, and F1 scores for each developmental stage within each tissue. n indicates the total number of samples per class (pooled across cross-validation folds).

Tissue	Stage	n	Precision	Recall	F1
Muscle	infant	194	0.939	0.959	0.949
Muscle	early childhood	76	0.838	0.816	0.827
Muscle	pre-pubertal	90	0.871	0.822	0.846
Muscle	post-pubertal/adult	401	0.973	0.980	0.976
Brain	pre-pubertal (<150d)	28	0.885	0.821	0.852
Brain	post-pubertal ($\geq$ 150d)	15	0.706	0.800	0.750
Liver	infant	87	0.965	0.954	0.960
Liver	early childhood	58	0.887	0.948	0.917
Liver	pre-pubertal	83	0.910	0.855	0.882
Liver	post-pubertal/adult	81	0.904	0.926	0.915
Blood	pre-pubertal (<150d)	45	0.957	0.978	0.967
Blood	post-pubertal ($\geq$ 150d)	33	0.969	0.939	0.954
Lung	pre-pubertal (<150d)	102	0.960	0.931	0.945
Lung	post-pubertal ($\geq$ 150d)	27	0.767	0.852	0.807

Supplementary Table 3: Functional annotations of cross-species developmental markers. Annotations compiled from UniProt, GeneCards, and literature review. FC = log₂ fold change (adult/infant); positive values indicate higher expression in adults, negative values indicate higher expression in infants. Pig FC from pig GTEx data (GEO: GSE257558); Human FC from Schaiter *et al.* (2024). Genes ordered by model importance within each functional module. “—” indicates the gene did not pass cross-species criteria (FDR < 0.10, |log₂ FC| > 0.5) but is included based on its known biological role.

Gene	Developmental Function	Pig FC	Human FC
<i>Module 1: Neuromuscular Maturation and Growth Signalling</i>			
ACHE	Functions as the key enzyme at the neuromuscular junction and as a structural adhesion molecule guiding neuronal growth during development ¹ .	−3.34	−0.53
NRK	Embryonic Ste20-type kinase that promotes actin polymerisation via cofilin phosphorylation and activates JNK signalling during skeletal muscle formation; expression restricted to embryonic and early postnatal development ² .	−8.56	−2.45
KREMEN1	Negative regulator of Wnt/ β -catenin signalling, partnering with Dkk1 to control cell survival and ensure proper patterning during postnatal development ^{3?} .	−1.69	−1.43
IGF2BP2	RNA-binding protein that promotes myoblast proliferation via translational control of growth-related transcripts (c-Myc, Igf1r, Sp1); postnatal downregulation required for terminal myogenic differentiation ⁴ .	−2.19	−2.03
SORCS2	Molecular switch directing trafficking of signalling receptors to regulate the balance between neuronal growth and pruning during neuromuscular maturation ⁵ .	−4.67	−0.95
ETV5	FGF-responsive PEA3-subfamily ETS transcription factor that promotes stem/progenitor cell self-renewal and myogenic differentiation in cooperation with MEF2; highly expressed during active postnatal myogenesis ⁶ .	−2.80	−2.24
IGF2BP3	Oncofetal RNA-binding protein that promotes myoblast proliferation and differentiation via IGF2–PI3K/Akt signalling; pronounced postnatal silencing reflects the conserved transition from fetal growth programs ⁴ .	−4.85	−1.23
CCDC8	Core component of the 3M complex (with CUL7 and OBSL1) that regulates microtubule dynamics, genome integrity, and growth hormone signalling during rapid postnatal growth; consistent with severe growth restriction in 3-M syndrome ⁷ .	−4.44	−3.14
<i>Module 2: Structural Refinement and ECM Remodelling</i>			
TCAF1	Regulates TRPM8 ion channel trafficking and Ca ²⁺ signalling; may support calcium-dependent processes during early postnatal muscle growth and metabolic maturation ⁸ .	−4.72	−1.02
FGFRL1	Acts as a “decoy” receptor dampening excessive FGF signalling to ensure proper formation of critical structures including the diaphragm and skeleton ⁹ .	−3.03	−2.10
WDR1	Cofactor of cofilin-mediated actin filament disassembly, essential for sarcomere assembly and myofibril organisation during postnatal muscle growth; higher expression in immature muscle reflects intensive cytoskeletal remodelling during myofibrillogenesis ¹⁰ .	−1.69	−0.52
DLK1	Crucial regulator of the myogenic program that inhibits myoblast proliferation while promoting differentiation; high infant expression declines as the muscle fiber population matures ^{11,12} .	−9.15	−2.88
FSCN1	Actin-bundling protein essential for formation of cellular protrusions during cell migration and tissue morphogenesis; highly expressed in embryonic somite-derived tissues, with postnatal downregulation as migratory programs complete ¹³ .	−4.28	−1.95
POSTN	Secreted matricellular protein that mediates ECM remodelling, collagen fibrillogenesis, and connective tissue maturation during postnatal myofiber development; expression declines as tissue architecture stabilises ¹⁴ .	−7.97	−2.93
FREM2	Basement membrane extracellular matrix protein of the FRAS/FREM complex maintaining epithelial–mesenchymal adhesion during organogenesis; postnatal downregulation reflects transition to collagen VII-dominated mature basement membranes ¹⁵ .	−4.58	−1.05
ADAMTS2	Procollagen N-proteinase that cleaves propeptides from fibrillar procollagens to initiate collagen fibrillogenesis; high infant expression reflects intensive ECM biosynthesis during rapid postnatal growth ¹⁶ .	−4.25	−1.83
COL11A2	Component of type XI collagen regulating collagen fibril organisation in cartilage and developing connective tissue; high infant expression reflects active skeletal morphogenesis and ECM assembly ¹⁷ .	−8.35	−0.70
TRIM54	Muscle-specific E3 ubiquitin ligase (MuRF3) involved in sarcomere Z-disc assembly; expression increases postnatally as the mature contractile apparatus is established.	—	—
<i>Module 3: Metabolic Transition</i>			
ME1	Malic enzyme 1, catalysing oxidative decarboxylation of malate to pyruvate; a key source of NADPH for lipid biosynthesis. Expression increases during development, reflecting metabolic maturation.	2.59	0.64
MAOB	Mitochondrial outer membrane enzyme catalysing oxidative deamination of monoamines; postnatal upregulation reflects developmental maturation of oxidative metabolism and the glycolytic-to-oxidative fiber-type shift.	2.55	0.80
PLA2G4E	Calcium-dependent N-acyltransferase synthesising N-acylphosphatidylethanolamines (endocannabinoid precursors); species-discordant expression may reflect divergent intramuscular lipid metabolism programs.	2.32	−1.25

Supplementary Table 4: Stage-by-stage expression trajectories of cross-species developmental markers. Mean TPM values across porcine developmental stages for genes with conserved pig-human developmental patterns. Stage boundaries: infant (0–20 d), early childhood (21–59 d), pre-pubertal (60–149 d), post-pubertal (150–365 d), adult (>365 d). “Peak Change” indicates the developmental transition with the largest fold change. Genes are grouped by functional module and ordered by model importance within each module.

Gene	infant	early	pre-pub	post-pub	adult	Peak Change
<i>Module 1: Neuromuscular Maturation and Growth Signalling</i>						
<i>ACHE</i>	17.5	7.6	3.4	1.3	1.7	infant→early
<i>NRK</i>	128.2	4.2	8.1	1.3	0.3	infant→early
<i>KREMEN1</i>	30.3	13.2	9.4	5.2	9.4	infant→early
<i>IGF2BP2</i>	11.3	1.5	1.9	1.0	2.5	infant→early
<i>SORCS2</i>	14.4	3.0	2.2	0.6	0.6	infant→early
<i>ETV5</i>	34.2	16.7	7.5	5.5	4.9	early→pre-pub
<i>IGF2BP3</i>	21.8	1.7	2.1	0.4	0.7	infant→early
<i>CCDC8</i>	177.4	28.2	25.4	13.4	8.2	infant→early
<i>Module 2: Structural Refinement and ECM Remodelling</i>						
<i>TCAF1</i>	78.9	8.3	8.1	3.8	3.0	infant→early
<i>FGFRL1</i>	77.4	15.6	14.2	4.3	9.5	infant→early
<i>WDR1</i>	202.9	125.6	95.0	51.9	63.1	pre-pub→post-pub
<i>DLK1</i>	338.8	3.9	6.4	0.7	0.6	infant→early
<i>FSCN1</i>	123.7	32.5	18.9	6.2	6.4	infant→early
<i>POSTN</i>	602.4	44.1	54.3	2.5	2.4	pre-pub→post-pub
<i>FREM2</i>	33.1	4.4	5.5	1.5	1.4	infant→early
<i>ADAMTS2</i>	51.1	16.6	16.0	6.4	2.7	infant→early
<i>COL11A2</i>	44.6	1.5	1.0	0.3	0.1	infant→early
<i>TRIM54</i>	78.3	351.1	149.4	377.9	144.4	infant→early
<i>Module 3: Metabolic Transition</i>						
<i>ME1</i>	2.8	4.0	3.9	6.9	16.9	post-pub→adult
<i>MAOB</i>	5.7	24.2	10.8	15.7	33.4	infant→early
<i>PLA2G4E</i>	0.1	1.4	0.7	0.6	0.5	infant→early

Supplementary Table 5: Within-project fold-change validation across developmental contrasts. Developmental fold-changes ($\log_2$ scale) were computed within single-project muscle cohorts—each spanning four developmental stages under uniform laboratory protocols—and correlated with full-dataset estimates across three stage comparisons (all using infant as baseline). Pooled within-project fold-changes were computed by averaging across both projects before correlating with full-dataset estimates. “All genes” includes all 31,908 expressed genes; “Cross-species” includes the 66 genes passing pre-specified criteria ($\text{FDR} < 0.10$, $|\log_2 \text{FC}| > 0.5$) in both pig and human. CI = 95% bootstrap confidence interval (1,000 iterations). Per-project results are available in the analysis repository. Full-dataset sample sizes: infant $n = 194$, early childhood $n = 76$, pre-pubertal $n = 90$, post-pubertal $n = 396$ (excludes 5 adult samples merged into post-pubertal/adult in the 4-class scheme).

Comparison	Gene set	Full dataset		Within-project		Pooled r [95% CI]	p -value
		n_{Inf}	n_{later}	n_{Inf}	n_{later}		
infant vs. early childhood	All genes	194	76	106	12	0.669 [0.66–0.68]	$< 10^{-300}$
	Cross-species					0.761	1.3×10^{-13}
infant vs. pre-pubertal	All genes	194	90	106	27	0.806 [0.80–0.82]	$< 10^{-300}$
	Cross-species					0.965	6.9×10^{-39}
infant vs. post-pubertal	All genes	194	396	106	10	0.799 [0.79–0.81]	$< 10^{-300}$
	Cross-species					0.978	5.6×10^{-45}

References

- [1] Richard L. Rotundo. Expression and localization of acetylcholinesterase at the neuromuscular junction. *J Neurocytol*, 32(5):743–766, June 2003. ISSN 1573-7381. doi: 10.1023/B:NEUR.0000020621.58197.d4. URL <https://doi.org/10.1023/B:NEUR.0000020621.58197.d4>.
- [2] M. Kanai-Azuma, Y. Kanai, M. Okamoto, Y. Hayashi, H. Yonekawa, and K. Yazaki. Nr1: a murine X-linked NIK (Nck-interacting kinase)-related kinase gene expressed in skeletal muscle. *Mech Dev*, 89(1-2):155–159, December 1999. ISSN 0925-4773. doi: 10.1016/s0925-4773(99)00193-8.
- [3] Masako Osada, Emi Ito, Hector A. Fermin, Edwin Vazquez-Cintron, Tadmiri Venkatesh, Roland H. Friedel, and Mark Pezzano. The Wnt signaling antagonist Kremen1 is required for development of thymic architecture. *Clin Dev Immunol*, 13(2-4):299–319, 2006. ISSN 1740-2522. doi: 10.1080/17402520600935097.
- [4] Zhizhong Li, Jason A. Gilbert, Yunyu Zhang, Minsi Zhang, Qiong Qiu, Krishnan Ramanujan, Tea Shavlakadze, John K. Eash, Annarita Scaramozza, Matthew M. Goddeeris, David G. Kirsch, Kevin P. Campbell, Andrew S. Brack, and David J. Glass. An HMGA2-IGF2BP2 axis regulates myoblast proliferation and myogenesis. *Dev Cell*, 23(6):1176–1188, December 2012. ISSN 1878-1551. doi: 10.1016/j.devcel.2012.10.019.
- [5] Pernille Bogetofte Thomasen, Alena Salasova, Kasper Kjaer-Sorensen, Lucie Woloszczuková, Josef Lavický, Hande Login, Jeppe Tranberg-Jensen, Sergio Almeida, Sander Beel, Michaela Kavková, Per Qvist, Mads Kjolby, Peter Lund Ovesen, Stella Nolte, Benedicte Vestergaard,

- Andreea-Cornelia Udrea, Lene Niemann Nejsum, Moses V. Chao, Philip Van Damme, Jan Krivanek, Jeremy Dasen, Claus Oxvig, and Anders Nykjaer. SorCS2 binds progranulin to regulate motor neuron development. *Cell Reports*, 42(11):113333, November 2023. ISSN 2211-1247. doi: 10.1016/j.celrep.2023.113333. URL <https://www.sciencedirect.com/science/article/pii/S2211124723013451>.
- [6] J. M. Taylor, E. E. Dupont-Versteegden, J. D. Davies, J. A. Hassell, J. D. Houlé, C. M. Gurley, and C. A. Peterson. A role for the ETS domain transcription factor PEA3 in myogenic differentiation. *Mol Cell Biol*, 17(9):5550–5558, September 1997. ISSN 0270-7306. doi: 10.1128/MCB.17.9.5550.
- [7] Dan Hanson, Philip G. Murray, James O’Sullivan, Jill Urquhart, Sarah Daly, Sanjeev S. Bhaskar, Leslie G. Biesecker, Mars Skae, Claire Smith, Trevor Cole, Jeremy Kirk, Kate Chandler, Helen Kingston, Dian Donnai, Peter E. Clayton, and Graeme C. M. Black. Exome Sequencing Identifies *CCDC8* Mutations in 3-M Syndrome, Suggesting that *CCDC8* Contributes in a Pathway with *CUL7* and *OBSL1* to Control Human Growth. *The American Journal of Human Genetics*, 89(1):148–153, July 2011. ISSN 0002-9297. doi: 10.1016/j.ajhg.2011.05.028. URL <https://www.sciencedirect.com/science/article/pii/S0002929711002217>.
- [8] Dimitra Gkika, Loic Lemonnier, George Shapovalov, Dmitri Gordienko, Céline Poux, Michela Bernardini, Alexandre Bokhobza, Gabriel Bidaux, Cindy Degerny, Kathye Verreman, Basma Guarmit, Mohamed Benahmed, Yvan de Launoit, Rene J. M. Bindels, Alessandra Fiorio Pla, and Natalia Prevarskaya. TRP channel-associated factors are a novel protein family that regulates TRPM8 trafficking and activity. *J Cell Biol*, 208(1):89–107, January 2015. ISSN 1540-8140. doi: 10.1083/jcb.201402076.
- [9] Ruth Amann, Stefan Wyder, Anne M. Slavotinek, and Beat Trueb. The FgfrL1 receptor is required for development of slow muscle fibers. *Dev Biol*, 394(2):228–241, October 2014. ISSN 1095-564X. doi: 10.1016/j.ydbio.2014.08.016.
- [10] Shoichiro Ono. Functions of actin-interacting protein 1 (AIP1)/WD repeat protein 1 (WDR1) in actin filament dynamics and cytoskeletal regulation. *Biochem Biophys Res Commun*, 506(2):315–322, November 2018. ISSN 1090-2104. doi: 10.1016/j.bbrc.2017.10.096.
- [11] Jolena N. Waddell, Peijing Zhang, Yefei Wen, Sanjay K. Gupta, Aleksey Yevtodiyenko, Jennifer V. Schmidt, Christopher A. Bidwell, Ashok Kumar, and Shihuan Kuang. Dlk1 is necessary for proper skeletal muscle development and regeneration. *PLoS One*, 5(11):e15055, November 2010. ISSN 1932-6203. doi: 10.1371/journal.pone.0015055.
- [12] Ditte Caroline Andersen, Jorge Laborda, Victoriano Baladron, Moustapha Kassem, Søren Paludan Sheikh, and Charlotte Harken Jensen. Dual role of delta-like 1 homolog (DLK1) in skeletal muscle development and adult muscle regeneration. *Development*, 140(18):3743–3753, September 2013. ISSN 1477-9129. doi: 10.1242/dev.095810.

- [13] Josephine C. Adams. Roles of fascin in cell adhesion and motility. *Curr Opin Cell Biol*, 16(5): 590–596, October 2004. ISSN 0955-0674. doi: 10.1016/j.ceb.2004.07.009.
- [14] Cansu Ozdemir, Uğur Akpulat, Parisa Sharafi, Yılmaz Yıldız, Ilyas Onbaşlar, and Cetin Kocaefe. Periostin is temporally expressed as an extracellular matrix component in skeletal muscle regeneration and differentiation. *Gene*, 553(2):130–139, December 2014. ISSN 1879-0038. doi: 10.1016/j.gene.2014.10.014.
- [15] Daiji Kiyozumi, Nagisa Sugimoto, and Kiyotoshi Sekiguchi. Breakdown of the reciprocal stabilization of QBRICK/Frem1, Fras1, and Frem2 at the basement membrane provokes Fraser syndrome-like defects. *Proc Natl Acad Sci U S A*, 103(32):11981–11986, August 2006. ISSN 0027-8424. doi: 10.1073/pnas.0601011103.
- [16] A. Colige, S. W. Li, A. L. Sieron, B. V. Nusgens, D. J. Prockop, and C. M. Lapière. cDNA cloning and expression of bovine procollagen I N-proteinase: a new member of the superfamily of zinc-metalloproteinases with binding sites for cells and other matrix components. *Proc Natl Acad Sci U S A*, 94(6):2374–2379, March 1997. ISSN 0027-8424. doi: 10.1073/pnas.94.6.2374.
- [17] Mei Sun, Eric Y. Luo, Sheila M. Adams, Thomas Adams, Yaping Ye, Snehal S. Shetye, Louis J. Soslowsky, and David E. Birk. Collagen XI regulates the acquisition of collagen fibril structure, organization and functional properties in tendon. *Matrix Biol*, 94:77–94, December 2020. ISSN 0945-053X. doi: 10.1016/j.matbio.2020.09.001. URL <https://pmc.ncbi.nlm.nih.gov/articles/PMC7722227/>.