

J/ ψ production from ^4He

Electron beam energy: 11 GeV

Luminosity: 1.2×10^{37} eN cm⁻² s⁻¹

Target: 15-cm liquid ^4He ($\sim 2\%$ radiation length)

Acceptance: $p > 300$ MeV, $6^\circ < \theta < 29^\circ$, full azimuthal

Cross section model: grids “duke-jpsi-02-19-2020” with and without the restriction “p” (initial nucleon in ^4He) < 300 MeV

Photoproduction:

Bremsstrahlung photon with target as the radiator, select $6.2 < E_\gamma < 9.2$ GeV

Subthreshold cut: $E_\gamma < 8.2$ GeV

Note: E_γ is known in simulation


Purpose:

Divide the subthreshold region into two bins [6.2, 7.2] and [7.2, 8.2] GeV of the photon energy


J/ ψ production from ^4He

Simulated cases


| Label | γ energy (GeV) | initial “p” |
|--------------------------|-----------------------|-------------|
| γ -nc-[6.2,7.2] | [6.2, 7.2] | No cut |
| γ -c300-[6.2,7.2] | [6.2, 7.2] | < 300 MeV |
| γ -nc-[7.2,8.2] | [7.2, 8.2] | No cut |
| γ -c300-[7.2,8.2] | [7.2, 8.2] | < 300 MeV |




γ -nc-[6.2,7.2]




γ -nc-[6.2,7.2]





γ -nc-[6.2,7.2]




γ -nc-[6.2,7.2]










γ -c300-[6.2,7.2]









γ -c300-[6.2,7.2]







γ -nc-[7.2,8.2]




γ -nc-[7.2,8.2]




γ -nc-[7.2,8.2]





γ -nc-[7.2,8.2]





γ -nc-[7.2,8.2]














γ -c300-[7.2,8.2]

