

Consequences of Regge behaviour

In Susy LFHQCD we obtain for the nucleon ($\tau = 3$) FF the beautiful result:

$$F(t) = \frac{1}{B(2, 1 - \alpha_0)} B(2, 1 - \alpha(t)) \quad (1)$$

where $\alpha(t) = \alpha_0 - \alpha' t$.

For vector current: $\alpha_0 = \frac{1}{2}$, for axial current $\alpha_0 = 0$, for both currents : $\alpha' = \frac{1}{4\lambda}$

From that we can deduce that the distribution function $q(x)$ can be expressed in a parametrization invariant way as:

$$q(x) = \frac{1}{B(2, 1 - \alpha_0)} (1 - w(x)) w(x)^{-\alpha(t)} w'(x) \quad (2)$$

For small x we assume $w(x) = Cx + O(x^2)$ This yields for the small x dependence:

$$q(x) \sim \frac{1}{B(2, 1 - \alpha_0)} (Cx)^{-\alpha(t)} C = \frac{1}{B(2, 1 - \alpha_0)} C^{1-\alpha_0-\alpha't} x^{-\alpha_0-\alpha't} \quad (3)$$

For small x Regge behaviour suggests :

$$q(x, t) \sim \beta_R(t) \left(\frac{x_0}{x} \right)^{\alpha(t)} \quad (4)$$

where $x_0 < 1$ is the value of x where Regge behaviour sets in. x is conventionally identified with the Bjorken $x = \frac{Q^2}{s+m^2+Q^2}$ Comparing the two expressions we obtain for the Regge residue the condition:

$$\beta_R(t) \left(\frac{x_0}{x} \right)^{-\alpha_0-\alpha't} = \frac{1}{B(2, 1 - \alpha_0)} C^{1-\alpha_0-\alpha't} x^{-\alpha_0-\alpha't} \quad (5)$$

$$\beta_R(t) = \frac{1}{B(2, 1 - \alpha_0)} C^{1-\alpha_0} x^{-\alpha_0} e^{-\log(C x_0) t/(4\lambda)} \quad (6)$$

For the vector current we obtain

$$\beta_V(t) = \frac{3}{4} \left(\frac{C_V}{x_0} \right)^{1/2} \exp[-\log(C_V x_0) t/(4\lambda)] \quad (7)$$

For the axial current:

$$\beta_A(t) = 2C_A \exp[-\log(C_A x_0) t/(4\lambda)] \quad (8)$$

If we assume that the Regge residues at $t = 0$ are approximately equal, then we have:

$$C_A \approx \frac{3}{8} \sqrt{\frac{C_V}{x_0}} \quad (9)$$

As a rough estimate for the t -dependence of the residue we assume that is similart to that of a FF, that is we assume that $C_V x_0 = e^{-n}$ with n in the range $2 \dots 4$, that is $x_0 \approx e^{-n}/C_V$ Then $C_A \approx \frac{3}{8} C_V e^{n/2}$

For the standard form we have: $w'(0, a) = e^{-a}$ and hence we expect wery roughly:

$$a_A = a_V - n/2 + 0.98 \quad (10)$$

That is we expect a_A to be near zero or even negative.